



Motor activities, Physical education and Sport
Interdisciplinary approaches and methodological implications

Series directed by DARIO COLELLA

5

Scientific Committee

Branislav Antala
Comenius University in Bratislava - Slovakia

Gheorghe Balint
University of Bacau - Romania

Antonio Borgogni
University of Bergamo

Attilio Carraro
University of Bolzano

Dario Colella
University of Salento, Lecce

Simone Digennaro
University of Cassino and Southern Lazio

Francesco Fischetti
University of Bari

Erika Gobbi
University of Urbino

Michael Kolb
University of Vienna - Austria

Stefania Pinnelli
University of Salento, Lecce

Italo Sannicandro
University of Foggia

Jana Vašíčková Palacky
University, Olomouc - Czech Republic

FÉDÉRATION INTERNATIONALE D'ÉDUCATION PHYSIQUE ET SPORTIVE



Game in Physical Education, Physical Activities and Sport of Children and Youth: Researches – Best Practices – Situation

Editors

Dario Colella
Domenico Monacis
Branislav Antala
Sergii Ivashchenko
Dario Novak





EUROPE

Publication Game in Physical Education, Physical Activities and Sport of Children and Youth. Researches-Best Practices-Situation

Editors **Dario Colella** (University of Salento, Lecce, Italy); **Domenico Monacis** (Pegaso University Telematic, Naples, Italy); **Branislav Antala**, (Comenius University in Bratislava, Slovakia); **Sergii Ivashchenko** (University of Kyiv, Ukraine) and **Dario Novak** (University of Zagreb, Croatia)

Manuela Ladogana (University of Foggia, Italy): linguistic editing and revision publication prepared with a contribution on the funds of University of Foggia, Department of Humanistic Studies, Italy

Reviewers Antonio Borgogni (Italy); antonio.borgogni@unibg.it
Athos Trecroci (Italy); athos.trecroci@unimi.it
Attilio Carraro (Italy); attilio.carraro@unibz.it
Cristiana D'Anna (Italy); cristiana.danna@unipegaso.it
František Seman (Slovakia); frantisek.seman@uniba.sk
Jelena Petrovic (Serbia); jelenailicpetrovic@gmail.com
Lubor Tománek (Slovakia); lubor.tomanek@uniba.sk
Stefan Kolimechkov (UK); dr.stefan.kolimechkov@gmail.com

Note No part of this publication may be reproduced without the prior permission of the Authors

ISBN volume 979-12-5568-095-6

2023 © by Pensa MultiMedia®

73100 Lecce • Via Arturo Maria Caprioli, 8 • Tel. 0832.230435

www.pensamultimedia.it

Table of contents

- 9 **Introduction**
Dario Colella, Domenico Monacis, Branislav Antala, Sergii Ivashchenko, Dario Novak

RESEARCHES

- 17 Oana-Maria Păstac
The benefits of using Gamification in teaching English as a Foreign Language
- 29 Wang Shiyue, Garry Kuan, Zhao Mengyuan, Kueh Yee Cheng, Zhou Ke
Research on Rehabilitation Strategy of Exceptional Child by using Chinese Traditional Games
- 45 Govindasamy Balasekaran, Peggy Boey, Ng Yew Cheo
Best Research Practices in Physical Education, Physical Activities and Sport Among Singaporean Children and Youths
- 57 Andreea Popescu, Daniela-Denisa Mitroi, Eduard-Leonard Guță
Evaluation of the postural rehabilitation process at school children flat foot by analysing the distribution of plantar pressure
- 65 John Jeswenny Fresshila, Chin Ngien-Siong, Kueh Yee Cheng, Garry Kuan
The Effect of Small-Sided Games Intervention on the Primary School Students
- 77 John Jeswenny Fresshila, Chin Ngien-Siong, Garry Kuan, Teo Boon Sian
Understanding Expectancy Beliefs and Values in Small-sided Games among Chinese Primary School Children
- 91 Wee Eng Hoe, Victor Kung, Ler Hui Yin, Chin Ngien Siong
Are social support and level of physical activity determinants of exercise adherence among university students?

- 107 Jose Francisco Mora Núñez, Manuel Guerrero Zainos
Benefits of formative sports chair in times of confinement
- 117 Sergii Ivashchenko, Natalya Kovalchuk, Branislav Antala
Game style of Physical Education and its influence on Motor Activity of children in Ukraine
- 129 Antonio Eugenio Rivera-Cisneros, José Guadalupe Montaña-Corona, Teresa Melchor-Moreno, Fernando Avila-González, Manuel Guerrero-Zainos, Néstor Rangel, Raymundo Romo, Aurelia Gutiérrez-Ríos, Jorge Manuel Sánchez-González, Oscar Ontivero
Evolution of game, body mass, metabolic status, and motricity in four decades of Mexican childrens

BEST PRACTICES

- 141 Dania Aspasia, Karolina Barzouka, Konstantinos Sotiropoulos, Ioannis Ktistakis, Nikolaos Apostolidis
The use of game-based approaches as playful learning in physical education teacher education programs
- 153 Salfou Ilboudo, Brigitte Nana, Delphine Rouamba, Daouda Ouattara, Florent Songnaba
Contraintes de la pratique régulière de l'activité physique et sportive dans la ville de Ouagadougou
- 165 Camelia-Daniela Plăstoi, Ilie Mihai, Simona Sgonde, Roxana Bodgan
Sustainable perspective through the contribution of winter sports activities for pupils from inclusive centers
- 175 Stefania Cazzoli, Lorenzo Sciascia, Federico Abate Daga
The role of multistation exercises in physical activity exposure during a curricular primary school lesson for learning and training by game approach: a pilot study
- 185 Ilaria Tosi, Francesco Casolo, Daniele Coco, Dario Cipani
Movement and Active Breaks Let's Play Together!
- 199 Amalia Tinto, Marta Campanella
Playing with rhythm

- 207 Cristiana D'Anna, Filippo Gomez Paloma
Observing from a biopsychosocial perspective during play activities in physical education: good practices from the *EduFiBES* project
- 225 Zhou Yali, Garry Kuan, Yee Cheng Kueh, Huang Zhijian, Ren Wenxuan
Application of games-based tennis in physical education for children
- 251 Dario Colella, Cristina d'Arando, Domenico Monacis
The Game to promote physical literacy and sports literacy. Physical education teacher training through teaching styles and the variability of practice
- 271 Jiarun Wu, Garry Kuan, Lou Hu, Hu Xiaoyu, Sabo Abdulwali, Kueh Yee Cheng
A Feasibility Study on Using Sports Games to Enhance Mental Toughness of Chinese Adolescent
- 287 Eduard Feroyan
The use of games in teaching sport swimming to children 4-7 years old
- 301 Zhang Haiyan, Garry Kuan, Kueh Yee Cheng, Luo Yi
Traditional Games in Physical Exercise of 3-6-year-old Children in China: Educational Value-Application Strategy
- 321 Elvis Mehid, Garry Kuan, Chin Ngien Siong
Iban Community of Sarawak Traditional Games For Enhancing Physical Activity

SITUATION

- 335 Eduarda Eugenia Dias de Jesus, Valeska Iliencko Villela Souto, Fabricio Faitarone Brasilino, Alexandre Rosa, Pedro Jorge Cortes Morales
Physical exercise and teenagers' Self-Determination: a theoretical-reflective review
- 343 Elena Romanato, Giampaolo Santi, Attilio Carraro
Competitive games with children and youth in physical education and sport in Italy
- 367 Clive Palmer, Andrew Sprake, David Grecic
Harnessing the power of '*game*' in Physical Education: towards the CARE Curriculum

- 387 Mouhamed Dit Momar Talla Ndongo, Djibril Seck, Hameth Dieng
Physical and Sports Education (PSE) in Senegal: a critical sociology of a school discipline in critical situation
- 409 Jaume Bantulà Janot
Presence of the play in Physical Education in the new educational curricula of Catalonia
- 427 Chatkamon Singnoy, Tanida Julvanichpong
Folk Gamed for Physical Education in Thailand
- 439 Italo Sannicandro
The games in Long-Term Athletic Development (LTAD) for youth sports: exploratory learning experience and opportunities to increase physical efficiency
- 459 Dhirveenraj Raj Kumar, Garry Kuan
Traditional Games for Enhancing Physical Activity Among Children and Youth in Suburban Area
- 469 Ngien-Siong Chin, Eng-Hoe Wee, Chatkamon Singhoy, Elvis Mehid
Traditional Games in the Malaysian Education Curriculum for Physical Education
- 479 Shuai Ying, Wang Shaoshen, Garry Kuan, Yee Cheng Kueh, Liu Xian
Traditional Chinese Game in Physical Education, Physical Activities and Sport for Children and Youth in China
- 501 Zhao Mengyuan, Wang Shiyue, Garry Kuan, Kueh Yee Cheng, Zhou Ke
Effectiveness of mini-basketball games for children: A Systematic Review
- 513 Wan Zhen Lee, Garry Kuan, Muhammad Hafiz Hanafi, Yee Cheng Kueh
Effect of exergaming in physical activity among children and adolescent: A review
- 527 **Conclusions and suggestions for future research**
Dario Colella, Domenico Monacis, Branislav Antala, Sergii Ivashchenko, Dario Novak
- 533 **List of Contributors**
- 537 **Index of Authors**

The role of multistation exercises in physical activity exposure during a curricular primary school lesson for learning and training by game approach: a pilot study

Stefania Cazzoli

School of Exercise and Sport Sciences SUSIM, and Department of Philosophy and Education Science DFE, University of Turin, Italy

Lorenzo Sciascia

School of Exercise and Sport Sciences SUSIM, Adapted Training and Performance-Research Group, University of Turin, Italy

Federico Abate Daga

Department of Clinical and Biological Sciences, University of Turin, Italy

Abstract

Background: The benefits of physical activity on primary school children are well reported. Nevertheless, the percentage of children meeting the minimum daily amount of recommended moderate-to-vigorous physical activity (MVPA) declines continuously. Many countries worldwide promote projects in primary schools to sensitize children to physical activity to fight against this phenomenon. However, most of the interventions focused on the available time dedicated to physical activity and not on the quality of time usage. Thus, this study aimed to identify the role of exercise set-up on the actual time spent by active moving during PA intervention using the «multistation exercise» method. **Methods:** Twenty children from the third grade of a primary school were recruited to perform 60 minutes of extra-curricular PA intervention. Children performed three different exercise formats, and active time spent by moving or exercising was calculated by video recording the intervention. **Results:** The final sample size was 12 primary school children (weight 26.83 ± 4.26 kg; height 125.83 ± 6.07 cm, BMI 16.87 ± 1.804 kg/m²). The mean active time spent moving or exercising was 50.75 ± 6.46 seconds in the one-station format and 100.13 ± 10.78 seconds in the two-station format, and 148.71 ± 7.26 seconds in the four-stations format. Repeated measure ANOVA showed a significant solid difference among the three formats ($p < 0.001$). Bonferroni Post Hoc showed a considerable difference between one and two workstation formats ($p < 0.001$; +97% of active time), between two and four workstation forms ($p < 0.001$; +49% of active time) and between one and four workstation set-ups ($p < 0.001$; +193% of active time) **Conclusions.** The multistation exercise method may be an excellent opportunity to optimize PA intervention in primary school children.

Keywords: *Physical education in primary school, Active time, Exercise organization, Exercise management, Physical education teacher.*

Introduction

Nowadays, the benefits of physical activity on children are well reported. Many studies declared that an appropriate amount of physical activity during childhood helps children improve cardiovascular fitness, body composition, and insulin sensitivity and prevent medical conditions such as overweight and obesity (Carrel et al., 2005; D W Harsha, 1995; Errisuriz et al., 2018). Moreover, regular physical activity can also provide multiple benefits to school performance (de Greeff et al., 2018). Despite this, the percentage of children meeting the minimum daily amount of recommended moderate-to-vigorous physical activity (MVPA) declines between 6–8 years (76.1%) and 9–11 years (64,7%) (Dobbins et al., 2013). In addition, the increased emphasis on academic success and the common belief that some subjects are more important than others in school careers have reduced physical education time in schools (Davison & Birch, 2001; Murray & Ramstetter, 2013). However, physical education interventions at school are the best way to reach many children across demographic groups (Weedon et al., 2022). Therefore, it has become a popular method to target physical education and fitness development on a large scale (Errisuriz et al., 2018; Fakhouri et al., 2013). Many countries all over the world encourage physical activity (PA) classes to try to guarantee the minimum daily amount of MPVA for their children. However, most of the PA interventions mainly focused on the general amount of time spent on physical activity (i.e. 10 hours of intervention) without considering the actual active time that children spent moving themselves. A 60-minute lesson full of movement, plays and motor task is entirely different from the same intervention but characterised by long sitting time due to game or rules exposition or waiting for your turn to play. Despite this, no data are available about the efficacy of PA intervention management on physical activity exposure. Thus, this study aims to identify the role of exercise set-up on the actual time spent by active moving during PA intervention using the «multistation exercise» method. This approach is utilised in a team sport or activities with many participants because it optimises time, space and the coach's control (Singh et al., 2018). We hypothesise that this approach could be extended to children's classes, which could help manage the intervention better and upgrade the time spent by active moving.

Research Question

Is the time spent in Physical Activity interventions really «active» for primary school children? Efficacy of the multistation exercise approach to increase the active time during PA intervention in primary school children.

Materials and methods

Study design and ethical considerations

The present study was conducted in a primary school in the metropolitan area of Turin (North-western Italy). Children from the third grade of the Italian primary school system were asked to perform a «one-shot» of extra physical activity. Participation was voluntary-based and extra-curricular. Parents should give their consent by signing an informed consent statement. Children without parents' approval were excluded from this study. The Institutional Research Bioethics Committee of the University of Turin approved this study (number of licenses: 0258948)

Participants

Twenty children from the third grade of a primary school in the metropolitan area of Turin were recruited for this study. Participants should be eight years old and regularly attend third-grade classes to be included in this study. In addition, parents must give written permission to participate by signing informed consent and an image release form. Finally, all participants should not suffer from any physio-psychological pathology (medically certified), have been severely injured in the last three months before the study (i.e. broken arm or leg) or be considered unfit to practice any physical activities because of physical or cognitive disabilities.

Procedures

Children considered eligible for this study were invited to participate in a 60-minute of additional physical activity intervention on a Saturday morning in the local sports centre, where an outdoor futsal pitch was booked and dedicated to this study. The futsal pitch is meanly 36-42 meters long and 18-22 meters large. These measures are stringent to a typical Italian school gym, so the futsal pitch was considered the best choice to replicate the usual space of movement that children have in a school gym. After a 10-minute warm-up, the children were ready to start the intervention. The exercise focus was on motor and cognitive

performance abilities. Children had to run along the track with or without a ball and pay attention to the teacher's colour call. Each colour was related to a specific task or target, so children had to remember the association between the colour and the correct motor task.

The exercise was administered in three different formats to reach the aim of this study. Each form was 10 minutes long, and five minutes of rest were observed between each trial to guarantee a complete recovery and avoid that results may be affected by the incoming fatigue. The first format involved only one station (starting/ending point). Thus, while one child performed the exercise, the remaining participants waited, standing in a row. Conversely, the second format involved two symmetrical stations (starting/ending points). Therefore, the group was divided into two rows, and two children could perform the exercise simultaneously. Furthermore, only two markers were added to the existing exercise track to create this new format. Finally, the third format involved four symmetrical stations (starting/ending points). Therefore, the group was divided into four rows, and two children could perform the exercise simultaneously. Four new cones and markers were added to the exercise track to create this new format. In conclusion, children observed five minutes of final debriefing with the PE teacher, and then the intervention was cleared. The PA intervention was video recorded using an iPad Pro 11» (Apple Inc., Cupertino, California, USA). Then, the video was analysed using the open-source Kinovea version 0.9.5 (www.kinovea.org) to detect the real active time of every child in each exercise format.

Statistical Analysis

All data were analysed using the open-source JASP, version 0.16.4 (Jeffreys's Amazing Statistics Program., University of Amsterdam, the Netherlands). Descriptive statistics ((mean and standard deviation (SD)) were used to present participants' characteristics and data distribution. In addition, repeated measures ANOVA was used to assess differences in real active time spent during PA intervention between the three different exercise formats. Significance was assumed at $p < 0.05$.

Results

Twenty children were recruited for this study. However, eight were excluded from this study because they did not meet the inclusion criteria, or their parents did not sign written permission for participation. Therefore, the remaining

twelve children participated in this study and their data were collected for further analysis. Participants' anthropometric characteristics are represented in Table 1. The mean weight was 26.83 ± 4.26 kg; the mean height was 125.83 ± 6.07 cm, and the mean BMI was 16.87 ± 1.804 kg/m². The entire sample belonged to the third grade of the Italian primary school system, and all were male and eight years old when this study was realised.

Furthermore, descriptive data regarding the active time spent during PA intervention are shown in Table 2. Data from all formats were sampled on a 10-minute exercise. For the set-up with only one workstation, the mean active time during 10 minutes of exercise was 50.75 ± 6.46 seconds. Conversely, in the 2-station format, the mean active time during 10 minutes of exercise was 100.13 ± 10.78 seconds. Finally, in the four-station form (multistation), the mean active time during 10 minutes of exercise was 148.71 ± 7.26 seconds. Repeated measure ANOVA showed a significant solid difference among the three formats ($p < 0.001$). Bonferroni Post Hoc showed a considerable difference between one and two workstation formats ($p < 0.001$; +97% of active time), between two and four workstation forms ($p < 0.001$; +49% of active time) and between one and four workstation set-ups ($p < 0.001$; +193% of active time). Data from repeated measure ANOVA and Bonferroi Post Hoc are represented in Tables 3 and 4.

Discussion

This study aimed to identify the role of exercise set-up on the actual time spent by active moving during PA intervention using the «multistation exercise» method. This approach is commonly used in team sports training schedules to enhance performance and prevent injuries (Eils et al., 2010; Pérez-Silvestre et al., 2019) because it provides the opportunity to train the entire squad at the same time on specific tasks. Therefore, this approach seems to be an excellent method in PA intervention in primary school to optimise the active time spent by moving and physically exercising. The results of this study show that moving from one to two workstations in a ten-minute exercise improved 97% of the time paid by active moving or exercising. Furthermore, moving from two to four workstations increased 49% of time spent actively moving or exercising. Finally, the most significant gains were detected moving from one to four workstations, where the time expended actively moving or exercising increased by up to 193%. Thus, «multistation exercise» may reveal an intelligent PA intervention management able to raise the total active time spent by moving or exercising in optimal use of space and stuff. Only four cones and four markers were added to increase

from one to four workstations. Cones and markers are simple elements that are usually supplied in each school to increase up to 193% of the active time of each child. Unfortunately, no studies have investigated this topic the same way as this research did. Thus, a comparison of our results is not properly accessible. Most literature focuses on the active recess's crucial role (Murray & Ramstetter, 2013). It was estimated that an active recess with moderate to vigorous PA could significantly improve school aptitudes, creativity and cognitive flexibility (Pedro Ángel et al., 2021), reduce BMI (Thalken et al., 2021) and positively affect physical fitness and social factors (Chen et al., 2018; Haapala et al., 2014). Our findings agree with these studies underlining the need to improve the time spent actively moving during primary school. However, this research moves further and provides a way to manage PA interventions to optimise time and space during physical education lessons. Finally, a limitation of this study can be the small sample size. However, Italian primary school classes could guest up to 25 children. Thus, extensive samples must be obtained involving more than one grade and, consequently, more than one school. Therefore, as a pilot study, we decided to limit the intervention to one school and one grade. Future researchers should increase the sample size by using the multistation exercise method on large numbers of classes to confirm and improve these findings.

Conclusion

In conclusion, this study underlines the efficacy of the multistation exercise method in increasing the active time spent by moving or exercising during PA intervention. With some simple «tricks», it can quickly move from one to four stations guaranteeing children an increase of up to 193% of active time during PA intervention. These findings may be beneficial for primary school teachers to raise the quality of physical education lesson management.

Figures and Tables

Descriptive Statistics of anthropometric characteristics							
		WEIGHT (kg)		HEIGHT (cm)		BODY MASS INDEX (BMI) (kg/m ²)	
Valid		12		12		12	
Missing		0		0		0	
Mean		26.833		125.833		16.871	
Std. Deviation		4.260		6.073		1.804	
Minimum		21.000		113.000		14.640	
Maximum		34.000		139.000		20.450	

Table 1 - Anthropometric characteristics of the sample. Participants belonged to the third grade of the Italian primary school system, and all were eight years old when this study was realised. Mean weight was 26.83 ± 4.26 kg; mean height was 125.83 ± 6.07 cm and mean BMI was 16.87 ± 1.804 kg/m²

Descriptive Statistics of real active time spent during PA interventions							
		1 WORKSTATION		2 WORKSTATIONS		4 WORKSTATIONS	
Valid		12		12		12	
Missing		0		0		0	
Mean		50.750*		100.125*		148.708*	
Std. Deviation		6.455*		10.773*		7.216*	
Minimum		43.900*		87.200*		129.600*	
Maximum		64.900*		123.800*		157.300*	

*seconds

Table 2 - Descriptive representation of real active time spent during PA interventions. For the one-station format, the mean active time during 10 minutes of exercise was 50.75 ± 6.46 seconds.

Conversely, in the 2-station format, the mean active time during 10 minutes of exercise was 100.13 ± 10.78 seconds. Finally, in the four-station format (multistation), the mean active time during 10 minutes of exercise was 148.71 ± 7.26 seconds

Repeated Measure Anova - Within Subjects Effects					
Cases	Sum of Squares	df	Mean Square	F	<i>p</i>
RM Factor 1	57576.264	2	28788.132	404.451	<.001
Residuals	1565.923	22	71.178		

Note. Type III Sum of Squares

Table 3 - Repeated Measure ANOVA, Within Subject Effects. A significant difference was detected among different formats

Post Hoc Comparisons – with Bonferroni Method								
		95% CI for Mean Difference						
		Mean Difference	Lower	Upper	SE	t	P _{bonf}	
1 WORK-STATION	2 WORK-STATIONS	-49.375	-58.300	-40.450	3.444	-14.335	< .001	***
	4 WORK-STATIONS	-97.958	-106.883	-89.033	3.444	-28.441	< .001	***
2 WORK-STATIONS	4 WORK-STATIONS	-48.583	-57.508	-39.658	3.444	-14.106	< .001	***

*** p < .001

Note. P-values and confidence intervals were adjusted for comparing a family of 3 estimates (confidence intervals were corrected using the Bonferroni method)

Table 4 - Post Hoc comparison with Bonferroni Method. A significant difference was detected between one and two workstation formats (p< 0.001), between two and four workstation forms (p<0.001) and between one and four workstation set-ups (p<0.001)

References

- Carrel, A. L., Clark, R Randall, Peterson, S. E., Nemeth, B. A., Sullivan, J., & Allen, D. B. (2005). *Improvement of Fitness, Body Composition, and Insulin Sensitivity in Overweight Children in a School-Based Exercise Program A Randomized, Controlled Study.*
- Chen, W., Hammond-Bennett, A., Hypnar, A., & Mason, S. (2018). Health-related physical fitness and physical activity in elementary school students. *BMC Public Health, 18*(1). <https://doi.org/10.1186/s12889-018-5107-4>

- D W Harsha. (1995). The benefits of physical activity in childhood. *The American Journal of The Medical Sciences*, 310, S109–S113.
- Davison, K. K., & Birch, L. L. (2001). Childhood overweight: A contextual model and recommendations for future research. *Obesity Reviews*, 2(3), 159–171. <https://doi.org/10.1046/j.1467-789x.2001.00036.x>
- de Greeff, J. W., Bosker, R. J., Oosterlaan, J., Visscher, C., & Hartman, E. (2018). Effects of physical activity on executive functions, attention and academic performance in preadolescent children: a meta-analysis. In *Journal of Science and Medicine in Sport* (Vol. 21, Issue 5, pp. 501–507). Elsevier Ltd. <https://doi.org/10.1016/j.jsams.2017.09.595>
- Dobbins, M., Husson, H., Decorby, K., & Larocca, R. L. (2013). School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. In *Cochrane Database of Systematic Review*, 2. John Wiley and Sons Ltd. <https://doi.org/10.1002/14651858.CD007651.pub2>
- Eils, E., Schröter, R., Schröderr, M., Gerss, J., & Rosenbaum, D. (2010). Multistation proprioceptive exercise program prevents ankle injuries in basketball. *Medicine and Science in Sports and Exercise*, 42(11), 2098–2105. <https://doi.org/10.1249/MSS-0b013e3181e03667>
- Errisuriz, V. L., Golaszewski, N. M., Born, K., & Bartholomew, J. B. (2018). Systematic Review of Physical Education-Based Physical Activity Interventions Among Elementary School Children. In *Journal of Primary Prevention* (Vol. 39, Issue 3, pp. 303–327). Springer New York LLC. <https://doi.org/10.1007/s10935-018-0507-x>
- Fakhouri, T. H. I., Hughes, J. P., Brody, D. J., Kit, B. K., & Ogden, C. L. (2013). Physical activity and screen-time viewing among elementary school-aged children in the United States from 2009 to 2010. *JAMA Pediatrics*, 167(3), 223–229. <https://doi.org/10.1001/2013.jamapediatrics.122>
- Haapala, H. L., Hirvensalo, M. H., Laine, K., Laakso, L., Hakonen, H., Kankaanpää, A., Lintunen, T., & Tammelin, T. H. (2014). Recess physical activity and school-related social factors in Finnish primary and lower secondary schools: Cross-sectional associations. *BMC Public Health*, 14(1). <https://doi.org/10.1186/1471-2458-14-1114>
- Murray, R., & Ramstetter, C. (2013). The crucial role of recess in school. *Pediatrics*, 131(1), 183–188. <https://doi.org/10.1542/peds.2012-2993>
- Pedro Ángel, L. R., Beatriz, B. A., Jerónimo, A. V., & Antonio, P. V. (2021). Effects of a 10-week active recess program in school setting on physical fitness, school aptitudes, creativity and cognitive flexibility in elementary school children. A randomised-controlled trial. *Journal of Sports Sciences*, 39(11), 1277–1286. <https://doi.org/10.1080/02640414.2020.1864985>
- Pérez-Silvestre, Á., Albert-Lucena, D., Gómez-Chiguano, G. F., Plaza-Manzano, G., Pecos-Martín, D., Gallego-Izquierdo, T., Martín-Casas, P., & Romero-Franco, N. (2019). Six weeks of multistation program on the knee proprioception and performance of futsal players. *Journal of Sports Medicine and Physical Fitness*, 59(3), 399–406. <https://doi.org/10.23736/S0022-4707.18.08141-0>

- Singh, N., Gupta, R., & Mahalakshmi, V. N. (2018). Multistation exercises: a combination of problem-based learning and team-based learning instructional design for large-enrollment classes. *Adv Physiol Educ*, 42, 424–428. <https://doi.org/10.1152/advan.00023.2018>.
- Thalken, J., Massey, W. v., Szarabajko, A., Ozenbaugh, I., & Neilson, L. (2021). From policy to practice: Examining the role of recess in elementary school. *Public Health in Practice*, 2, 100091. <https://doi.org/10.1016/j.puhip.2021.100091>
- Weedon, B. D., Liu, F., Mahmoud, W., Burden, S. J., Whaymand, L., Esser, P., Collett, J., Izadi, H., Joshi, S., Meaney, A., Delextrat, A., Kemp, S., Jones, A., & Dawes, H. (2022). Declining fitness and physical education lessons in UK adolescents. *BMJ Open Sport and Exercise Medicine*, 8(1). <https://doi.org/10.1136/bmjsem-2021-001165>.