



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A survey on the milk chemical and microbiological quality in dairy donkey farms located in NorthWestern Italy

This is the author's manuscript Original Citation: Availability: This version is available http://hdl.handle.net/2318/1507557 since 2017-07-27T14:05:50Z Published version: DOI:10.1016/j.foodcont.2014.08.019 Terms of use: Open Access Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright

(Article begins on next page)

protection by the applicable law.





This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in FOOD CONTROL, 50, 2015, 10.1016/j.foodcont.2014.08.019.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), 10.1016/j.foodcont.2014.08.019

The publisher's version is available at: http://linkinghub.elsevier.com/retrieve/pii/S095671351400468X

When citing, please refer to the published version.

Link to this full text: http://hdl.handle.net/2318/1507557

This full text was downloaded from iris - AperTO: https://iris.unito.it/

A survey on the milk chemical and microbiological quality in dairy donkey farms located in NorthWestern Italy

Laura Cavallarin¹, Marzia Giribaldi¹*, Maria de los Dolores Soto- Del Rio², Gandolfo Barbarino³, Maria Silvia Gennero⁴, Tiziana Civera²

¹ Istituto di Scienze delle Produzioni Alimentari, Consiglio Nazionale delle Ricerche, Largo

Braccini 2, 10095 Grugliasco, Italy. laura.cavallarin@ispa.cnr.it; marzia.giribaldi@ispa.cnr.it

² Dipartimento di Scienze Veterinarie, Università di Torino, Largo Braccini 2, 10095 Grugliasco,

 $Italy.\ \underline{mariadelosdolores.sotodelrio@unito.it;}\ \underline{tiziana.civera@unito.it}$

³ Regione Piemonte, Direzione Sanità, Settore Prevenzione e Veterinaria, Via Lagrange 24, 10123 Torino, Italy. <u>gandolfo.barbarino@regione.piemonte.it</u>

⁴ Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle d'Aosta, via Bologna 148,

10154 Torino, Italy. mariasilvia.gennero@izsto.it

* Corresponding author. Tel.: +39.011.670.9231 Fax: +39.011.670.9297

email address: marzia.giribaldi@ispa.cnr.it

The authors disclose any conflict of interest.

All authors have participated to both research and article preparation. All authors have approved the

final version of the article.

The present study received no financial support.

Abstract

There is a growing interest in donkey's milk as food for sensitive consumers, such as infants with cow's milk protein allergy and elderly people. The aim of this study was to carry out a survey on the dairy donkeys farming in Piedmont, Italy. The research was conducted in order to analyze the farm characteristics as well as the chemical and microbiological quality of milk. All the farms were small-sized, family-run, and, in most cases, animals were farmed semi-extensively. The donkey milk from Piedmont farms was characterized by a protein content around 1.5 g/100 mL and a fat content lower than 0.1 g/100 mL. Lysozyme activity was considerably higher than that reported in raw cow milk. The milk microbiological profile greatly differed among the farms. Milk sampled in the farm that performed hand milking showed total viable counts significantly lower than milk collected in the farms equipped with automatic milking. Samples were tested for several pathogens and negative results were observed, except for the detection of *B. cereus* in one sample. The survey provided useful data for the laying down of recent regional regulation for the production and commercialization of donkey's milk. The results of the survey indicate that further research is needed in order to define the best management and nutritional strategies for the improvement of the quali-quantitative production of dairy donkeys.

Key words: donkey farming, donkey milk, microbiological quality, protein profile, lysozyme

Abbreviations:

DM: donkey milk; LAB: lactic acid bacteria; PVDF: polyvinylidene fluoride; TVC: total viable count.

1. Introduction

There is a growing interest in donkey farming for milk production, since donkey's milk (DM) is considered a valid alternative foodstuff in terms of clinical tolerability, palatability and nutritional adequacy for children affected by cow's milk protein allergy (Iacono et al., 1992; Mansueto et al., 2013; Monti et al., 2007; Monti et al., 2012).

Since 2005, the number of published papers on DM has increased to almost 30 per year, the great majority by Italian researchers. Although data covering the potential of DM use as a substitute of cow's milk for allergic individuals are available, information regarding donkey farm management and some aspects of milk production are limited, such as the study of its main microbial communities. The majority of surveys on DM quality were carried out in the South of Italy, mainly on Martina Franca (D'Alessandro, De Petro, Claps, Pizzillo, & Martemucci, 2009; D'Alessandro & Martemucci, 2007; D'Alessandro, Martemucci, Jirillo, & Leo, 2011; Martemucci & D'Alessandro, 2012) and Ragusana breeds (Alabiso, Giosuè, Alicata, Mazza, & Iannolino, 2009; Giosuè, Alabiso, Russo, Alicata, & Torrisi, 2008). One very recent report was published on Amiata donkey milk chemical composition (Martini, Altomonte, & Salari, 2014). Some data are also available for Chinese donkeys (Guo et al., 2007) and for Balkan Littoral-Dinaric breed (Ivanković et al., 2009; Sarić et al., 2012). The literature indicates that fat content in DM range from 0.3 to 1.8 g/100 mL, while the protein content is reported to be less variable, with values from 1.4 to 1.8 g/100 mL (Medhammar et al., 2012). DM is also known to contain high concentration of lysozyme, if compared to other mammalian milks. Donkey's lysozyme, similarly to cow's, displays antibacterial activity against a vast number of bacteria (Conte, Foti, Malvisi, Giacopello, & Piccinini, 2012; Zhang, Zhao, Jiang, Dong, & Ren, 2008; Šarić et al., 2012), affecting directly the composition of the milk microbiome (Sarno, Santoro, Di Palo, & Costanzo, 2012). Very recently, it has been shown that the addition of jenny milk during cheese-making reduced the number of coliforms in the resulting cheese (Cosentino, Freschi, Paolino, & Valentini, 2013). Further compositional aspects of DM, that are not considered in the present study, were extensively reviewed in two recent reports by Salimei & Fantuz (2012) and Medhammar and coworkers (2012).

Low bacterial counts are generally reported in DM, in comparison to milk and dairy products from other animals (Pilla, Daprà, Zecconi, & Piccinini, 2010; Salimei et al., 2004; Sarno et al., 2012). Previous reports have shown that the DM microbiota is mainly composed by lactic acid bacteria (Zhang et al., 2008; Šarić et al., 2012). Nevertheless, undesirable species bacteria, such as *Streptococcus* spp., *Staphylococcus* spp. and coliforms, have been also detected in DM (Pilla et al., 2010; Sarno et al., 2012).

The aim of this research is to provide the first overview of the chemical and microbiological characteristics of milk produced in dairy donkey farms in the North West of Italy. This outline provided a significant contribution to the recent publication of a regional regulation for the production and commercialization of DM (Regione Piemonte Direzione Sanità Settore Prevenzione e Veterinaria, 2013).

2. Materials and Methods

2.1. Farm description and milk sampling

Five donkey dairy farms located in the North Western Italy, in Piedmont, were surveyed during this study, and a questionnaire was filled in for each farm. Data and samples were collected during Autumn 2012 and Spring 2013. The on-farm survey included a face-to-face interview with the farm manager. We developed a questionnaire, which consisted of semi-closed questions, including topics related to farm management and husbandry practices.

The DM samples were obtained from a different number of jennies and by different milking practices, depending on farm management. In each surveyed farm, one liter of DM was sampled for

chemical and biochemical analyses, while six replicate DM aliquots (300 mL each) were collected in sterile bottles for microbiological screening.

Immediately after collection, DM samples were transported, refrigerated and either used fresh (as for microbiological analyses), stored within 6 hours from the collection at -20 °C, or lyophilized.

2.2. Milk chemical and biochemical analyses

The DM samples were analyzed for gross energy, dry matter, crude protein and ether extract contents. In addition, the lysozyme activity of each sample was determined using a fluorescence-based assay (EnzCheck[®] Lysozyme Assay Kit, Life Technologies Italia, Monza, Italy).

Dry matter content was measured on DM using a gravimetric method (Baldini, Fabietti, Giammarioli, Onori, Orefice, & Stecchini, 1996). Gross energy content was determined in excess oxygen by adiabatic bomb calorimeter (Mod. 700, IKA GmbH & Co., Staufen, Germany), using benzoic acid as a reference (26.454 MJ/kg). The nitrogen content in milk samples was estimated by Kjeldahl-based block digestion method (AOAC Official Method 991.20, 2000), using a 2020 Tecator Digestor (VWR International Pbi, Milano, Italy), a Kjeltec-System 1002 Distilling Unit (Foss Italia, Padova, Italy) and a 655 Dosimat automatic titrimeter (Metrohm Italiana, Origgio, Italy). The total fat content of DM samples was measured gravimetrically on ether extract using a Soxhtraction device (VWR International Pbi), following manufacturer's instructions. All the analyses were performed on lyophilized DM samples in triplicate.

Lysozyme activity assay was conducted by EnzCheck[®] Lysozyme Assay Kit following manufacturer's instruction in triplicate on 500-fold diluted DM samples, using a fluorescence microplate reader (Victor 3D, Perkin Elmer, Waltham, USA), equipped with fluorescein filters (485 nm emission and 535 nm excitation).

2.3. Milk protein profile

DM samples were accurately mixed and vortexed at room temperature in order to ensure homogeneity, then 100 μ L from each sample were taken up to 1 mL with milliQ water and vortexed thoroughly again. One aliquot (100 μ L) of each diluted sample was mixed with the same volume of

NuPAGE® LDS Sample Buffer (2X diluted) (Life Technologies Italia), containing 50mM dithiothreitol (Sigma Aldrich, Milano, Italy) as reducing agent, and warmed to 70°C for 10 minutes. Each sample solution (10 μ L) was then loaded onto a 10-well, NuPAGE[®] Bis-Tris mini gels (12% polyacrylamide, 1mm width). One lane on each gel was loaded with 5 μ L of Mark12[®] Unstained Standard (Life Technologies Italia).

The proteins were separated on a Novex Mini-cell (Life Technologies Italia) filled with cold 1X NuPAGE[®] MES SDS Running Buffer (Life Technologies Italia), at 200V. The gels were then stained with colloidal Coomassie Blue staining (Candiano et al., 2004) and digitized with a ImageScanner device (Amersham Pharmacia, now GE Healthcare Life Sciences, Uppsala, Sweden) at 300 dpi.

Two bands were excised with a sterile scalpel and passively eluted from the gel pieces onto PVDF membranes, as previously described (Reuter et al., 2005) to be subjected to N-terminal sequencing. The membranes were then microsequenced on a Procise 492 protein sequencer (Applied Biosystems, now Life Technologies Italia). All the chemicals used in the procedure were from Life Technologies Italia. The N-terminal amino acid sequences were searched with the MS-Homology software package on NCBI non-redundant database (NCBInr2013.6.17).

2.4. Milk microbiological analyses

For this part of the study, a third sampling was carried out during the 2014 Spring season.

Two aliquots of each DM sample were collected and stored in sterile conditions at 4°C. One of them was analyzed the day after milking, while the second one was examined 5 days after the sampling.

The total viable count (TVC) was performed onplate count agar, using the inclusion method. One mL of undiluted samples and of each 10⁻¹, 10⁻² and 10⁻³dilutions in a mixture of physiological saline solution and peptone (85:15 v:v) (OXOID LTD, Basingstoke, Hampshire, England) were included in the plate count agar (OXOID LTD) and plates were incubated for 72 hours at 30°C before counting. The results were then expressed as cfu/mL (UNI EN ISO 4833: 2004). Lactic acid bacteria (LAB) colonies, grown on MRS agar (OXOID LTD) at 30°C for 48 hours in anaerobiosis were counted (ISO/FDIS 15214:1998). According to ISO 6579:2002, for the detection of *Salmonella* spp., BPW enrichment media (OXOID LTD) was used for an incubation of 24 hours at 37°C, a second incubation was performed with Rappaport media (OXOID LTD) (41°C for 24 hrs) and with MKTTn (OXOID LTD) at 37°C for 24 hrs. Both incubation products were finally grown on XLD agar medium (OXOID LTD) for 24 hrs at 37°C.

For *E. coli*, EN/ISO 16649/2:2001 regulation parameters were followed, growing in TBX agar (OXOID LTD) at 44°C for 24 hrs. Regarding *E.coli* O157, the enrichment media TSBm (OXOID LTD) was utilized, incubated at 41°C for 24 hours. Dynabeads anti-*E. coli* O157 (Invitrogen Dynal AS, Oslo, Norway) were used to selectively capture these bacteria. *E. coli* O157 was grown on TBX agar (OXOID LTD) at 41°C for 24 hrs and on CT-SMAC (OXOID LTD) at 37°C for 24 hrs.

Listeria monocytogenes presence was evaluated by enriching with Demi-Fraser (Sifin, Berlin, Germany) and Fraser media (Sifin) incubated at 37°C for 24 hours (EN/ISO 11290/1:1996), a subsequent growth on OCLA and PALCAM agar mediums (OXOID LTD) was done for both enriching media for 48 hrs at 37 °C.

For *B. cereus*, the Bacillus agar base medium (OXOID LTD) was used, incubating at 37°C for 24 hrs (EN/ISO 7932:2005).

The detection of *Campylobacter* spp, (Bacteriological Analytical Manual - FDA (2001)) consisted in an initial enrichment with Bolton Broth media (OXOID LTD), an incubation of 41°C for 24 hours in anaerobiosis and a final growth on CCD agar (OXOID LTD) in the same conditions but for 48 hrs.

7

All the analysis were done in technical triplicate.

2.5. Statistical analysis

Significant differences in total viable count (log10 cfu/mL) among the farms were assessed by ANOVA at p \leq 0.05, and classes of uniformity were grouped according to Tukey's post-hoc test. Significant differences in lactic acid bacteria count (log10 cfu/mL) among the farms were assessed by Kruskal-Wallis non parametric test at $\alpha \leq 0.05$, and classes of uniformity were grouped according to Dunnet-C post-hoc test, not assuming equal variances. The statistical significance of increase in TVC and LAB counts from day 0 to day 5 were evaluated by Student T test (not shown).

2.6 PCR for confirmation of Bacillus cereus

DNA was extracted by boiling procedure from broth and amplified as previously reported (Park, Kim, Kim, Kim & Kim, 2007). The PCR product was visualized in a 2% agarose gel (Invitrogen, Carlsbad, USA) in the Gel Doc XR+ system (BIO-RAD Italia, Segrate, Italy).

3. Results and discussion

3.1. Farm characteristics

Data regarding farm management and husbandry practices are reported in Table 1. All the farms were small-sized, family-run, and recently began the donkey farming activity. In most cases, animals were farmed semi-extensively, partially grazing and partially fed *ad libitum* with on-farm produced hay. Farms A, C and E were located in agricultural areas. Farm B was located in a mountain rural area. Farm D, which was located in a hilly area, did not produce or sell milk during the surveyed timeframe. In all the farms surveyed, donkeys spent around 12 h/d on pasture. All farms, except one, used machine milking and milked the jennies once a day. The mean milk yield

was around 1 L/day/animal. All the farms sold the milk directly to families of cow's milk protein allergic children. Some of the producers also allocated part of the milk for cosmetic production.

3.2. Donkey milk chemical characterization

The average quality of DM produced in Piedmont farms was assessed by evaluating its chemical characteristics, in Autumn and Spring. Table 2 reports the mean values recorded for each assayed parameter in the two different samplings, as well as their overall means. Milk was characterized by a mean protein content around 1.5 g/100mL and a mean fat content lower than 0.1 g/100mL. Farm C milk was the only sample displaying a slightly higher fat concentration, ranging from 0.2 to 0.4 g/100mL, depending on the season. Fat is the most variable component of DM, ranging from 0.1 g/100mL (Salimei et al., 2004) to 1.8 g/100mL (Guo et al., 2007) in different reports, being affected from both lactation and foaling season (Cosentino et al., 2013; Giosuè et al., 2008; Ivanković et al., 2009; Salimei et al., 2004). The Piedmontese DM showed very low fat concentration, even below the lower limit reported in literature. The low DM fat concentration and its consequent low energy content are the main limits to its use in nutrition of children allergic to cow's milk protein, in their first year of life, since recommended dietary allowances may not be reached, unless adequate supplementation is provided (D'Auria, Mandelli, Ballista, Di Dio, & Giovannini, 2011). Future perspectives on DM research thus include studies aimed at setting up new nutritional strategies for the lactating donkey, in order to increase the DM energetic value. On the other hand, this feature makes DM an hypo-caloric and highly digestible food for consumers with specific dietary requirements, such as athletes and elderly people.

The observed values for milk protein content are in accordance to the results reported for Italian DM. Milk was reported to range between 1.2 g/100mL (Cosentino et al., 2013) and 1.9 g/100mL (Giosuè et al., 2008), and it has been shown to be affected by the lactation stage, the lactation and foaling season (D'Alessandro et al., 2009; Giosuè et al., 2008; Ivanković et al., 2009; Martemucci & D'Alessandro, 2012; Salimei et al., 2004; Tidona et al., 2014).

The DM produced in Piedmont farms was characterized by a mean lysozyme activity of 162 U/ μ L, very high in comparison to other dairy species, as also reported by Conte et al. (2012). Such an elevated lysozyme activity may explain the low incidence of mastitis in jennies, that usually follows physical injuries to the glands or drying off (Conte, Piccinini, Daprà, & Gagliano, 2006).

3.3. Donkey milk protein profile characterization

The mono-dimensional gradient electrophoresis pattern of the proteins contained in milk samples from the different farms is reported in Figure 1. The NuPAGE® gels coupled with MES SDS Running Buffer were chosen in order to enhance the separation of medium-low molecular weight proteins, and to allow a reliable and easy comparison of the protein patterns from the DM samples collected in the different farms. Small differences in the profile of the different DM samples could be seen. Milk collected from Farm B displayed a slightly different protein pattern, showing two protein bands, characterized by an apparent different abundance compared to the other milk samples, especially in the Spring sampling. The two protein bands from Farm B milk sample were subjected to N-terminal sequencing in order to be identified. Results are detailed in Table 3. The two bands represented a mixture of beta lactoglobulin 1 and alpha s2 casein B (band 1) and beta lactoglobulin 1 (band 2). Differential protein glycosylation of beta lactoglobulin and phosphorylation of alpha s2 casein, already observed on the same proteins in previous DM proteomic characterization (Bertino et al., 2010), may play a role in the observed changes in protein bands. It is interesting to note that Farm B was the only farm breeding exclusively Martina Franca donkeys. Polymorphism of protein isoelectrofocusing bands was already observed in Ragusana breed by Criscione et al. (2009), which reported the absence of other DM proteins (alpha s1 casein and beta lactoglobulin II) in some individuals. They also reported the absence of alpha s2 casein in all DM samples. In a very recent study, Tidona et al. (2014) investigated the relevance of DM protein profile individual variability in determining the size of casein micelles and *in vitro* milk protein digestibility. They reported that the size of casein micelles was not modified by the absence of alpha s1 casein, while it was partially correlated with total protein content. They also reported that the absence of beta lactoglobulin II increased the rate of gastric digestion of the lactoglobulin fraction in DM. As alpha casein and beta lactoglobulin isoforms are considered as major allergens in cow milk, the presence/absence of these protein bands may have an impact on the residual allergenic potential of DM, as already reported by Tidona et al. (2014). A deeper insight into DM protein profiles from different breeds may shed light on this feature.

3.4. DM microbiological quality

The occurrence of known milk pathogens in DM was investigated by searching for *Salmonella* spp., *E. coli*, *E. coli* O157, *Listeria monocytogenes, Bacillus cereus* and *Camplylobacter* spp. The results were negative for all the farms at day 0 as well as at day 5, with the exception of farm A in the 2014 Spring sampling, being positive only for B. cereus (1.3*102 cfu/mL). The presence of B. cereus in milk has been previously reported, with seasonal fluctuations in raw and processed cow milk (Bartoszewicz, Hansen, & Swiecicka, 2008). Preliminary studies on DM have shown the presence of *B. cereus*, with counts similar to the ones found in this study (Scatassa et al., 2011).

The total viable and lactic acid bacteria counts obtained for each of the farms on the three samplings are shown in Table 4. The Farm A presented the highest TVC in this study (Table 4). Different microbiological tests and sequencing analyses were performed for some isolates of this farm (data not shown), showing *Pseudomonas* spp. as the group having the highest occurrence, thus suggesting a possible contamination issue.

On the other hand, farm D presented the lowest values for the TVC (Table 4). This farm was run extensively. The animals were free to pasture and were hand milked, not regularly, in contrast with the standard management of the other surveyed farms (Table 1). Moreover, Farm D displayed the lowest TVC value after 5 days, in contrast to farms using automatic milkers, also being the only farm to show a not significant increase of TVC following cold storage, after Student T test comparison of data at day 0 and day 5 (data not shown). Similar results were observed in a previous

study on goat milk in Spain (Delgado-Pertiñez et al., 2003): the authors found that hand milking determined a significant reduction of bacteria in the milk, in comparison to farms that use mechanic procedures. This observation might be due to the fact that hand milking results in a more suitable control of the hygienic management of the farm, as well as of milk collection and conservation. It is also important to mention that in all the studied farms, the milkers were adaptations of machines for goats or cows.

The overall mean for the three samplings of the TVC at day 0 was 2.4*10⁵cfu/mL. This value is one order of magnitude higher than previously reported (Chiavari et al., 2005; Coppola et al., 2002; Salimei et al., 2004; Zhang et al., 2008). Regarding the LAB count, the overall mean number was 1.6*10⁴ cfu/mL (Table 4). This LAB value is consistent with counts recently reported by Carminati et al. (2014) in DM. The LAB count increase in DM during cold storage was not significant, as assessed by Student T test comparison of data at day 0 and day 5 (data not shown), for any of the surveyed farms, thus indicating a possible effect of lysozyme antimicrobial activity against LAB. Gram-positive bacteria, such as LAB, possess a cell wall that can be hydrolyzed by lysozyme (Salazar & Asenjo, 2007), thus causing inhibition of LAB development and, as a consequence, decreasing their abundance in milk.

4. Conclusion

The profile of the donkey milk collected in Piedmont resulted to be quite similar among the surveyed farms, from a chemical point of view. The milk was characterized by a very low fat content, which may indicate the need for implementation of new nutritional protocols for the lactating donkeys, to increase its energetic value. The microbiological content of milk showed significant variations among surveyed farms, indicating a possible effect of the adopted milking practice. A better training of the farm operators is needed for an appropriate and efficient use of the

milking machinery. Besides, there is a need for the implementation of specific instruments to milk donkeys. An improvement in these aspects might help in reducing milk contamination, increasing the hygienic and sanitary properties of donkey milk.

References

- Alabiso, M., Giosuè, C., Alicata, M. L., Mazza, F., & Iannolino, G. (2009). The effects of different milking intervals and milking times per day in jennet milk production. *Animal*, 3(4), 543-547.
- AOAC Official Method 991.20 (1991). Nitrogen (Total) in Milk (15th ed.). In Official Methods of Analysis of AOAC International, Arlington, VA:USA.
- Baldini, M., Fabietti, F., Giammarioli, S., Onori, R., Orefice, L., & Stecchini, A. (1996). Rapporti ISTISAN 96/34: Metodi di analisi per il controllo chimico degli alimenti. Available at http://www.iss.it/binary/publ2/cont/Rapporto%2096-34.1140450878.pdf
- Bartoszewicz, M., Hansen, B. M., & Swiecicka, I. (2008). The members of the *Bacillus cereus* group are commonly present contaminants of fresh and heat-treated milk. *Food Microbiology*, 25(4), 588-596.
- Bertino, E., Gastaldi, D., Monti, G., Baro, C., Fortunato, D., Perono Garoffo, L., et al. (2010).
 Detailed proteomic analysis on DM: insight into its hypoallergenicity. *Frontiers in Bioscience (Elite Edition)*, 2,526-536.
- Candiano, G., Bruschi, M., Musante, L., Santucci, L., Ghiggeri, G. M., Carnemolla, B., et al. (2004). Blue silver: a very sensitive colloidal Coomassie G-250 staining for proteome analysis. *Electrophoresis*, 25(9), 1327-1333.

- Carminati, D., Tidona, F., Fornasari, M.E., Rossetti, L., Meucci, A., & Giraffa, G. (2014).
 Biotyping of cultivable lactic acid bacteria isolated from donkey milk. Lett Appl Microbiol. doi: 10.1111/lam.12275.
- Chiavari, C., Coloretti, F., Nanni, M., Sorrentino, E., & Grazia, L. (2005). Use of donkey's milk for a fermented beverage with lactobacilli. *Lait*, *85*(6), 481-490.
- Conte, F., Foti, M., Malvisi, M., Giacopello, C., & Piccinini, R. (2012). Valutazione dell'azione antibatterica del lisozima del latte d'asina. Considerazioni igienico - sanitarie. *Large Animal Review*, 18(1), 13-16.
- Conte, F., Piccinini, R., Daprà, V., & Gagliano, V. (2006, September). *Evaluation of lysozyme in asses' milk*. Presented at the 60th Convegno Nazionale S.I.S.Vet., Terrasini (BA), Italy.
- Coppola, R., Salimei, E., Succi, M., Sorrentino, E., Nanni, M., Ranieri, P., et al. (2002). Behaviour of Lactobacillus rhamnosus strains in ass's milk. *Annals of Microbiology*, *52*(1), 55-60.
- Cosentino, C., Paolino, R., Freschi, P., & Calluso, A. M. (2013). Short communication: Jenny milk production and qualitative characteristics. *Journal of Dairy Science*, *95*(6), 2910-2915.
- Criscione, A., Cunsolo, V., Bordonaro, S., Guastella, A. M., Saletti, R., Zuccaro, A., et al. (2009). Donkeys' milk protein fraction investigated by electrophoretic methods and mass spectrometric analysis. *International Dairy Journal*, *19*(4), 190-197.
- D'Alessandro, A. G., & Martemucci, G. (2007). Influence of milking number and frequency on milk production in Martina Franca breed asses. *Italian Journal of Animal Science*, 6(SUPPL. 1), 643-645.
- D'Alessandro, A. G., De Petro, R., Claps, S., Pizzillo, M., & Martemucci, G. (2009). Yield and quality of milk and udder health in Martina Franca ass: Effects of daily interval and time of machine milking. *Italian Journal of Animal Science*, 8(SUPPL. 2), 697-699.
- D'Alessandro, A. G., Martemucci, G., Jirillo, E., & Leo, V. D. (2011). Major whey proteins in donkey's milk: Effect of season and lactation stage. Implications for potential dietary

interventions in human diseases. *Immunopharmacology & Immunotoxicology, 33*(2), 259-265.

- D'Auria, E., Mandelli, M., Ballista, P., Di Dio, F., & Giovannini, M. (2011). Growth impairment and nutritional deficiencies in a cow's milk-allergic infant fed by unmodified donkey's milk. *Case Reports in Pediatrics*,2011,1-4.
- Delgado-Pertiñez, M., Alcalde, M. J., Guzmán-Guerrero, J. L., Castel, J. M., Mena, Y., & Caravaca,
 F. (2003). Effect of hygiene-sanitary management on goat milk quality in semi-extensive
 systems in Spain. *Small Ruminant Research*, 47(1), 51-61.
- EN/ISO 11290/1. (1996). Microbiology of food and animal feeding stuffs. Horizontal method for the detection and enumeration of Listeria monocytogenes - Part 1: detection method.
 Geneva: International Organization for Standardization.
- EN/ISO 16649/2. (2001). Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of beta-glucuronidase-positive Escherichia coli - Part 2: Colony-count technique at 44°C using 5-bromo-4-chloro-3-indolyl beta D-glucuronide. Geneva: International Organization for Standardization.
- EN/ISO 16654. (2003). Microbiology of food and animal feeding stuffs. Horizontal method for the detection and enumeration of Escherichia coli O157:H7. Geneva: International Organization for Standardization.
- EN/ISO 7932. (2005). Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of presumptive Bacillus cereus Colony-count technique at 30°C. Geneva: International Organization for Standardization.

FDA. (2001). Bacteriological analytical manual. Arlington, VA: AOAC International.

Giosuè, C., Alabiso, M., Russo, G., Alicata, M. L., & Torrisi, C. (2008). Jennet milk production during the lactation in a Sicilian farming system. *Animal*, 2(10), 1491-1495.

- Guo, H. Y., Pang, K., Zhang, X. Y., Zhao, L., Chen, S. W., Dong, M. L., & Ren, F. Z. (2007). Composition, physiochemical properties, nitrogen fraction distribution, and amino acid profile of donkey milk. *Journal of Dairy Science*, 90(4), 1635-1643.
- Iacono, G., Carroccio, A., Cavataio, F., Montalto, G., Soresi, M., & Balsamo, V. (1992). Use of ass' milk in multiple food allergy. *Journal of Pediatric Gastroenterology & Nutrition*, 14(2), 177-181.
- ISO 6579. (2002). *Microbiology of food and animal feeding stuffs. Horizontal method for detection of Salmonella spp.* Geneva: International Organization for Standardization.
- ISO/FDIS 15214. (1998). Microbiology of food and animal feeding stuffs Horizontal method for the enumeration of mesophilic lactic acid bacteria – Colony count technique at 30°C.
 Geneva: International Organization for Standardization.
- Ivanković, A., Ramljak, J., Štulina, I., Antunac, N., Ivan, B., Kelava, N., & Konjacić, M. (2009). Characteristics of the lactation, chemical composition and milk hygiene quality of the Littoral-Dinaric ass. *Mljekarstvo*, 59(2), 107-113.
- Mansueto, P., Iacono, G., Taormina, G., Seidita, A., D'Alcamo, A., Adragna, F., et al. (2013). Ass's milk in allergy to cow's milk protein: A review. *Acta Medica Mediterranea*, 29(2), 153-160.
- Martemucci, G., & D'Alessandro, A. G. (2012). Fat content, energy value and fatty acid profile of donkey milk during lactation and implications for human nutrition. *Lipids in Health and Disease*, 11, 113.
- Martini, M., Altomonte, I., & Salari, F. (2014). Amiata donkeys: Fat globule characteristics, milk gross composition and fatty acids. *Italian Journal of Animal Science*, *13*(1), 123-126.
- Medhammar, E., Wijesinha-Bettoni, R., Stadlmayr, B., Nilsson, E., Charrondiere, U. R., &
 Burlingame, B. (2012). Composition of milk from minor dairy animals and buffalo breeds:
 A biodiversity perspective. *Journal of the Science of Food and Agriculture*, 92(3), 445-474.

- Monti, G., Bertino, E., Muratore, M. C., Coscia, A., Cresi, F., Silvestro, L., et al. (2007). Efficacy of donkey's milk in treating highly problematic cow's milk allergic children: an in vivo and in vitro study. *Pediatric Allergy & Immunology*, *18*(3), 258-264.
- Monti, G., Viola, S., Baro, C., Cresi, F., Tovo, P. A., Moro, G., et al. (2012). Tolerability of donkey's milk in 92 highly-problematic cow's milk allergic children. *Journal of Biological Regulators & Homeostatic Agents*, 26(3 Suppl), 75-82.
- Park, S. H., Kim, H. J., Kim, J. W., Kim, T. W., & Kim, H. Y. (2007). Simultaneous detection and identification of *Bacillus cereus* group bacteria using multiplex PCR. *Journal of Microbiology and Biotechnology*, 17(7), 1177-1182.
- Pilla, R., Daprà, V., Zecconi, A., & Piccinini, R. (2010). Hygienic and health characteristics of donkey milk during a follow-up study. *Journal of Dairy Research*, 77(4), 392-397.
- Qureshi, A. S., & Enbergs, H. (2012). Studies on the lysozyme activity in the milk of female donkeys (Equus asinus) in relation to reproductive physiology. *The Journal of Animal & Plant Sciences*, 22(Suppl. 2), 70-74.
- Regione Piemonte Direzione Sanità Settore Prevenzione e Veterinaria. *Indicazioni operative per la produzione e commercializzazione di latte crudo di asina*. Available at http://www.regione.piemonte.it/governo/bollettino/abbonati/2013/29/attach/dddb200000461 _830.pdf.
- Reuter, A., Fortunato, D., Garoffo, L. P., Napolitano, L., Scheurer, S., Giuffrida, M. G., et al.
 (2005). Novel isoforms of Pru av 1 with diverging immunoglobulin E binding properties identified by a synergistic combination of molecular biology and proteomics. *Proteomics*, 5(1), 282-289.
- Salazar, O., Asenjo, J. A. (2007) Enzymatic lysis of microbial cells. *Biotechnology Letters*, 29(7), 985-994.
- Salimei, E., & Fantuz, F. (2012). Equid milk for human consumption. *International Dairy Journal*, 24(2), 146-152.

- Salimei, E., Fantuz, F., Coppola, R., Chiofalo, B., Polidori, P., & Varisco, G. (2004). Composition and characteristics of ass's milk. *Animal Research*, *53*(1), 67-78.
- Šarić, L. Ć., Šarić, B. M., Mandić, A. I., Torbica, A. M., Tomić, J. M., Cvetković, D. D., & Okanović, D. G. (2012). Antibacterial properties of Domestic Balkan donkeys' milk. *International Dairy Journal*, 25(2), 142-146.
- Sarno, E., Santoro, A. M. L., Di Palo, R., & Costanzo, N. (2012). Microbiological quality of raw donkey milk from Campania region. *Italian Journal of Animal Science*, 11(3), 266-269.
- Scatassa, M.L., Carrozzo, A., Ducato, B., Giosuè, C., Miraglia, V., & Arcuri L. (2010). Bacillus cereus: isolamento in latte d'asina. Abstract XX Convegno A.I.V.I. Qualità dell'ambiente e sicurezza alimentare, Copanello di Stalettì (CZ), 69.
- Tidona, F., Criscione, A., Devold, T. G., Bordonaro, S., Marletta, D., & Vegarud, G. E. (2014).Protein composition and micelle size of donkey milk with different protein patterns: Effects on digestibility. *International Dairy Journal*, *35*(1), 57-62.
- UNI EN ISO:4833. (2004). Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of microorganisms. Colony-count technique at 30°C. Geneva: International Organization for Standardization.
- Zhang, X. Y., Zhao, L., Jiang, L., Dong, M. L., & Ren, F. Z. (2008). The antimicrobial activity of donkey milk and its microflora changes during storage. *Food Control*, 19(12), 1191-1195.

Figure captions

Figure 1. NuPAGE[®] (12% Bis –Tris; MES buffer) total protein profile of donkey milk, sampled from different farms (A-E) in the North Western Italy, in different sampling seasons (Panel A: Autumn; Panel B: Spring). Numbers: band 1 and band 2 (see Table 3 for identification). Colloidal Coomassie Brilliant Blue

- 458 stained. Std: Mark12[®] Unstained Standard markers (Life Technologies). Gels were run in reducing and
- 459 denaturing conditions.

	Farm A	Farm B	Farm C	Farm D	Farm E
Farm area (ha)	35	12	10	10	42
Altitude above sea level (m)	194	1.110	395	600	183
Total donkeys ^a (no.)	65	63	45	150	48
Jennies ^a (no.)	45	40	40	70	32
Milking jennies ^a (no.)	7-10	7-10	8-10	30-33	6
Herd breed	Crossbreds	Martina Franca	Crossbreds	Crossbreds	Martina Franca,
					Ragusana, crossbreds
Total milk sold (L/year)	1200	2000	2000	0	1000
Milking practice	Automatic in milking	Automatic in milking room	Automatic in cowshed	Hand milking	Automatic in milking
	room				room
Farming type	Semi-extensive	Semi-extensive	xtensive Semi-extensive		Semi-extensive
Feed	Grazing - Hay	Hay - Bread – Protein Grazing - Hay		Grazing - Hay	Grazing - Hay
		supplementation			
Milk use	Food -cosmetics	Food	Food -cosmetics	-	Food -cosmetics

Table 1. Farm characteristics as from face-to-face questionnaire to farm managers

^a counted during the visits

	Season	Dry matter (g/100mL)	Crude proteins (g/100mL)	Total lipids (g/100mL)	Gross energy (kcal/100mL)	Lysozyme activity (U/µL)
Farm A	Autumn	8.77 ± 0.03	1.35 ± 0.04	0.02 ± 0.01	32.7 ± 0	166 ± 21
	Spring	8.69 ± 0.01	1.49 ± 0.02	0.01 ± 0.00	35.1 ± 0	140 ± 29
Farm B	Autumn	8.79 ± 0.02	1.41 ± 0.09	0.02 ± 0.01	35.5 ± 0	180 ± 23
	Spring	9.18 ± 0.03	1.53 ± 0.01	0.06 ± 0.00	37.9 ± 0	153 ± 28
Farm C	Autumn	9.80 ± 0.01	1.68 ± 0.01	0.40 ± 0.03	44.2 ± 0	181 ± 21
	Spring	8.87 ± 0.02	1.55 ± 0.01	0.22 ± 0.01	36.8 ± 0	194 ± 32
Farm D	Autumn	8.75 ± 0.02	1.75 ± 0.08	0.06 ± 0.02	35.7 ± 0	154 ± 34
	Spring	9.05 ± 0.00	1.39 ± 0.00	0.08 ± 0.01	38.0 ± 0	154 ± 13
Farm E	Autumn	8.35 ± 0.03	1.28 ± 0.02	0.02 ± 0.01	33.6 ± 0	155 ± 25
	Spring	8.72 ± 0.00	1.40 ± 0.06	0.02 ± 0.00	34.4 ± 0	140 ± 19
Overall	mean	8.89	1.48	0.09	36.3	162

Table 2. DM chemical composition. Values represent mean values \pm standard deviations (n=3), and the overall means of data collected in the surveyed farms.

	N-term sequence	Identified protein	Accession (NCBI)	Source organism	Theoretical mass (kDa)
Band 1	T-N-I-P-Q-T-M-Q	Beta lactoglobulin 1	P13613.1	Equus asinus	18.53
	K-H-E-I-K-X-V-S	Alpha s2 casein B precursor	CAX65660	Equus asinus	16.72
Band 2	T-N-I-P-Q-T-M-Q	Beta lactoglobulin 1	P13613.1	Equus asinus	18.53

Table 3. N-terminal sequencing and protein identification of bands 1 and 2 from Figure 1.

		Total viable count (TVC)		Lactic acid bacteria count (LAB)		
		Day 0	Day 5	Day 0	Day 5	
Farm A	Mean	5.77 ^a	7.32 ^a	4.34 ^a	5.15 ^a	
	SD	0.45	0.13	0.40	0.10	
	Median	5.85	7.30	4.23	5.11	
Farm B	Mean	4.51 ^{ab}	6.22 ^a	3.61 ^a	4.00 ^{ab}	
	SD	0.66	0.52	0.10	0.31	
	Median	4.18	6.40	3.57	4.00	
Farm C	Mean	5.27 ^a	7.39 ^a	4.01 ^a	4.78 ^{ab}	
	SD	0.42	0.30	0.89	1.22	
	Median	5.11	7.46	3.65	4.20	
Farm D	Mean	2.84 ^b	3.64 ^b	2.69 ^a	2.62 ^{ab}	
	SD	0.96	1.52	1.18	1.15	
	Median	2.72	4.04	1.95	1.95	
Farm E	Mean	4.05 ^{ab}	6.76 ^a	2.34 ^a	2.31 ^b	
	SD	1.12	1.26	0.66	0.62	
	Median	4.00	7.41	1.95	1.95	
Overall	Mean	5.38	7.21	4.20	5.19	
	Median	4.86	7.00	3.57	4.00	

Table 4. DM microbiological analysis results. The numbers represent mean values and standard deviations (n=3) for the three samplings (log cfu/mL).

^{a-b} Different superscript letters in a column indicate significant differences according to Tukey's post-hoc test for TVC and to Dunnet-C test for LAB.

Figure 1

