



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Angiotensin II induces tumor necrosis factor-alpha expressionand release from cultured human podocytes

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/100646

since 2017-06-19T16:27:27Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

The final publication is available at Springer via http://dx.doi.org/10.1007/s00011-011-0412-8

Angiotensin II induces Tumor Necrosis Factor-α expression and release from cultured human podocytes

Arianna Carolina Rosa¹, Lorenza Rattazzi¹, Gianluca Miglio¹, Massimo Collino¹and Roberto Fantozzi¹

¹Department of Anatomy, Pharmacology and Forensic Medicine, University of Turin, Turin, Italy

Running title: Effects of Ang II on TNF- α production by human podocytes

Correspondence: Arianna Carolina Rosa, PhD,

Department of Anatomy, Pharmacology and Forensic Medicine, University of Turin, Corso Raffaello 33, 10125, Turin, Italy Phone: +390116707955 Fax: +390116707688 e-mail: ariannacarolina.rosa@unito.it

ABSTRACT

Objective High level of both Angiotensin (Ang) II and Tumor Necrosis Factor (TNF)- α heve been implicated in the pathogenesis of glomerular injury by affecting podocytes. The aim of this study was to investigate the Ang II-TNF- α relationship in human podocytes.

Methods Immortalized podocytes were exposed to Ang II for 6 days in the absence or presence of either losartan or PD123,319 (both at 100 nM), AT₁ and AT₂ receptor antagonists, respectively.

Results Ang II, after at least 72 h of repeated treatment, increased basal *TNFA* expression and cytokine release with a biphasic pattern and the maximum response at 10 nM. Losartan dampened the Ang II effects on TNF- α production throughout all the experimental period, demonstrating an AT₁ contribution. PD123,319 affected the second TNF- α production peak showing even an AT₂ receptor contribution. Moreover, Ang II causes TNFR1 and TNFR2 over-expression in a time-dependent manner. The functional interaction between Ang II and TNF- α has been demonstrated when the pro-proliferative effect of Ang II was antagonized by a neutralizing TNF- α antibody. *Conclusions* In conclusion, our results point out a functional interaction between Ang II and TNF- α

and indicate this cytokine as a mediator in Ang II long-term pathoadaptive podocytes changes.

Keywords podocytes, Tumor Necrosis Factor- α , angiotensin II

Clinical studies have shown that the angiotensin-converting-enzyme inhibitors (ACEi) and the angiotensin II-receptor blockers (ARB) exert significantly nephroprotective effects by decreasing proteinuria and disease progression both in patients affected by diabetic and non diabetic nephropathies [1, 2]. These data suggest a relevant contribution of the angiotensin (Ang) II signal to the development of glomerular dysfunction. Podocytes are terminally differentiated epithelial cells [3] and constituents of the glomerular filtration barrier (GFB). Their dysfunction and loss have been implicated in the onset and progression of glomerular diseases. These cells express all the components of the renin-angiotensin system (RAS) and are a major source of intraglomerular Ang II [4]. Ang II production results in being enhanced in podocytes following their exposure to mechanical stretch and hyperglycemia [5, 6]. Moreover, the expression of both AT₁ and AT₂ Ang II receptors on podocyte plasma membrane suggests that could be targeted by Ang II [7]. It has been demonstrated in podocytes that Ang II promotes actin rearrangement, tight junction protein ZO-1 and nephrin protein down-regulation, thus leading to proteinuria, as well as an increase of vascular endothelial growth factor (VEGF) and transforming growth factor (TGF)-synthesis and, finally, it triggers apoptosis [8, 9]. Since podocyte response to Ang II may be a complex mixture of both direct and indirect effects, a further elucidation is warranted, particularly by testing the effects of a prolonged cell exposure to the vasopeptide with the aim of mimicking the clinical condition of a chronic exposure to high level of Ang II.

The proinflammatory cytokine tumor necrosis factor (TNF)- α has been reported to participate in the pathogenesis of glomerular injury. In particular, high levels of TNF- α mRNA were found in the glomeruli of rodents and patients with diabetic nephropathy [10, 11], and this overexpression has been associated with the development of renal injury, leading to microalbuminuria [12, 13]. TNF- α is expressed and released not only by infiltrating leukocytes but also by resident cells, including endothelial, mesangial and tubular epithelial cells [14]. Among the many deleterious effects, TNF- α in the kidney promotes an inflammatory response and the accumulation of extracellular matrix thus

reducing the glomerular blood flow and damaging the GFB [15]. Recently, Bruggeman et al. (2011) have demonstrated that TNF- α , through the TNFR2-NF- κ B pathway, contributes to the proliferative and pro-inflammatory phenotype of diseased podocytes [16]. The detrimental role of TNF- α in the pathogenesis of glomerular injury has been confirmed by the attempt to use TNF- α antagonists to treat glomerular diseases [17]. Infliximab, a chimeric monoclonal antibody against TNF- α , has been reported to significantly decrease albuminuria in a rat model of diabetic nephropathy [18].

In vitro studies suggest that a functional linkage between Ang II and TNF- α may exist, having been demonstrated that Ang II induces TNF- α expression in endothelial and mesangial cells [19, 20]. However, at the best of our knowledge, whether Ang II could promote TNF- α production by podocytes remains still an unsolved question. As the high level of both Ang II and TNF- α have been reported in chronic progressive glomerular diseases, where podocytes dysfunction plays a critical role (i.e., hypertensive nephropathy, diabetic nephropathy, classic focal segmental glomerulosclerosis) [15, 21-23], the aim of this study was to investigate the Ang II-TNF- α relationship in human podocytes.

MATERIALS AND METHODS

Drugs and chemicals

Ang II, TNF- α , losartan, PD123,329 and all other reagents were from Sigma-Aldrich (Milan, Italy). Ang II receptor antagonists were dissolved in dimethylsulfoxide, TNF- α in water, as indicated by manufacturers' instructions. Final drug concentrations were obtained by dilution of stock solutions in the experimental buffers. The final concentration of organic solvent was less than 0.1%, which has no effect on cell viability. In all experiments un-stimulated cells served as controls.

Cell cultures

Primary cultures of human podocytes were kindly provided by Prof. G. Camussi (Department of Internal Medicine and Center for Molecular Biotechnology, University of Turin, Turin, Italy). Podocytes were characterized and cultured as previously reported [24, 25].

Citokine release

TNF- α release was evaluated by measuring cytokine concentration in the experimental medium by the Instant ELISA system (Bender MedSystems, GmbH, W., Austria) following the manufacturers' instructions. All experiments were performed with 30×10^3 cells/well as initial cell density and results were corrected for the number of cells (cell growth) and are expressed as percent change over the control.

RT-PCR and quantitative real-time PCR analysis

The analyses were performed as previously described [25, 26]. Briefly, RT-PCR amplifications were performed in 25 μ l reaction mixtures containing 1.0 μ l of cDNA, 2.5 μ l of 10× buffer, 1.0 μ l of 50 mM MgCl₂, 0.20 μ l of 25 mM dNTPs mix, 0.05 U of Euro*Taq* DNA polymerase (Euroclone,

Milan, Italy), and 2.5 nM of each primer. PCR amplicons were resolved in an ethidium bromidestained agarose gel (2%-3%) by electrophoresis, and signals were quantified using NIH Image J 1.41 software. β -*actin* expression was used as an internal control (Table 1). Real-time PCR experiments were performed in 25 µl reaction mixtures containing 10 ng of cDNA template, the Power SYBR® Green PCR Master Mix and the AmpliTaq Gold® DNA Polymerase LD (Applied Biosystems). Relative quantization of the products was performed using a 48-well StepOneTM Real Time System (Applied Biosystems). For all real-time PCR analyses, β -actin mRNA was used to normalize RNA inputs (Table 1).

Western blot analysis

Cell culture dishes were washed with phosphate-buffered saline (PBS) before adding ice-cold lysis buffer (20 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% SDS, 1% Triton X-100, 1% sodium deoxycholate, 5 mM EDTA, 1 µl/ml protease inhibitors, 0.1 mM ZnCl₂, and 1 mM Ph-Me-Sul-Flu). Cell lysates were processed to determine protein concentrations using a BCA protein assay (Pierce Biotechnology Inc. Rockford, IL, USA) following the manufacturers' instruction. Samples containing 20 µg of protein were subjected to SDS-PAGE using a 10% gel. Proteins were transferred to a PVDF membrane (Millipore, Bradford, USA), which was incubated with Super Block blocking buffer (Pierce Biotechnology Inc.). AT₁R, TNFR1 and TNFR2 were detected following incubation with mouse monoclonal antibodies (Santa Cruz Biotechnology, CA, USA). ATR2 was detected following incubation with rabbit polyclonal antibody (Santa Cruz Biotechnology), according to the manufacturer's instructions. To confirm the homogeneity of the proteins loaded, the membranes were stripped and incubated with an anti-β-actin monoclonal antibody (Sigma-Aldrich). The membranes were overlaid with Western Lightning Chemiluminescence Reagent Plus (PerkinElmer Life Science, Norwalk, CT, USA) and then exposed to Hyperfilm ECL film (Amersham Biosciences, Piscataway, NJ, USA).

Cell proliferation assay

Podocytes (3000 cells/well) were exposed to Ang II 10 nM for 6 days according to an everyday medium changing protocol. In some experiments, at the day 3, a polyclonal goat anti-human IgG blocking TNF- α or a nonspecific polyclonal goat IgG (Santa Cruz Biotechnology) were added. The proliferative response was determined by counting the viable cells in a hemacytometer by the trypan blue exclusion test, blind by an observer.

Data analysis

Results are expressed as means \pm SEM of at least three experiments. One-way analysis of variance (ANOVA) followed by the post hoc Dunnet's multiple comparison and Student –Newman-Keuls tests were used to determine significant differences between means. Differences were considered statistically significant when P < 0.05.

RESULTS

Effects of Ang II on TNF-a production by human podocytes

Our cell line posses both AT₁ and AT₂ receptors, as measured at both mRNA and protein level (Figure 1A and 1B). Their activation by Ang II stimulates TNF- α production as shown in Figure 2A and 2B. Cells were treated with Ang II (10 nM, 1-6 days) according to a daily exposure protocol and both *TNFA* expression and TNF- α release were measured (see materials and methods). The cytokine release showed a biphasic pattern with a first peak at the forth day and a second one at the sixth: medium TNF- α concentration was 14.9±3.0 pg/ml at day 0, 37.9±3.7 pg/ml at the forth day (about +150%; *P*<0.01) and 49.2±5.2 pg/ml at the sixth day (about +230%; *P*<0.01 *vs* both day 0 and forth day). The same pattern was displayed by *TNFA* but with a ~24 h time-lag (Figure 2A). Ang II concentration-dependent curve (0.03-100 nM) disclosed a sigmoid shape with the maximum effect achieved from 10 nM and the EC₅₀ at 0.43±0.02 nM (Figure 2B). These results demonstrate

that Ang II modulates TNF- α release by human podocytes in a time- and concentration-dependent manner. These effects are regulated at the gene level.

Effects of losartan and PD123,319 on cytokine production induced by Ang II

To evaluate whether AT_1 and AT_2 receptors contribute differently to Ang II response, podocytes were treated with the vasopeptide (10 nM, 1-6 days) in the presence or absence of selective AT_1 or AT_2 receptor antagonists, losartan (AT_1) or PD123,319 (AT_2), both at 100 nM (Figure 3). Only losartan was able to abolish Ang II effect on TNF- α release throughout all the experimental period. On the contrary, PD123,319 did not affect the first peak, but antagonized the second one (Figure 3). From these data it appears that Ang II response is differently and selectively mediated by both Ang II receptors. The same results were recorded by testing gene expression (data not shown).

Effects of Ang II on TNF-a receptor expression

A further link between Ang II and TNF- α in human podocytes was discovered when the vasopeptide effect on TNF- α receptor expression was investigated at both mRNA and protein level. As previously, cells were exposed to 10 nM Ang II up to 6 days (Figure 4A and 4B). Ang II upregulated TNF- α receptors expression in a time-dependent manner starting respectively from the first day for TNFR2 and the third one for TNFR1.

Effects of Ang II on human podocytes proliferation

To evaluate whether the stimulatory agonism of Ang II on TNF- α production by human podocytes may have functional effects, the proliferative podocyte response to Ang II was tested. In comparison to medium alone, Ang II daily exposure increased cell growth rate: the cell number was significantly from the fourth day (Figure 5A). This effect was blunted by an α TNF- α neutralizing antibody (at the third day; Figure 5B), thus suggesting that the Ang II induced proliferative response is mediated by TNF- α .

DISCUSSION

In this study we reported, in a concentration- and time-dependent manner, that Ang II induces TNF- α production by human podocytes. These results suggest that Ang II could activate a paracrine/autocrine loop centred on TNF- α . A similar positive effect of Ang II has been described in other kidney cells, such as cells from isolated tubules of rat medullary thick ascending limb and mesangial cells [19, 20], extending the evidence to the human podocytes. In our experiments Ang II, after at least 72 h of repeated treatment, increased basal TNFA expression and cytokine release with a biphasic pattern. Ang II-induced TNF- α release required a repeated exposure to the vasopeptide and it is related to the gene transcription. The analysis of the two kinetic profiles highlights a ~24 h-delay of release in comparison to gene expression, likely due to the multi-step process leading to mature TNF- α secretion [27]. Ang II up-regulates TNF- α in a concentrationdependent manner (EC₅₀s 0.43 ± 0.02 nM) with the maximum response at 10 nM. Notably, these concentrations are within the intrarenal range reported to be ~one thousand times higher than the circulating one [28, 29]. Furthermore, our protocol, consisting of repeated stimulations, intended to mimic a chronic in vivo podocyte exposure to the vasopeptide. The everyday replacement of the medium with or without Ang II was planned to overcome Ang II catabolism [30]. Velez et al. (2007) have demonstrated that podocytes not only synthesize and release Ang II, but also convert it into the bioactive Ang (1-7) and Ang III [31]. The formation of these active metabolites has to be considered while discussing the data. However, as losartan, a selective AT₁ antagonist, abolished the Ang II effects, the contribution of Ang (1-7), acting on the mas receptor [32], can be excluded. Even Ang III contribution can be rejected, because, even if it activates AT₁ and AT₂ receptors with an affinity quite close to Ang II [32], its half life is ~5 fold shorter than the one of the parent peptide [31]. Thus, we may conclude that the observed effects are mainly due to Ang II. Taken these data together, our experimental conditions might be relevant to elucidate at least one of the podocytes pathoadaptive changes, TNF- α up-regulation, caused by a prolonged exposure to high concentration of Ang II, in the same way as it happens in chronic progressive glomerular disease where podocyte dysfunction plays a critical role (i.e., hypertensive nephropathy, diabetic nepfropathy, classic focal segmental glomerulosclerosis) [15, 21-23].

Previous findings indicate that in rat renal cortex [22, 33] and in human renal proximal tubular epithelial cells [34] the interaction between Ang II and TNF- α is mediated by AT₁ receptors. These evidences have been confirmed by using a selective AT₁ antagonist, such as losartan. Drug treatment dampened all the Ang II effects on TNF- α gene expression and release, throughout the entire experimental period. However, our results show that even AT₂ receptors contribute to evoke the effects measured in the last period of the experimental observation, thus indicating a delayed involvement of theirs. This suggests that while the first peak reflects the AT₁ receptor contribution on cytokine production, the second one seems to be driven by a more complex signaling.

Another original result of this study is that Ang II, besides evoking TNF- α production, causes TNFR1 and TNFR2 over-expression, which might underlie an increased cell responsiveness to the cytokine. Actually, a functional interaction between Ang II and TNF- α has been demonstrated in our study when the pro-proliferative effect of Ang II was antagonized by a neutralizing TNF- α antibody. This experiment has been planned only to ascertain the cytokine role, being not able to elucidate the relative contribution of the two TNF- α receptor types. However, the kinetic data depicted in figure 4 indicate a more rapid and more intense modulation by Ang II of the TNFR2 expression in comparison to that of TNFR1. Interestingly, while TNFR1 has been implicated in cell death processes [14, 27], recent data indicate the TNF- α -TNFR2 axis as a candidate pathway contributing to the development of a proliferative and pro-inflammatory podocyte phenotype,

associated with proliferative podocytopathies [16]. Ore data are derived from cultured immortalized human podocytes that are proliferating cells. Therefore these observations are not in keeping with that podocytes are postmitotic terminally differentiated cells with little or no capacity for regenerative replication [35]. However, proliferating podocytes have been observed in experimental models of glomerular injury [36], which may be explained admitting that some podocytes reengage the cell cycle as an adaptive response to injury in an attempt to mitigate their loss [37]. Further, indirect evidence for podocyte proliferation in human diabetes stems from the increased numbers of podocytes that are detected in the urine long before any reduction in glomerular podocyte numbers [38, 39]. In our study, we have demonstrated that our model leads to an Ang II induced proliferation of human podocytes. A critical aspect of our model is the use of immortalized podocytes, that are a cell type which is able to proliferate by itself. While recognizing this experimental bias, we have to stress that an increase in proliferation rate was observed in the presence of Ang II, as well as its reduction after α TNF- α . Our results, consistently with previous data indicate that TNF- α could act as a mitogenic stimulus on podocytes [16] and suggest that the TNF- α release and the TNFRs expression, could mediate the proliferative effects induced by Ang II on podocytes.

In conclusion, our results point out a functional interaction between Ang II and TNF- α and indicate this cytokine as a mediator in Ang II long-term pathoadaptive podocytes changes. These results get more insight into the pathophysiological Ang II role in podocytes dysfunction leading to glomerular disease.

ACKNOWLEDGMENTS

This work was supported by a grant from the Italian Ministry of Education, University, and Research (*Programmi di ricerca scientifica di rilevanza nazionale*, PRIN 2007), the Regione Piemonte (2008 and 2009), and the University of Turin. We thank Prof. G. Camussi and Dr. C. Grange of the Department of Internal Medicine and Center for Molecular Biotechnology, University of Turin (Italy) for providing human immortalized podocytes.

Part of this study was presented at the 16th World Congress on Basic and Clinical Pharmacology, July 17-23, 2010, Cophenaghen, Denmark; and was published in abstract form on *Basic Clin Pharmacol Toxicol*, 2010, 107 (Suppl. 1), 548.

The specific contributions to the work is listed as follow: performed the research A.C.R., L.R. and G.M.; designed the research study A.C.R. and R.F.; analyzed the data A.C.R., L.R., M.C. and G.M.; wrote the paper A.C.R., G.M. and R.F.

CONFLICT OF INTEREST

None.

TITTLES AND LEGENDS

Figure 1. Basal AT_1 and AT_2 receptor expression by human podocytes. Representative pictures of *AGTR1* and *AGTR2* mRNAs detected by RT-PCR (A) or AT_1 and AT_2 proteins by Western blot analysis (B).

Figure 2. Effects of Ang II on TNF- α production by human podocytes. (A) Podocytes were exposed to Ang II. Experimental medium and Ang II were replaced every day. Cells were processed for quantitative real-time PCR analysis and cytokine concentration in the supernatants were measured by ELISA. Data are expressed as percent change over the control (untreated cells; day 0). (B) Podocytes were exposed to increasing Ang II concentrations (0.03 – 100 nM) for 4 days and TNF- α release was measured by ELISA. Data are expressed as percentage of the maximal effect. Data are the mean ± SEM of three independent experiments performed in duplicate. **P*<0.01 *vs*. control and #*P*<0.01 *vs*. first peak.

Figure 3. Effects of losartan and PD123,319 on TNF- α production induced by Ang II. Podocytes were exposed to Ang II (10 nM, 1-6 days) in presence of losartan (100 nM) or PD123,319 (100 nM). Experimental medium, Ang II and antagonists were replaced every day. Cytokine concentration in the supernatants was measured by ELISA. Results are expressed as percent change over the control (untreated cells; day 0). Data are the mean \pm SEM of three independent experiments performed in duplicate. **P*<0.01 *vs*. control.

Figure 4. Effects of Ang II on TNFR1 and TNFR2 receptor expression. Podocytes were exposed to Ang II. Experimental medium and Ang II were replaced every day and cells were processed for quantitative real-time PCR analysis (A) or Western blot analysis, pictures shown are representative

of three independent experiments (B). Results are expressed as percent change over the control (untreated cells; day 0). Data are the mean \pm SEM of three independent experiments performed in duplicate. **P*<0.01 *vs*. control.

Figure 5. Effects of Ang II on proliferative response of human podocyte. (A) Cells were cultured in the absence or presence of Ang II (10 nM, 1-6 days) Experimental medium and Ang II were replaced every day. (B) Cells were cultured for 4 days, on the day 3 α TNF- α or nonspecific polyclonal goat IgG were added. Results are expressed as cells/well. Data are the mean ± SEM of three independent experiments performed in triplicate. *P<0.01 vs. control and #P<0.01 vs. α TNF- α .

REFERENCES

- 1. Benigni A, Cassis P, Remuzzi G. Angiotensin II revisited: new roles in inflammation, immunology and aging. EMBO Mol Med; 2:247-57.
- 2. Ruggenenti P, Cravedi P, Remuzzi G. The RAAS in the pathogenesis and treatment of diabetic nephropathy. Nat Rev Nephrol; 6:319-30.
- 3. Shankland SJ. Cell cycle regulatory proteins in glomerular disease. Kidney Int 1999; 56:1208-15.
- 4. Liebau MC, Lang D, Bohm J, Endlich N, Bek MJ, Witherden I, et al. Functional expression of the renin-angiotensin system in human podocytes. Am J Physiol Renal Physiol 2006; 290:F710-9.
- 5. Shankland SJ. The podocyte's response to injury: role in proteinuria and glomerulosclerosis. Kidney Int 2006; 69:2131-47.
- 6. Miceli I, Burt D, Tarabra E, Camussi G, Perin PC, Gruden G. Stretch reduces nephrin expression via an angiotensin II-AT(1)-dependent mechanism in human podocytes: effect of rosiglitazone. Am J Physiol Renal Physiol; 298:F381-90.
- 7. Wang L, Flannery PJ, Spurney RF. Characterization of angiotensin II-receptor subtypes in podocytes. J Lab Clin Med 2003; 142:313-21.
- 8. Durvasula RV, Petermann AT, Hiromura K, Blonski M, Pippin J, Mundel P, et al. Activation of a local tissue angiotensin system in podocytes by mechanical strain. Kidney Int 2004; 65:30-9.
- 9. Ding G, Reddy K, Kapasi AA, Franki N, Gibbons N, Kasinath BS, et al. Angiotensin II induces apoptosis in rat glomerular epithelial cells. Am J Physiol Renal Physiol 2002; 283:F173-80.
- 10. Nakamura T, Fukui M, Ebihara I, Osada S, Nagaoka I, Tomino Y, et al. mRNA expression of growth factors in glomeruli from diabetic rats. Diabetes 1993; 42:450-6.
- 11. Mahmoud RA, el-Ezz SA, Hegazy AS. Increased serum levels of interleukin-18 in patients with diabetic nephropathy. Ital J Biochem 2004; 53:73-81.
- 12. Kalantarinia K, Awad AS, Siragy HM. Urinary and renal interstitial concentrations of TNFalpha increase prior to the rise in albuminuria in diabetic rats. Kidney Int 2003; 64:1208-13.
- 13. Navarro JF, Mora C, Maca M, Garca J. Inflammatory parameters are independently associated with urinary albumin in type 2 diabetes mellitus. Am J Kidney Dis 2003; 42:53-61.
- 14. Vielhauer V, Mayadas TN. Functions of TNF and its receptors in renal disease: distinct roles in inflammatory tissue injury and immune regulation. Semin Nephrol 2007; 27:286-308.
- 15. Sanchez-Nino MD, Benito-Martin A, Goncalves S, Sanz AB, Ucero AC, Izquierdo MC, et al. TNF superfamily: a growing saga of kidney injury modulators. Mediators Inflamm; 2010.
- 16. Bruggeman LA, Drawz PE, Kahoud N, Lin K, Barisoni L, Nelson PJ. TNFR2 interposes the proliferative and NF-kappaB-mediated inflammatory response by podocytes to TNF-alpha. Lab Invest; 91:413-25.
- 17. Marasa M, Kopp JB. Monoclonal antibodies for podocytopathies: rationale and clinical responses. Nat Rev Nephrol 2009; 5:337-48.
- 18. Moriwaki Y, Inokuchi T, Yamamoto A, Ka T, Tsutsumi Z, Takahashi S, et al. Effect of TNF-alpha inhibition on urinary albumin excretion in experimental diabetic rats. Acta Diabetol 2007; 44:215-8.
- 19. Ferreri NR, Escalante BA, Zhao Y, An SJ, McGiff JC. Angiotensin II induces TNF production by the thick ascending limb: functional implications. Am J Physiol 1998; 274:F148-55.
- 20. Nakamura A, Johns EJ, Imaizumi A, Niimi R, Yanagawa Y, Kohsaka T. Role of angiotensin II-induced cAMP in mesangial TNF-alpha production. Cytokine 2002; 19:47-51.

- 21. Crowley SD, Frey CW, Gould SK, Griffiths R, Ruiz P, Burchette JL, et al. Stimulation of lymphocyte responses by angiotensin II promotes kidney injury in hypertension. Am J Physiol Renal Physiol 2008; 295:F515-24.
- 22. Siragy HM, Awad A, Abadir P, Webb R. The angiotensin II type 1 receptor mediates renal interstitial content of tumor necrosis factor-alpha in diabetic rats. Endocrinology 2003; 144:2229-33.
- 23. Ma L, Fogo AB. Role of angiotensin II in glomerular injury. Semin Nephrol 2001; 21:544-53.
- 24. Doublier S, Ruotsalainen V, Salvidio G, Lupia E, Biancone L, Conaldi PG, et al. Nephrin redistribution on podocytes is a potential mechanism for proteinuria in patients with primary acquired nephrotic syndrome. Am J Pathol 2001; 158:1723-31.
- 25. Miglio G, Rosa AC, Rattazzi L, Grange C, Collino M, Camussi G, et al. The subtypes of peroxisome proliferator-activated receptors expressed by human podocytes and their role in decreasing podocyte injury. Br J Pharmacol; 162:111-25.
- 26. Miglio G, Rosa AC, Rattazzi L, Collino M, Lombardi G, Fantozzi R. PPARgamma stimulation promotes mitochondrial biogenesis and prevents glucose deprivation-induced neuronal cell loss. Neurochem Int 2009; 55:496-504.
- 27. Bemelmans MH, van Tits LJ, Buurman WA. Tumor necrosis factor: function, release and clearance. Crit Rev Immunol 1996; 16:1-11.
- 28. Kobori H, Nangaku M, Navar LG, Nishiyama A. The intrarenal renin-angiotensin system: from physiology to the pathobiology of hypertension and kidney disease. Pharmacol Rev 2007; 59:251-87.
- 29. Siragy HM, Howell NL, Ragsdale NV, Carey RM. Renal interstitial fluid angiotensin. Modulation by anesthesia, epinephrine, sodium depletion, and renin inhibition. Hypertension 1995; 25:1021-4.
- 30. Schling P. Expression of angiotensin II receptors type 1 and type 2 in human preadipose cells during differentiation. Horm Metab Res 2002; 34:709-15.
- 31. Velez JC, Bland AM, Arthur JM, Raymond JR, Janech MG. Characterization of reninangiotensin system enzyme activities in cultured mouse podocytes. Am J Physiol Renal Physiol 2007; 293:F398-407.
- Fyhrquist F, Saijonmaa O. Renin-angiotensin system revisited. J Intern Med 2008; 264:224-36.
- 33. Rompe F, Artuc M, Hallberg A, Alterman M, Stroder K, Thone-Reineke C, et al. Direct angiotensin II type 2 receptor stimulation acts anti-inflammatory through epoxyeicosatrienoic acid and inhibition of nuclear factor kappaB. Hypertension; 55:924-31.
- 34. Kagawa T, Takao T, Horino T, Matsumoto R, Inoue K, Morita T, et al. Angiotensin II receptor blocker inhibits tumour necrosis factor-alpha-induced cell damage in human renal proximal tubular epithelial cells. Nephrology (Carlton) 2008; 13:309-15.
- 35. Pavenstadt H, Kriz W, Kretzler M. Cell biology of the glomerular podocyte. Physiol Rev 2003; 83:253-307.
- 36. Macconi D, Sangalli F, Bonomelli M, Conti S, Condorelli L, Gagliardini E, et al. Podocyte repopulation contributes to regression of glomerular injury induced by ACE inhibition. Am J Pathol 2009; 174:797-807.
- 37. Herman-Edelstein M, Thomas MC, Thallas-Bonke V, Saleem M, Cooper ME, Kantharidis P. Dedifferentiation of immortalized human podocytes in response to transforming growth factor-beta: a model for diabetic podocytopathy. Diabetes; 60:1779-88.
- 38. Yamaguchi Y, Iwano M, Suzuki D, Nakatani K, Kimura K, Harada K, et al. Epithelialmesenchymal transition as a potential explanation for podocyte depletion in diabetic nephropathy. Am J Kidney Dis 2009; 54:653-64.

39. Nakamura T, Ushiyama C, Suzuki S, Hara M, Shimada N, Ebihara I, et al. The urinary podocyte as a marker for the differential diagnosis of idiopathic focal glomerulosclerosis and minimal-change nephrotic syndrome. Am J Nephrol 2000; 20:175-9.











Table 1 Supplemental material

PCR primers		Amplicon	Denaturation	Anneali
		size(bp)		
TNFA	F:5'-CAATGGCGTGGAGCTGAGAGA-3'	344	95°C, 60 s	62°C, 6
	R:5'-CCAAAGTAGACCTGCCCAGAC-3'			
Tnfr1	F:5'-TCTATGCCCGAGTCTCAACC-3'	244	94°C, 30 s	55°C, 3
	R:5'-GGTGAGGGACCAGTCCAATA-3'			
Tnfr2	F:5'-AACTGGGTTCCCGAGTGCTTG-3'	629	94°C, 30 s	64°C, 3
	R:5'-AGTGCTGGGTTCTGGAGTTGG-3'			
AGTR1	F:5'-AGCCAAATCCCACTCAAACCT-3'	120	94°C, 30 s	55°C, 3
	R:5'-TCGAACATGTCACTCAACCTCA-3'			
AGTR2	F:5'-AAGAAGAAATCCCTGGCAAGC-3'	302	94°C, 30 s	55°C, 3
	R:5'-CCACGGCCTTGCTCTTGTT-3'			
β -actin	F:5'-TGACGGGGTCACCCACACTGTGCCCATCTA-3'	660	95°C, 45 s	60°C, 4
	R:5'-CTAGAAGCATTTGCGGTGGACGATGGAGGG-3'			