

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Peri-Adriatic mélanges and their evolution in the Tethyan realm

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/101216> since

*Published version:*

DOI:10.1080/00206810902949886

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITA' DEGLI STUDI DI TORINO

**This is an author version of the contribution published on:**

Questa è la versione dell'autore dell'opera:

*Festa et al. (2010) - International Geology Review, v.52 (4-6), 369-406,*  
doi:10.1080/00206810903557704

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://www.tandfonline.com/loi/tigr20>

# **Peri-Adriatic mélanges, and their evolution in the Tethyan realm**

<sup>1-3</sup>Andrea Festa, <sup>2</sup>Gian Andrea Pini, <sup>3</sup>Yildirim Dilek, <sup>1</sup>Giulia Codegone, <sup>1</sup>Livio Vezzani,  
<sup>4</sup>Francesca Ghisetti, <sup>5</sup>Claudio Corrado Lucente & <sup>6</sup>Kei Ogata

1 Dipartimento di Scienze della Terra – Università di Torino, Via Valperga Caluso 35, 10125 Torino, Italy

2 Dipartimento di Scienze della Terra e Geologico-Ambientali, Università di Bologna, Via Zamboni 67, 40127  
Bologna, Italy

3 Department of Geology, Miami University, Oxford, OH 45056, USA

4 TerraGeoLogica, 55 Mansfield Avenue, Christchurch 8014, New Zealand

5 Servizio Tecnico Bacini Conca e Marecchia, Regione Emilia-Romagna, Via Rosaspina 7, 47900 Rimini

6 Dipartimento di Scienze della Terra – Università di Parma, V.le G.P. Usberti, 157/A - Campus Universitario,  
43100 Parma, Italy

Corresponding author:

Dr. A. Festa

[andrea.festa@unito.it](mailto:andrea.festa@unito.it)

Submitted to:

International Geology Review

Special Issue: Eastern Mediterranean Geodynamics

Revised

Ms.

## Abstract

In the peri-Adriatic region, mélanges represent a significant component of the Apennine and Dinaride-Albanide-Hellenide orogenic belts as well as ancient and present day accretionary wedges. Different mélange types in this broad region provide an excellent case study to investigate the mode and nature of main processes (tectonic, sedimentary and diapiric) involved in mélange formation in contrasting geodynamic settings. We present a preliminary subdivision and classification of the peri-Adriatic mélanges based on several years of field studies on chaotic rock bodies, including detailed structural and stratigraphic analyses. Six main categories of mélanges are distinguished on the basis of the processes and geodynamic settings of their formation. These mélange types are spatially and temporally associated with: 1) extensional tectonics, 2) passive margin evolution, 3) strike-slip tectonics, cases, there appears to have been a strong interplay and some overlap between tectonic, sedimentary, and diapiric processes during mélange formation; however, in highly deformed regions it is still possible to distinguish those mélanges that formed in different geodynamic environments and their main processes of formation. This study shows that a strong relationship exists between mélange-forming processes and the paleogeographic settings and conditions of mélange formation. Given the differences in age, geographic location and evolutionary patterns, we document the relative importance of mélanges and broken formations in the tectonic evolution of the peri-Adriatic mountain belts.

Key words: Mélange, olistostromes, broken formation, tectonosomes, mud diapirs, subduction processes, subduction channels, obduction, allochthonous nappes, Apennines, Dinarides-Albanides-Hellenides.

## Introduction

In the peri-Adriatic region (Fig. 1), mélanges are common as part of the Apennines and Dinaride-Albanide-Hellenide orogenic belts and accretionary wedges. Their formation and incorporation into these orogenic systems played a significant role in the Mesozoic-Cenozoic tectonic evolution of the central Mediterranean region.

The term “mélange” was first coined by Greenly (1919) to describe a unit of highly disrupted rocks in North Wales (Anglesey Island). After its reintroduction by Hsü (1968), the term mélange has been applied worldwide to indicate chaotic, block-in-matrix rocks. In particular, the term has been used to indicate rock units associated with the subduction of oceanic crust in the circum-Pacific areas, such as the west coast of US, New Zealand, and Japan. Since its application to the circum-Mediterranean region, the term has been applied to different kinds of chaotic rock types containing a wide range of characteristics (see Camerlenghi and Pini, 2009 and references therein), that result from protracted cycles of rifting, drifting, oceanic subduction, continental collision and intracontinental deformation, different from the almost exclusively subduction-related Pacific mélanges. Moreover, the term mélange frequently flanks or substitutes for more traditional terms, such as wildflysch, schistes à blocs, agglomerates, argille scagliose, and olistostromes (see Camerlenghi and Pini, 2009 and references therein). The large majority of these indicate chaotic, mostly block-in-matrix rocks, related to a sedimentary origin, a complex sedimentary-tectonic origin, and/or to mud diapirism (Hsü, 1968, 1973; Abbate et al., 1970; Elter and Trevisan, 1973; Naylor, 1981; Castellarin et al., 1998; Pini, 1999; Festa et al., 2005; Trümpy, 2006; Dela Pierre et al., 2007; Camerlenghi and Pini, 2009).



Therefore, in the circum-Mediterranean mountain chains, the term mélanges has been frequently applied to rock units of sedimentary origins (see below), whereas in the Pacific realm the term has been always considered as related to a tectonic origin (i.e., Meschede et al., 1999; Chang et al., 2000; Wakabayashi, 2004; Ikesawa et al., 2005) with notable exceptions (Williams et al., 1984; Cowan, 1985; Barber et al., 1986; Brandon, 1989; Horton and Rast, 1989; Brown, 1990; Cousineau, 1998; Yamamoto et al., 2000). Notably, some of these, non-Mediterranean exceptions used Mediterranean mélanges as paradigmatic examples (see, e.g. Hsü, 1965; 1974; Cowan, 1985).

In this paper, we describe the mélange occurrence in the Apennine and Dinaride-Albanide-Hellenide mountain belts, both of which are part of the broad Alpine orogenic system. Given the differences in age, geographic location, and evolutionary patterns, we propose a new categorization of these mélanges, taking into account their geodynamic environments and processes of formation. We also document their relative importance in the evolution of these mountain belts and in the Tethyan system.

## Contribution of Apenninic geology to the mélange concept

Bianconi (1840) described the mesoscopic scaly texture of block-in-matrix rocks of the Ligurian nappe in the northern Apennines as “argille scagliose” (scaly clays). This term was then progressively extended to all the tectonic, tectono-stratigraphic and sedimentary chaotic rock bodies of the Apennines, also named *mélanges (s.l.)* after Greenly (1919), that are characterized by a block-in-matrix arrangement at the outcrop scale (Signorini, 1946; Azzaroli, 1948; Merla, 1952; Ogniben, 1953; Maxwell, 1959; Elter, 1960). Different chaotic units, including both Jurassic (Ligurian units) ophiolites and Mesozoic-Cenozoic sedimentary successions originally deposited on oceanic and/or continental crust of the Adria microplate (Subligurian, Sicilide, Tuscan, Marche-Umbria, Molise-Sannio, Lagonegro units), were recognized and described in different stratigraphic and structural context of the Apenninic thrust-and-fold belt.

The argille scagliose were considered to be the products of sedimentary processes deformed in response to tectonic events (orogenic landslides; Merla, 1952; Signorini, 1956). Another descriptive term, “argille brecciate” (brecciated clays), was proposed by Ogniben (1953) and Rigo de Righi (1956) to describe the block-in-matrix fabric at the millimeter-to centimeter scale of chaotic sediments in southern Apennines and Sicily. In contrast to argille scagliose, argille brecciate has been interpreted as the product of mass-wasting processes (debris flows and debris avalanches, see Camerlenghi and Pini; 2009 for major details).

Argille scagliose, argille brecciate and chaotic rock bodies with block-in-matrix fabric, recognized as sedimentary bodies within layered coherent sedimentary marine successions, were defined in Sicily as “olistostromes” by Flores (1955; 1956). This term, from the Greek “ (to slide) and “ (bed), was used to indicate different products of mass-gravitational transport (i.e., Beneo, 1956; Rigo de Righi, 1956; Abbate et al., 1970; 1981; Elter and Trevisan, 1973; Ricci Lucchi, 1975) including block slides (olistoliths), avalanches, debris flows and iper-concentrated flows (Camerlenghi and Pini, 2009). These differ from the argille scagliose on the basis of their brecciated matrix consisting of rounded to angular polymictic millimeter-to centimeter-scale clasts of claystone, random distributed throughout the clayey matrix. The scaly fabric is often present and may overprint but not completely obliterate the brecciated texture (Bettelli and Panini, 1985).

The chaotic appearance of the argille scagliose, argille brecciate and olistostromes in outcrops led to the proliferation of different terms such as Ligurian *mélange*, chaotic complex, undifferentiated complex, *complesso caotico eterogeneo* (heterogeneous chaotic complex) or *caotico eterogeneo* (heterogeneous chaotic) to describe the same chaotic successions in different sectors of the Apennines chain (see, e.g., Abbate and Sagri, 1970). Some of these terms have been used synonymously with the term olistostrome in regional-scale maps (e.g. Boni et al., 1970, Bellinzona et al., 1971; Boccaletti and Coli, 1982). As a consequence, a conceptual linkage between olistostromes, argille scagliose and other terms that describe rocks with a block-in-matrix fabric has always been improperly maintained or never been excluded (see discussion in Camerlenghi and Pini, 2009 for major details). While the application of these terms in the Apennines was changing, many of these concepts were exported and favorably applied to disrupted rock bodies of different ages exposed in different geodynamic contexts and stratigraphic-structural positions elsewhere (see for example, Hsü, 1965; Raymond, 1984; Cowan, 1985; Orange and Underwood, 1995).

Starting in the 1980s, chaotic rock bodies and disrupted rocks of the Apennines have been distinguished (Bettelli and Panini, 1985, 1987; Castellarin et al., 1986; Castellarin and Pini, 1987; Pini, 1987; Camerlenghi and Pini, 2009 and references therein) into: a) strongly deformed units (tectonosomes or broken formations) displaying a prevailing block-in-matrix fabric, in which part of the same coherent stratigraphic unit can be recognized and mapped; b) sedimentary block-in-matrix rocks (or olistostromes or sedimentary *mélanges*) related to different mass-wasting sedimentary processes (debris flows and avalanches, sliding-gliding of blocks), with the possible contribution of mud volcanoes and diapirs. These sedimentary bodies have a classic block-in-brecciated-matrix fabric, with the matrix made up of clays supporting millimeter-scale clasts (brecciated matrix, see Swarbrick and Naylor, 1980; Abbate et al., 1981 and references in Camerlenghi and Pini, 2009).

### **Regional setting of the peri-Adriatic region**

During late Mesozoic-Cenozoic, the Mediterranean region (Fig. 1) experienced complex subduction and associated collisional events, resulting in the development of different fold-and-thrust belts (Dilek, 2006). The Apennine and Dinaride-Albanide-Hellenide mountain belts are the product of this subduction-collision history and presently are separated by the Apulia-Adriatic foreland.

The Apennines consist of an east-to northeast vergent fold-and-thrust belt developed after the Late Cretaceous – Early Cenozoic closure of the Ligurian ocean (Alpine Tethys) (Fig. 2a) and the convergence between

the continental margin of the European plate (Corsica-Sardinia), to the west, and of the Adria microplate (of African affinity), to the east (i.e., Boccaletti et al., 1980; Dewey et al., 1989; Castellarin, 1994;).

During pre-collisional eo- and meso-alpine episodes of convergence (Vai and Castellarin, 1993; Pini, 1999; Vescovi et al., 1999) the Ligurian units, which represent the most internal paleogeographic domain (Fig. 2) adjacent to the European passive margin, were deformed and incorporated in the Late Cretaceous - middle Eocene accretionary wedge related to oceanic ("B" type) subduction (i.e., Marroni et al., 2001; Bortolotti et al., 2005). These units are the remnants of the Ligurian ocean (Alpine Tethys) and consist of Mesozoic to lower Tertiary sedimentary successions and subordinate Jurassic ophiolitic rocks (portion of oceanic crust). They represent the more far-travelled units of the Apennine belt and currently occupy the highest structural position in the chain.

In the Northern Apennines, the Ligurian units are the innermost and uppermost units of the Ligurian nappe (Elter, 1975; Bortolotti et al., 2001; Cerrina Feroni et al., 2002) that is represented in a more external and lower position by the Subligurian units. These units originated from the thinned continental margin of the Adria microplate (Figs. 2b, 3 and 4).

In the southern Apennines (Figs. 2 and 6), the Ligurian units (Ligurian complex *sensu* Ogniben, 1969) are classically subdivided into three main units of Upper Jurassic to lower Miocene age (see Bonardi et al., 1988; Patacca and Scandone, 2007 for major details). The two lower units respectively consist of slightly metamorphic rocks (Frido unit) followed by a metapelitic mélangé (Episcopia-San Severino mélangé *sensu* Patacca and Scandone, 2007) with serpentinite, granulite, amphibolite, granitoid and marble blocks (e.g., Spadea, 1982; Patacca and Scandone, 2007). The uppermost unit is non-metamorphic and consists of pillow basalts overlain by upper Jurassic radiolarites, shales, and quartz arenites (Timpa delle Murge Formation), black shales (Crete nere Formation) and alternating calcareous-siliciclastic turbidites (Saraceno Formation).

The Sicilide units, which also occupy the highest position in the southern Apennine thrust sheet, were originally deposited in a more external Tethyan basin (Figs. 2b) resting on basement of unknown character (thinned continental and/or oceanic crust; i.e. Patacca and Scandone, 2007).

During neo-Alpine deformation (late Oligocene – early Pleistocene) thrust accretion across the Apulia-Adriatic continental margin is shown by the east-northeastward migration of the Ligurian and Subligurian units. In the Northern Apennines, these units, which form a detached thrust sheet, tectonically overlie the Miocene sedimentary succession of Tuscan and Romagna-Marche-Umbria units (Figs. 3 and 4). In the southern Apennines (Figs. 6 and 7), Ligurian and Sicilide units tectonically overlie a complex imbricate orogenic stack consisting of carbonate platform (Lazio-Abruzzi and Campania-Lucania units) and pelagic basins of the Adriatic margin (Molise, Sannio and Lagonegro units).

In both the northern and southern Apennines, the timing of eastward migration and shortening is strongly constrained by the discordant deposition of eastward-younging top-thrust basins (middle-late Eocene – Pliocene, Figs. 1, 5 and 6), known as Epiligurian basins in the northern Apennines (Fig. 5), onto the inner Ligurian, Subligurian and Sicilide units and the outer accretionary wedge of the thrust belt (Ghisetti et al., 2003). Deposition of top-thrust basins and Epiligurian units occurred simultaneously with the east and northeast migration of the thrust-and-fold belt that, as a consequence, strongly controlled the shape and sedimentation of these basins (Ori and Friend, 1984).

Miocene – early Pliocene thrust accretion across the Apulia – Adriatic continental margin was accompanied by Tyrrhenian back-arc extension (i.e., Mazzoli and Helman, 1994).

The northwest-trending Dinaride – Albanide - Hellenide orogenic belt lies to the east of the Apulia-Adriatic foreland (Figs. 1 and 2). It is characterized by a double-vergent structural architecture bounded by west-vergent thrust faults in the external zone (to the west) and east-verging faults in the internal zone (to the east, Fig. 8a) (Dilek, 2006 and reference therein).

The west-directed thrust faults are the result of the Tertiary tectonic collision (Fig. 2b) between the Adria microplate (Apulia-Adriatic foreland) and Eurasia (Pelagonian platform), whereas the east-directed thrust faults are associated with Latest Jurassic-Early Cretaceous ophiolite emplacement (Dilek et al., 2005; Dilek, 2006).

Ophiolites are distributed along a northwest-trending belt (Figs. 1 and 8a) from the Kosovo region in the north, through Albania and Greece, to western Turkey, where they join the Tauride belt (Ghikas, 2007; Dilek et al., 2008; Ghikas et al., this issue). The northwest trending belt formed through the closure of several Tethyan basins during northeastward migration of Gondwana and the opening of the Atlantic Ocean (Dilek et al., 2005; Smith and Rassios, 2003). During the Middle Triassic – Jurassic, the Pindos basin, located at the northern margin of the Neotethys Ocean and bounded by Pelagonia to the east and Apulia to the west, rifted (Dilek et al., 2007; Sharp and Robertson, 2006). Extensive carbonate shelves developed on both of these microcontinents during subsidence along

the margins of the basin (Dilek et al. 2005). In the Middle Jurassic (Fig. 2a), the Pindos basin began to close, with the formation of a west-dipping intra-oceanic (Casey and Dewey, 1984) subduction zone. Westward subduction continued until the arrival and partial subduction of Pelagonia (Smith and Rassios, 2003) during the Early Cretaceous (Sharp and Robertson, 2006). This continued until the Tertiary oblique collision of Apulia with Pelagonia, completing the emplacement of the western Dinaride-Albanide-Hellenide ophiolite (Dilek et al. 2007; Ghikas et al., this issue). The collision-induced deformation propagated westward as an oblique convergence between Adria and Eurasia that produced a thin-skinned fold-and-thrust belt, consisting of Eocene-Quaternary sedimentary successions (Fig. 2b). Thus, along the external thrust front of the Dinaride-Albanide-Hellenide mountain belt strain partitioning occurred in and across a broad dextral shear zone in the Balkan Peninsula. This promoted the local collapse of the orogenic belt in well-developed transtensional zones (Dilek and Koçiu, 2004; Dumurdzanov et al., 2005; Dilek, 2006).

Postcollisional magmatism is widespread in the Dinarides and the Hellenides with the occurrence in the northern Greece (Voras mountains) and Macedonia of low-K calc-alkaline volcanic rocks (ca. 5.0-4.0 Ma) and shoshonitic volcanism (ca. 2.5-1.5 Ma) (Pe-Piper and Piper, 2002).

### **Geodynamic environment and processes of formation of the peri-Adriatic mélanges**

The peri-Adriatic region, provides several excellent examples and a complete case history that allow us to consider how mélanges related to different geodynamical situations during the complex geologic history of the eastern Mediterranean, as well as how different processes (tectonic, sedimentary and diapiric) were involved in their formation. Thus, we propose a preliminary subdivision and classification of the peri-Adriatic mélanges. Although many examples are preserved elsewhere in this complex region, the proposed classification scheme only applies to select examples from different sectors of the Apennine and Dinaride-Albanide-Hellenide mountain belts.

#### **1) Extensional tectonics-related mélanges**

This group refers exclusively to mélanges that formed from sedimentary, en-mass processes (mass-wasting). Bodies from debris avalanches and flows, single-slide blocks and groups of slide blocks are commonly referred to as megabreccia, olistoliths, and olistolith fields or swarms, respectively (Castellarin, 1972, 1982; Bernoulli, 2001; Camerlenghi and Pini, 2009 and references therein). The most notable examples are the Norian-Jurassic and, in part, Cretaceous megabreccias of the Southern Alps (Castellarin, 1972; Bosellini et al., 1977) and Northern Apennines (see, among many others, Castellarin et al.; 1978, Cecca et al., 1981; Castellarin, 1982; Fazzuoli et al., 1985; Bernoulli, 2001; Galluzzo and Santantonio, 2002) and the Upper Cretaceous megabreccias of the Apulia foreland in Central-Southern Apennines (Maiella, Gargano and Adriatic off-shore; see Bernoulli, 2001 and reference therein; Fig. 6). Other examples include the megabreccias and olistoliths of the Cretaceous Calcirudite a Rudiste Fm. in the Gran Sasso area (i.e., Ghisetti and Vezzani, 1986; Ghisetti and Vezzani, 1998; Bernoulli, 2001; Fig. 6). In each case, the mélanges are deposited in pelagic limestones. They have an intrabasinal composition, and their main sources are the already cemented carbonate platform rocks (frequently slightly older than the age of deposition of bodies), with some contribution of contemporaneous, slumped pelagic limestones.

In the Western Hellenides, the Upper Triassic-Lower Jurassic Agios Nicolas Formation. (see Naylor and Harle, 1976) consists of pebbly mudstones and boulders deposited by debris flows and sliding into the shale basins, triggered by extensional faulting during rifting. The mudstone matrix contains pebbles, cobbles, and boulders of sandstone (arenite and wacke) derived from the Pelagonian hinterland (Ghikas et al., this issue). A comparable example is the proto-Advella mélange, emplaced by gravitational-movement on the eastern margin of the pre-Apulian foreland (see Ghikas et al., this issue).

Sedimentary bodies associated with extension are thought to be related to either 1) normal faults dissecting and controlling the margins of carbonate platforms (Castellarin, 1972, 1982; Eberli, 1987; Accordi and Carbone, 1988; Eberli et al., 1993; Vecsei, 1991, 1998; Bosellini et al., 1993; De Alteris and Aiello; 1993; Graziano, 2000; Rusciadelli, 2005) during and after rifting (Naylor and Harle, 1976; Fazzuoli et al., 1985; Marchegiani et al., 1999; Bernoulli, 2001), or 2) the geometry of the carbonate platform margin (scaloped margins, see Bosellini, 1998).

#### **2) Passive margin (?)-related mélanges**

Passive margins instability may trigger the progressive deformation of basin slopes, such as in the case of the carbonate and siliciclastic fine-grained turbidites and mudstones of the Early Cretaceous Palombini Shales (locally defined Ronchi argillites), which constitutes the lowest horizon of the stratigraphic succession of the Modino mass-wasting complex (Modino “basal” complex, De Libero, 1998; Pini et al., 2004; Fig. 4). The Ronchi argillites were

deformed at various degrees, from in-situ folding and boudinage to slump-slide structures, evolving down flow to cohesive debris flows. The latter led to complete stratal disruption and development of a block-in matrix fabric. Sediments were deformed when poorly or non-consolidated and show plastic deformation of clasts with enhanced fluidal structures of the matrix (Pini et al., 2004; Camerlenghi and Pini, 2009).

Similar examples are described in the Othris Mountains (in Greece) by Smith et al. (1979). Here, late Jurassic- early Cretaceous radiolarian siltstones were deformed by debris flows and slumping related to instabilities of the passive margin (Smith et al., 1979). The ophiolitic *mélange* of Albania (Shallo, 1990) also formed during the late Jurassic-early Cretaceous continental margin evolution, due to development of slides triggered by tectonic and sedimentary events.

All of these bodies are mainly monomictic, recycling the sediments of the same formation. The major problem with this interpretation concerns the triggering mechanism. These deformed bodies are found in the Early Cretaceous, after rifting concluded, and in a passive margin setting at the edge of the thinned Adria continental margin or at the transition to the oceanic realm (see, e.g., De Libero, 1998; Pini et al., 2004). Many processes could have triggered the slides in these situations, as well as the Late Cretaceous to early Tertiary “extensional” olistostromes in the Adria-based units (Tuscan, Umbrian domains). Some possibilities include: 1) tectonic reactivation of extensional Jurassic structures, 2) the influence of pre-existing submarine topography, 3) deep ocean processes, and/or 4) extension throughout the Cretaceous and early Tertiary (Fazzuoli et al., 1985; Montanari et al., 1989; Marchegiani et al., 1999; Bernoulli, 2001). An alternative explanation considers the onset of Early Cretaceous compression related to subduction at the northern edge of the Ligurian ocean, as suggested by some paleogeographic reconstructions (Schettino and Scotese, 2002).

### 3) Strike-slip tectonics-related *mélanges*

The more classic example in the international literature of the Mediterranean area are the Moni and Arakapas *mélange* in Cyprus (Robertson, 1977; Krylov et al., 2005). They are mud-rich, block-in-matrix bodies, which are considered to be of both tectonic and sedimentary origin. The Arakapas *mélange*, (Krylov et al., 2005) is an olistostrome developed along fault scarps during Cenomanian-Campanian strike-slip activity in a fore-arc basin transform zone (Troodos zone). The younger, Tertiary, stage of the Moni *mélange* (Neo-Moni, Krylov et al., 2005) is a tectonic *mélange* related to the activity of a transpressional wrench fault, reactivating olistostromes associated with subduction (Robertson, 1977).

In the Northern Apennines, the lower portion of the Eastern Ligurian sedimentary succession, the so-called basal complexes of the Helminthoid flysch (see Abbate et al., 1970; Bortolotti et al., 2001; Marroni et al., 2001; Cerrina Feroni et al., 2002) (Fig. 2b) are characterized by Late Cretaceous cohesive debris flows with a “brecciated” argillaceous matrix (classic olistostromes) mostly sourced in Early Cretaceous Ligurian sediments (Palombini shales). These, in turn, host large-scale (from hundred of meters to several kilometres in length) ophiolitic blocks and continental margin fragments (Marroni et al., 1998). Marroni et al. (2001) interpret these bodies as products of submarine landslides or a portion of the sea floor that was uplifted by synsedimentary transpressive faults, acting in the retroarc basin of the paleo-Apenninic accretionary wedge. Detrital evidence from the oceanic crust has been used as a paleogeographic key in discriminating between “western” and “eastern” External Ligurian units (Marroni et al., 2001).

The lower part of the Messinian *mélange*, of the northern Tertiary Piedmont Basin (Torino Hill and Monferrato, Figs. 1 and 9), represents another example of strike-slip tectonic *mélanges* (Festa et al., 2005; Dela Pierre et al., 2007; Festa, 2009). Regional scale wrench-faulting (e.g. Rio Freddo deformation zone *sensu* Piana and Polino, 1995, Fig. 1) dismembered of the previously coherent evaporitic succession and produced a tectonic disrupted unit.

Other comparable examples are exposed along the Villalvernia- Varzi line (Fig. 1), a regional and polyphase deformational zone that juxtaposed the Tertiary Piedmont Basin and the northern Apennines during late Oligocene to late Miocene. Here, strike-slip movements mixed three different chaotic rock bodies (Festa, unpublished data) originally deposited at the base of the Epi-Ligurian units (Calcari di Zebedassi *Auct.*, Upper Cretaceous in age, see Boni et al., 1970) and as olistostromes within the Epi-Ligurian units (lower and upper Miocene olistostromes, Complesso caotico and Complesso caotico indifferenziato *sensu* Boni et al., 1970), thus producing a new strike-slip tectonic *mélange*.

The Thitonian ophiolitic *mélange* of Albania (Shallo, 1990; Rassios and Dilek, 2009, Fig. 8) may also have been associated with strike-slip tectonics during the late Jurassic - early Cretaceous. Its general trend and structure crosscut different ophiolitic (i.e. Kashnjet-Gornisique-Kcire) and/or continental margin successions (i.e.



Barmash-Leskovik area). It is characterized by zones of elongated blocks set within an argillitic-foliated matrix (Shallo, 1990).

#### **4) (Oceanic crust) Subduction-related mélanges**

These types of mélangé could be subdivided on the basis of their formational processes as:

##### ***4a - mass-wasting deposits at the wedge front***

Mass-wasting deposits are widely abundant in the Bocco Shales, above the Internal Ligurian units (Figs. 4 and 5). These Paleocene (Danian) bodies overlay a complete stratigraphic succession. The lower term is a complete sequence of oceanic crust (Marroni and Pandolfi, 2001). Very thick (hundreds of meters), dismembered slide blocks (olistoliths) are present together with bodies from mud-rich cohesive debris flows (olistostromes), very coarse to coarse turbidites, thin-bedded turbidites, and mudstones. Slide blocks and olistostromes are composed of the same Jurassic ophiolite sequence and the related Late Jurassic to early Paleocene sedimentary sequence, which outcrops below the Bocco Shales. The largest blocks consist of contorted thin-bedded turbidites of the same Bocco Shales, interpreted as slide blocks from a steep slope. According to Marroni and Pandolfi (2001), slide blocks and olistostromes of the Bocco Shales were generated by episodic instability of the front of the Paleocene accretionary wedge due to the subduction of reactivated faults offsetting the oceanic crust and causing relief in the lower plate. In modern subduction systems, subducting sea-mounts and ridges cause slope oversteepening and front-of-wedge instability, resulting in gigantic submarine landslides (e.g., Hikurangi margin, continental margin of New Zealand, Collot et al., 2001; Costarica continental margin, von Heune et al., 2003; Hühnerbach et al., 2005). Mud diapirism, which has been suggested as a triggering mechanism for similar deposits in the Western Alps (Di Giulio, 1992), has been discounted by the lack of diapiric structures (Marroni and Pandolfi, 2001).

The Late Cretaceous pre-flysch successions of the Internal Ligurian (Fig. 5) units are characterized by large scale (several tens of meters thick) mud-rich olistostromes, such as the classic Forcella Pass olistostrome in the Val Lavagna Shales (Bortolotti et al., 2004), that are mainly composed by older, Early Cretaceous rocks (Palombini Shales), and ophiolitic clasts (Abbate et al., 1970). The leading mechanisms of transport and deposition were cohesive debris flows acting on already consolidated, tectonized and disrupted units. These bodies episodically occur as meter to tens of meter thick sedimentary bodies in well-bedded, normal successions. Considering the Late Cretaceous geodynamic setting and the examples of the younger Bocco Shales, the hypothesis of instability of a passive margin (Naylor, 1981) can be rejected (Bortolotti et al., 2004) in favour of instability on a tectonically active slope at the front of the paleo Apenninic accretionary wedge, as suggested by Görler and Reutter (1968), but more definitive conclusions have not been reached.

##### ***4b - Tectonic mélanges and broken formations***

A large number of stratally-disrupted units in the External Ligurian units (Fig. 5) belong to this group, after the so-called basal complexes, chaotic complexes, or argille scagliose of the Northern Apennines have been distinguished in either Epiligurian olistostromes (see Type 6), or stratally disrupted stratigraphic units (broken formation or tectonosomes, Pini, 1987; Bettelli and Panini, 1987; Castellarin and Pini, 1987 and following papers, see references in Pini, 1999 and Camerlenghi and Pini, 2009). The latter, defined as broken formations and tectonosomes formed from the interaction of different subduction-related processes: layer-parallel boudinage in an early stage of subduction by fluid overpressure and lithostatic loading (Bettelli and Vannucchi, 2003), progressive oblate-type boudinage and transposition of sedimentary layers by protracted sheath folding, associated with large scale thrusting (Vannucchi and Bettelli, 2002; Bettelli and Vannucchi, 2003), block stacking by both large scale and mesoscopic-scale duplexing and thrusting (Pini, 1999; Cowan and Pini, 2001).

Due to their progressive deformational origin, the finite style and fabric of these mélanges depends on the rheology of the rocks involved, that is from the mechanical characters of the multilayer, which reflect the main depositional environment and the composition (see Bettelli and Vannucchi, 2003). Different deformational styles within diverse stratigraphic intervals in the same sequence may be, therefore, explained by the rheology of the multilayer, but can also be related to the superposition of different tectonic phases (see Pini, 1999), namely the Upper Cretaceous, eo-alpine phases and the Eocene, meso-alpine, or Ligurian phase (see, among many others, Vai and Castellarin, 1993; Pini, 1999; Vescovi et al., 1999; Daniele and Plesi, 2000; Bortolotti et al., 2001; Marroni et al., 2002; Catanzariti et al., 2007).

Different intensity as of stratal disruption are recognizable in the Northern Apennines broken formations-tectonosomes. The “broken formations” in Fig. 5 are Neocomian to Campanian basal complexes, detached from the Ligurian Helminthoid flysches, that contain structures ranging from contorted, highly-folded and boudinaged beds to

complete stratal disruption (Vannucchi and Bettelli, 2002; Bettelli and Vannucchi, 2003 and references therein). The tectonosomes of Pini (1999) correspond to an entire Ligurian-type succession (Fig. 5), ranging in age from Neocomian to lower Eocene (Sillaro-Samoggia structural unit, see Cerrina Feroni et al., 2002), which is more severely deformed, with a prevailing complete stratal disruption (Pini, 1987; 1999, Conti, 1987).

The Frido Formation (late Jurassic – early Cretaceous) of the Ligurian units (Fig. 6-7) is a tectonic *mélange* that represents the remnants of the outcropping deeper portion of the accretionary wedge of the Southern Apennines. It underwent collision with the Apulia passive margin during the early Miocene (Knot, 1994), producing a series of thrust sheets with different lithological features and metamorphic overprinting that records subduction of oceanic crust and deformation during HPLT metamorphism (Laurita et al., 2009 and reference therein).

In Greece (Vourinos Mountains., see Ghikas et al, this issue) and Albania (Dilek et al., 2005, 2008) meter-to kilometre-scale, exotic blocks of ultramafic rocks and ophiolites are seen in an early Cretaceous sedimentary *mélange* (i.e., Vourinos sub-ophiolitic *mélange*) resting on the Pelagonian platform succession (Fig. 8). The Vourinos sub-ophiolitic *mélange* rests tectonically on the Pelagonian microcontinental margin. Its internal structure and stratigraphy is reminiscent of a block-in-matrix sedimentary *mélange* representing initially a broken formation that formed in a continental slope-rise setting (Ghikas et al., this issue). However, this *mélange* was subsequently tectonically incorporated into the Pelagonian margin during the obduction of the Vourinos ophiolite in the Middle Jurassic during which it was pervasively metamorphosed and tectonized.

### 5) **Collision-related *mélanges*.**

In the Apennines, no *mélanges* have been conclusively attributed to this because the timing of the end of subduction and the inception of intracontinental deformation varies greatly throughout paleogeographic extent of the Ligurian ocean (see Marroni and Treves, 1998; Cibin et al., 2001; Cerrina Feroni et al., 2002). There are also multiple interpretations regarding the paleogeographic positions of the diverse Ligurian units (see, e.g., Daniele and Plesi, 2000; Bortolotti et al., 2001; Argnani et al., 2004). The late Eocene olistostromes at the base of the Epiligurian succession (Figs. 4 and 5) above the Ligurian nappe (Type 6c) may represent this type of *mélange*. The Advella *mélange* in the Western Hellenides, for example, could represent deformation that has occurred since the Eocene as a result of continental collision (Fig. 8) between the pre-Apulian foreland and Pelagonia (Ghikas et al., this issue). These polygenic *mélange*, originally emplaced by sedimentary processes (see 1. Extensional tectonic-related *mélanges*), is characterized by a block-in-matrix fabric with blocks ranging in age from Mid-Triassic to Cretaceous, and was emplaced onto Cretaceous-Eocene shelf and turbidite deposits of pre-Apulia (Kostopoulos, 1988; Jones and Robertson, 1991; Rassios and Moores, 2006). It contains rocks associated not only with the initial rifting of Pelagonia and pre-Apulia, but also with the development of a mature carbonate platform, the deposition of thick shallow-marine turbidite and detrital sequences, and the emplacement of the Pindos ophiolite (Jones and Robertson, 1991; Ghikas et al., this issue). The Advella *mélange*, therefore, represents a much more complete history of the Pindos basin than the Vourinos *mélange*.

### 6) **Intracontinental deformation, "obduction" related *mélanges***

This group is by far the most common type observed in the Peri-Adriatic *mélanges*. It is related to obduction of the accretionary wedge over the continental crust and of the "oceanic" nappe translation.

#### *6a - Sub-nappe *mélanges**

##### 6a1- precursory olistostromes

Considered among the classic olistostromes of the Apennines (Abbate et al., 1970, 1981), these sedimentary bodies contain a typical block-in-brecciated-matrix fabric (type A olistostromes), or are aggregates of individual blocks, supported or not supported by a brecciated matrix (Type B and C, respectively, Lucente and Pini, 2003). These bodies were deposited by cohesive debris flows and/or blocks avalanches (Figs. 3 and 10) in migrating foredeep basins (see, for the northern Apennines, Lucente and Pini, 2008). Defined as precursory olistostromes by Elter and Trevisan (1973), these bodies have been described elsewhere at the front of tectonic *mélanges*, facing lateral or frontal ramps or at the front of accretionary wedges and/or nappe systems (Tuscan, Umbrian, Lazio-Abruzzi and Molise foredeep successions, see, e.g., Abbate et al., 1970, 1981; Sgrosso, 1988; Pescatore et al., 2000; Cowan and Pini, 2001; Pini et al., 2004; Vezzani et al., 2004; 2009; Festa et al., 2006; Lucente and Pini, 2008; Patacca and Scandone, 2007; Camerlenghi and Pini, 2009). In the northern Apennines, olistostromes are present in all stage of the migrating foredeep complex, from the early Oligocene Macigno foredeep to the middle-late Miocene Marnoso-arenacea foredeep, and in the Messinian to Pliocene front-Apenninic deposits (Figs. 3 and 10) (see, e.g.,

Abbate et al., 1970, 1981; Ricci Lucchi, 1986; Conti, 1987; Ricci Lucchi and Vai, 1994; Cornamusini, 2001; Roveri et al., 2002; Landuzzi, 2004; Lucente and Pini, 2003, 2008). They are the result of the collapse of the wedge front, resedimenting Ligurian and Subligurian rocks and deposits from the wedge-top Epiligurian basins (Lucente and Pini, 2008 and references therein). They are often associated with intrabasinal mass-wasting deposits, consisting of sediments from the margins of the foredeep basins close to the front-of-the-wedge (inner slopes) and the same deposits of the basin plains (Lucente and Pini, 2003, 2008).

In the Central and Southern Apennines, for example, chaotic rock bodies of argille scagliose have been emplaced in the foredeep sediments at the front of tectonic *mélange* of the Sicilide units (Figs. 6 and 11). These olistostromes, which are interbedded in the different foredeep successions from the inner peri-Tyrrhenian region (upper Tortonian Frosinone Fm., see Cosentino et al., 2002) to the outer peri-Adriatic region of the Matese platform area (Pietraraja Flysch, upper Tortonian – lower Messinian in age, see Sgrosso, 1988; Vezzani et al., 2004, 2009; Festa et al., 2006; Patacca and Scandone, 2007), testify to the eastward migration of the Sicilide units.

Different factors are able to trigger these sediments failures. In fact, earthquakes at active margins, rather than oversteepening slope angles, are the most efficient factor in increasing stresses in weakened sediments and triggering failures (Maltman, 1994; Camerlenghi and Pini, 2009).

#### 6a2 - Olistostromal carpet at the base of a nappe

This group is conceptually related to the previous one, since it comes from the protracted activity of debris flows and avalanches at the front of an advancing nappe or tectonic *mélange* (Figs. 4 and 11). This coalescent carpet of olistostromes has been recognized at the base of the Ligurian nappe in the Bologna area (Landuzzi, 2004; Pini et al., 2004; Camerlenghi and Pini, 2009) as the lowest part of a system of imbricated stacks of normal bedded units, tectonosomes, tectonic *mélanges*, and olistostromes at the base of the Ligurian nappe (Pini, 1987, 1993; Bettelli and Panini, 1992; Landuzzi, 2004). This widely extending composite lithosome, see Fig. 5, is directly connected to the precursory olistostromes, as suggested by Lucente and Pini (2008) (Fig. 10), and consists of extrabasinal (Ligurian-Subligurian) and intrabasinal (wedge-top basins, basin margin and foredeep) blocks and bodies.

This olistostromal carpet, some of the related large-scale mass wasting complexes, such as the Modino mass-wasting complex (Figs. 4 and 10), and part of the stack of units at the base of the Ligurian nappe, have been grouped together in the Sestola-Vidiciatico unit and interpreted as an equivalent of a subduction channel by Vannucchi et al. (2008). These two interpretations are not in disagreement. The origin of the Sestola-Vidiciatico unit from submarine landslides at the front of the Ligurian nappe is not excluded, at least in part, by Remitti et al. (2007) and Vannucchi et al. (2008). They do provide examples of tectonized olistostromes. The Sestola-Vidiciatico unit (Fig. 3), or the part coinciding with the olistostromal carpet, is therefore to be considered a mixed tectono-sedimentary *mélange*, because: 1) the exotic blocks are supplied by mass-wasting episode at the front of the wedge, sourcing rocks from the front of the wedge (Ligurian-Subligurian) and from the wedge-top basins (Epiligurian), and also involving the slope and part of the basin plain, as in the case of Monte Modino complex (Lucente et al., 2006; Lucente and Pini, 2008); and 2) part of the exotic blocks may derived from erosion at the base of the nappe (Vannucchi et al., 2008). These intimate interplay of tectonic and sedimentary processes closely recalls observations from the front of modern convergent wedges during erosion (see, e.g., Choconat et al., 2002; Ranero and von Heune, 2002; von Heune et al., 2003), and confirms the validity of the Sestola-Vidiciatico as a fossil equivalent of a subduction channel, even if it is related to the emplacement of an “oceanic” nappe atop an intracontinental stacking of units, above the continental crust of Adria (Boccaletti et al. 1990; Castellarin et al., 1992; Barchi et al., 2001; Carmignani et al., 2001; Argnani et al. 2003).

The Cretaceous sedimentary *mélange* exposed in the Othris Mountain (Greece) (see Smith et al., 1979), is a debris flow that was deposited just in front of the advancing ophiolitic thrust sheet. The east-ward displacement of the thrust sheet tectonically deformed the *mélange* that, located in the lowest part of a system of embricated stack, represented an olistostromal carpet enveloping exotic blocks of ultramafic rocks and ophiolites (up to 200 meters wide).

#### 6a3 - tectonic *mélange* and tectonic-sedimentary *mélanges* at a base of a nappe (6a1 plus 6a3)

In other areas of the circum-Mediterranean region, the concentration of olistolithes and olistostromes below the base of allochthonous nappes is emblematic (e.g. Ankara *mélange*, see Bailey and Mc Callian, 1950, 1953).

In the Central and Southern Apennines (Fig. 6), the Flysch Rosso (late Cretaceous-early Miocene) and the Sicilide argille scagliose (Late Oligocene – Early Miocene) represent notable examples of tectonic-sedimentary *mélanges* at the base of the Molise units (Figs. 7 and 11) and of the Mt. Moschiatturo Klippe in the

Sannio-Molise region (Vezzani et al., 2004; 2009; Festa et al., 2006). In spite of an intense stratal disruption that led locally to a block-in-matrix fabric, the stratigraphy (alternating varicoloured clayey marls, grey-reddish marls and siliceous calcarenites), can still be recognized and mapped. It is characterized by a structurally-ordered block-in-matrix fabric. In the matrix, pervasive scaly fabric is present and often associated to mesoscale S-C shear zones, defined by centimeter-to decimeter-sized lithons.

The whole internal structure consists of fault-bounded bodies ranging from decimeter (Mt. Moschiatturo)-to kilometer (Molise units) scale. Tectonic shearing, acting at the base of the units, was probably the main factor in disrupting the originally coherent succession. Tectonic loading probably triggered overpressure conditions at the base of the unit, as fluids expelled upwards by the clayey matrix produced brittle deformation in the calcareous interlayers (Festa et al., 2006). Locally upward rising of overpressured and underconsolidated sediments and mud diapirs, as is frequently imaged in seismic reflection profiles in Plio-quadernary sediments above the Apenninic frontal thrust, either north of Gargano, or in the Gulf of Taranto, is not excluded (Roure et al., 1991).

#### **6b - *intra-nappe mélanges*** (related to deformation during nappe translation)

These types of *mélange* are subdivided on the basis of the processes involved in their formation. These types are:

##### 6b1 - Sedimentary

Submarine mass-wasting bodies, dominantly composed of only interbasinal sediments, have been described as olistostromes by Jacobacci (1963), Abbate et al. (1970; 1981) and Elter and Trevisan (1973).

Some notable examples include the Breccia della Renga (Serravallian – Tortonian, see, for example, Cipollari and Cosentino, 1995) at Mt. Simbruini (Fig. 6) and the Breccie di San Massimo (Late Tortonian – Lower Messinian, see Ghisetti and Vezzani, 1998, Fig. 6) at the front of the Matese platform (Central Apennines). These lenticular chaotic rock bodies are composed only of intrabasinal sediments, emplaced by submarine mass-wasting phenomena (i.e., Ghisetti and Vezzani, 1986; Vezzani et al., 2004; 2009; Festa et al., 2006). In southern Apennines, the Gorgoglione Flysch (Langhian – Serravallian) is characterized by channelled polygenetic calcareous conglomerates and sandstones, olistostromes of Liguride and Sicilide units (Sauro and Basento rivers, see Carbone and Lentini, 1990) and calcareous olistoliths (Mt. Carruozzo), up to a kilometer wide (Vezzani et al., 2009) (Fig. 6).

##### 6b2 - Tectonic and/or tecto-sedimentary

The Coscogno *mélange* in the Northern Apennines is a rather puzzling unit, that has been recognized and mapped by Bettelli et al. (2002). Exhumed intermixed chunks of Ligurian, Epiligurian and Subligurian units comprise this *mélange*. According to Bettelli et al. (2004), these lithosomes of different paleogeographic origin represent thrust splays with contractional contacts. This *mélange* should represent a tectonic *mélange* developed deep inside the Ligurian nappe (Bettelli and Panini, 1992; Capitani, 1993). The exhumation of the *mélange*, and the involvement of the Epiligurian deposits(?) occurred in the late Oligocene-early Miocene, and has been attributed to either out-of-sequence thrust dissecting the entire Ligurian nappe, transpressional activity along strike-slip faults, or to normal faulting (Capitani, 1993). The scarce continuity of the outcrops does not allow for a determination of the nature of the contacts inside and around the body, so an alternative interpretation of these *mélanges*, as separated, stacked Ligurian and Subligurian units and subunits that are exposed in a tectonic window, has been proposed (Cerrina Feroni et al., 2002).

The outer accretionary wedge of the central-southern Apennines (Mt. Frentani *mélange*, Vezzani et al., 2004, 2009; Festa et al., 2006; Torrente Calaggio Fm., Patacca and Scandone, 2007 and the Falda di Metaponto *Auct.* Figs. 6 and 7) is marked by a 10-15 km wide belt of *intra-nappe* tecto-sedimentary *mélange*. It consists of a block-in-matrix arrangement with intraformational block coming from Cenozoic Molise units, the Messinian evaporitic succession and late Miocene – lower Pliocene foredeep deposits (Ghisetti et al., 2003; Vezzani et al., 2004, 2009; Festa et al., 2006; Patacca and Scandone, 2007). The blocks range in size from decimeters-to hundred of meters. The matrix mainly consists of brecciated matrix with the locally overprinting of a pervasive scaly cleavage. The emplacement and deformation of these chaotic bodies (Fig. 11) is consistent with slope and debris avalanches at the external frontal thrust that produced mass-gravity deposits during tectonic deformation (Ghisetti et al., 2003). The rapid burial provided by the deposition of the early-middle Pliocene top-thrust and foredeep clastic Mutignano successions (late Pliocene – early Pleistocene) could have caused pore-fluid dissipation and promoted the upward rise of overpressured chaotic sediments, exceeding hydrostatic pressure (mud diapirs; diapiric *mélange*) as suggested by some outcrops north of Atessa (early – late Pliocene). Later deformational stages of the already emplaced *mélange* are associated with folding and thrusting of the Apulia platform.

### **6c - Epi-nappe mélanges**

These types of mélange are subdivided, based of their formational processes into three categories. These categories are:

#### 6c1 – Sedimentary

One of the most notable examples of this type of mélanges is the Messinian mélange in the episutural Tertiary Piedmont Basin (Figs. 1 and 9). It represents a sedimentary body of mixed rocks, resulting from dismemberment of the originally coherent stratigraphic succession (Dela Pierre et al., 2002; 2007; Irace, 2004; Festa et al., 2005; Irace et al., 2005; Festa, 2009). This succession consisted of pre-evaporitic Tortonian - lower Messinian hemipelagic marls and shallow-water Messinian gypsum deposited in response to the Mediterranean salinity crisis. Despite interpretations (Dela Pierre et al., 2002; 2003; Irace, 2004; Irace et al., 2005) that have related the Messinian mélange (Valle Versa chaotic complex *sensu* Dela Pierre et al., 2002) to gravity-driven phenomena triggered by intra-Messinian tectonics, Festa et al. (2005; 2009), Dela Pierre et al. (2007) and Festa (2009) have shown that Messinian mélange has a more complex stratigraphic and structural arrangement. It is a composite chaotic unit formed from different bodies of mixed rocks from a variety of processes (tectonic, gravitational and diapiric), each of which can be inferred to represent a mélange-forming process. Tectonic disrupted units, gravity-driven sedimentary units and diapiric-disrupted units of mixing rocks (*sensu* Festa, 2009) correspond to type 3 (strike-slip tectonic-related mélange), type 6c1 (epi-nappe sedimentary mélange) and type 6c3 (epi-nappe diapiric mélange), respectively, of this paper.

Several olistostromal bodies characterize the episutural, wedge-top Epiligurian deposits (Figs. 3, 4, 5 and 10) and dominantly consist of mud-rich, block-in-brecciated-matrix deposits related to cohesive debris flows, which may also host large blocks (meters to tens of meter thick) of both well-stratified and stratally-disrupted Ligurian units. Generally speaking, the main components are from previously lithified and deformed External Ligurian units, although blocks and detritus from the contemporaneous and slightly older Epiligurian deposits are also present.

The dimension of the bodies spans from some hundreds of square meters to some tens of square kilometers of areal extent and from few meters to as much as 300-400 meters in thickness (Pini, 1987, 1999). In the largest bodies, thick (some tens of centimeters to several meters) zones of banding and fluidal structures in the matrix at meter to centimeter-scale are present at the base and inside the bodies (Pini, 1999) and display simple shear-related flow structures that geometrically resemble mylonitic fabrics. These thick olistostrome bodies can be, therefore, considered as either stacks of anastomosing smaller bodies (in analogy with examples in Cyprus, see Swarbrick and Naylor, 1980), or the result of progressive, multiphase emplacement of large bodies with independent relative motion of discrete masses along shear zones.

Their stratigraphic distribution is almost continuous through the entire Epiligurian stratigraphic record (from late Eocene to Pliocene olistostromes, see, e.g., 1:50.000 Italian Geological Map, Bologna sheet). They seem to be more frequent and more widely distributed at late Eocene base of the Epiligurian succession (Figs. 3, 5 and 10), marked by the Baiso breccias, and the lower Miocene Canossa-Val Tiepido olistostrome (Bettelli et al., 1987). Some, very laterally extended and hundred of meters thick chaotic sedimentary bodies, such as the early Oligocene Specchio megabody, have not been compared with the olistostromes yet, because of a more “coherent” aspect at the outcrop due to the higher abundance of rafting blocks and slabs.

Frequently, the overlaying sediments sink inside the olistostromes, with viscous flow of the block-in-matrix rocks around the “normal” bedded Epiligurian strata, as well as the basal contacts of the olistostrome are almost vertical and seem to “pierce” their substratum of tectonosome units. One possible explanation can be a post-depositional reactivation (as mud- diapirs?) of the olistostromes, induced by their underconsolidated state and high pressure of pore fluids. The magnitude of these phenomena, already suggested for the Bologna foothills (see, e.g., Pini, 1999; Pini et al, 2004) deserves further study.

#### 6c2 - Tecto-sedimentary

Some notable examples are most of the Sicilide units (late Cretaceous – early Miocene) exposed in the Central - Southern Apennines which are mainly emplaced by tectonic and subordinate sedimentary processes. They are well-exposed in the Campobasso sector (Molise, Figs. 6 and 7) where a large (more than thousand km<sup>2</sup>) and thick (up to 2000 meters) nappe (“Coltre sannitica” *sensu* Selli, 1962, Sannio unit *sensu* Patacca et al., 1992) tectonically overlies the Molise units (Fig. 7). It consists mainly of a block-in-matrix fabric that lacks total stratal disruption (Vezzani et al., 2004; 2009; Festa et al., 2006). Large blocks (up to few km<sup>2</sup> wide), which preserve the original stratigraphic succession, show preferential distributions that are aligned with main shear zones. The

emplacement of this huge nappe is consistent with the tectonic movement above the Apennine thrust sheet from the inner to the outer portions. During this movement, local mass-wasting phenomena and gravitational sliding were triggered by tectonic instability.

### 6c3 - Diapiric

In some cases, the epi-nappe mélanges are (or are suitable to be) reactivated as diapirs. As previously described (see 6c1), this is documented for the Messinian mélange of the Tertiary Piedmont Basin (Festa et al., 2005; Dela Pierre et al., 2007; Festa, 2009, Figs. 1 and 9). Here, diapiric bodies (up to tens of meter wide) pierce the gravity-driven and tectonically disrupted units of the Messinian mélange, causing their partial reorganization. Loading from deposition of gravity-driven chaotic sediments, the presence of low permeability layers (primary evaporites), and strike-slip faulting may have caused the Messinian mélange to locally reactivate as a diapir (Festa, 2009).

Other comparable examples in different sectors of the Tertiary Piedmont Basin and Northern and Southern Apennines are still being examined (Codegone, Festa and Pini, unpublished data).

### **Discussion and conclusions**

The examples of mélanges described in this paper represent only a few of numerous excellent examples preserved in the peri-Adriatic region. We have proposed a preliminary, subdivision and classification of the peri-Adriatic mélanges, with the perspective of investigating some classical mélange problems. The main problems we have examined are: (1) whether a relationship exists between different types of mélanges and their tectonic paleogeographic settings, and (2) in which geotectonic setting(s) mélanges are most abundant and influential.

**Obducted ophiolitic and ophiolite-bearing submarine nappes:** The largest number of mélanges occurs at the base of submarine allochthonous nappes, in the examples discussed here and in the other circum-Mediterranean chains (see Camerlenghi and Pini, 2009). Particularly, the most abundant cases are related to ophiolitic or ophiolite-bearing nappes, already obducted, and overriding and moving onto a continental plate and a foreland basin system. The main question regarding this group of mélanges is their origin, specifically regarding the contributions of sedimentary vs. tectonic processes at their onset and evolution. The different methodological approaches and geodynamic visions of various authors have caused several interpretative problems, but most of these sub-nappe mélanges are derived from the interplay of both tectonic and sedimentary processes. Sedimentary mass-wasting processes at the front of the nappes supply a considerable amount of materials to the mélanges throughout their formation and evolution, through the time of nappe translation. This is recorded by the presence of precursory olistostromes in the sedimentary record of migrating foreland basins, at least in the Apennines and the Dinarides-Albanides-Hellenides as described in this paper, in the Alps (wildflysch units, Camerlenghi and Pini, 2009 and references therein) and in Turkey (Camerlenghi and Pini, 2009). Tectonic processes, however, play a significant role when elements from the base of the nappe are incorporated into the basal mélange through possible frictional-erosional processes (Jurewicz et al., 2007), in a similar way to subduction erosion (von Huene and Ranero, 2003), or to folding-thrusting of the basal contact of the nappe (see Catalano et al., 2007) related to out-of-sequence thrusts (see Castellarin and Pini, 1989).

**Mélanges at the base of accretionary prisms and subduction channels:** Mélanges associated with subduction of oceanic crust (accretionary wedges) appear to be an entirely quite different situation. The circum-Adriatic, fossil sedimentary mélanges are only concentrated in peculiar situation and along certain stratigraphic horizons (see the example of the Paleocene Bocco Shales of the Internal Ligurian units, Northern Apennines). They do not have an ubiquitous and continuous distribution in time and space through the sedimentary record. Accretionary-wedge-related sedimentary mélanges are therefore not so common, they are outnumbered by the “continental” sub-nappe mélanges, and their volumetric relevance is scarce (see Fig. 5).

This seems to be consistent with the actualistic scenario, suggested by the examples of the Pacific convergent margins. The concentration of mass-wasting bodies in this setting is scarce, and they are very small in comparison to other geodynamical settings, such as passive margins (see discussion in Camerlenghi and Pini, 2009); with the notable exception of the situation of tectonic erosion (see, e.g., Choconat et al., 2002; Ranero and von Heune, 2002; von Heune et al., 2004) and the collapse of front of the wedge due to subduction of relevant lower plate irregularities (seamounts, volcanoes, normal faults, etc., see, e.g., Collot et al., 2001; von Huene et

al., 2004; Hühnerbach et al., 2005). The largest fossil mass-wasting deposits are in fact the expression of similar geotectonic situation as evidence by the Bocco Shales (Marroni and Pandolfi, 2001).

In this geodynamical setting the tectonic *mélanges* should prevail. In the circum-Adriatic chains, several disrupted formation have been recently interpreted as related to subduction processes (Pini, 1999; Vannucchi and Bettelli, 2002; Bettelli and Vannucchi, 2003). However, these rock units are severely to completely disrupted stratigraphic units that always maintain their lithostratigraphic and chronologic identities (broken formations, tectonosomes), and not true *mélanges*. Broken formations-tectonosome do not contain exotic blocks, and are conceptually more similar to the “normal” bedded, but strongly deformed basal complexes of the Internal Ligurian units and some of the External Ligurian units (Levi et al., 2007). The “normal” bedded basal complexes, moreover, have not been considered as Argille scagliose or *mélanges* at all, but they display zone of higher stratal disruption (see Levi et al., 2006) comparable with the zone of moderate disruption of the units labelled as “broken formation” in Fig. 5.

Generally speaking, considering the basal complexes of the Ligurian units as a whole and with the only exclusion of the sedimentary *mélanges* at the base of the External units (Fig. 5), there is a progressive grading in the intensity of stratal disruption between the normal bedded units and the broken formations and tectonosomes. The only *mélanges*, referred to as merely tectonic that have been reported in this paper, always show characteristic features that suggest a possible contribution of sedimentary processes in including the exotic blocks or defining certain fabric elements. One can wonder if solely tectonic *mélanges* can really be found in the circum-Adriatic realm.

**Mud-diapiric reactivation of sedimentary and/or tectonic *mélanges*:** Another important point is the existence of *mélanges* related to mud diapirs and volcanoes. All the examples we describe are related to the reactivation of previously deposited sedimentary and/or tectono-sedimentary *mélanges* as mud diapirs. Two main points about this group of *mélanges* can be discussed: do they occur in a particular geodynamic setting, and if so what is the role of this setting in producing *mélanges*?

The occurrence or rare but well-exposed examples of mud diapirs or minor mud-diapiric phenomena in different geodynamic setting (see, for example, the sub-nappe Flysch Rosso, the intra-nappe Mt. Frentani *mélange*, the epi-nappe Messinian *mélange* of the Tertiary Piedmont Basin and the Epiligurian olistostromes) suggests that their formation is not strictly related to a particular setting. On the contrary, fluid overpressure caused by the quick loading provided by deposition of gravity-driven sediments (or olistostromes) or by the emplacement of tectonic nappe, the presence of low-permeable clayey layers and faulting, represent necessary factors to form mud diapirs. These conditions are easily reached in the environment and mechanism in which *mélanges* form. Moreover, deep seated olistostromes, or water-rich, tectono-sedimentary *mélanges* are probably the major candidates in supplying mud diapirs, as observed in the Gulf of Cadiz and the Alboran Sea (see Camerlenghi and Pini, 2009 for major details). High pore-fluid content and a very high porosity remain inside mass-wasting deposits long after their emplacement.

Tectonosomes and strongly deformed and dewatered tectonic *mélanges* should be less prone to generating mud diapirs. However, pseudo-diapiric anticlines and ridges have been tentatively recognized and mapped in the Po-Plain border of the Northern Apennines (Pini, 1987, 1993, 1999) and in the Southern Apennines.

*Mélanges*, therefore, preserve a complex deformational history due to the superposition of different and often linked processes. During this long history, diapirism can play an important role in re-shaping the *mélange* bodies, changing and complicating the contacts with the normal, well bedded formations and, basically, in exhuming the *mélanges* (see the case of Coscogno *mélange*).

We conclude that tectonic, sedimentary, and diapiric processes commonly work together with often an intimate, mutual interplay and linkage in the formation of *mélanges*. Yet, it is still possible to distinguish *mélanges* formed in different geodynamic environments and the main processes of their formation. The different case studies we have examined in the peri-Adriatic region suggest that a close relationship exists between different types of *mélanges* and their tectonic paleogeographic settings. Moreover, the complex evolutionary history of *mélanges*, generally involving the superposition of tectonic, sedimentary and diapiric processes and the reactivation of previously-formed *mélanges*, allow us to better understand the role of *mélanges* in the complex evolution of the peri-Adriatic orogens and accretionary wedges.

## **Acknowledgments**

This work derives from several years of field studies on chaotic rock bodies of the northern Apennines (G.A. Pini, C.C. Lucente and K. Ogata), southern Apennines (A. Festa, F. Ghisetti and L. Vezzani), Tertiary Piedmont Basin (A. Festa and G. Codegone) and Dinarides, Albanides and Hellenides (Y. Dilek). It also presents evidence and interpretations that have resulted from subsequent work and from new unpublished research. We are greatly indebted to A. Camerlenghi, (ICREA, Institutió Catalana de Recerca i Estudis Avançats, Barcelona, Spain), D.S. Cowan (Department of Earth and Space Sciences, University of Washington, Seattle, USA), A. Castellarin (Dipartimento di Scienze della Terra e Geologico-Ambientali, Università di Bologna), Annie Rassios (IGME-Kozani, Greece), and Alan G. Smith (University of Cambridge, U.K.) for fruitful discussions and for their support during the course of this research. Our melange studies were funded by MIUR (Ministero dell'Università e della Ricerca) grants to P. Clari (2005-043300) and G.A. Pini (2003-040755 and 2005-045211 grants) and Consiglio Nazionale delle Ricerche, Istituto di Geoscienze e Georisorse, Torino, and by research grants from the NATO science for Peace Programme, Institute of Geology and Mineral Exploration (IGME) of Greece, and Miami University to Y. Dilek.



## REFERENCES

- Abbate, E., Bortolotti, V., and Passerini, P., 1970, Olistostromes and olistoliths. In: Sestini, G., ed., Development of the Northern Apennines Geosyncline, *Sedimentary Geology*, v. 4, p. 521-557.
- Abbate, E., Bortolotti, V., and Sagri, M., 1981, Excursion No. 5: Olistostromes in the Oligocene Macigno formation (Florence area), Introduction: an approach to olistostrome interpretation. In: Ricci Lucchi, F., ed., International Association of Sedimentologist, Second European Regional Meeting, Bologna, 1981, Excursion Guidebook, Tecnoprint, Bologna, p. 165-185.
- Abbate, E., and Sagri, M., 1970, The eugeosynclinal sequence. In: Sestini, G., ed., Development of the Northern Apennines Geosyncline, *Sedimentary Geology*, v. 4, p. 521-557.
- Accordi, G., and Carbone, F., 1988, Sequenze carbonatiche meso-cenozoiche: Quaderni Ricerca Scientifica, v. 114/115, p. 11-92.
- Argnani, A., Barbacini, G., Bernini, M., Camurri, F., Ghielmi, M., Papani, G., Rizzini, F., Rogledi S., and Torelli, L., 2003, Gravity tectonics driven by Quaternary uplift in the Northern Apennines: insights from the La Spezia-Reggio Emilia geo-transect: *Quaternary International*, v. 101-102, p. 13-26.
- Argnani, A., Fontana, D., Stefani, C., and Zuffa, G.G., 2004, Late Cretaceous Carbonate Turbidites of the Northern Apennines: Shaking Adria at the Onset of Alpine Collision: *Journal of Geology*, v. 112, p. 251-259.
- Azzaroli, A., 1948, Osservazioni sulla tettonica dell'alta Val di Nievole e delle zone adiacenti della catena di monte Albano: *Bollettino della Società Geologica Italiana*, v. 67, p. 143-159.
- Bailey, E.B., and McCallien, W.J., 1950, The Ankara Mélange and the Anatolian Thrust: *Nature*, v. 166, p. 938-940.
- Bailey, E.B., and McCallien, W.J., 1953, Serpentine lavas, the Ankara mélange and the Anatolian thrust: *Transaction Royal Society Edinburgh*, v. 6. No. 2, p. 403-442.
- Barber, A.J., Tjokrosapoetro, S., and Charlton, T.R., 1986, Mud volcanoes, shale diapirs, wrench faults and mélanges in accretionary complexes, Eastern Indonesia: *American Association of Petroleum Geologists Bulletin*, v. 70, p. 1729-1741.
- Barchi, M., Landuzzi, A., Minelli, G., and Piali, G., 2001, In: Vai, G.B., and Martini, I.P., eds., Anatomy of a Mountain Belt: The Apennines and Adjacent Mediterranean Basins, Kluwer Academic Publishers, Dordrecht, p. 215-254.
- Bellinzona, G., Boni, A., Braga, G., and Marchietti G., 1971, Note Illustrative della Carta Geologica d'Italia alla scala 1:100.000, Foglio 71 "Voghera": Servizio Geologico d'Italia. Nuova Tecnica Grafica, Roma, pp. 121.
- Beneo, E., 1956, Accumuli terziari da risedimentazione (olistostroma) nell'Appennino centrale e frane sottomarine; estensione tempo-spaziale del fenomeno: *Bollettino del Servizio Geologico Italiano*, v. 78, p. 291-319.
- Bernoulli, D., 2001, Mesozoic-Tertiary carbonate platforms, slope and basins of the external Apennines and Sicily. In: Vai, G.B., and Martini, I.P., eds., Anatomy of an Orogen: the Apennines and Adjacent Mediterranean Basins, Kluwer Academic Publishers, Great Britain, p. 307-326.
- Bettelli, G., Bonazzi, U., Fazzini, P., and Panini, F., 1987, Schema introduttivo alla geologia delle Epiliguridi dell'Appennino Modenese e delle aree limitrofe. *Memorie della Società Geologica Italiana*, v. 39, p. 215-244.
- Bettelli, G., and Panini, F., 1985, Il mélange sedimentario della Val Tiepido (Appennino modenese)- composizione litologica, distribuzione areale e posizione stratigrafica: *Atti della Società dei Naturalisti e Matematici di Modena*, v. 115, p. 91-106.
- Bettelli, G., and Panini, F., 1987, I mélanges dell'Appennino settentrionale dal T. Tresinaro al T. Sillaro: *Memorie della Società Geologica Italiana*, v. 39, p. 187-214.
- Bettelli, G., and Panini, F., 1992, Liguridi, mélanges e tettoniti nel complesso caotico lungo la "linea del Sillaro" (Appennino Settentrionale, Province di Firenze e Bologna): *Memorie Descrittive della Carta Geologica d'Italia*, v. 46, p. 387-415.
- Bettelli, G., Panini, F. and Capitani, S., 2002, Carta Geologico-Strutturale dell'Appennino emiliano sudorientale. In: *Atti del terzo seminario sulla cartografia geologica*, Bologna, 2002, p. 47-52.
- Bettelli, G., and Vannucchi, P., 2003, Structural style of the offscraped Ligurian oceanic sequences of the Northern Apennines: new hypothesis concerning the development of melange block-in-matrix fabric: *Journal of Structural Geology*, v. 25, p. 371-388.
- Bianconi, G., 1840, Storia naturale dei terreni ardenti, dei vulcani fangosi, delle sorgenti infiammabili, dei pozzi idropirici e di altri fenomeni geologici operati dal gas idrogeno e della origine di esso gas. Marsigli, Bologna, pp. 164.

- Bigi, G., Cosentino, D., Parotto, M., Sartori, R., and Scandone, P., 1990, Structural Model of Italy, scale 1:500,000, Sheet 1, Geodynamic project, C.N.R., Ed. S.EL.CA. Firenze.
- Boccaletti, M., Calamita F., Deiana, G., Gelati, R., Massari F., Moratti, G., and Ricci Lucchi F., 1990, Migrating foredeep-thrust belt system in the northern Apennines and southern Alps: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 77, p. 3-14.
- Boccaletti, M., and Coli, M. (eds.), 1982, Carta Strutturale dell'Appennino Settentrionale. CNR Progetto Finalizzato Geodinamica, Publication No. 429-1982, 1:250.000 scale geological map, 4 sheets, Ed. S.EL.CA. Firenze.
- Boccaletti, M., Coli, M., Decandia, F.A., Giannini, E., and Lazzarotto, A., 1980, Evoluzione dell'Appennino settentrionale secondo un nuovo modello strutturale: Memorie della Società Geologica Italiana, v. 21, p. 359-373
- Bonardi, G., Amore, F.O., Ciampo, G., De Capoa, P., Miconnet, P., and Perrone, V., 1988, Il complesso ligure auct.: stato delle conoscenze e problemi aperti sulla sua evoluzione pre-appenninica ed i suoi rapporti con l'Arco Calabro: Memorie della Società Geologica Italiana, v. 41, p. 17-35.
- Boni, A., Casnedi, R., Anfossi, G., Bruno, G.M., Motta, E., Perotto, G., and Zezza, F., 1970, Note Illustrative della Carta Geologica d'Italia alla scala 1:100.000, Fogli 69 e 70 "Asti-Alessandria": Servizio Geologico d'Italia, Poligrafica e Cartevalori, Ercolano (Napoli), pp. 64.
- Bortolotti, V., Cortesogno, L., Gaggero, L., Lahondere, D., Marroni, M., Molli, G., Montanini, A., Pandolfi, L., Principi, G., Rossi, P., Saccano, E., Treves, B., and Tribuzio, R., 2004, Northern Apennine and Corsica ophiolites: the oceanic lithosphere of the ligure-piemontese basin and its transition to the Adria continental margin (Italy): Memorie descrittive della Carta Geologica d'Italia, v. 63, p. 27, 1-12.
- Bortolotti, V., Marroni, M., Pandolfi, L., and Principi, G., 2005, Mesozoic to Tertiary tectonic history of the Mirdita ophiolites, northern Albania: The Island Arc, v. 14, p. 471-493.
- Bortolotti, V., Principi, G., and Treves, B., 2001, Ophiolites, Ligurides and the tectonic evolution from spreading to convergence of a Mesozoic Western Tethys segment. In: Vai, G.B., and Martini, I.P., eds., Anatomy of an Orogen: the Apennines and Adjacent Mediterranean Basins, Kluwer Academic Publishers, Great Britain, p. 151-164.
- Bosellini, A., 1998, Scalloped versus faulted carbonate margins and the origin of basinal megabreccias: Memorie della Società Geologica Italiana, v. 53, p. 63-74.
- Bosellini, A., Castellarin, A., Rossi, P.L., Simboli, G., and Somnavilla, E., 1977, Schema sedimentologico e stratigrafico per il Trias medio della Val di Fassa ed aree circostanti (Dolomiti Centrali): Giornale di Geologia, v. 42, p. 83-108.
- Bosellini, A., Neri, C., and Luciani, V., 1993, Platform margin collapses and sequence stratigraphic organization of carbonate slopes: Cretaceous-Eocene, Gargano Promontory, Southern Italy: Terra Nova, v. 5, p. 282-297.
- Brandon, M.T., 1989, Deformational styles in a sequence of olistostromal mélanges, Pacific Rim Complex, western Vancouver Island, Canada: Geological Society of America Bulletin, v. 101, p. 1520-1542.
- Brown, K.M., 1990, The nature and hydrogeologic significance of mud diapirs and diatremes for accretionary systems: Journal of Geophysical Research, v. 95, p. 8969-8982.
- Camerlenghi, A., and Pini, G.A., 2009, Mud volcanoes, olistostromes and Argille scagliose in the Mediterranean region: Sedimentology, v. 56, p. 319-365.
- Capitani, M., 1993, Analisi macrostrutturale della fascia di deformazioni trasversali dei torrenti Lavino e Samoggia (Province di Bologna e Modena). Ph.D Thesis, Università di Bologna, pp. 104.
- Carbone, S., and Lentini, F., 1990, Migrazione neogenica del sistema catena-avampaese nell'Appennino meridionale: problematiche paleogeografiche e strutturali: Rivista Italiana di Paleontologia e Stratigrafia, v. 96, No. 2-3, p. 271-296.
- Carmignani, L., Decandia, F.A., Disperati, L., Fantozzi, P.L., Kligfield, R., Lazzarotto, A., Lotta, D., and Meccheri, M., 2001, Inner Northern Apennines. In: Vai, G.B., and Martini, I.P., eds., Anatomy of an Orogen: the Apennines and Adjacent Mediterranean Basins, Kluwer Academic Publishers, Great Britain, p. 197-214.
- Casey, J., and Dewey, J., 1984, Initiation of subduction zones along transform and accreting plate boundaries, triple-junction evolution, and forearc spreading centres; implications for ophiolitic geology and obduction. In: Gass, I., Lippard, S., and Shelton, A., eds., Ophiolites and Oceanic Lithosphere, Geological Society, London, Special Publication, v. 13, p. 269-290.
- Castellarin, A., 1972, Evoluzione paleotettonica sin-sedimentaria del limite tra "piattaforma veneta" e "bacino lombardo", a Nord di Riva del Garda: Giornale di Geologia, v. 2, No. 38, p. 11-212.

- Castellarin, A., 1982, Tettonica sinsedimentaria e associazioni di facies carbonatiche (con principali riferimenti al Giurassico sudalpino). In: Castellarin, A., and Vai, G.B., eds., Guida alla Geologia del Sudalpino Centro-Orientale, SGI-Pitagora Tecnoprint, Bologna, p. 23-40.
- Castellarin, A., 1994, Strutturazione eo-mesoalpina dell'Appennino Settentrionale attorno al "nodo ligure". In: Capozzi, R., and Castellarin, A., eds., Studi preliminari all'acquisizione dati del profilo CROP 1-1A La Spezia-Alpi orientali, Studi Geologici Camerti, Volume Speciale 1992/2, p. 99-108.
- Castellarin, A., Cantelli, L., Fesce, A.M., Mercier, J.L., Picotti, V., Pini, G.A., Prosser, G., and Selli, L., 1992, Alpine compressional tectonics in the Southern Alps: relationships with the N-Apennines: *Annales Tectonicae*, v. 4, p. 62-94.
- Castellarin, A., Colacicchi, R., and Praturlon, A., 1978, Fasi distensive, trascorrenze e sovraccorrimenti lungo la "linea Ancona-Anzio", dal Lias medio al Pliocene: *Geologica Romana*, v. 17, p. 161-189.
- Castellarin, A., Eva, C., and Capozzi, R., 1994, Tomografie sismiche e interpretazione geologica profonda dell'Appennino Settentrionale Nord-occidentale: Studi Geologici Camerti, Volume Speciale 1992/2, p. 85-98.
- Castellarin, A., and Pini, G.A., 1987, L'arco del Sillaro: la messa in posto delle Argille Scagliose al margine appenninico padano (Appennino bolognese): *Memorie della Società Geologica Italiana*, v. 39, p. 127-142.
- Castellarin, A., Pini, G.A., Crestana, G., and Rabbi, E., 1986, Caratteri strutturali mesoscopici delle Argille Scagliose dell'Appennino bolognese: *Memorie di Scienze Geologiche*, v. 38, p. 459-477.
- Castellarin, A., Selli, L., Picotti, V., and Cantelli, L., 1998, Tettonismo e diapirismo medio Triassico delle Dolomiti: *Memorie della Società Geologica Italiana*, v. 53, p. 145-169.
- Catalano S., Grasso G., Mazzoleni P., Monaco C., and Tortrici L., 2007, Intracontinental tectonic mélange in Southern Apennines: *Terra Nova*, v. 19, p. 287-293.
- Catanzariti, R., Ellero, A., Levi, N., Ottria, G., and Pandolfi, L., 2007, Calcareous nannofossil biostratigraphy of the Antola Unit succession (Northern Apennines, Italy): new age constraints for the Late Cretaceous Helminthoid Flysch: *Cretaceous Research*, v. 28, p. 841-860.
- Cecca, F., Cresta, S., Giovagnoli, M.C., Manni, R., Mariotti, N., Nicosia, U., and Santantonio, M., 1981, Tithonian "Ammonitico Rosso" near Bolognola (Marche-Central Apennines): a shallow nodular limestone. In: Farinacci, A., and Elmi, S., eds., Rosso Ammonitico Symposium Proceedings, Edizioni Tecnoscienza-Roma, p. 91-112.
- Cerrina Feroni, A., Ottria, G., Martinelli, P., and Martelli, L., 2002, Structural Geological Map of the Emilia-Romagna Apennines, 1:250,000 Scale, Ed. S.EL.CA. Firenze.
- Chang, C.-P., Angelier, J., and Huang, C.-Y., 2000, Origin and evolution of a mélange: the active plate boundary and suture zone of the Longitudinal Valley, Taiwan: *Tectonophysics*, v. 325, p. 43-62.
- Choconat, P., Cadet, J.P., Lallemand, S.J., Mazzotti, S., Nouzé, H., Fouchet, C., and Foucher, J.P., 2002, Slope instabilities and gravity processes in fluid migration and tectonically active environment in the eastern Nankai accretionary wedge (KAIKO-Tokai'96 cruise): *Marine Geology*, v. 187, p. 193-202.
- Cibin, U., Spadafora, E., Zuffa, G.G., and Castellarin, A., 2001, Continental collision history from arenites of episutural basins in the Northern Apennines, Italy: *Geological Society of America Bulletin*, v. 113, p. 4-19.
- Cipollari, P., and Cosentino, D., 1995, Miocene unconformities in the central Apennines: geodynamic significance and sedimentary basin evolution: *Tectonophysics*, v. 252, p. 375-389.
- Collot, J.-Y., Lewis, K., Lamarche, G., and Lallemand, S., 2001, The giant Ruatoria debris avalanche on the northern Hikurangi margin, New Zealand: result of oblique seamount subduction: *Journal of Geophysical Research*, v. 106, p. 19271-19297.
- Conti, S., 1987, Studio geologico della geotraversa Grosseto-Val Marecchia tra le valli del Savio e del Foglia (Appennino romagnolo-marchigiano). Ph.D Thesis, University of Bologna, pp. 150.
- Cornamusini, G., 2001, The early depositional phases of the northern Apennine foredeep-thrust belt system: implications from the "Macigno costiero" (Late Oligocene, Italy): *Ofioliti*, v. 26, p. 263-274.
- Cosentino, D., Cipollari, P., Di Donato, V., Sgrosso, I., and Sgrosso, M., 2002, The Volsci Range in the kinematic evolution of the northern and southern Apennine orogenic system: *Bollettino della Società Geologica Italiana*, Volume Speciale No. 1, p. 209-218.
- Cousineau, P.A., 1998, Large-scale liquefaction and fluidization in the Cap Chat Mélange, Quebec Appalachians: *Canadian Journal of Earth Sciences*, v. 35, p. 1408-1422.
- Cowan, D.S., 1985, Structural styles in Mesozoic and Cenozoic melanges in the western Cordillera of North America: *Geological Society of America Bulletin*, v. 96, p. 451-462.

- Cowan, D.S., and Pini, G.A., 2001, Disrupted and chaotic rock units in the Apennines. In: G.B. Vai and I.P. Martini, eds., *Anatomy of a Mountain Belt: The Apennines and Adjacent Mediterranean Basins*, Kluwer Academic Publishers, Dordrecht, p. 165-176.
- Daniele, G., Plesi, G., 2000, The Ligurian Helminthoid Flysch units of the Emilian Apennines: stratigraphic and petrographic features, palaeogeographic restoration and structural evolution: *Geodinamica Acta*, v. 13, p. 313-333.
- De Alteris, G., and Aiello, G., 1993, Stratigraphy and tectonics offshore of Puglia (Italy, Southern Adriatic sea): *Marine Geology*, v. 113, p. 197-212.
- Dela Pierre, F., Clari, P., Cavagna, S., and Bicchi, E., 2002, The Parona chaotic complex: a puzzling record of the Messinian (late Miocene) events in Monferrato (NW Italy): *Sedimentary Geology*, v. 152, p. 289-311.
- Dela Pierre, F., Festa, A., and Irace, A., 2007, Interaction of tectonic, sedimentary and diapiric processes in the origin of chaotic sediments: an example from the Messinian of the Torino Hill (Tertiary Piedmont Basin, NW Italy): *Geological Society of America Bulletin*, v. 119, p. 1107-1119.
- Dela Pierre, F., Piana, F., Fioraso, G., Boano, P., Bicchi, E., Forno, M.G., Violanti, D., Clari, P., and Polino, R., 2003, Note illustrative della Carta Geologica d'Italia alla scala 1:50,000. Foglio 157 Trino: APAT, Dipartimento Difesa del Suolo, pp. 147.
- De Libero, C.M., 1998, Sedimentary vs. tectonic deformation in the "Argille Scagliose" of Mt. Modino (Northern Apennines): *Giornale di Geologia*, v. 56, p. 143-166.
- Dercourt, J., Zonenshain, L.P., Ricou, L.E., Kazmin, V.G., Le Pichon, X., Knipper, A.L., Granjacquet, C., Sbertshikov, I.M., Geyssant, J., Lepvrier, C., Pechersky, D.H., Boulin, J., Sibuet, J.C., Savostin, L.A., Sorokhtin, O., Westphal, M., Bazhenov, M.L., Lauer, J.P., and Biju-Duval, B., 1986, Geological evolution of the Tethys belt from the Atlantic to the Pamirs since the Lias: *Tectonophysics*, v. 123, p. 241-315.
- Dewey, J.F., Helman, M.L., Turco, E., Hutton, D.W.H., and Knott, S.D., 1989, Kinematics of the western Mediterranean. In: Coward, M.P., Dietrich, D., and Park, R.G., eds., *Alpine Tectonics*, Geological Society London, Special Publication, v. 45, p. 265-283.
- Di Giulio, A., 1992, The evolution of the western Ligurian Flysch Units and the role of mud diapirism in ancient accretionary prisms (Maritime Alps, Northwestern Italy): *Geologische Rundschau*, v. 81. No. 3, p. 655-668.
- Dilek, Y., 2006, Collisional tectonics of the Mediterranean region: Causes and consequences. In: Dilek, Y., and Pavlides, S., eds., *Postcollisional Tectonics and Magmatism in the Mediterranean Region and Asia*, Geological Society of America Special Paper, v. 409, p. 1-13.
- Dilek, Y., Furnes, H., and Shallo, M., 2007, Suprasubduction zone ophiolite formation along the periphery of Mesozoic Gondwana: *Gondwana Research*, v. 11, p. 453-475.
- Dilek, Y., Furnes, H., and Shallo, M., 2008, Geochemistry of the Jurassic Mirdita Ophiolite (Albania) and the MORB to SSZ evolution of a marginal basin oceanic crust: *Lithos*, v. 100, p. 174-209.
- Dilek, Y., and Koçiu, S., 2004, Neotectonics, seismology and the mode of Late Cenozoic crustal deformation in the Albanides, Balcan Peninsula: *Geological Society of America Abstracts with programs*, v. 36, No. 5, p. 52-53.
- Dilek, Y., Shallo, M., and Furnes, H., 2005, Rift-drift, seafloor spreading, and subduction tectonics of Albanian ophiolites: *International Geology Review*, v. 47, p. 147-176.
- Dumurdzanov, N., Serafimovski, T., and Burchfiel, B.D., 2005, Cenozoic tectonics of Macedonia and its relation to the South Balkan extensional regime: *Geosphere*, v. 1, No. 1, p. 1-22.
- Eberli, G.P., 1987, Carbonate turbidite sequences deposited in rift-basins of the Jurassic Tethys Ocean (eastern Alps, Switzerland): *Sedimentology*, v. 34, p. 363-388.
- Eberli, G.P., Bernoulli, D., Sanders, D.G.K., and Vecsei, A., 1993, From aggradation to progradation: the Maiella platform (Abruzzi, Italy). In: Simo, A.J., Scott, R.W., and Masse, J.P., eds., *Cretaceous Carbonate Platforms*, American Association of Petroleum Geologists Memories, v. 56, p. 213-232.
- Elter, P., 1960, I lineamenti tettonici dell'Appennino a Nord Ovest delle Apuane: *Bollettino della Società Geologica Italiana*, v. 79, p. 273-308.
- Elter, P., 1975, L'ensemble ligure: *Bulletin de la Société Géologique de France*, v. 17, p. 984-997.
- Elter, P., and Trevisan, L., 1973, Olistostromes in the tectonic evolution of the Northern Apennines. In: De Jong, K.A., and Scholten, R., eds., *Gravity and Tectonics*, John Wiley and Sons, New York, p. 175-188.

- Fazzuoli, M., Ferrini, G., Pandeli, E., and Sguazzoni, G., 1985, Le formazioni giurassico-mioceniche della falda toscana a nord dell'Arno: Considerazioni sull'evoluzione sedimentaria: Memorie della Società Geologica Italiana, v. 30, p. 159-201.
- Festa, A., 2009, Tectonic, sedimentary, and diapiric formation of the Messinian melange: Tertiary Piedmont Basin (northwestern Italy). In: Wakabayashi, J., and Dilek, Y., eds., *Melanges: Processes of formation and societal significance*, Geological Society of America Special Paper, in press.
- Festa, A., Dela Pierre, F., and Irace, A., 2005, Role of tectonic deformation in the genesis of the Messinian chaotic sediments of the Northern – Tertiary Piemonte Basin (Torino Hill and Monferrato unit): *Rendiconti della Società Geologica Italiana, Nuova Serie*, v. 1, p. 105-106.
- Festa, A., Dela Pierre, F., Piana, F., Fioraso, G., Lucchesi, S., Boano, P., and Forno M.G., 2009, Note Illustrative della Carta Geologica d'Italia alla scala 1:50.000. Foglio 156 Torino Est, APAT, Dipartimento Difesa del Suolo, pp. 138, in press.
- Festa, A., Ghisetti, F., and Vezzani, L., 2006, Carta Geologica del Molise (Scala 1:100.000) - Note Illustrative. Litografia Geda - Nichelino (TO), pp. 104. ISBN 88-902635-0-4. with CD-Rom.
- Flores, G., 1955, Discussion in: Beneo, E., 1955, *Les résultats des études pour la recherche pétrolifère en Sicilie*. Proceedings, Fourth World Petroleum Congress, Rome, Section 1/A/2, p. 121-122.
- Flores, G., 1956, The results of the studies on petroleum exploration in Sicily: *Discussion, Bollettino del Servizio Geologico d'Italia*, v. 78, p. 46-47.
- Galluzzo, F., and Santantonio, M., 2002, The Sabina Plateau: a new element in the Mesozoic palaeogeography of Central Apennines: *Bollettino della Società Geologica Italiana, Volume Speciale No. 1*, p. 561-588.
- Ghikas, A.C., 2007, Structure and tectonics of a subophiolitic mélange (Zavordas mélange) of the Vourinos ophiolite (Greece) and kinematics of ophiolite emplacement. Unpublished Msc Thesis, Miami University, Oxford, OH.
- Ghikas A.C., Dilek, Y., and Rassios, A.E., 2009, Structure and Tectonics of Sub-ophiolitic Mélanges in the Western Hellenides (Greece) and Implications for Ophiolite Emplacement Tectonics: *International Geology Review*, this issue.
- Ghisetti, F., and Vezzani L., 1986, Assetto geometrico ed evoluzione strutturale della catena del Gran Sasso tra Vado di Siella e Vado di Corno: *Bollettino della Società Geologica d'Italia*, v. 105, p. 131-171.
- Ghisetti, F., and Vezzani, L., 1998, Segmentation and tectonic evolution of the Abruzzi-Molise thrust belt (central Apennines, Italy): *Annales Tectonicae*, v. 12, p. 97-112.
- Ghisetti, F., Vezzani L., and Festa, A., 2003, The exhumed Miocene-Pliocene accretionary wedge of the central Apennines (Italy). The new 1:100.000 geological map of Molise: Geological Society of New Zealand Annual General Meeting, Abstract, 90, Dunedin (NZ), 1-4 December 2003.
- Görler, K., and Reutter, K.J., 1968, Entstehung und Merkmale der Olistostrome: *Geologische Rundschau*, v. 57, p. 484-514.
- Graziano, R., 2000, The Aptian-Albian of the Apulia carbonate platform (Gargano Promontory, southern Italy): evidence of palaeoceanographic and tectonic controls on the stratigraphic architecture of the platform margin: *Cretaceous Research*, v. 21, p. 107-126.
- Greenly, E., 1919, *The Geology of Anglesey*, Vols I and II: *Memoirs of Geological Survey*. HM Stationary Office, London, pp. 1-388, 390-980.
- Horton, J.W. Jr. and Rast, N. (eds.), 1989, Mélanges and Olistostromes of the U.S. Appalachians: Geological Society of America Special Paper, v. 228, p. 275.
- Hsü, K.J., 1965, Franciscan rocks of Santa Lucia Range, California, and the "argille scagliose" of the Apennines, Italy: a comparison in style of deformation: *Geological Society of America Abstracts for 1965*, p. 210-211.
- Hsü, K.J., 1968, Principles of melanges and their bearing on the Franciscan-Knoxville paradox: *Geological Society of America Bulletin*, v. 79, p. 1063-1074.
- Hsü, K.J., 1973, Mesozoic evolution of California Coast ranges; a second look. In: DeJong, K.A., and Scholten, R., eds., *Gravity and Tectonics*, John Wiley and Sons, New York, p. 379-396.
- Hsü, K.J., 1974, Melanges and their distinction from olistostromes. In: Dott Jr., R.H., and Shaver, R.H., eds., *Modern and Ancient Geosynclinal Sedimentation*, SEPM, Special Publication, v. 19, p. 321-333.
- Hühnerbach, V., Masson, D.G., Bohrmann, G., Bul, J.M., and Weinrebe, W., 2005, Deformation and submarine landsliding caused by seamount subduction beneath the Costa Rica continental margin - new insights from high resolution sidescan sonar data. In: Hodgson, D.M., and Flint, S.S., eds., *Submarine Slope*

- Systems: Processes and Products, Geological Society of London, Special Publication, v. 244, p. 195-205.
- Ikesawa, E., Kimura, G., Sato, K., Ikehara-Ohmori, K., Kitamura, Y., Yamaguchi, A., Ejiie, K., and Hashimoto, Y., 2005, Tectonic incorporation of the upper part of oceanic crust to overriding plate of a convergent margin: an example from the Cretaceous-early Tertiary Mugi Melange, the Shimanto Belt, Japan: *Tectonophysics*, v. 401, p. 217-230.
- Irace, A., 2004, *Il Messiniano Piemontese: nuovi dati da due aree campione*. PhD Thesis. Torino, Italy, University of Torino, pp. 167.
- Irace, A., Dela Pierre, F., and Clari, P., 2005, "Normal" and "chaotic" deposits in the Messinian Gessoso Solfifera Fm. at the north-eastern border of the Langhe domain (Tertiary Piedmont Basin): *Bollettino della Società Geologica Italiana, Volume Speciale No. 4*, p. 77-85.
- Jacobacci, A., 1963, Frane sottomarine nelle formazioni geologiche. Interpretazione dei fenomeni olistostromici e degli olistoliti nell'Appennino ed in Sicilia: *Bollettino del Servizio Geologico Italiano*, v. 86, p. 65-85.
- Jones, G., and Robertson, A.H.F., 1991, Tectono-stratigraphy and evolution of the Mesozoic Pindos Ophiolite and related units, Northwestern Greece: *Journal of the Geological Society of London*, v. 148, p. 267-288.
- Jurewicz, E., Giren, B., and Steller, J., 2007, Cavitation erosion - a possible cause of the mass loss within thrust zones in the Tatra Mts., Poland: *Acta Geologica Polonica*, v. 57, p. 305-323.
- Knot, S.D., 1994, Structure, kinematics and metamorphism in the Liguride Complex, southern Apennines, Italy: *Journal of Structural Geology*, v. 16, p. 1107-1120.
- Kostopoulos, D., 1988, *Geochemistry, petrogenesis, and tectonic setting of the Pindos Ophiolite, NW Greece*. PhD Thesis, University of Newcastle upon Tyne.
- Krylov, K.A., Kaleda, K.G., and Bragin N.Yu., 2005, Chaotic Rock Complexes in South-Western and Central Cyprus. In: Krasheninnikov, V.A., Hall, J.K., Hirsch, F., Beniamini, C., and Flexer, A., eds., *Geological Framework of the Levant, Volume I: Cyprus and Syria*, p. 127-134, Historical Productions-Hall, Jerusalem.
- Landuzzi, A., 2004, Sin-depositional advancement of the Liguride allochthon in the Miocene foredeep of the Western Romagna Apennines (Italy). In: Pasquaré, G., and Venturini, C., eds., *Mapping Geology in Italy: Case Studies from Different Crustal Levels*, Ed. S.EL.CA. Firenze.
- Laurita, S., Cavalcante, F., Belviso, C., and Prosser G., 2009, The Liguride Complex of the Pollino area (Southern Apennines): tectonic setting and preliminary mineralogical data: *Rendiconti online della Società Geologica Italiana*, v. 5, p. 111-114.
- Levi, N., Ellero, A., Ottria, G., and Pandolfi, L., 2006, Polyorogenic deformation history recognized at very shallow structural levels: the case of the Antola Unit (Northern Apennine, Italy): *Journal of Structural Geology*, v. 28, p. 1694-1709.
- Lucente, C.C., and Pini, G.A., 2003, Anatomy and emplacement mechanism of a large submarine slide within the Miocene foredeep in the Northern Apennines, Italy: a field perspective: *American Journal of Science*, v. 303, p. 565-602.
- Lucente, C.C., and Pini, G.A., 2008, Basin-wide mass-wasting complexes as markers of the Oligo-Miocene foredeep-accretionary wedge evolution in the Northern Apennines, Italy: *Basin Research*, v. 20, p. 49-71, DOI: 10.1111/j.1365-2117.2007.00344.x.
- Lucente, C.C., Pini, G.A., and De Libero, C.M., 2006, The role of large scale sedimentary failures at the wedge front-foredeep boundary: a case study from the northern Apennines, Italy: *GeoActa*, v. 5, p. 113-128.
- Maltman, A., 1994, Introduction and overview, in: Maltman, A., ed., *The geological deformation of sediments*, London, Chapman & Hall, p. 1-35.
- Marchegiani, L., Bertotti, G., Cello, G., Deiana, G., Mazzoli, S., and Tondi, E., 1999, Pre-orogenic tectonics in the Umbria-Marche sector of the Afro-Adriatic continental margin: *Tectonophysics*, v. 315, p. 123-143.
- Marroni, M., Cerrina Feroni, A., di Biase, D., Ottria, G., Pandolfi, L., and Taini, A., 2002, Polyphase folding at upper structural levels in the Borbera valley (Northern Apennines, Italy): implications for the tectonic evolution of the linkage area between Alps and Apennines: *Comptes Rendus de l'Academie des Sciences Geosciences*, v. 334, p. 565-572.
- Marroni M., Molli G., Montanini A., and Tribuzio R., 1998, The association of continental crust rocks with ophiolites (northern Apennine, Italy): implications for the continent-ocean transition: *Tectonophysics*, v. 292, p. 43-66.

- Marroni, M., Molli, G., Ottria, G., and Pandolfi, L., 2001, Tectono-sedimentary evolution of the External Liguride units (Northern Apennines, Italy): insight in the pre-collisional history of a fossil ocean-continent transition zone: *Geodinamica Acta*, v. 14, p. 307-320.
- Marroni, M., and Pandolfi, L., 2001, Debris flow and slide deposit at the top of the Internal Liguride ophiolitic sequence, Northern Apennines, Italy: a record of frontal tectonic erosion in a fossil accretionary wedge: *The Island Arc*, v. 10, p. 9-21.
- Marroni, M., and Treves, B., 1998, Hidden terranes in the northern Apennines, Italy: a record of Late Cretaceous-Oligocene transpressional tectonics: *Journal of Geology*, v. 106, p. 149-162.
- Maxwell, J.C., 1959, Turbidite, tectonic and gravity transport, Northern Apennine Mountains, Italy: *American Association of Petroleum Geologists Bulletin*, v. 43, p. 2701-2719.
- Mazzoli, S., and Helman, M., 1994, Neogene patterns of relative plate motions for Africa-Europe: some implications for recent central Mediterranean tectonics: *Geologische Rundschau*, v. 83, p. 464-468.
- Merla, G., 1952, *Geologia dell'Appennino settentrionale*: *Bollettino della Società Geologica Italiana*, v. 70, p. 95-382.
- Meschede, M., Zweigel, P., Frisch, W., and Völker, D., 1999, Mélange formation by subduction erosion: the case of the Osa melange in southern Costa Rica: *Terra Nova*, v. 11, p. 141-148.
- Montanari, A., Chan, L.S., and Alvarez, W., 1989, Synsedimentary tectonics in the Late Cretaceous–Early Tertiary pelagic basin of the Northern Apennines, Italy. In: Crevello, P.D., Wilson, J.L., Sarg, J.F., and Read, J.F., eds., *Controls on Carbonate Platform and Basin Development*, Society of Economic Paleontologists and Mineralogists, Special Publication, v. 44, p. 379-399.
- Naylor, M.A., 1981, Debris flows (olistostromes) and slumping on a distal passive continental margin: the Palombini limestone-shale sequence of the northern Apennines: *Sedimentology*, v. 28, p. 837-852.
- Naylor, M.A., and Harle, T., 1976, Palaeogeographic significance of rocks and structures beneath the Vourinos ophiolite, northern Greece: *Journal of the Geological Society, London*, v. 132, p. 667-676.
- Ogniben, L., 1953, “Argille scagliose” ed “argille brecciate” in Sicilia: *Bollettino del Servizio Geologico Italiano*, v. 75, p. 279-289.
- Ogniben, L., 1969, Schema introduttivo alla geologia del confine calabro-lucano: *Memorie della Società Geologica Italiana*, v. 8, p. 453-763.
- Orange, D.L., and Underwood, M.B., 1995, Patterns of thermal maturity as diagnostic criteria for interpretation of mélanges: *Geology*, v. 23, No. 1, p. 1144-1148.
- Ori, G.G., and Friend, P.F., 1984, Sedimentary basins formed and carried piggy-back on active thrust sheets: *Geology*, v. 12, p. 475-478.
- Patacca, E., and Scandone, P., 2007, *Geology of the Southern Apennines*: *Bollettino della Società Geologica Italiana, Volume Speciale No. 7*, p. 75-119.
- Patacca, E., Scandone, P., Bellatalla, M., Perilli, N., and Santini, U., 1992, La zona di giunzione tra l'arco appenninico settentrionale e l'arco appenninico meridionale nell'Abruzzo e nel Molise. In: Tozzi, M., Cavinato, G.P., and Parotto, M., eds., *Studi preliminari all'acquisizione dati del profilo CROP 11 Civitavecchia-Vasto, AGIP-CNR-ENEL, Studi Geologici Camerti, Volume Speciale 1991-2*, p. 417-441.
- Pe-Piper, G., and Piper, D.J.W., 2002, The igneous rocks of Greece, the anatomy of an orogen. *Beiträge zur Regionalen Geologie der Erde, Gebrüder Borntraeger, Berlin-Stuttgart, Germany*, pp. 573.
- Pescatore, T., Di Nocera S., Matano, F., and Pinto, F., 2000, L'unità del Fortore nel quadro della geologia del settore orientale dei Monti del Sannio (Appennino meridionale): *Bollettino della Società Geologica Italiana*, v. 119, p. 587-601.
- Piana, F., and Polino, R., 1995, Tertiary structural relationships between Alps and Apennines: the critical Torino Hill and Monferrato area, Northwestern Italy: *Terra Nova*, v. 7, p. 138-143.
- Pini, G.A., 1987, *Studio geologico e analisi strutturale delle “argille scagliose” del Pedepennino bolognese*. PhD Thesis, University of Bologna, Bologna, pp. 247.
- Pini, G.A., 1993, Geological map of the Bologna area foothills, scale 1:50000. CTF, Grafiche STEP, Parma.
- Pini, G.A., 1999, Tectonosomes and olistostromes in the Argille Scagliose of the Northern Apennines, Italy: *Geological Society of America Special Paper*, v. 335, p. 73.
- Pini, G.A., Lucente, C.C., Cowan, D.S., De Libero, C.M., Dellisanti, F., Landuzzi, A., Negri, A., Tateo, F., Del Castello, M., Morrone, M., and Cantelli, L., 2004, The role of olistostromes and argille scagliose in the structural evolution of the Northern Apennines. In: Guerrieri, L., Rischia, I., and Serva, L., eds., *Field Trip Guidebooks, 32nd IGC Florence 20–28 August 2004. Memorie Descrittive della Carta Geologica d'Italia*, 63, B13, p. 1-40.

- Ranero, C.R., and von Heune, R., 2002, Subduction erosion along the Middle America convergent margin: *Nature*, v. 404, p. 748-752.
- Rassios, A.E., and Dilek, Y., 2009, Rotational deformation in the Jurassic Mesohellenic ophiolites, Greece, and its tectonic significance: *Lithos*, v. 108, p. 207-223.
- Rassios, A.E., and Moores, E., 2006, Heterogeneous mantle complex, crustal processes, and Obduction kinematics in a unified Pindos-Vourinos ophiolitic slab (northern Greece). In: Robertson, A., and Mountrakis, D., eds., *Tectonic Development of The Eastern Mediterranean Region*, Geological Society, London, Special Publications, v. 260, p. 237-266.
- Raymond, L.A. (ed.), 1984, *Melanges: their nature, origin and significance*. Geological Society of America Special Paper, v. 198, p. 170.
- Remitti, F., Vannucchi, P., and Bettelli, G., 2007, Internal structure and tectonic evolution of an underthrust tectonic mélange: the Sestola-Vidiciatico tectonic unit of the Northern Apennines, Italy: *Geodinamica Acta*, v. 20/1-2, p. 37-51.
- Ricci Lucchi, F., 1975, Miocene paleobiogeography and basin analysis in the Periadriatic Apennines. In: Squyres, C., ed., *Geology of Italy, Earth Science Society Lybian Arabian Republic*, PESL Editor, Castelfranco, p.129-236.
- Ricci Lucchi, F., 1986, The Oligocene to Recent foreland basin of the Northern Apennines. In: Allen, P.A., and Homewood, P., eds., *Foreland Basins*, International Association of Sedimentologists, Special Publication, v. 8, p. 105-139.
- Ricci Lucchi, F. and Vai, G.B., 1994, A stratigraphic and tectonofacies framework of the “calcari a Lucina” in the Apennine Chain, Italy: *Geo-Marine Letters*, v. 14, p. 210-218.
- Rigo de Righi, M., 1956, Olistostromi neogenici in Sicilia: *Bollettino della Società Geologica Italiana*, v. 75, p. 185-215.
- Robertson, A.H.F., 1977, The Moni Melange, Cyprus; an olistostrome formed at a destructive plate margin: *Journal of Geological Society of London*, v. 133, p. 447-466.
- Roure, F., Casero, P., and Vially, R., 1991, Growth processes and melange formation in the Southern Apennines accretionary wedge: *Earth and Planetary Science Letters*, v. 102, p. 395-412.
- Roveri, M., Ricci Lucchi, F., Lucente, C.C., Manzi, V., and Mutti, E., 2002, Stratigraphy, facies and basin fill history of the Marnoso-Arenacea Formation. In: Mutti, E., Ricci Lucchi, F., and Roveri, M., eds., *Turbidite Workshop, 64th EAGE Conference, Excursion Guidebook, III/1-15*. Copy & Press, Parma.
- Rusciadelli, G., 2005, The Maiella escarpment (Apulia platform, Italy): geology and modeling of the Upper Cretaceous scalloped erosional platform margin: *Bollettino della Società Geologica Italiana*, v. 124, p. 661-673.
- Schettino, A., and C.R. Scotese, 2002, Global kinematic constraints to the tectonic history of the Mediterranean region and surrounding areas during the Jurassic and Cretaceous. In: Rosenbaum, G., and Lister, G.S., eds., *Reconstruction of the evolution of the Alpine-Himalayan orogen*, *Journal of the Virtual Explorer*, v. 7, p. 147-166.
- Selli, R., 1962, Il Paleogene nel quadro della geologia dell'Italia centro- meridionale: *Memorie della Società Geologica Italiana*, v. 3, p. 737-789.
- Sgrosso, I., 1988, Nuovi elementi per un più articolato modello paleogeografico nell'Appennino centro-meridionale: *Memorie della Società Geologica Italiana*, v. 41, p. 225-242.
- Shallo, M., 1990, Ophiolitic mélange and flyschoidal sediments of the Tithonian–Lower Cretaceous in Albania: *Terra Nova*, v. 2, p. 476-483.
- Sharp, I., and Robertson, A., 2006, Tectonic-sedimentary evolution of the western margin of the Mesozoic Vardar Ocean: evidence from the Pelagonian and Almopias Zones, northern Greece. In: Robertson, A., and Mountrakis, D., eds., *Tectonic Development of The Eastern Mediterranean Region*, Geological Society, London, Special Publication, v. 260, p. 373-412.
- Signorini, R., 1946, Autoctonia e alloctonia dei terreni dell'Appennino centrale e settentrionale: *Atti dell'Accademia Nazionale dei Lincei*, v. 8, No. 1, p. 99-106.
- Signorini, R., 1956, Tipi strutturali di scendimento e Argille Scagliose: *Bollettino della Società Geologica Italiana*, v. 75, p. 69-89.
- Smith, A.G., and Rassios, A.E., 2003, The evolution of idea for the origin and emplacement of the western Hellenic ophiolites. In: Dilek, Y., and Newcomb, S., eds., *Ophiolite Concept and the Evolution of Geological Thought: Geological Society of America Special Paper*, v. 373, p. 337-350.



- Smith, A.G., Woodcock, N.H., and Naylor, M.A., 1979, The structural evolution of a Mesozoic margin, Othris Mountains, Greece: *Journal of the Geological Society, London*, v. 136, p. 589-603.
- Spadea, P., 1982, Continental rocks associated with ophiolites in Lucanian Apennines (southern Italy): *Ophioliti*, v. 2, No. 3, p. 501-522.
- Swarbick, R.E., and Naylor, M.A., 1980, The Kathikas mélanges, SW Cyprus: late Cretaceous submarine debris flows: *Sedimentology*, v. 27, p. 63-78.
- Trümpy, R., 2006, Geologie der Iberger Klippen und ihrer Flysch-Unterlage: *Eclogae Geologicae Helvetiae*, v. 99, p. 79-121.
- Vai, G.B., and Castellarin, A., 1993, Correlazione sinottica delle unità stratigrafiche nell'Appennino Settentrionale. In: Capozzi, R., and Castellarin, A., eds., *Studi preliminari all'acquisizione dati del profilo CROP 1-1a La Spezia-Alpi orientali*, Studi Geologici Camerti, Volume Speciale No. 2, p. 171-185.
- Vannucchi, P., and Bettelli, G., 2002, Mechanism of subduction accretion as implied from the broken formations in the Apennines, Italy: *Geology*, v. 30, p. 835-838.
- Vannucchi, P., Remitti, F., and Bettelli, G., 2008, Geological record of fluid flow and seismogenesis along an erosive subducting plate boundary: *Nature*, v. 451, No. 7, p. 699-703
- Vecsei, A., 1991, Aggradation and progradation eines karbonatplattform- randes: kreide mis mittleres tertiar der montagna della Maiella, Abruzen. PhD Thesis, Mitteilungen des Geologischen Institutes der Eidgenossischen Technischen Hochschule und der Universität, Zurich, 294, pp. 171.
- Vecsei, A., 1998, Bioclastic sediment lobes on a supply dominated Upper Cretaceous carbonate platform margin, Montagna della Maiella, Italy: *Sedimentology*, v. 45, p. 473-487.
- Vescovi, P., Fornaciari, E., Rio, D., and Valloni, R., 1999, The Basal Complex Stratigraphy of the Helmintoid Monte Cassio Flysch: a key to the Eoalpine tectonics of the Northern Apennines: *Rivista Italiana di Paleontologia e Stratigrafia*, v. 105, No. 1, p. 101-128.
- Vezzani, L., Ghisetti, F., and Festa, A., 2004, Carta geologica del Molise (scala 1:100.000), Ed. S.EL.CA. Firenze.
- Vezzani, L., Ghisetti, F., and Festa, A., 2009, Geological-structural map of the central-southern Apennines (Italy) (1:250,000 scale), 2 sheets, Ed. S.EL.CA. Firenze.
- Von Heune, R., Ranero, C.R., and Watts, P., 2003, Tsunamigenic slope failure along the Middle America Trench in two tectonic setting: *Marine Geology*, v. 203, p. 303-327.
- Wakabayashi, J., 2004, Contrasting settings of serpentinite bodies, San Francisco Bay area, California; derivation from the subduction plate vs. mantle hanging wall?: *International Geology Review*, v. 46, p. 1103-1118.
- Williams, P.R., Pigram, C.J., and Dow, D.B., 1984, Melange production and the importance of shale diapirism in accretionary terrains: *Nature*, v. 309, p. 145-146.
- Yamamoto, Y., Otha, Y., and Ogawa, Y., 2000, Implication for the two-stage layer-parallel faults in the context of the Isu forearc collision zone: examples from the Miura accretionary prism, Central Japan: *Tectonophysics*, v. 325, p. 133-144.

## FIGURE CAPTIONS

**Figure 1** – Distribution of main mélanges and broken formations in the peri-Adriatic region. The map has been compiled after a large number of citations quoted in the text and on the basis of the following maps (Bigi et al., 1990; Cerrina Feroni et al., 2002; Vezzani et al., 2009, and various sheets of the 1:50.000 scale Geological Map of Italy).

**Figure 2** – Paleogeographic sketch of the Apennines and Dinarides-Albanides-Hellenides at different stages: (A) the Alpine Neo-Tethys during middle-late Jurassic (modified after Dercourt et al., 1986); (B) the paleogeography at the end of the Mesozoic tectonic phase during Late Eocene – early Oligocene (modified after Castellarin, 1994).

**Figure 3** - Structural-stratigraphic diagram of the Northern Apennines showing the main structural units, the lithostratigraphic groups and the paleogeographic domains (modified after Camerlenghi and Pini, 2009). A particular emphasis is given to the distribution of sedimentary mélanges (olistostromes) and broken formation (tectonosomes). Block letters indicate stratigraphic and paleogeographic names; structural units are in italics.

**Figure 4** - Simplified, speculative cross section of the Northern Apennines and the adjoining Po Plain. Liberally based on: Castellarin et al. (1994) and Argnani et al. (2003).

**Figure 5** - Geological map of the Northern Apennines showing the main bodies of mélanges described in the text. The map is after Pini (1999) and Lucente and Pini (2008). Main sources for the mélange distribution are in: Boccaletti and Coli, (1982), Pini (1993), Marroni et al. (2001), Marroni and Pandolfi (2001), Bettelli et al. (2002), Cerrina Feroni et al. (2002); various sheets of the 1:50.000 scale Geological Map of Italy.

**Figure 6** – Geological-structural map of the Central-Southern Apennines (after Vezzani et al., 2009). Abbreviations indicate the type of mélange described in the text and its location.

**Figure 7** – Cross section of the Central-Southern Apennines (after Vezzani et al., 2009) showing the distribution of the main mélanges described in the text. Location in Fig. 6.



**Figure 8** – Simplified geological map of Albanide-Hellenide mountain belt showing the distribution of the Mesohellenic ophiolite belt and other major tectonic zones. Cross section showing the distribution of the main mélanges (modified after Rassios and Dilek, 2009).

**Figure 9** – Scheme (not in scale) showing the stratigraphic and structural relationships between strike-slip tectonics-related mélanges (3), epi-nappe sedimentary mélanges (6c1) and epi-nappe diapiric mélanges (6c3) in the Torino Hill (Tertiary Piedmont Basin). Modified after Festa (2009).


**Figure 10** - Distribution of the large scale mass-wasting complexes, m.w.c., (I to X) in the SW to NE migrating foreland basin system of the Northern Apennines, from late Oligocene to late Miocene. Adapted from Lucente and Pini (2008).

**Figure 11** – Schematic cross sections showing the emplacement of the Sicilide units in the Molise-Sannio region, Central-Southern Apennines (modified after Ghisetti et al., 2003). During the first tectonic stage (A), Sicilide units represented a tecto-sedimentary epi-nappe mélange (6c2). Slope and debris avalanches at the external frontal thrust (B) of the Sicilide units produced an intra-nappe tecto-sedimentary mélange (6b2). Later tectonic stages (C) deformed the already emplaced mélange.





**Figure 2**



BS = Bocco Shales and equivalents  
 Lf = Ligurian flysches (Helminthoid and Paleocene-Eocene flysches)  
 om = Casanova and Mt. Ragola complexes (ophiolitic mélanges)  
 bc = Ligurian pre-flysch successions ("basal complexes")  
 cs = Triassic-Jurassic substratum from a continental margin (Adria?)  
 op = ophiolites  
 Cc = Canetolo complex  
 ccs = carbonate and carbonate-clastic succession  
 ev = Triassic evaporites  
 Hb = Hercinian basement and continental deposits


-  Strataly disrupted stratigraphic sequences (tectonosomes)
-  Olistostromes, olistoliths complexes, slide blocks
-  Precursory olistostromes and olistostromal carpet
-  Turbiditic infills of foreland basins
-  Contact of uncertain, debated nature (tectonic vs. sedimentary origin)

Figure 3





Figure 4



- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li> Front-Appenninic deposits</li> <li> Sedimentary mélanges (olistostromes) in the Epiligurian successions</li> <li> Epiligurian successions and Tertiary Piedmont Basin</li> <li> Crevasco-Voltaggio and Western Alps metamorphic units</li> <li> External Ligurian units: highly disrupted units (tectonosomes)</li> <li> External Ligurian units: moderate to highly disrupted units (broken formations)</li> <li> Sedimentary mélanges at the base of the External Ligurian units</li> </ul> | <ul style="list-style-type: none"> <li> External Ligurian units</li> <li> Sedimentary mélanges atop the Internal Ligurian units</li> <li> Internal Ligurian units: Gottero and associated units</li> <li> Internal Ligurian units: Bracco and Colli Tavarone ophiolitic units</li> <li> Subligurian units</li> <li> Sub-nappe mélanges</li> <li> Mass wasting deposits and olistostromal carpet</li> </ul> | <p>Adria-related units: Umbrian units</p> <ul style="list-style-type: none"> <li> Romagna-Marche-Umbria fold-and-thrust belt</li> <li> Cervarola-Falserona unit</li> </ul> <p>Tuscan units</p> <ul style="list-style-type: none"> <li> Tuscan nappe</li> <li> Tuscan metamorphic units</li> </ul> <ul style="list-style-type: none"> <li> Stratigraphic contact</li> <li> Normal fault</li> <li> Strike-slip fault</li> <li> Thrust fault</li> </ul> |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|




Figure 6






Figure 7





Figure 8




Figure 9




Figure 10




Figure 11