

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

State of art and future perspectives of biological control of postharvest fruit diseases

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/101450> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in

Spadaro D., Gullino M.L. (2004) – State of art and future perspectives of biological control of postharvest fruit diseases. International Journal of Food Microbiology, 91 (2), 185-194.

DOI: 10.1016/S0168-1605(03)00380-5

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [10.1016/S0168-1605(03)00380-5]

23 **STATE OF ART AND FUTURE PERSPECTIVES OF BIOLOGICAL CONTROL OF**
24 **POSTHARVEST FRUIT DISEASES**

25

26 Davide Spadaro and Maria Lodovica Gullino*

27 Di.Va.P.R.A. – Plant Pathology, University of Torino, Via L. da Vinci 44, 10095 Grugliasco
28 (TO), Italy.

29

30 **Abstract**

31 Synthetic fungicides, when admitted, are the primary means to control postharvest diseases of
32 fruits. Biological control using antagonists has emerged as one of the most promising
33 alternatives to chemicals. During the last twenty years, several biocontrol agents have been
34 widely investigated against different pathogens and fruit crops. Many biocontrol mechanisms
35 have been suggested to operate on fruit including antibiosis, parasitism, induced resistance
36 and competition. Trying to extend the use of the biofungicides, there have been deep studies
37 on application of antagonists mixes, preharvest use, and integration with chemical and
38 physical means of protection. The formulation and application methods are key issues for the
39 efficacy and final success of the commercial product. The new molecular techniques are
40 useful tools in the characterisation of the microorganisms and enhancement of their biocontrol
41 capabilities, through genetic engineering. Although a huge number of scientific papers
42 published on biological control, this method at the moment should be viewed as an important
43 component of an integrated disease management scheme if a significant and permanent
44 reduction of pesticide use is our goal. Anyway scientists, growers and consumers alike must
45 accept the fact that BCA's are usually not as effective as pesticides.

* Corresponding author: Davide Spadaro, Di.Va.P.R.A. – Plant Pathology, University of Torino, Via L. da Vinci
44, 10095 Grugliasco (TO), Italy; phone: +39-6708696; fax: +39-6708541, email: spadaro@agraria.unito.it

46

47 **Keywords:** antagonist, fruit, mechanism of action, postharvest disease.

48

49 **1. Introduction**

50

51 Fruits are an important part of the human diet, because they supply essential nutrients such as
52 vitamins, minerals, and they are important to human health and well-being, for their contents
53 in antioxidants and anticancer substances. An increasing awareness by consumers that diet
54 and health are linked has resulted in a greater consumption of fruits. At the same time,
55 consumers are also more concerned about the safety of the fruits they eat, and they ask for
56 food free from pesticide residues, toxins and pathogens.

57 Losses due to pests and diseases attacks on fruit in field and during storage, transit, and
58 commercialisation steps, before reaching the consumer, are not easily assessed, but can result
59 in 25% of the total production in industrialised countries (Harvey, 1978). In developing
60 countries damages are often higher, exceeding 50%, because of the lack of adequate storage
61 structures (Eckert and Ogawa, 1985). The high water content of plant products, such as fruit,
62 is one of the features that makes them more susceptible to pathogens attack, since they are in
63 orchard (Harvey, 1978). Another factor favourable to pathogenic fungi, particularly to the
64 necrotrophic ones, is the presence during storage on the plant organs of wounds, often
65 produced during harvest and transport of fruit, which represent an ideal way of access for
66 microorganisms.

67 Synthetic fungicides, when admitted, are the primary means to control postharvest diseases.
68 However, several reasons, such as the growing public concern over the human health
69 conditions and the environmental pollution associated with pesticide usage in orchards
70 (Wilson and Wisniewski, 1994), the development of fungicide resistant strains of postharvest

71 pathogens (Romano et al., 1983; Spotts and Cervantes, 1986) and the lack of reregistration of
72 some of the most effective fungicides (Gullino and Kuijpers, 1994) have encouraged the
73 search of alternative approaches.

74 Biological control well fits with the concept of sustainable agriculture, because it mostly
75 exploits natural cycles with zero or reduced environmental impact. Among the biological
76 strategies adoptable in postharvest, the induction of resistance in the fruit, the use of plant or
77 animal products with a fungicidal activity, and, above all, the application of antagonistic
78 microorganisms can be considered. Biological control using antagonists (Wilson and
79 Wisniewski, 1994) has emerged as one of the most promising alternatives, either alone or as
80 part of an integrated pest management to reduce pesticide use.

81 The postharvest environment represents a particular sector for the development of biological
82 measures. Peculiar difficulties are present in the control of postharvest diseases (Chalutz and
83 Droby, 1998): the disease control level required is extremely high (also 95-98%); the
84 nutritional safety imposes special care to the direct use of living microorganisms on food
85 products; the potential market to employ a biofungicide expressly developed for postharvest
86 use is relatively small. On the other side, the possibilities of success for postharvest biological
87 means can be numerous (Chalutz and Droby, 1998): the storage conditions partially
88 controlled, such as temperature and humidity, can switch the host-pathogen-antagonist
89 equilibrium towards the antagonist and the laboratory trials and results have a higher
90 possibility to be transferred into practice; the application site of the antagonist, which is the
91 fruit, is limited, permitting an increase of the biocontrol agent (BCA) efficacy and avoiding
92 the presence of some interfering factors; finally the high value of fruit can justify a treatment
93 with a product relatively expensive.

94 During the last 20 years, several biocontrol agents have been exploited and widely
95 investigated against different postharvest fungal pathogens (*Alternaria*, *Botrytis*,

96 *Colletotrichum*, *Monilia*, *Penicillium*, *Rhizopus* spp.) on different host species. Many of the
97 first studies were aimed at the study of the mode of action and the evaluation of the efficacy
98 of some potential biocontrol bacteria, such as *Brevibacillus subtilis*, producer of antibiotics
99 (Pusey et al., 1986). The consideration that the application of bacterial antagonists on the fruit
100 was not commercially acceptable, at least in the short period, brought to switch the interest on
101 antagonists using modes of action different from antibiosis. Wilson and Wisniewski (1994)
102 indicated the characteristics of an ideal antagonist: genetic stability, efficacy at low
103 concentrations and against a wide range of pathogens on various fruit products, simple
104 nutritional requests, survival in adverse environmental conditions, growth on cheap substrates
105 in fermenters, lack of pathogenicity for the host plant and lack of production of metabolites
106 potentially toxic for humans, resistance to the most frequently used pesticides, compatibility
107 with other chemical and physical treatments. Yeasts, between the potential BCAs, possess a
108 good number of the mentioned features and, during the last years, research focused on the
109 selection and study of these microorganisms.

110 At present, three products containing *Pseudomonas syringae* Van Hall, active against
111 *Botrytis*, *Penicillium*, *Mucor* and *Geotrichum* spp., named Bio-Save 100, Bio-Save 110 and
112 Bio-Save 1000 and commercialised by EcoScience Corp. (Janisiewicz and Jeffers, 1997), and
113 a product containing *Candida oleophila* Montrocher, effective against *Botrytis* and
114 *Penicillium* spp., named Aspire and commercialised by Ecogen Inc. (Hofstein et al., 1994),
115 are available on the market for postharvest protection. Other yeast species have been
116 extensively tested and could be registered in relatively short times. Among the
117 microorganisms under development, there are antagonistic strains belonging to the species
118 *Aureobasidium pullulans* (Ippolito et al., 2000), *Candida saitoana* (El-Ghaouth et al., 1998),
119 *Candida sake* (Teixido et al., 1998) and *Metschnikowia pulcherrima* (Spadaro et al., 2001a).

120

121 **2. Mechanisms of action**

122

123 Information on the mechanisms of action for most of the antagonists investigated is still
124 incomplete, because of the difficulties encountered during the study of the complex
125 interactions between host, pathogen, antagonist and others microorganisms present in the site
126 of interaction. However a good understanding of the mode of action is essential to develop
127 appropriate formulation and methods of application, and to obtain registration.

128 Several possible biocontrol mechanisms have been suggested to operate against post-harvest
129 rots on fruit including antibiosis, parasitism or direct interaction with the pathogen
130 (extracellular hydrolases), induced resistance and competition for nutrients and space (Droby
131 and Chalutz, 1994).

132

133 *2.1. Antibiosis*

134 Some of the most active bacteria are producer of antibiotics, whose action, at least partially,
135 influences these BCA's effectiveness. *B. subtilis* was able to produce iturin, a powerful
136 antifungal peptide (Gueldner et al., 1988) but also gramicidin S (Edwards and Seddon, 2001),
137 *Pseudomonas cepacia* showed to synthesise pyrrolnitrin, that used alone can control *Botrytis*
138 *cinerea* and *Penicillium expansum* attacks on pome fruit (Janisiewicz et al., 1991). Also from
139 these observation, it was possible to develop phenylpyrroles fungicides (Gehmann et al.,
140 1990). Often, over antibiosis, other mechanisms are present: *Penicillium digitatum* strains
141 resistant to pyrrolnitrin are still inhibited by *P. cepacia* (Smilanick and Denis-Arrue, 1992).

142 The main concerns, related to the use of antibiotics in food products, regard the selection of
143 human pathogens resistant to these chemicals and the likely rapid development of resistance
144 in fruit pathogens. Even if antibiotic producers are able to control wound infections occurred
145 before antagonist application or latent infections, at the moment there are not such antagonists

146 registered for use on food.

147

148 2.2 Competition

149 Others selected microorganisms, particularly yeasts, act mainly establishing with the pathogen
150 a competition for space and/or utilisation of some nutrients (Piano et al., 1997; Filonow,
151 1998; Spadaro et al., 2001a). Yeasts can successfully compete with the pathogen, inhibiting
152 its growth but often leaving it alive. Also some bacteria act for competition, for example
153 *Enterobacter cloacae* against *Rhizopus stolonifer* on peaches (Wisniewski et al., 1989). In the
154 competition for space yeasts are helped by the formation of an extracellular polysaccharide
155 capsule that can promote adhesion to fruit surface (Andrews et al., 1994). Competition for
156 nutrients was demonstrated for *Pichia guilliermondii* against *P. digitatum* cocultivated on
157 synthetic means (Droby et al., 1989): the addition of exogenous nutrients resulted in a reduced
158 efficacy, because antagonists offered better results when nutrients were scarce. A rapid
159 multiplication and colonisation by antagonist cells in the wound was elucidated in various
160 interactions (Droby et al., 1989; Smilanick and Denis-Arrue, 1992; Piano et al., 1997). Recent
161 studies on the repartition of radiolabelled glucose between the antagonist yeasts
162 *Sporobolomyces roseus* or *Cryptococcus laurentii* and the pathogen *B. cinerea* point out a
163 strong sugar assumption by the BCAs, that blocks fungus conidial germination for nutrients
164 deprivation (Filonow, 1998). In fruit wounds, nutrients competition is likely extended to other
165 nutrients, such as nitrogen compounds low concentrated. Janisiewicz et al. (2000) recently have
166 developed a non-destructive method using tissue culture plates with cylinder inserts
167 containing defusing membrane at one end to study competition for nutrients without
168 competition for space.

169

170 2.3 Parasitism

171 Antagonist and pathogen can interact also through a direct parasitism. Wisniewski et al.
172 (1991) observed a strong adhesion *in vitro* of *P. guilliermondii* antagonist cells to *B. cinerea*
173 mycelium, perhaps due to a lectin link. Such adhesion is blocked by exposure to compounds
174 able to alter the protein integrity and the respiration process. Moreover *P. guilliermondii*
175 shows an high activity of β -1,3-glucanase enzyme, that could be associated to pathogen link
176 and result in the degradation of the fungal cell walls (Jijakli and Lepoivre, 1998). Also
177 *Aureobasidium pullulans* in apple wounds produces extracellular exochitinase and β -1,3-
178 glucanase, that could play a role in the biocontrol activity (Castoria et al., 2001). El-Ghaouth
179 et al. (1998) found, through ultrastructural and cytochemical studies, that *Candida saitoana*
180 yeast cells, when cultivated together with *B. cinerea* mycelium, link fungal hyphae that show
181 cytological damages, such as papillae and other protuberances in the cell wall, and
182 degeneration of the cytoplasm. Finally some yeasts, such as *S. roseus* and *C. laurentii*, when
183 applied, are able to reduce, on apple carposphere, the conidial adhesion and germination of *B.*
184 *cinerea*, that is favoured by butyl acetate, volatile aroma produced by the fruit (Filonow,
185 2001).

186

187 *2.4 Induced resistance*

188 Some BCAs can interact with the host tissue, particularly the wounds, increasing the
189 cicatrisation processes (Droby and Chalutz, 1994). Several antagonistic yeasts are as much
190 effective as applied before pathogen inoculation. This observation has brought to suppose that
191 yeast cell application could induce resistance processes in the fruit skin. Some *Candida*
192 strains are able to cause chemical and osmotic changes in apple tissues, favouring antagonist
193 settlement (McLaughlin et al., 1990). A *P. guilliermondii* strain stimulates the production of
194 ethylene, an hormone able in grapefruit to activate the phenylalaninammonium-lyase
195 (Wisniewski et al., 1991), enzyme involved in the phenols, phytoalexins and lignins synthesis.

196 A phytoalexins accumulation (scoparon and scopoletin) was noted in citrus fruit treated with
197 yeast cell (Rodov et al., 1994). In addition to control decays, *A. pullulans* can cause on apple
198 fruit a transient increase in β -1,3-glucanase, chitinase, and peroxidase activities starting 24 h
199 after treatment (Ippolito et al., 2000).

200

201 **3. Extension of use**

202

203 Potential biocontrol agents often have some significant limitations: they have a reduced action
204 range, because they partially act on specific hosts against well defined pathogens on particular
205 environmental conditions. A method to select antagonist with a broader spectrum of action,
206 preferably for commercial development, is the carrying out of initial efficacy tests on various
207 pathogens and fruit species (Wilson et al., 1993; Lima et al., 1999).

208

209 *3.1 Antagonist mixtures*

210 In the enhancement of a biocontrol system, work could focuses on a promising approach,
211 which is the development of antagonist mixtures. There is a superior probability to have an
212 effective biological control with a mixture of several antagonists, complementary and non
213 competitive, that with only one. Such mixture has some peculiar advantages (Janisiewicz,
214 1998): over the broadening of the spectrum of action (different fruits, cultivars and ripening
215 stages), it can increase the efficacy (less biomass necessary) and offer a higher reliability,
216 permitting a reduction in the application times and treatment costs. Moreover, it permits the
217 combination of different genetic traits, avoiding genetic engineering. The evaluation of casual
218 combinations of antagonists isolated from apple fruits and leaves resulted in the selection of a
219 mixture of *P. syringae* and *S. roseus*, superior to both BCAs applied alone in the control of
220 the *Penicillium* rot (Janisiewicz and Bors, 1995).

221

222 3.2 Preharvest use

223 One of the major obstacles to the development of postharvest BCAs is their inability to
224 control previously established infections, such as latent infections. Field application of the
225 BCAs may enable early colonisation of the fruit surfaces, protecting from these infections
226 (Ippolito and Nigro, 2000). To be successful in preharvest application, potential antagonists
227 should be able to tolerate low-nutrient availability, UV rays, high temperature and dry
228 conditions. Leibinger et al. (1997) applied in preharvest a mixture of a yeast *A. pullulans* and
229 a bacterium (*B. subtilis*) obtaining a level of control of *P. expansum* and *B. cinerea* on apple
230 similar to fungicides. Teixido et al. (1998) applied unmodified and low water activity tolerant
231 cells of *Candida sake* before harvest to control blue mould of apples, obtaining similar
232 disease control.

233

234 3.3 Integrated use

235 Biological means of control can not at the moment solve all the problems of postharvest rots
236 during fruit storage and they must be considered instruments to be used in combination with
237 other methods of control in an integrated vision of disease management. For example,
238 biocontrol agents can be combined with waxes and fungicides applied not only in post but
239 also in pre-harvest (Pusey, 1994). In some laboratory and semi-commercial trials the efficacy
240 of the BCA *P. guilliermondii* was consistently increased by the addition of small
241 concentrations of imazalil or thiabendazole, reaching the levels of control of the fungicide
242 alone (Droby et al., 1993). Antagonists, if employed together with fungicides, have to be
243 tolerant or they request the development of resistant strains. Yeasts generally are tolerant to
244 many of the fungicides used in postharvest: *Metschnikowia pulcherrima* (Spadaro et al.,
245 2001a) is tolerant to relative high concentrations of benzimidazoles (benomyl and

246 thiabendazole) and dicarboximides (vinchlozolin and procymidone). Moreover
247 microorganisms need to survive to thermal, hygrometry and atmospheric storage conditions.
248 The efficacy of *M. pulcherrima* against *B.cinerea* and *P. expansum* is increased when fruits
249 are stored at low temperatures optimal for host, respect to storage at 23°C, because the
250 antagonist is more adaptable than the pathogens to cold temperatures (Spadaro et al., 2001a).
251 Among the strategies evaluated during the last years, it is to remember the combination of
252 biological treatments with other techniques alternative to chemicals: thermotherapy (Barkai-
253 Golan and Phillips, 1991), ultraviolet rays (Chalutz et al., 1992), animal and plant natural
254 products (Aharoni et al., 1993), calcium infiltrations (Janisiewicz et al.,1998), sodium
255 bicarbonate (Teixido et al., 2001) or ethanol (Spadaro et al., 2001b).
256 Calcium chloride infiltrations combined to antagonist application on apples increases control
257 of *P. expansum* after 3 and 6 months of storage at 1°C, compared to simple biological
258 treatment (Janisiewicz et al., 1998). A even more integrated approach, experienced on “Gala”
259 apples (Conway et al., 1999) consists of a heat treatment (4 days at 38°C), followed by a
260 calcium chloride (2%) infiltration and a *Pseudomonas syringae* cell suspension application.
261 The result of the three treatments carried out together was a reduction of the Penicillium rots
262 (91%) higher respect to calcium, antagonist, or heat treatments applied alone.
263 The efficacy of *Pantoea agglomerans* for the control of green mould was improved when
264 combined with sodium bicarbonate, or baking soda (Teixido et al., 2001), a common food
265 additive, allowed with no restrictions for many application by European and North American
266 regulations, and listed as approved ingredients on products labelled as “organic”. The
267 combination of sodium bicarbonate, or ethanol, or acybenzolar-S-methyl with *M. pulcherrima*
268 cell suspension and heat treatment was also studied by Spadaro et al. (2001b). Ethanol occurs
269 naturally in fruit and many other food products, and its toxic effects on spores of fungal
270 pathogens have been reported: the flesh treated with the alcohol was significantly firmer and

271 injury to the fruit did not occur.

272 It is possible to conclude that the possibilities to apply antagonists in the context of integrated
273 control are very extensive.

274

275 **4. Improvement of biocontrol agents**

276

277 *4.1 Formulation and application*

278 Differently from soilborne or open field pathogens, where a disease control of 70-80% can be
279 considered satisfying, postharvest disease control requests a higher level of efficacy and more
280 consistent results. Even considering the most effective BCA's studied until now, they rarely
281 reach the levels of efficacy of the fungicides. It is therefore necessary an increase in their
282 antagonistic activity to achieve a practical use.

283 A rapid, efficient and cheap mass production of the antagonist, generally by fermentation, is a
284 key issue. The efficacy of many antagonists of wound pathogens is directly related to the
285 number of antagonist propagules inoculated (Hofstein et al., 1994), so that a really simple way
286 to increase the effectiveness is the application of a higher number of cells. To scale-up a
287 laboratory fermentation process to an industrial level, it is fundamental to find the nitrogen
288 and carbon sources that provide maximum biomass production and minimum cost of media,
289 whilst maintaining biocontrol efficacy. Costa et al. (2001) have studied yeast extract, dry beer
290 yeast, sucrose, and molasses as possible substrate for the production of the biocontrol agent *P.*
291 *agglomerans*.

292 A correct formulation can be decisive in the improvement of the efficacy and extension of the
293 product shelf-life, facilitating the storage for periods of time commercially acceptable
294 (Janisiewicz and Jeffers, 1997). The application of adjuvants can protect and stimulate the
295 establishing of the antagonist on the host surface. The addition of calcium salts increases the

296 activity of several antagonistic yeasts (Janisiewicz et al., 1998). The addition of glycerol and
297 trealose to the culture means augmented the osmotic tolerance and control capability of *C.*
298 *sake* against *P. expansum* on apple (Janisiewicz, 1998). Sodium alginate,
299 carboxymethylcellulose and chitosan, are adhesivants and can be added to yeast cell
300 suspension, to increase the activity of the formulate. The three adhesivants have been added to
301 a strain of *M. pulcherrima* (Piano et al., 1998) significantly increasing the efficacy against
302 grey rot on apple. Chitosan has also a fungistatic activity demonstrated against the main
303 postharvest pathogens of strawberry (El-Ghaouth et al., 1992). Recently, El-Ghaouth et al.
304 (2000a) developed a biocontrol product called "bioactive coating" consisting of a unique
305 combination of an antagonist with glycolchitosan, a chemically-modified chitosan. The
306 bioactive coating made it possible to exploit the antifungal property of glycolchitosan and the
307 biological activity of the antagonist. Moreover, when applied as a pretreatment, sodium
308 carbonate enhanced the efficacy of the bioactive coating (El-Ghaouth et al., 2000b).

309 Also application systems have a fundamental role in the determination of the final results. The
310 coatings most often applied to citrus fruit contain shellac, which is a purified product of the
311 hardened resinous secretion of the scale insect *Kerria lacca*. Having developed coating
312 formulations before biological control, it is important to test the suitability of such products
313 for antagonistic applications. McGuire (2000) found that, after 3 or 4 months storage, *C.*
314 *oleophila* survival was more efficacious when applied in the shellac than when applied by
315 preliminary immersion and subsequent drying of fruit prior to shellacking. On the contrary,
316 Chalutz and Droby (1998) had noted that the application of antagonistic yeasts in aqueous
317 suspension before waxing results in an efficacy much more consistent respect to a unique
318 application of antagonists and wax.

319 Another issue to face in the commercial production of biocontrol agents is the storage, that
320 should be as long as possible. The BCA's should have a storability of at least 6 months and

321 preferably of 2 years (Pusey, 1994). Abadias et al. (2001a) found that freezing at -20°C was
322 the best method to preserve the viability of *C. sake* cells after freeze drying. Survival of the
323 cells was higher using 10% skim milk as a protectant, and increased by using other
324 appropriate protectants, such as lactose, glucose, fructose or sucrose. Skimmed milk was also
325 the best rehydration medium with 1% peptone (Abadias et al., 2001b). However the efficacy
326 of freeze-dried cells was significantly lower than that of fresh cells.

327 The transition from fungicide use to employment of biological means will be as easier as
328 more flexible are the biofungicides.

329

330 4. 2 *The nutritional environment*

331 Temperature, humidity and gas composition in warehouse are optimised to guarantee a high
332 fruit quality during storage and should not be changed but it is possible to manipulate the
333 chemical environment to favour the antagonist. The addition of nutrients preferably
334 metabolised by the antagonist and difficultly by the pathogen was suggested in several
335 antagonist-pathogen interactions (Janisiewicz, 1998). The application of two nitrogen
336 compounds, L-asparagine and L-proline (Janisiewicz et al., 1992), and an analogue of sugar,
337 2-deoxy-D-glucose (Janisiewicz, 1994) showed a consistent increase of the control of *P.*
338 *expansum* on apple. Both aminoacids stimulated *P. expansum* germination but slowed
339 mycelial growth. Also L-glutamine added to *M. pulcherrima* showed a direct influence on the
340 yeast, because its application with the antagonist contributed to reduce Botrytis rot, while
341 without yeast, it was ineffective (Piano et al., 1998). The sugar analogue 2-deoxy-D-glucose
342 could be a useful additive to antagonistic microorganisms, provided that it has a fungicidal
343 action on the major postharvest pathogens of apple and peach fruit (El-Ghaouth et al., 1997)
344 and that the antagonist is resistant to its toxic effects. Recently the sugar analogue was found
345 to be compatible with the antagonistic yeast *C. saitoana* and effective against apple and citrus

346 fruit decay (El-Ghaouth et al., 2000c).

347 Nutritional composition can influence not only density and competitiveness of the population,
348 but also the production of metabolites crucial in many control systems, such as antibiotics
349 (Gueldner et al., 1988) and cell wall degrading enzymes (Wisniewski et al., 1991). The form
350 and concentration of nitrogen and carbon sources can therefore be important factors in the
351 synthesis and secretion of key compounds in the biocontrol mechanism.

352

353 *4.3 Manipulation of antagonists*

354 It is possible to increase the ecological competence of microorganisms manipulating
355 antagonists with techniques of mutagenesis (physical or chemical) or of sexual recombination,
356 through protoplast fusion or continued culture.

357 Because nitrogen seems to be a limiting substance, it was noted that biocontrol increases
358 adding some aminoacids. Therefore biocontrol strains with a superior capability of
359 exploitation of nitrogen compounds present, or with a higher transport or metabolism rate of
360 the limiting factor, could be developed (Janisiewicz, 1998). Some phenolic compounds,
361 among them benzoic or chlorogenic acid, present at the wound sites, influence negatively the
362 colonisation by yeasts; it is therefore conceivable trying to obtain strains resistant to these
363 phenolic compounds (Bizeau et al., 1989). Mutants that use new substrates, not metabolised
364 by the pathogen, to provide a nutritional advantage, could also be induced (Janisiewicz,
365 1998).

366 With genetic engineering techniques, features for the exploitation of plant products or
367 additives applied together with the antagonist, for fungicide resistance, for carposphere
368 colonisation in storage conditions, fro synthesis of compounds favouring antagonism
369 (antibiotics or siderophores), could be transferred into the potential antagonistic
370 microorganisms (Pusey, 1994). Because one mechanism involved in the biocontrol of

371 postharvest fruit pathogens is mycoparasitism, lytic enzymes such as chitinases (Chernin et
372 al., 1997), proteases and glucanases (De la Cruz et al., 1995) produced by bacterial and fungal
373 microorganisms, could be inserted in the potential antagonists to improve the degradation of
374 the pathogen cell walls, resulting in death or inhibition of growth of the attacked fungus. For
375 biocontrol of soilborne pathogens, for example, a chitinase gene, isolated from *Serratia*
376 *marcescens*, was introduced into the endophytic bacterium *Pseudomonas fluorescens* to
377 improve the control of *Rhizoctonia solani* on bean (Downing and Thomson, 2000).

378 Another idea could be the insertion of the gene for amylase under constitutive promoter in
379 some BCA's, because many fruits are rich in amid and antagonists could use effectively this
380 carbon source having a consistent advantage. First experiments of transformation have been
381 successful: *M. pulcherrima* was transformed with the green fluorescent protein gene (Nigro et
382 al., 1999), *C. oleophila* was transformed with β -glucuronidase gene (Chand-Goyal et al.,
383 1998), and histidine auxotrophs of *C. oleophila* were transformed with *HIS3*, *HIS4* and *HIS5*
384 genes (Chand-Goyal et al., 1999). All these studies were accomplished only to obtain variants
385 of the antagonistic strains with a genetically stable marker to expedite studies on the ecology
386 of the yeasts on the fruit surface, but are really useful to put successive insertions of useful
387 genes right. In all cases the transformed antagonists maintained the biocontrol capability.

388

389 **5. Molecular characterisation**

390

391 New molecular technologies improving the preliminary selection of biocontrol agents and the
392 monitoring of field applied antagonists may greatly facilitate the selection and screening of
393 yeast isolates having a superior antagonistic activity. Morphological and cultural
394 characteristics alone are not sufficient to distinguish between strains of the same species and
395 do not allow the exploitation of any possible naturally available genetic variability. Moreover

396 the effectiveness of the preharvest strategy depends to a large extent upon the survival of the
397 antagonists in competition with other microorganisms on the surface of fruits. A study of the
398 ecology of the BCA's is therefore necessary but often complicated by the existence of other
399 microorganisms on the fruit that are morphologically similar to the antagonistic strain.
400 Molecular approaches can help to differentiate a biocontrol agent from an epiphytic
401 microorganism. Between the technologies already used to obtain DNA fingerprintings the
402 random amplified polymorphic DNA (RAPD) and the arbitrarily primed PCR (AP-PCR) are
403 to remember (Schena et al., 1999; 2000). Both these methods are based on the amplification
404 of random genomic DNA fragments by arbitrarily selected PCR primers and have the major
405 disadvantage that they are very sensitive to the reaction conditions, DNA quality and PCR
406 temperature profiles, which limit their applications.

407 New techniques more robust and reliable, such as amplified fragment length polymorphism
408 (Vos et al., 1995; De Barros et al., 1999), based on the selective PCR amplification of
409 restriction fragments from a total digest of genomic DNA, and PCR amplification of the ITS
410 regions of the ribosomal DNA (Masih et al., 2001) can be applied on potential biocontrol
411 agents.

412

413 **6. Conclusions**

414

415 From this review, it appears that some significant progress has been made toward biological
416 and integrated control of postharvest diseases on fruits. Some biofungicides are already on the
417 market in a few countries, and will probably become more widely available as they are
418 registered in more areas. Other BCA should reach the market soon. Postharvest conditions
419 provide an ideal niche for BCAs since they are less subject to sudden climate changes, and are
420 often equipped with a sophisticated system of climate control.

421 It is unrealistic to assume that perfect conditions for the development of BCA's will always
422 prevail in the warehouse, and as a result, biofungicides will rarely stand alone as a complete
423 measure of disease control under all conditions. For this reason, scientists, growers and
424 consumers alike must accept the fact that BCA's are usually not as effective as pesticides. The
425 success of biological control greatly depends in a change of mentality of the consumer, not
426 anymore willing to have fruit with satisfying exterior aspects, but with an inner quality.
427 Biological control should be viewed at the moment as an important if not essential component
428 of an integrated disease management scheme if a significant and permanent reduction of
429 pesticide use is our goal.

430

431 **Acknowledgements**

432

433 The manuscript has been written by the first author and revised by the second one. Work
434 supported by a grant from Italian Ministry for Environment and Territory.

435

436 **References**

437

438 Abadias, M., Benabarre, A., Teixido, N., Usall, J., Vinas, I., 2001a. Effect of freeze drying
439 and protectants on viability of the biocontrol yeast *Candida sake*. Int. J. Food Microbiol.
440 65, 3, 173-182.

441 Abadias, M., Teixido, N., Usall, J., Benabarre, A., Vinas, I., 2001b. Viability, efficacy, and
442 storage stability of freeze-dried biocontrol agent *Candida sake* using different protective
443 and rehydration media. J. Food Prot. 64, 6, 856-861.

444 Aharoni, Y., Copel, A., Fallik, E., 1993. Hinokitiol (b-thujaplicin), for postharvest decay
445 control on "Galia" melons. N. Z. J. Crop Hortic. Sci. 21, 165-169.

446 Andrews, J.H., Harris, R.F., Spear, R.N., Lau, G.W., Nordheim, E.V., 1994. Morphogenesis
447 and adhesion of *Aureobasidium pullulans*. *Can. J. Microbiol.* 40, 6-17.

448 Barkai-Golan, R., Phillips, D. J., 1991. Postharvest heat treatment of fresh fruits and vege-
449 tables for decay control. *Plant Dis.* 75, 1085-1089.

450 Bizeau, C., Ponchant, F., Moreau, M., 1989. Evolution of yeast flora on cider apples of
451 Brittany. *Proc. 7th Int. Symp. on Yeasts, Perugia, Italia.* *Yeast*, vol. 5. John Wiley &
452 Sons, Ltd., Chichester, UK, 491-494.

453 Castoria, R., De Curtis, F., Lima, G., Caputo, L., Pacifico, S., De Cicco, V., 2001.
454 *Aureobasidium pullulans* (LS-30) an antagonist of postharvest pathogens of fruits: study
455 on its modes of action. *Postharvest Biol. Technol.* 22, 7-17.

456 Chalutz, E., Droby, S., 1998. Biological control of postharvest disease. In: Boland, G.J.,
457 Kuykendall, L.D. (Eds.), *Plant-Microbe Interactions and Biological Control* Marcel
458 Dekker, New York, USA, pp. 157-170.

459 Chalutz, E., Droby, S., Wilson, C.L., Wisniewski, M.E., 1992. UV-induced resistance to
460 postharvest diseases of citrus fruit. *Phytochem. Phytobiol. B.: Biol.* 15, 367.

461 Chand-Goyal T., Eckert J.W., Droby, S., Atkinson, K., 1998. A method for studying the
462 population dynamics of *Candida oleophila* on oranges in the grove, using a selective
463 isolation medium and PCR technique. *J. Microbiol. Res.* 153, 3, 265-270.

464 Chand-Goyal, T., Eckert, J.W., Droby, S., Glickmann, E., Atkinson, K., 1999. Transformation
465 of *Candida oleophila* and survival of a transformant on orange fruit under field
466 conditions. *Curr. Genet.* 35, 51-57.

467 Chernin, L.S., De La Fuente, L., Sobolev, V., Haran, S., Vorgias, C.E., Oppenheim, A.B.,
468 Chet, I., 1997. Molecular cloning, structural analysis, and expression in *Escherichia coli*
469 of a chitinase gene from *Enterobacter agglomerans*. *Appl. Environm. Microbiol.* 63, 834-
470 839.

471 Conway, W.S., Janisiewicz, W.J., Klein, J.D., Sams, C.E., 1999. Strategy for combining heat
472 treatment, calcium infiltration, and biological control to reduce postharvest decay of
473 “Gala” apples. HortScience 34, 4, 700-704.

474 Costa, E., Teixido, N., Usall, J., Atares, E., Vinas I., 2001. Production of the biocontrol agent
475 *Pantoea agglomerans* strain CPA-2 using commercial products and by-products. Appl.
476 Microbiol. Biotechnol. 56, 367-371.

477 De Barros, L., Rainieri, S., Henschke, P.A., Langridge, P., 1999. AFLP fingerprinting for
478 analysis of yeast genetic variation. Int. J. Syst. Bacteriol. 49, 915-924.

479 De La Cruz, J., Pintor-Toro, J.A., Benitez, T., Llobell, A., Romero, L.C., 1995. A novel endo-
480 1,3-glucanase, BGN 13.1, involved in the mycoparasitism of *Trichoderma harzianum*. J.
481 Bacteriology 177, 6937-6945.

482 Downing, K.J., Thomson, J.A., 2000. Introduction of the *Serratia marcescens* *chiA* gene into
483 an endophytic *Pseudomonas fluorescens* for the biocontrol of phytopathogenic fungi.
484 Can. J. Microbiol. 46, 4, 363-369.

485 Droby, S., Chalutz, E., 1994. Mode of action of biocontrol agents of postharvest diseases. In:
486 Wilson, C.L., Wisniewski, M.E. (Eds.), Biological Control of Postharvest Diseases.
487 Theory and Practice. CRC Press, Boca Raton, USA, pp. 63-75.

488 Droby, S., Chalutz, E., Wilson, C.L., Wisniewski, M.E., 1989. Characterization of the
489 biocontrol activity of *Debaryomyces hansenii* in the control of *Penicillium digitatum* on
490 grapefruit. Can. J. Microbiol. 35, 794-800.

491 Droby, S., Hofstein, R., Wilson, C.L., Wisniewski, M.E., Fridlender, B., Cohen, L., Weiss, B.,
492 Timar, D., Chalutz, E., 1993. Pilot testing of *Pichia guilliermondii*: a biocontrol agent of
493 postharvest diseases of citrus fruit. Biol. Control 3, 47-52.

494 Eckert, J.W., Ogawa, J.M., 1985. The chemical control of postharvest diseases: subtropical
495 and tropical fruits. Annu. Rev. Phytopathol. 23, 421-454.

496 Edwards, S.G., Seddon, B., 2001. Mode of antagonism of *Brevibacillus brevis* against
497 *Botrytis cinerea* in vitro. J. Appl. Microbiol. 91, 4, 652-659.

498 El-Ghaouth, A., Arul, J., Grenier, J., Asselin, A., 1992. Antifungal activity of chitosan on two
499 postharvest pathogens of strawberry fruits. Phytopathology 82, 398-402.

500 El-Ghaouth, A., Smilanick, J.L., Brown, G.E., Ippolito, A., Wisniewski, M., Wilson, C.L.,
501 2000a. Application of *Candida saitoana* and glycolchitosan for the control of postharvest
502 diseases of apple and citrus fruit under semi-commercial conditions. Plant Dis. 84, 243-
503 248.

504 El-Ghaouth, A., Smilanick, J.L., Wilson, C.L., 2000b. Enhancement of the performance of
505 *Candida saitoana* by the addition of glycolchitosan for the control of postharvest decay
506 of apple and citrus fruit. Postharvest Biol. Technol. 19, 103-110.

507 El-Ghaouth, A., Smilanick, J.L., Wisniewski, M., Wilson, C.L., 2000c. Improved control of
508 apple and citrus fruit decay with a combination of *Candida saitoana* and 2-deoxy-D-
509 glucose. Plant Dis. 84, 249-253.

510 El-Ghaouth, A., Wilson, C.L., Wisniewski, M., 1998. Ultrastructural and cytochemical
511 aspects of the biological control of *Botrytis cinerea* by *Candida saitoana* in apple fruit.
512 Phytopathology 88, 282-291.

513 El-Ghaouth A., Wilson C.L., Wisniewski M., 1997. Antifungal activity of 2-deoxy-D-glucose
514 on *Botrytis cinerea*, *Penicillium expansum*, and *Rhizopus stolonifer*. Ultrastructural and
515 cytochemical aspect. Phytopathology 88, 772-779.

516 Filonow, A.B., 2001. Butyl acetate and yeasts interact in adhesion and germination of *Botrytis*
517 *cinerea* conidia in vitro and in fungal decay of golden delicious apple. J. Chem. Ecol. 27,
518 831-844.

519 Filonow, A.B., 1998. Role of competition for sugars by yeasts in the biocontrol of gray mould
520 of apple. Biocontrol Sci. Technol. 8, 243-256.

521 Gehmann, K., Nyfeler, R., Leadbeater, A.J., Nevill, D., Sozzi, D., 1990. CGA 173506: a new
522 phenylpyrrole fungicide for broad spectrum disease control. Proc. Brighton Crop Prot.
523 Conf., Pests and Diseases, 399-406.

524 Gueldner, R.C., Reilly, C.C., Pusey, R.L., Costello, C.E., Arrendale, R.F., Cox, R.H.,
525 Himmelsbach, D.S., Crumley, F.G., Cutler, H.G., 1988. Isolation and identification of
526 iturins as antifungal peptides in biological control of peach brown rot with *Bacillus*
527 *subtilis*. J. Agric. Food Chem. 36, 366-370

528 Gullino, M.L., Kuijpers, L.A.M., 1994. Social and political implications of managing plant
529 diseases with restricted fungicides in Europe. Annu. Rev. Phytopathol. 32, 559-579.

530 Harvey, J.M., 1978. Reduction of losses in fresh market fruits and vegetables. Annu. Rev.
531 Phytopathol. 16, 321-341.

532 Hofstein, R., Friedlender, B., Chalutz, E., Droby, S., 1994. Large scale production and pilot
533 testing of biocontrol agents of postharvest diseases. In: Wilson, C.L., Wisniewski, M.
534 (Eds.), Biological Control of Postharvest Diseases – Theory and Practice. CRC Press
535 Inc., Boca Raton, FL, USA, pp. 89-100.

536 Ippolito, A., El-Ghaouth, A., Wilson, C.L., Wisniewski, M., 2000. Control of postharvest
537 decay of apple fruit by *Aureobasidium pullulans* and induction of defense responses.
538 Postharvest Biol. Technol. 19, 265-272.

539 Ippolito, A., Nigro, F., 2000. Impact of preharvest application of biological control agents on
540 postharvest diseases of fresh fruits and vegetables. Crop Prot. 19, 610-619.

541 Janisiewicz, W.J., 1994. Enhancement of biocontrol of blue mold with nutrient analog 2-
542 deoxy-D-glucose on apples and pears. Appl. Environ. Microbiol. 60, 2671-2676.

543 Janisiewicz W.J., 1998 – Biocontrol of postharvest diseases of temperate fruits – Challenges
544 and opportunities. In: Boland, G.J., Kuykendall, L.D. (Eds.), Plant-Microbe Interactions
545 and Biological Control. Marcel Dekker, New York, USA, pp. 171-198.

546 Janisiewicz, W. J., Bors, R., 1995. Development of a microbial community of bacterial and
547 yeast antagonists to control wound-invading postharvest pathogens. Appl. Environ.
548 Microbiol. 61, 3261-3267.

549 Janisiewicz, W.J., Conway, W.S., Glenn, D.M., Sams, C.E., 1998. Integrating biological
550 control and calcium treatment for controlling postharvest decay of apples. HortScience
551 33, 1, 105-109.

552 Janisiewicz, W.J., Jeffers, S.N., 1997. Efficacy of commercial formulation of two
553 biofungicides for control of blue mold and gray mold of apples in cold storage. Crop Prot.
554 16, 629-633.

555 Janisiewicz, W.J., Tworkoski, T.J., Sharer, C., 2000. Characterizing the mechanism of
556 biological control of postharvest diseases on fruits with a simple method to study
557 competition for nutrients. Phytopathology 90, 1196-1200.

558 Janisiewicz, W.J., Usall, J., Bors, B., 1992. Nutritional enhancement of biocontrol of blue
559 mold on apples. Phytopathology 82, 1364-1370.

560 Janisiewicz, W.J., Yourman, L., Roitman, J., Mahoney, N., 1991. Postharvest control of blue
561 mold and gray mold of apples and pears by dip treatments with pyrrolnitrin, a metabolite
562 of *Pseudomonas cepacia*. Plant Dis. 75, 490-494.

563 Jijakli, M.H., Lepoivre, P., 1998. Characterization of an exo-b-1,3-glucanase produced by
564 *Pichia anomala* strain K, antagonist of *Botrytis cinerea* on apples. Phytopathology 88, 4,
565 335-343.

566 Leibinger, W., Breuker, B., Hahn, M., Mengden, K., 1997. Control of postharvest pathogens
567 and colonization of the apple surface by antagonistic microorganisms in the field.
568 Phytopathology 87, 1103-1110.

569 Lima, G., Arru, S., De Curtis, F., Arras, G., 1999. Influence of antagonist, host fruit and
570 pathogen on the biological control of postharvest fungal diseases by yeasts. J. Ind.

571 Microbiol. Biot. 23, 223-229.

572 Masih, E.I., Slezack-Deschaumes, S., Marmaras, I., Barka, E.A., Vernet, G., Charpentier, C.,
573 Adholeya, A., Paul, B., 2001. Characterisation of the yeast *Pichia membranifaciens* and
574 its possible use in the biological control of *Botrytis cinerea*, causing the grey mould
575 disease of grapevine. FEMS Microbiol. Lett. 202, 2, 227-232.

576 McGuire, R.G., 2000. Population dynamics of postharvest decay antagonists growing
577 epiphytically and within wounds on grapefruit. Phytopathology 90, 1217-1223.

578 McLaughlin, R.J., Wisniewski, M.E., Wilson, C.L., Chalutz, E., 1990. Effects of inoculum
579 concentration and salt solutions on biological control of postharvest diseases of apple
580 with *Candida* sp.. Phytopathology 80, 456-461.

581 Nigro, F., Finetti Sialer, M.M., Gallitelli, D., 1999. Transformation of *Metschnikowia*
582 *pulcherrima* 320, biocontrol agent of storage rot, with the green fluorescent protein gene.
583 J. Plant Pathol. 81, 3, 205-208.

584 Piano, S., Cerchio, F., Migheli, Q., Gullino, M.L., 1998. Effetto di diverse sostanze
585 sull'attività dell'antagonista *Metschnikowia pulcherrima* 4.4 contro *Botrytis cinerea* e
586 sulla sua sopravvivenza sulle mele. Atti Giornate Fitopatologiche, 3-7 maggio 1998, 495-
587 500.

588 Piano, S., Neyrotti, V., Migheli, Q., Gullino, M.L., 1997. Characterization of the biocontrol
589 capability of *Metschnikowia pulcherrima* against *Botrytis* postharvest rot of apple.
590 Postharvest Biol. Technol. 11, 131-140.

591 Pusey, P.L., 1994. Enhancement of biocontrol agents for postharvest diseases and their
592 integration with other control strategies. In: Wilson, C.L., Wisniewski, M.E. (Eds.),
593 Biological Control of Postharvest Diseases. Theory and Practice. CRC Press, Boca
594 Raton, WV, USA, pp. 77-88.

595 Pusey, P.L., Wilson, C.L., Hotchkiss, M.W., Franklin, J.D., 1986. Compatibility of *Bacillus*

596 *subtilis* for postharvest control of peach brown with commercial fruit waxes, dicloran and
597 cold-storage conditions. Plant Dis. 70, 622-626.

598 Rodov, V., Ben-Yehoshua, S., D'hallewin, G., Castina, T., Farg, D., 1994. Accumulation of
599 the phytoalexins, scoparone and scopoletine in citrus fruit subjected to various
600 postharvest treatments. Int. Symp. Natural Phenols in Plant Resistance, Acta Hortic. 381,
601 517.

602 Romano, M.L., Gullino, M.L., Garibaldi, A., 1983. Evaluation of the sensitivity to several
603 fungicides of post-harvest pathogens in North-western Italy. Mede. Fac. Landoubouww.
604 Gent 48, 591-602.

605 Schena, L., Ippolito, A., Zahavi, T., Cohen, L., Droby, S., 2000. Molecular approaches to
606 assist the screening and monitoring of postharvest biocontrol yeasts. Eur. J. Plant Pathol.
607 106, 681-691.

608 Schena, L., Ippolito, A., Zahavi, T., Cohen, L., Nigro, F., Droby, S., 1999. Genetic diversity
609 and biocontrol activity of *Aureobasidium pullulans* isolates against postharvest rots.
610 Postharvest Biol. Technol. 17, 189-199.

611 Smilanick, J.L., Denis-Arrue, R., 1992. Control of green mold of lemon with *Pseudomonas*
612 species. Plant Dis. 76, 481-485.

613 Spadaro, D., Piano, S., Duverney, C., Gullino, M.L., 2001b. Use of microorganisms, heat
614 treatment, and natural compounds against Botrytis rot on apple. Proc. 2nd International
615 Conference on the alternative control methods against plant pests and diseases, Lille,
616 France, 4-7 marzo 2002, in press.

617 Spadaro, D., Vola, R., Piano, S., Gullino, M.L., 2001a. Mechanisms of action and efficacy of
618 four isolates of the yeast *Metschnikowia pulcherrima* active against postharvest
619 pathogens on apples. Postharvest Biol. Technol. 15, in press.

620 Spotts, R.A., Cervantes, L.A., 1986. Population, pathogenicity, and benomyl resistance of

621 *Botrytis* spp., *Penicillium* spp., and *Mucor piriformis* in packinghouses. Plant Dis. 70,
622 106-108.

623 Teixido, N., Usall, J., Palou, L., Asensio, A., Nunes, C., Vinas, I., 2001. Improving control of
624 green and blue molds of oranges by combining *Pantoea agglomerans* (CPA-2) and
625 sodium bicarbonate. Eur. J. Plant Pathol. 107, 7, 685-694.

626 Teixido, N., Vinas, I., Usall, J., Magan, N., 1998. Control of blue mold of apples by
627 preharvest application of *Candida sake* grown in media with different water activity.
628 Phytopathology 88, 960-964.

629 Vos, P., Hogers, R., Bleeker, M., Reijans, M., Van De Lee, T., Hornes, M., Frijters, A., Pot,
630 J., Peleman, J., Kuiper, M., Zabeau, M., 1995. AFLP: a new technique for DNA
631 fingerprinting. Nucleic Acids Res. 23, 21, 4407-4414.

632 Wilson, C.L., Wisniewski, M.E., Droby S., Chalutz E., 1993. A selection strategy for
633 microbial antagonists to control postharvest disease of fruits and vegetables. Scientia
634 Hort. 53, 183-189.

635 Wilson, C.L., Wisniewski, M.E., 1994. Biological Control of Postharvest Diseases. Theory
636 and Practice. CRC Press, Boca Raton, USA.

637 Wisniewski, M.E., Biles, C., Droby, S., Mclaughlin, R., Wilson, C.L., Chalutz, E., 1991.
638 Mode of action of the postharvest biocontrol yeast, *Pichia guilliermondii*. I.
639 Characterization of attachment to *Botrytis cinerea*. Physiol. Mol. Plant P. 39, 245-258.

640 Wisniewski, M.E., Wilson, C.L., Hershherger, W., 1989. Characterization of inhibition of
641 *Rhizopus stolonifer* germination and growth by *Enterobacter cloacae*. Can. J. Bot. 67,
642 2317-2323.