AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Self-Renewal and Multipotency Coexist in a Long-Term Cultured Adult Rat Dental Pulp Stem Cell Line: An Exception to the Rule?

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/109817> since 2022-01-12T14:09:27Z

Published version:

DOI:10.1089/scd.2012.0141

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Sprio AE; Di Scipio F; Raimondo S; Salamone P; Pagliari F; Pagliari S;
Folino A; Forte G; Geuna S; Di Nardo P; Berta GN. Self-Renewal and
Multipotency Coexist in a Long-Term Cultured Adult Rat Dental Pulp Stem
Cell Line: An Exception to the Rule?. STEM CELLS AND
DEVELOPMENT. 21 (18) pp: 3278-3288.
DOI: 10.1089/scd.2012.0141

The publisher's version is available at:

<http://online.liebertpub.com/doi/abs/10.1089/scd.2012.0141>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/109817>

Self-renewal and multipotency coexist in a long-term cultured adult rat dental pulp stem cell line: an exception to the rule?

A.E. Sprio¹, F. Di Scipio¹, S. Raimondo^{1,2}, P. Salamone¹, F. Pagliari^{3,4}, S. Pagliari⁵, A. Folino¹, G. Forte⁵, S. Geuna^{1,2}, P. Di Nardo^{3,4,6,#}, G.N. Berta^{1,6,#}

Affiliation:

¹University of Turin, Department of Clinical and Biological Sciences, 10043, Orbassano, Turin, Italy,

²University of Turin, Neuroscience Institute Cavalieri Ottolenghi, 10043, Orbassano, Turin, Italy

³University of Rome "Tor Vergata", Department of Internal Medicine, Laboratory of Molecular and Cellular Cardiology, 00133, Rome, Italy

⁴BioLink Institute, Link Campus University, Rome, Italy

⁵National Institute for Materials Science (NIMS), International Research Center for Materials Nanoarchitectonics (MANA), Tsukuba, Ibaraki 305–0044, Japan

⁶National Institute of Cardiovascular Research, 40126, Bologna, Italy (INRC)

[#]These Authors contributed equally to this work

Running title: Adult stem cell model characterization

keywords: dental pulp stem cell; neural crest; cell model; long-term culture

Address correspondence to:

Giovanni N. Berta, PhD, DMD

Department of clinical and Biological Sciences

University of Turin,

reg Gonzole 10, 10043 Orbassano (TO)

Italy

E-mail: giovanni.berta@unito.it

Summary

The stemness state is characterized by self-renewal and differentiation properties. However, stem cells are not able to preserve these characteristics in long-term culture because of the intrinsic fragility of their phenotype easily undergoing senescence or neoplastic transformation. Furthermore, although isolated from the same original tissue using similar protocols, adult stem cells can display dissimilar phenotypes and important cell clone/species contamination. Finally, the lack of a clear standardization contributes to complicate the comprehension about the stemness condition. In this context, cell lines displaying a particularly stable phenotype must be identified in order to define one or multiple benchmarks against which other stem cell lines could be reliably assessed. The present paper demonstrates that it is possible to isolate from the rat dental pulp a stem cell line (MUR-1) that does not display neoplastic transformation in long-term culture. MUR-1 cells stably express a broad range of stemness markers and are able to differentiate into adipogenic, osteogenic, chondrogenic, neurogenic and cardiomyogenic lineages independently of the culture passages. Moreover, serial in vitro passages have not changed their immunophenotype, proliferation capacity or differentiation potential. The uniqueness of these characteristics candidates MUR-1 as a model to reliably improve the understanding of the mechanisms governing the stem cell fate in the same as well as in other stem cell populations.

Introduction

The interest around stem cells has increased over the years proportionally to new findings. Although postulated about one century ago [1], stem cells were revealed only in the 1963 [2]. The discovery of “*cluster of hæmatopoietic cells*” within the bone marrow paved the way to the first stem cells transplant [3] starting the regenerative medicine era.

In the last decade, stem cells have been identified in almost all body tissues [4], from which they have been isolated on the basis of their intrinsic characteristics of self-renewal and potency. Among others, adult mesenchymal stem cells (MSC) have attracted great attention for their ability to generate different cell lineages [5-7], fueling the assumption that they could be the panacea to regenerate all injured tissues. Nevertheless, the clinical worthiness of MSC to repair injured organs remains a chimera. In fact, owing to the poor knowledge of some MSC intrinsic features together with the inadequacy of current handling and engraftment protocols, clinically reliable cell treatments are still lacking.

In mammals, MSC are segregated in particular tiny regions (niches) in which the microenvironment governs their quiescence, self-renewal and/or commitment [8]. Each niche is thus responsible for MSC features and behavior in terms of growth and differentiation potential. In bone marrow, MSC represent only a small percentage (0.5-10% depending on the anatomical site) of the total cell population [8-10] and are often hardly identifiable for their morphological and biochemical camouflage with the differentiated surroundings [8,11]. Moreover, even if correctly isolated, MSC are not able to retain their stemness condition when cultured in vitro for long-term [12], very likely because of the lack of niche control. Consistently, stem cell primary cultures undergo senescence and their capabilities of self-renewal and multi-potency are faded by an innate drift towards commitment linked to in vitro conditions [13,14]. On the other hand, MSC, when apparently not affected by senescence after prolonged in vitro culture, may acquire cytogenetic abnormalities prompting unwanted neoplastic transformations [15].

Furthermore, although isolated from the same tissue following identical protocols, primary MSC cultures can display dissimilar phenotypes, corresponding to different stages of cell immaturity along the differentiation pathway [16-18] and/or important contamination problems [13,19-24]. Consistently, similarly to the Hindu “Blind men and an elephant” tale [25], different investigators have described the same stem cell population using a plethora of often non-specific “Cluster of Differentiations” markers. As result, the lack of standardized procedures in stem cell research [26] confuses and complicates the comprehension about the stemness condition, consequently frustrating the regenerative

medicine philosophy itself. On these evidences, it is of paramount importance to identify populations that can be used as benchmarks to validate further advancements in stem cell research. In this context, the aim of the present study has been to verify the phenotypic stability in long-term culture and the capability to differentiate upon specific stimuli of a stem cell population (MUR-1, from latin Mòrsus/Múrsus = muzzle) isolated from the dental pulp of rat continuously growing incisors. Indeed, the results have shown that MUR-1 are adult mesenchymal stem cells characterized by geno/phenotypical stability and non-tumorigenicity in long-term culture. These characteristics candidate them as a benchmark in stemness/differentiation studies as well as to assess the potency of other stem cell populations.

Materials and Methods

Cell Isolation and Purification

Dental pulp stem cells (DPSC) were isolated from mandible incisive teeth of five months old male Wistar rats (Charles River, Monza, Italy). The dental pulp was gently removed through the apical foramen. The resulting tissue was washed twice in phosphate buffered saline (PBS) and centrifuged for 5 min at 1000 rpm; subsequently, the dental pulp was digested, washed, and seeded as previously described [27]. After 24 h, non-adherent cells were removed while the remaining cells were cultured in RPMI-1640 supplemented with 10% fetal calf serum (FCS), 100 U/ml penicillin G, 40 µg/ml gentamicin sulfate and 2.5 µg/ml amphotericin B (all from Sigma-Aldrich, St Louis, MO, USA) at 37°C in a humidified 5% CO₂ atmosphere. The final clone selection was conducted by serial dilutions in 96-well plates. Only wells enclosing homogeneous population were considered for further expansion. Among them, one clone (named MUR-1) was chosen for subsequent experimentations.

MUR-1 Culture Conditions and Cryopreservation

Cells were cultured in 75 cm² TPP[®] flasks (TPP AG, Switzerland) as above described, without any feeder-layer. Cells were harvested twice a week with 0.25% trypsin solution (Sigma-Aldrich). Standard 1:4 dilution ratio was used for cell expansion while MUR-1 cryopreservation was performed in the same culture medium added of 10% DMSO.

FACS Analysis, Cell Growth Kinetics, and Colony-Forming Units Fibroblastoid Assay

Cell volume and morphological complexity was assessed by a FACScalibur (Beckton Dickinson Italia SpA, Buccinasco, Italy) instrument, after suspension at the density of 5x10⁵ cells/ml. Doubling time (DT) of cell line was calculated as previously described [28] in cells (1x10⁵) seeded into 25 cm² flasks and cultured in complete medium. The clonogenicity of DPSC was determined by means of the Colony-Forming Units Fibroblastoid (CFU-F). Cells were seeded in 25 cm² flasks at the density of 2, 5, or 10 cells/cm². After two weeks, the adherent cells were toluidine blue-stained, and macroscopic colonies were counted. All experiments were performed in triplicate.

Light and Electron Microscopy

MUR-1 cell morphology in adherent conditions was investigated in cells cultured on coverslips (DelChimica, Napoli, Italy), stained with haematoxylin/eosin, and observed by light microscopy. For electron microscopy, cells were processed as previously described [29].

Karyotype Analysis

Chromosome analysis was performed every 6 months of culture adopting a conventional G-banding karyotype protocol. Briefly, cells undergoing active division were blocked at metaphase by colchicine and actinomycin D, detached from the growth surface by trypsin–EDTA, and subsequently swollen by exposure to hypotonic KCl solution. Cells were then fixed in methanol/acetic acid solution, glass-mounted and finally metaphase dishes were identified by light microscope and photographed. Chromosomes were identified according to the Committee for a Standardized Karyotype of *Rattus norvegicus* [30].

mRNA and Protein Expression Analysis

Total RNA was extracted with TRIreagent solution (Sigma-Aldrich) according to manufacturer's instructions. One microgram of RNA was purified from contaminating DNA through a Deoxyribonuclease I kit and retrotranscribed with Random Hexamer Primer according to M-MuLV Reverse Transcriptase manufacturer's instructions (all reagents were from Fermentas International Inc., Burlington, Canada). PCR reactions were carried out using 2.5 µl cDNA, 2.5 µl 10X PCR Buffer II (Roche Applied Science, Indianapolis, USA), 1.5 µl 25 mM MgCl₂, 0.5 µl 10 mM dNTP (Fermentas), 0.75 µl of each specific 10 µM primer (Sigma-Genosys), and 0.05 U/µl Taq DNA Polymerase (Fermentas) to a final volume of 25 µl. Specific primer sequences are summarized in Table 1. Amplification products were separated by agarose gel electrophoresis. DNA bands were stained with ethidium bromide and digitally captured by Kodak ImageStation. As positive controls, cDNAs extracted from brain (for Thy1, cKit, Oct4, Stella, SSEA1, Tubb3, and Sox2), foetal brain (MEF2C), heart (GATA4, ACTC1, CX43, and ADRB1), pancreas (Islet1), lung (ADRB2, Fragilis1, ALPL, and HLA-DR), and liver (HDAC6) were used.

Protein expression was assessed by immunocytochemistry analysis. MUR-1 cells were cultured on poly-D-lysine-coated coverslips and then incubated with specific primary antibodies overnight at 4°C and then with appropriate secondary antibodies for 1 h at room temperature. Primary antibodies are reported in Table 2. Image acquisition of CD90,

CD117, YAP1, vimentin, Nkx2.5, CD45, and CD34 expression was carried out with a LSM 510 confocal laser microscopy system (Zeiss, Jena, Germany), while FLK1, alpha smooth muscle actin (α SMA), cMyc, CD105, and GATA-4 expression was visualized under a Leica DRMB microscope equipped with a digital camera.

Differentiation Culture Conditions

The adipogenic differentiation was induced by growing MUR-1 cells in RPMI-1640 containing 10% FBS, 1.7 μ M insulin, 1 μ M dexamethasone, and 0.5 mM methylisobutylxanthine (Sigma-Aldrich) [31]. After 5 days of induction, cells were fixed with 4% paraformaldehyde for 20 min, washed with PBS, incubated with Nile Red (Sigma-Aldrich) for 10 min to emphasize lipid vesicles occurrence [32] and then observed by confocal laser microscope. For electron microscopy, differentiated cells were processed as previously described [29].

Osteogenic differentiation was induced in MUR-1 cells cultured in complete medium supplemented with 10 mM β -glycerophosphate, 0.05 mM ascorbic acid, and 100 nM dexamethasone [33]. After 3 weeks, osteogenic differentiation was evaluated by alkaline phosphatase staining kit (BioOptica Milano SpA, Milano, Italy).

MUR-1 chondrogenic differentiation was induced by complete medium supplemented with 10 ng/ml TGF- β 3, 0.025 mM ascorbic acid, and 100 nM dexamethasone [34]. After 3 weeks, differentiated cells were highlighted with toluidine blue staining as described elsewhere [35].

MUR-1 neuro-glial differentiation was induced by pre-treating cells with 1 mM β -mercaptoethanol in complete medium for 24 h and then with 35 ng/ml retinoic acid for 72 h. Subsequently, MUR-1 cells were grown in complete medium supplemented with 5 ng/ml PDGF (Platelet-derived Growth Factor), 10 ng/ml bFGF (basic Fibroblast Growth Factor), 252 ng/ml GGF (Glial Growth Factor), and 14 μ M Forskolin for 2 weeks [36]. The induction of neuro-glial phenotype was confirmed by specific immunostaining for neuron growth-associated protein 43 (GAP43 - Goat, 1:100, Santa Cruz Biotechnology, Santa Cruz, CA, USA), glial fibrillary acidic protein (GFAP - Mouse, 1:50, DakoCytomation, Dako Italia SpA, Milano, Italy) and S100 protein (Rabbit, 1:400, Sigma-Aldrich). MUR-1 cells without differentiation factors were prepared in the same conditions and used as negative controls. Myocardial differentiation was induced through direct co-culture of MUR-1 with murine neonatal cardiomyocytes in complete medium for 7 days. Neonatal cardiomyocytes were obtained as described elsewhere [37] and plated at the concentration of 5×10^4 cells/cm²

onto fibronectin (2 µg/ml)-laminin (0.2%) pre-coated glass chamber slides (BD Biosciences, San Jose, CA). Twenty-four hours later, MUR-1 cells, previously labeled with 1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate (Dil, Vybrant™, Molecular Probes, Life Technologies Italia, Monza, Italy), were seeded onto cardiomyocytes at a 1:5 ratio. MUR-1 without cardiac cells and viceversa were prepared in the same conditions and used as controls. The induction of myocardial differentiation was assessed by evaluating sarcomeric α -actinin (Mouse, 1:700, Sigma-Aldrich), connexin 43 (CX43) (Rabbit, 1:100, Sigma-Aldrich) and Troponin I (Rabbit, 1:50, Santa Cruz Biotechnology) expression.

Colony Formation and Tumorigenicity Assay

Soft-agar assay was carried out to evaluate the possibility that MUR-1 cells could undergo anchorage-independent growth. Briefly, 1×10^5 MUR-1 cells were seeded into 10 mm Petri dish in 0.3% agar-complete medium. Cells were cultured for 21 days and then colony formation assessed by means of light microscopy. MUR-1 tumorigenicity was evaluated in vivo by injecting 1.0 or 10×10^6 cells between the shoulder blades in six male Wistar rats randomly divided into two groups. Animals were followed for 60 days, evaluating whether masses were developing in the injection area. At 10, 30 and 60 days, one rat for each group was anesthetized and sacrificed. Tissues surrounding the injection area were excised and processed for histological analysis by standard haematoxylin/eosin stain. The whole experimentation was performed according to the Ethic Committee of the University of Turin, Italy.

Statistical Analysis

DT and CFU-F were expressed as arithmetical mean $\pm$ standard deviation. DT results were analyzed with a two-tailed paired t-test, whereas one-way ANOVA and Newman-Keuls Multiple Comparison Test (for post-ANOVA comparisons) were used to evaluate the statistical significance of the CFU-F differences. All analyses were carried out with GraphPad Prism version 5.00 (GraphPad Software, San Diego, CA, USA, www.graphpad.com), with $P < 0.05$ as the significance cut-off.

Results

CFU-F, Growth Kinetic, Karyotype and Morphological Characteristics

Under light microscopy (Fig. 1A), adherent MUR-1 cells showed a fibroblastoid shape; after haematoxylin/eosin staining (Fig. 1B), they displayed an irregular and eosinophilic cytoplasm with central round nucleus. Electron microscopy revealed a pale, eccentric and irregularly shaped nucleus with one or more nucleoli located near the MUR-1 perinuclear cisternae (Fig. 1C,D). Chromatin was spread throughout the nucleus except for a thin dense layer located immediately inside the perinuclear cisternae (Fig. 1D,E). The cytoplasm was rich in mitochondria and ribosomes predominantly associated with endoplasmic reticulum (Fig. 1F,G). Yet, ultrastructural observation showed that the irregularities of the plasma membrane seen by light microscopy were due to small pseudopodia located all around the cells (Fig. 1C,H).

MUR-1 cells were characterized by an estimated doubling time of 25.4 ± 2.2 and 23.2 ± 1.8 h at the exponential growth phase, when assessed after thirty (P30) and two-hundred (P200) culturing passages, respectively (P30 vs. P200, $P=0.120$). The frequency of CFU-F was proportional to the cell seeding density ($P=0.0028$ and $P=0.001$ at P30 and P200, respectively). The CFU-F corresponding to 2, 5, and 10 cells/cm² was 73.7 ± 14.4 , 179.7 ± 23.7 , and 250.0 ± 55.9 for MUR-1 at P30, and 77.0 ± 10.5 , 147.3 ± 26.8 , and 256.7 ± 43.8 for MUR-1 at P200, respectively (Fig. 2A). Results did not show any significant differences in terms of CFU-F between P30 and P200 passages. As evinced by FACS analysis, MUR-1 was a population with homogeneous dimensions and morphological complexity (Fig. 2B). During the entire cell culture time, G-banding cytogenetic analysis of MUR-1 cells did not reveal macroscopic chromosome alterations, if compared with the reference rat karyotype (Fig. 2C). In addition, cryopreservation did not affect MUR-1 characteristics, in accord to what already described [38].

MUR-1 Gene and Protein Expression

Analysis of mRNA expression (Fig. 3) revealed that MUR-1 cells transcribed several genes related to the embryonic/pluripotent undifferentiated status, as the transcription factor OCT4 (POU5F1), the membrane glycoprotein Thy-1 (CD90), the tyrosine-kinase receptor c-Kit (CD117), the developmental pluripotency-associated protein 3 (Stella), the stage-specific embryonic antigen 1 (SSEA1), the interferon-induced transmembrane protein 1

(Fragilis 1), and the histone deacetylase 6 (HDAC6), but not the SRY-box2 transcription factor (SOX2). Contextually, commitment transcripts were present: MUR-1 cells expressed mRNAs encoding for the transcription factors GATA4 and MEF2C, as well as the cytoskeleton components ACTC1 and tubulin β 3, the gap-junction CX43, and the receptor ADRB2, but they did not transcribe those for the ISL-1 transcription factor (islet-1), ADRB1 receptor, the non tissue specific alkaline phosphatase (ALPL), and the major histocompatibility complex HLA-DR.

At the protein level (Fig. 4), the expression of stem cell antigens CD90 and CD117 was further confirmed. Consistently, the presence of c-MYC transcription factor, TGF β RIII receptor Endoglin (CD105), FLK1 receptor (VEGFR2 or CD309), CX43, important transcription factors (NKX2.5, GATA4, and YAP1), intermediate filament vimentin, and α smooth muscle actin (α SMA) was highlighted. MUR-1 cells did not express typical blood lineage commitment markers, namely CD34 and CD45.

Plasticity of MUR-1 Cells

Adequately treated MUR-1 cells have been able to successfully differentiate in precursors of different histotypes, also after prolonged culture (P200). Following adipogenic stimuli, the presence of lipid-rich cells was detectable with the lipophilic Nile Red dye (Fig. 5A): indeed, ultrastructural observation indicated the presence of lipid droplets inside the cells (Fig. 5B). If compared with untreated MUR-1 cells (Fig. 5C), chondrogenic medium led to the formation of round-shaped cell clusters characterized by a basophilic cytoplasm (Fig. 5D). Similarly, osteogenic stimulation induced the aggregation of clusters of alkaline phosphatase-stained round-shaped cells surrounded by a dense extracellular matrix (Fig. 5E,F).

MUR-1 cells were also able to trans-differentiate into other cell types. If cultured with appropriate growth factors (see Materials and Methods section), MUR-1 may acquire neuronal morphologies including bipolar one (Fig. 5G), and express neuro-glial markers, such as GAP43 (Fig. 5H), GFAP (Fig. 5I), and S100 (Fig. 5J), within 2 weeks of incubation.

The co-culture of MUR-1 with neonatal cardiomyocytes induced cardiac differentiation as demonstrated by the expression of troponin I (Fig. 5K,L), CX43 (Fig. 5M,N) and α sarcomeric actinin (Fig. 5O,P), assembled in sarcomeric bands within the cells.

Soft-Agar and Isograft Assay

After more than 6 months of continued culture, MUR-1 cells were assessed for their tumorigenic potential. Both in vitro and in vivo tests showed no cells colony and mass development in the soft-agar test and in subcutaneously injected tissues, respectively. Also autoptic analysis of isograft areas of injection did not reveal the presence of any inflammatory infiltrates (data not shown).

Discussion

Dental pulp holds promise to be an interesting source of different adult stem cell populations generally characterized by an enhanced differentiation potential [39-41], although the gene/antigen expression profile is comparable to bone marrow stem cells [27]. Nevertheless, the actual potential of DPSC has been underestimated, since investigations have predominantly been focused on obtaining cells for in vivo dental/calcifying tissue recovery.

The present study demonstrates that, among DPSC, it is possible to isolate a cell line (MUR-1) displaying uncommon properties and behavior, even though morphological features strictly resemble those of MSC [27]. Ultrastructural analysis of MUR-1 cells reveals a finely dispersed euchromatin, presence of numerous mitochondria featuring tightly pleated cristae, and several ribosomes associated to the endoplasmic reticulum, clues of easy access to transcription factors, enhanced energy production and protein synthesis, respectively. Furthermore, MUR-1 cells transcribe and translate the classical CD90⁺/CD117⁺/CD34⁻/CD45⁻ profile without commitment to blood lineages, typically depicting the undifferentiated mesenchymal condition [7]. This expression pattern associates with that of both c-Myc and Oct4, which are essential to induce self-renewal in embryonic stem cells and are pivotal to sustain pluripotency [42]. Intriguingly, MUR-1 cells also express Stella antigen, further clue of self-renewal and multi/pluripotency. In fact, Stella has been observed, up to now, only in the embryo inner mass cells and in the germ cells line [43], where it acts as protector of the DNA demethylation, as demonstrated by its involvement in epigenetic reprogramming [44]. In addition, MUR-1 cells express the embryonic SSEA1 marker, which is also specific for neural crest cells [45,46], and Frigilis 1, which is mainly present in the mesoderm and the pharyngeal arch during the embryo development [47]. Finally, the expression of YAP1, marker of epithelium-mesenchyma transition [48], could represent the remnant of the ecto-mesenchymalization process underlying the dental pulp development. Indeed, the dental pulp originates from an unique mesenchyma-epithelium cross-talk, in which the epithelium of the first pharyngeal arch interacts with the neural crest, considered as the mesenchymal counterpart [49-51].

These evidences indicate that MUR-1 are characterized by a very early differentiation status, as also confirmed by their high proliferation rate and clonogenic potential. Notably, both phenomena remain unchanged up to the 200th culture passage. Since cells cultured in vitro for extended periods could escape senescence when affected by genetic aberrations and neoplastic transformation, MUR-1 cells have been repeatedly karyotyped

without any evidence of macroscopic chromosomal alterations. Furthermore, they were neither able to grow in vitro without adhesion-support nor to provoke mass development after injection in living experimental animals. Taken together, these evidences indicate that long-term cultured MUR-1 cells cannot be considered cancerous.

Beside these characteristics, MUR-1 cells translate both CD105 and Flk1 antigens strengthening the mesenchymal feature, though they are also typical endothelial markers [52,53]. Notably, albeit being negative for ADRB1, MUR-1 cells display a cardiac/vascular-like CD117⁺/Flk1⁺/NKX-2.5⁺/MEF2C⁺ profile, together with the low expression of endoderm gene GATA4. Such a pattern concurs to postulate a parenting with those cells driving the cardiogenesis during the embryo development [54,55]. In addition, CX43 is functionally organized in cell membranes, likewise the high motility neural crest cell subpopulation that arises in the endothelium of the aortic arch arteries and the septum between the aorta and the pulmonary artery [56], and portions of the cardiac conduction system [57]. These data suggest that MUR-1 cells could share a neural crest ancestral progenitor, while prone to differentiate to cardiomyocytes. Additionally, MUR-1 cells express ADRB2 that is involved in both adipo- and osteogenic differentiation [58,59], and vimentin, a marker of early neural phenotype [60], but also generally related to the mesenchymal status [61]. Finally, markers of differentiations, such as non-tissue-specific alkaline phosphatase or HLA-DR, have not been detected.

This broad profile of MUR-1 gene and antigen expression is reflected by their potency. Indeed, MUR-1 cells are able to adopt an adipocyte, chondrocyte and osteocyte phenotype, as evinced by the intracellular lipid vesicles development, the basophilic cytoplasm and alkaline phosphatase activity, respectively. These capabilities are substantially equivalent to those of stem cells from bone marrow. Furthermore, besides the above mentioned differentiating capabilities, MUR-1 cells can generate neuronal and glial precursors expressing typical markers, such as GFAP, GAP43 and S100, which are involved in neurite formation and elongation [62,63]. Finally, they can differentiate to cardiomyocytes expressing troponin I and α sarcomeric actinin in properly assembled Z-disks; conversely, CX43 expression and location remain as in undifferentiated MUR-1 cells. Notably, the MUR-1 extended differentiating potential remains unchanged independently of the cell passage and is not significantly affected by supplement/feeder-layer free culture conditions, such as when they are cultured in low-antioxidant high-glucose RPMI-1640. This is a very peculiar feature of MUR-1 cells, since the increment of

the Reactive Oxygen Species (ROS) level within stem cells is believed to be inversely correlated with the maintenance of their stemness state [64].

It can be speculated that MUR-1 peculiar characteristics could be related to the continuous growth of Myomorpha teeth during the entire animal life; however, adult stem cells deriving from other continuously growing tissues do not display a similar behavior. For example, when bone marrow-MSC are cultured in vitro, they lose the capability to self renew within 30-50 population-doublings [65,66].

All these observations indicate that the MUR-1 cells phenotype can be properly maintained even after isolation from their native niches. This is in apparent contrast with the notion that niche is crucial in regulating stem cells proliferation, self-renewal and differentiation potential [67]. It can be conjectured that, under some respect, culture conditions mimic the native niche environment.

In conclusion, MUR-1 is a cell population phenotypically very stable in conventional culture conditions and, at the same time, endowed with a very extensive differentiating potential that can be easily activated with specific stimuli. This makes MUR-1 a very reliable model to study the mechanisms regulating stemness state and to test biologically active substances of potential pharmacological interest. Furthermore, MUR-1 could represent a benchmark to assess the characteristics and potential of many other stem cell populations, whose phenotypic instability has so far hampered to deepen the knowledge about their behavior.

Acknowledgement

We are thankful to Dr. M. Pautasso and Mrs. L.M. Marinos for their technical supports. We thank Prof. A Ceccarelli for providing critical assistance. F. Di Scipio is supported by a grant of Compagnia di San Paolo, Turin, Italy.

Author Disclosure Statement

None of the authors has any disclosure to declare. No competing financial interests exist.

References

1. Maximow A. (1909). Der Lymphozyt als gemeinsame Stammzelle der verschiedenen Blutelemente in der embryonalen Entwicklung und im postfetalen Leben der Säugetiere. *Folia Haematologica* 125-134.
2. Becker AJ, EA McCulloch and JE Till. (1963). Cytological demonstration of the clonal nature of spleen colonies derived from transplanted mouse marrow cells. *Nature* 197:452-454.
3. Bach FH, RJ Albertini, P Joo, JL Anderson and MM Bortin. (1968). Bone-marrow transplantation in a patient with the Wiskott-Aldrich syndrome. *Lancet* 2:1364-1366.
4. Weissman IL. (2000). Translating Stem and Progenitor Cell Biology to the Clinic: Barriers and Opportunities. *Science* 287:1442-1446.
5. Baksh D, L Song and RS Tuan. (2004). Adult mesenchymal stem cells: characterization, differentiation, and application in cell and gene therapy. *J Cell Mol Med* 8:301-316.
6. Beltrami AP, D Cesselli, N Bergamin, P Marcon, S Rigo, E Puppato, F D'Aurizio, R Verardo, S Piazza, A Pignatelli, A Poz, U Baccarani, D Damiani, R Fanin, L Mariuzzi, N Finato, P Masolini, S Burelli, O Belluzzi, C Schneider and CA Beltrami. (2007). Multipotent cells can be generated in vitro from several adult human organs (heart, liver, and bone marrow). *Blood* 110:3438-3446.
7. Dominici M, K Le Blanc, I Mueller, I Slaper-Cortenbach, F Marini, D Krause, R Deans, A Keating, D Prockop and E Horwitz. (2006). Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 8:315-317.
8. Orford KW and DT Scadden. (2008). Deconstructing stem cell self-renewal: genetic insights into cell-cycle regulation. *Nat Rev Genet.* 9:115-128.
9. Gordon JI, GH Schmidt and KA Roth. (1992). Studies of intestinal stem cells using normal, chimeric, and transgenic mice. *FASEB J* 6:3039-3050.
10. Potten CS and RJ Morris. (1988). Epithelial stem cells in vivo. *J Cell Sci Suppl* 10:45-62.
11. Fuchs E and JA Segre. (2000). Stem cells: a new lease on life. *Cell* 100:143-155.
12. Vacanti V, E Kong, G Suzuki, K Sato, JM Canty and T Lee. (2005). Phenotypic changes of adult porcine mesenchymal stem cells induced by prolonged passaging in culture. *J Cell Physiol* 205:194-201.
13. Honda MJ, M Imaizumi, S Tsuchiya and C Morscheck. (2010). Dental follicle stem cells and tissue engineering. *J Oral Sci* 52:541-552.
14. Stenderup K, J Justesen, C Clausen and M Kassem. (2003). Aging is associated with decreased maximal life span and accelerated senescence of bone marrow stromal cells. *Bone* 33:919-926.
15. Tolar J, AJ Nauta, MJ Osborn, A Panoskaltsis Mortari, RT McElmurry, S Bell, L Xia, N Zhou, M Riddle, TM Schroeder, JJ Westendorf, RS McIvor, PC Hogendoorn, K Szuhai, L Oseth, B Hirsch, SR Yant, MA Kay, A Peister, DJ Prockop, WE Fibbe and BR Blazar. (2007). Sarcoma derived from cultured mesenchymal stem cells. *Stem Cells* 25:371-379.
16. Itzhaki-Alfia A, J Leor, E Raanani, L Sternik, D Spiegelstein, S Netser, R Holbova, M Pevsner-Fischer, J Lavee and IM Barbash. (2009). Patient characteristics and cell source determine the number of isolated human cardiac progenitor cells. *Circulation* 120:2559-2566.
17. Gronthos S, AC Zannettino, SE Graves, S Ohta, SJ Hay and PJ Simmons. (1999). Differential cell surface expression of the STRO-1 and alkaline phosphatase antigens on discrete developmental stages in primary cultures of human bone cells. *J Bone Miner Res* 14:47-56.
18. Stewart K, S Walsh, J Screen, CM Jefferiss, J Chainey, GR Jordan and JN Beresford. (1999). Further characterization of cells expressing STRO-1 in cultures of adult human bone marrow stromal cells. *J Bone Miner Res* 14:1345-1356.

19. Digirolamo CM, D Stokes, D Colter, DG Phinney, R Class and DJ Prockop. (1999). Propagation and senescence of human marrow stromal cells in culture: a simple colony-forming assay identifies samples with the greatest potential to propagate and differentiate. *Br J Haematol* 107:275-281.
20. Rossi DJ, CH Jamieson and IL Weissman. (2008). Stems cells and the pathways to aging and cancer. *Cell* 132:681-696.
21. Kerkis I, A Kerkis, D Dozortsev, GC Stukart-Parsons, SM Gomes Massironi, LV Pereira, AI Caplan and HF Cerruti. (2006). Isolation and characterization of a population of immature dental pulp stem cells expressing OCT-4 and other embryonic stem cell markers. *Cells Tissues Organs* 184:105-116.
22. Sloan AJ and RJ Waddington. (2009). Dental pulp stem cells: what, where, how? *Int J Paediatr Dent* 19:61-70.
23. Waddington RJ, SJ Youde, CP Lee and AJ Sloan. (2009). Isolation of distinct progenitor stem cell populations from dental pulp. *Cells Tissues Organs* 189:268-274.
24. Colter DC, R Class, CM DiGirolamo and DJ Prockop. (2000). Rapid expansion of recycling stem cells in cultures of plastic-adherent cells from human bone marrow. *Proc Natl Acad Sci U S A* 97:3213-3218.
25. Saxe JG, ed. *The poems of John Godfrey Saxe*. (1872). J. Osgood, Boston.
26. Di Nardo P and GC Parker. (2011). Stem cell standardization. *Stem Cells Dev* 20:375-377.
27. Gronthos S, M Mankani, J Brahimi, P Gehron Robey and S Shi. (2000). Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A* 97:13625-13630.
28. Sprio AE, F Di Scipio, P Ceppi, P Salamone, F Di Carlo, GV Scagliotti, M Papotti, A Ceccarelli and GN Berta. (2012). Differentiation-inducing factor 1 enhances 5-fluorouracil action on oral cancer cells inhibiting E2F1 and Thymidylate Synthase mRNAs accumulation. *Cancer Chemother Pharmacol* 69:983-989.
29. Audisio C, S Raimondo, S Nicolino, G Gambarotta, F Di Scipio, L Macrì, F Montarolo, MG Giacobini-Robecchi, P Porporato, N Filigheddu, A Graziani, S Geuna and I Perroteau. (2009). Morphological and biomolecular characterization of the neonatal olfactory bulb ensheathing cell line. *J Neurosci Methods* 185:89-98.
30. Committee for a standardized karyotype of *rattus norvegicus*. (1973). Standard karyotype of the Norway rat, *Rattus norvegicus*. *Cytogenet Cell Genet* 12:199-205.
31. Rim JS, RL Mynatt and B Gawronska-Kozak. (2005). Mesenchymal stem cells from the outer ear: a novel adult stem cell model system for the study of adipogenesis. *FASEB J* 19:1205-1207.
32. Fowler SD and P Greenspan. (1985). Application of Nile red, a fluorescent hydrophobic probe, for the detection of neutral lipid deposits in tissue sections: comparison with oil red O. *J Histochem Cytochem* 33:833-836.
33. Jaiswal N, SE Haynesworth, AI Caplan and SP Bruder. (1997). Osteogenic differentiation of purified, culture-expanded human mesenchymal stem cells in vitro. *J Cell Biochem* 64:295-312.
34. Barry F, RE Boynton, B Liu and JM Murphy. (2001). Chondrogenic differentiation of mesenchymal stem cells from bone marrow: differentiation-dependent gene expression of matrix components. *Exp Cell Res* 268:189-200.
35. Johnstone B, TM Hering, AI Caplan, VM Goldberg and JU Yoo. (1998). In vitro chondrogenesis of bone marrow-derived mesenchymal progenitor cells. *Exp Cell Res* 238:265-272.
36. Kingham PJ, DF Kalbermatten, D Mahay, SJ Armstrong, M Wiberg and G Terenghi. (2007). Adipose-derived stem cells differentiate into a Schwann cell phenotype and promote neurite outgrowth in vitro. *Exp Neurol* 207:267-274.
37. Pagliari S, AC Vilela-Silva, G Forte, F Pagliari, C Mandoli, G Vozzi, S Pietronave, M Prat, S Licocchia, A Ahluwalia, E Traversa, M Minieri and P Di Nardo. (2011). Cooperation of biological and mechanical signals in cardiac progenitor cell differentiation. *Adv Mater* 23:514-518.
38. Papaccio G, A Graziano, R d'Aquino, MF Graziano, G Pirozzi, D Menditti, A De Rosa, F Carinci and G Laino. (2006). Long-term cryopreservation of dental pulp stem cells (SBP-DPSCs) and their differentiated osteoblasts: a cell source for tissue repair. *J Cell Physiol* 208:319-325.
39. Huang GT, S Gronthos and S Shi. (2009). Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. *J Dent Res* 88:792-806.

40. Stevens A, T Zuliani, C Olejnik, H LeRoy, H Obriot, J Kerr-Conte, P Formstecher, Y Bailliez and RR Polakowska. (2008). Human dental pulp stem cells differentiate into neural crest-derived melanocytes and have label-retaining and sphere-forming abilities. *Stem Cells Dev.* 17:1175-1184.
41. Paino F, G Ricci, A De Rosa, R D'Aquino, L Laino, G Pirozzi, V Tirino and G Papaccio. (2010). Ecto-mesenchymal stem cells from dental pulp are committed to differentiate into active melanocytes. *Eur Cell Mater.* 20:295-305.
42. Takahashi K and S Yamanaka. (2006). Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 126:663-676.
43. Hayashi K, SM Chuva de Sousa Lopes, F Tang and MA Surani. (2008). Dynamic Equilibrium and Heterogeneity of Mouse Pluripotent Stem Cells with Distinct Functional and Epigenetic States. *Cell Stem Cell* 3:391-401.
44. Nakamura T, Y Arai, H Umehara, M Masuhara, T Kimura, H Taniguchi, T Sekimoto, M Ikawa, Y Yoneda, M Okabe, S Tanaka, K Shiota and T Nakano. (2007). PGC7/Stella protects against DNA demethylation in early embryogenesis. *Nat Cell Biol* 9:64-71.
45. Sieber-Blum M. (1989). SSEA-1 is a specific marker for the spinal sensory neuron lineage in the quail embryo and in neural crest cell cultures. *Dev Biol* 134:362-375.
46. Liu S, DA Wilcox, M Sieber-Blum and M Wong-Riley. (1990). Developing neural crest cells in culture: correlation of cytochrome oxidase activity with SSEA-1 and dopamine-beta-hydroxylase immunoreactivity. *Brain Res* 535:271-280.
47. Lange UC, M Saitou, PS Western, SC Barton and MA Surani. (2003). The fragilis interferon-inducible gene family of transmembrane proteins is associated with germ cell specification in mice. *BMC Dev Biol* 3:1.
48. Lian I, J Kim, H Okazawa, J Zhao, B Zhao, J Yu, A Chinnaiyan, MA Israel, LS Goldstein, R Abujarour, S Ding and KL Guan. (2010). The role of YAP transcription coactivator in regulating stem cell self-renewal and differentiation. *Genes Dev* 24:1106-1118.
49. Gilbert SF, ed. *Developmental Biology*. (2006). Sinauer Associates, Sunderland, MA.
50. Chai Y, X Jiang, Y Ito, PJ Bringas, J Han, DH Rowitch, P Soriano, AP McMahon and HM Sucov. (2000). Fate of the mammalian cranial neural crest during tooth and mandibular morphogenesis. *Development* 127:1671-1679.
51. Rothová M, J Feng, PT Sharpe, R Peterková and AS Tucker. (2011). Contribution of mesoderm to the developing dental papilla. *Int J Dev Biol* 55:59-64.
52. Jiang Y, BN Jahagirdar, RL Reinhardt, RE Schwartz, CD Keene, XR Ortiz-Gonzalez, M Reyes, T Lenvik, T Lund, M Blackstad, J Du, S Aldrich, A Lisberg, WC Low, DA Largaespada and CM Verfaillie. (2002). Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 418:41-49.
53. Guo H, B Fang, L Liao, Z Zhao, J Liu, H Chen, SH Hsu, Q Cui and RC Zhao. (2003). Hemangioblastic characteristics of fetal bone marrow-derived Flk1(+)CD31(-)CD34(-) cells. *Exp Hematol* 31:650-658.
54. Kattman SJ, ED Adler and GM Keller. (2007). Specification of multipotential cardiovascular progenitor cells during embryonic stem cell differentiation and embryonic development. *Trends Cardiovasc Med* 17:240-246.
55. Wu SM, Y Fujiwara, SM Cibulsky, DE Clapham, CL Lien, TM Schultheiss and SH Orkin. (2006). Developmental origin of a bipotential myocardial and smooth muscle cell precursor in the mammalian heart. *Cell* 127:1137-1150.
56. Waldo K, S Miyagawa-Tomita, D Kumiski and ML Kirby. (1998). Cardiac neural crest cells provide new insight into septation of the cardiac outflow tract: aortic sac to ventricular septal closure. *Dev Biol* 196:129-144.
57. Nakamura T, MC Colbert and J Robbins. (2006). Neural crest cells retain multipotential characteristics in the developing valves and label the cardiac conduction system. *Circ Res* 98:1547-1554.
58. Li H, C Fong, Y Chen, G Cai and M Yang. (2010). Beta-adrenergic signals regulate adipogenesis of mouse mesenchymal stem cells via cAMP/PKA pathway. *Mol Cell Endocrinol* 323:201-207.
59. Li H, C Fong, Y Chen, G Cai and M Yang. (2010). beta2- and beta3-, but not beta1-adrenergic receptors are involved in osteogenesis of mouse mesenchymal stem cells via cAMP/PKA signaling. *Arch Biochem Biophys* 496:77-83.

60. Cochard P and D Paulin. (1984). Initial expression of neurofilaments and vimentin in the central and peripheral nervous system of the mouse embryo in vivo. *J Neurosci* 4:2080-2094.
61. Hoppo T, H Fujii, T Hirose, K Yasuchika, H Azuma, S Baba, M Naito, T Machimoto and I Ikai. (2004). Thy1-positive mesenchymal cells promote the maturation of CD49f-positive hepatic progenitor cells in the mouse fetal liver. *Hepatology* 39:1362-1370.
62. Benowitz LI and A Routtenberg. (1997). GAP-43: an intrinsic determinant of neuronal development and plasticity. *Trends Neurosci.* 20:84-91.
63. Reeves RH, J Yao, MR Crowley, S Buck, X Zhang, P Yarowsky, JD Gearhart and DC Hilt. (1994). Astrocytosis and axonal proliferation in the hippocampus of S100b transgenic mice. *Proc Natl Acad Sci U S A* 91:5359-5363.
64. Hamanaka RB and NS Chandel. (2010). Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes. *Trends Biochem Sci* 35:505-513.
65. Bianco P, M Riminucci, S Gronthos and PG Robey. (2001). Bone marrow stromal stem cells: nature, biology, and potential applications. *Stem Cells* 19:180-192.
66. Bruder SP, N Jaiswal and SE Haynesworth. (1997). Growth kinetics, self-renewal, and the osteogenic potential of purified human mesenchymal stem cells during extensive subcultivation and following cryopreservation. *J Cell Biochem.* 64:278-294.
67. Seita J and IL Weissman. (2010). Hematopoietic stem cell: self-renewal versus differentiation. *Wiley Interdiscip Rev Syst Biol Med.* 2:640-653.