AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Clinical usefulness of peripheral blood lymphocyte subsets in canine lymphoma

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1120> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Clinical Usefulness of Peripheral Blood Lymphocyte Subsets in Canine Lymphoma

B. Miniscalco^{1*}, R. Guglielmino¹, E. Morello¹, A. Tarducci¹ and M. Geuna²

¹Department of Animal Pathology, University of Turin, Grugliasco (Turin);

²Laboratory of Tumor Immunology, I.R.C.C., Candiolo, Italy

*Correspondence: Dipartimento di Patologia Animale, Facoltà di Medicina Veterinaria, Via L. da Vinci, 44, 10095 Grugliasco (TO), Italy

E-mail: barbara.miniscalco@unito.it

Keywords: blood, dog, flow cytometry, immunophenotype, lymphoma

INTRODUCTION

The classification of canine lymphoid malignancies into groups based on cell of origin (B- or T-cell types) is possible with immunophenotypic analysis. A high frequency of B-cell lymphomas has been described; about 30% of lymphomas are of T-cell origin (Teske *et al.*, 1994a; Fournel-Fleury *et al.*, 1997; Grindem *et al.*, 1998). In human medicine the major diagnostic role of blood lymphocyte subset typing is the screening for non-Hodgkin's lymphomas with peripheral blood involvement and their differentiation from reactive lymphoproliferative conditions (Thalhammer-Scherrer *et al.*, 2000). The aim of this project is to evaluate the diagnostic efficacy of blood immunophenotyping in dogs with lymphoma in order to detect the blood involvement and the prognostic benefits.

MATERIALS AND METHODS

Forty-eight dogs, 32 males and 16 females, ranging from 3 to 14 years old, of varying breeds, were examined. The diagnosis of lymphoma was performed on cytological smears. The immunophenotype was detected from lymph node cellular suspensions or cavity fluids using flow cytometry (FACScan Becton Dickinson): 36 cases were CD21⁺ B-cell, 7 were CD4⁺ T-cell, 3 were CD8⁺ T-cell and 2 were CD4⁺CD8⁺ T-cell lymphomas. A complete blood count and a phenotypic analysis of blood lymphocyte subpopulations were performed. The following monoclonal antibodies, specific for canine leukocyte antigens, were used: CD21 (CA2.1D6), IgG F(ab')₂ and IgM, CD5 (DH3B), CD3 (CA17.2A12), CD4 (CA13.1E4), CD8 (CA9.JD3), CD45RA (CA4.1D3), CD49d (CA4.5B3) and CD45 (CA12.10C12). Except for CD5 (VMRD, Pulmann, Washington, USA) and IgG, IgM (Jackson ImmunoResearch, West Grove, PA, USA), all monoclonal antibodies were kindly provided by Dr P.F. Moore (U.C.

Davies, CA, USA). The results were compared to reference values of a control group of 50 healthy dogs (Guglielmino *et al.*, 1995).

RESULTS

In agreement with Jain (1993), the complete blood count showed a mild to modest anaemia in 46% of the dogs, thrombocytopenia in 44% and mild (18,300/ μ l) to extreme (342,000/ μ l) leukocytosis in 33%. Only 19% of cases exhibited lymphocytosis; lymphoblasts were present for 12 dogs (25%), and 15% were without lymphocytosis. In 10% of cases leukopenia was observed and lymphopenia was observed in 25% of cases. Five out of 36 (14%) B-cell lymphomas showed CD21⁺ B-cell lymphocytosis (ranging from 5520 to 115,670/ μ l); four cases showed a moderate increase of B lymphocytes without lymphocytosis. A CD8⁺ T-cell increase was observed in seven B-cell lymphomas. This is also noted for human patients with neoplastic disorders, where an increase in the so-called 'anti-tumour' cytolytic CD8⁺ T lymphocytes occurs (Quaglino *et al.*, 1993). Two out of seven CD4⁺ T-cell lymphomas showed lymphocytosis (17,680 and 45,050/ μ l) with an increase in CD4⁺ T lymphocytes. Five dogs (71%) exhibited an increased CD4/CD8 ratio: in three cases this was due to an absolute increase in CD4⁺ T cells. One dog showed an abnormal immunophenotype (TCD4⁺CD45⁻) both in the lymph node and in the peripheral blood. In two out of the three dogs with CD8⁺ T-cell lymphoma a mild increase in the absolute count of CD8⁺ T lymphocytes, without lymphocytosis, was noted; one of these showed the same abnormal immunophenotype as the lymph node (TCD8⁺CD45⁻). Extreme lymphocytosis (12,690 and 338,580/ μ l), due to neoplastic cells, was observed for two dogs with CD4⁻CD8⁻ T-cell lymphoma.

DISCUSSION

In dogs with lymphoma the total and differential leukocyte counts and the presence of atypical lymphocytes were very variable: indeed blood involvement was not always detectable on a simple complete blood count. The detection of neoplastic cells (5 T-cell and 4 B-cell lymphomas) with the same immunophenotype as the origin tumours, also without lymphocytosis and cells with atypical morphology, was possible using flow cytometric analysis of peripheral blood. The immunophenotype of the peripheral blood was able to provide more precise information about systemic involvement. In 75% (9/12) of T-cell lymphomas and in 25% (9/36) of B-cell lymphomas a systemic involvement was present, confirming a poorer prognosis for T-cell lymphomas as compared with B-cell lymphomas (Teske *et al.*, 1994b). Simultaneous immunophenotyping of blood, lymph node and bone marrow from dogs with lymphoma could represent an important diagnostic and prognostic tool. Our study supports the potential use of blood immunophenotyping as a sensitive tool in the

detection of leukaemic cells, facilitation of tumour staging and monitoring of remission or relapse.

ACKNOWLEDGEMENTS

This research was supported by a grant from the Italian MURST.

REFERENCES

- Fournel-Fleury, C., Magnol, J.P., Bricaire, P., Marchal, T., Chabanne, L., Delverdier, A., Bryon, P.A. and Felman, P., 1997. Cytohistological and immunological classification of canine malignant lymphomas. Comparison with human non-Hodgkin's lymphomas. *Journal of Comparative Pathology*, **117**, 35–59
- Grindem, C.B., Page, R.L., Ammerman, B.E. and Breitschwerdt, E.B., 1998. Immunophenotypic comparison of blood and lymph node from dogs with lymphoma. *Veterinary Clinical Pathology*, **27**(1), 16–20
- Guglielmino, R., Geuna, M., Tarducci, A., D'Angelo, A., Ru, G. and Bruatto, G., 1995. Analisi citofluorimetrica delle sottopopolazioni linfocitarie nel sangue di 50 cani sani. *Atti della Società Italiana delle Scienze Veterinarie*, **49**, 811–812
- Jain, N.C., 1993. *Essentials of Veterinary Hematology*, (Lea & Febiger, Philadelphia), 334–335
- Quaglino, D., De Pasquale, A. and Ginaldi, L., 1993. Hematological changes in malignant disease. *European Journal of Laboratory Medicine*, **1**, 23–33
- Teske, E., Wisman, P., Moore, P.F. and Van Heerde, P., 1994a. Histological classification and immunophenotyping of canine non-Hodgkin's lymphomas. Unexpected high frequency of T-cell lymphomas with B-cell morphology. *Experimental Hematology*, **22**, 1179–1187
- Teske, E., Van Heerde, P., Rutteman, G.R., Kurzman, I.D., Moore, P.F. and MacEwen, E.G., 1994b. Prognostic factors for treatment of malignant lymphoma in dogs. *Journal of American Veterinary Medical Association*, **205**, 1722–1728
- Thalhammer-Scherrer, R., Veitl, M., Exner, M., Schneider, B., Geissler, K., Simonitsch, I. and Schwarzingner, I., 2000. Role of immunological lymphocyte subset typing as a screening method for lymphoid malignancies in daily routine practice. *Cytometry*, **42**, 5–10