AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The admiR-able advances in cardiovascular biology through the zebrafish model system

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/119146> since

Published version:

DOI:10.1007/s00018-012-1181-4

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

The *admiR*-able advances in cardiovascular biology through the zebrafish model system

Dafne Gays · Massimo Mattia Santoro

Received: 11 July 2012 / Revised: 12 September 2012 / Accepted: 24 September 2012
© Springer Basel 2012

Abstract MicroRNAs are small non-coding RNAs endogenously expressed by all tissues during development and adulthood. They regulate gene expression by controlling the stability of targeted messenger RNA. In cardiovascular tissues microRNAs play a role by modulating essential genes involved in heart and blood vessel development and homeostasis. The zebrafish (*Danio rerio*) system is a recognized vertebrate model system useful to study cardiovascular biology; recently, it has been used to investigate microRNA functions during natural and pathological states. In this review, we will illustrate the advantages of the zebrafish model in the study of microRNAs in heart and vascular cells, providing an update on recent discoveries using the zebrafish to identify new microRNAs and their targeted genes in cardiovascular tissues. Lastly, we will provide evidence that the zebrafish is an optimal model system to uncover new microRNA functions in vertebrates and to improve microRNA-based therapeutic approaches.

Keywords MicroRNA · Zebrafish · Heart · Blood vessel · Development

MiRNAs: function and mechanism

MicroRNAs (e.g., miRs or miRNAs) are evolutionary conserved small non-coding RNAs, usually 21–25 nt long, that regulate gene expression at a post-transcriptional level.

MiRNAs act by impairing target mRNA expression through an antisense mechanism. They recognize partial or full complementary sequences that are present mainly in the 3'-UTR (untranslated region) of mRNAs and thereby inhibit protein synthesis [1, 2]. Based on their locations in the genome, microRNAs can be intergenic and so transcribed individually as single transcriptional units or intragenic (intronic or exonic) and then transcribed together with the genes in which they are located. Often, more miRNAs are located close by in the genome and form clusters; in this case they are transcribed in a single polycistronic unit that after cleavage releases single microRNAs [3, 4].

MiRNAs are transcribed by RNA polymerase II as a stem-loop precursor known as pri-miRNA. In the nucleus it is cleaved by a ribonuclease III enzyme called *Drosha* with its cofactor DiGeorge syndrome critical region 8 (DGCR8) into 60–100-nucleotides hairpins called pre-miRNA that can leave the nucleus through exportin 5. In the cytoplasm a second RNase, *Dicer*, removes the loop that generates a duplex containing the mature miRNA sequence and a 'passenger strand' miRNA sequence (called *star*). Subsequently, the miRNA duplex is recognized by the protein complex *Argonaute* to form the RNA-induced silencing complex (RISC). Thus the 'seed' region of mature miRNA (nucleotides 2–8) guides RISC on target mRNA(s). MiRNAs post-transcriptional regulation of gene expression can be achieved through different mechanisms: (1) degradation of the targeted mRNA and/or (2) translation repression. Recently, a role for miRNA in translational stimulation has also been proposed [2, 5–7]. While miRNAs commonly repress their mRNA targets, in rare cases it has been observed that miRNA stimulation of mRNA translation occurs [8].

MiRNAs have numerous high- and low-affinity targets and therefore have the potential to modulate multiple pathways. Often many of these targets can themselves

D. Gays · M. M. Santoro (✉)
Department of Biology, Biochemistry and Genetics,
Molecular Biotechnology Center, University of Turin,
Via Nizza 52, 10126 Turin, Italy
e-mail: massimo.santoro@unito.it

control the expression and function of miRNAs generating regulatory circuits [9–12]. In positive feedback circuits, miRNAs repress a repressor, leading to activation of transcription factors that activate miRNA expression. The regulatory loop formed between miR-1, HDAC4, and MEF2 exemplifies this form of regulation in skeletal muscle [13]. Here, repression of HDAC4 by miRNA-1 derepresses the activity of the MEF2 transcription factor, which in turn activates expression of miRNA-1 and other target genes. On the other hand, miRNAs can also establish negative feedback circuits by repressing activators of miRNA/mRNA expression. In a negative regulatory loop, miRNAs repress transcription factors that are required for miRNA expression, leading to decreased expression of miRNAs. The reciprocal regulatory interaction among a posterior Hox gene, *nob-1*, and miRNA-57 has been found to be essential for regulation of posterior cell fate determination in *C. elegans* [14]. On one hand, the Hox gene is required for normal activation of miRNA-57 expression, and on the other, the Hox gene functions as a direct target of and is repressed by the miRNA. Given the conservation of the two genes, a negative feedback loop between Hox and miRNA genes might be broadly used across species to regulate cell fate along the anteroposterior axis [14].

In addition, 3'-UTRs can contain binding sites for multiple miRNAs, allowing redundancy and cooperation between them. Collectively, these mechanisms establish a complex network of gene regulation that fine-tunes the biological processes [9, 15].

MiRNAs are widely expressed in almost all vertebrate tissues, and functional roles for miRNA during development have been extensively described [16, 17]. MiRNA expression has been linked to a wide range of physiologic processes, including development, differentiation, and cell proliferation. Therefore, it is likely that miRNAs participate in nearly every physiological process since miRNA expression is temporally and spatially regulated.

The role of miRNAs in shaping heart and vascular networks in vertebrates

The heart is the first organ to form and function during embryo development, and its continued contraction is essential for life. Processes that lead to heart development are very complex, and their alterations can result in congenital heart disease; functional abnormalities in the adult heart result in a variety of disorders, among which are arrhythmias, cardiomyopathies, heart failure, and sudden death [18, 19]. In vertebrates, the linear heart tube forms from the migration of bilateral populations of myocardial and endocardial cells and their fusion at the midline. Then, morphogenetic movements lead to the formation of a

looped, multichambered organ [20]. The heart has the function to pump blood in the vascular system. The vascular system is a highly organized network of arteries, capillaries, and veins that supplies virtually all body tissues, enabling an efficient exchange of oxygen and nutrients and removal of waste products. The vascular system is indispensable for embryonic development and adult life, and its aberrant alterations are associated with numerous diseases, including cancer, atherosclerosis, retinopathy, and stroke [21]. The generation of the vascular system starts with a process called vasculogenesis that gives rise to the primary vascular plexus of the embryo from which new vessels sprout in a process called angiogenesis [22]. During this phase, the vascular plexus progressively expands and remodels into a highly organized and stereotyped vascular network. The development and maturation of the cardiovascular system also require the recruitment of smooth muscle cells that stabilize vessels and regulate blood flow [23, 24]. After birth, angiogenesis still contributes to organ growth, but, during adulthood, most blood vessels remain quiescent until they receive physiopathological cues that stimulate endothelial cells (ECs) proliferation.

Cardiovascular formation and function are precisely controlled by a network of transcription factors closely intertwined with families of microRNAs that modulate multiple aspects of cardiovascular development, function, and dysfunctions [12]. The importance of microRNAs in the regulation of cardiovascular development and function has been demonstrated through earlier studies on vertebrate mutants of Dicer, the enzyme required for microRNA biogenesis [25–29]. The complete loss of Dicer function leads to severe developmental defects in both zebrafish and mouse. To overcome the early lethality, conditional Dicer mutant mice were created [30]. Cardiac-specific deletion of Dicer leads to cardiomyopathy, heart failure, and postnatal lethality [31]. Again, in mammals specific inactivation of Dicer in endothelial cells shows that endothelial microRNAs are important for postnatal angiogenic responses to a variety of stimuli, including VEGF, tumors, limb ischemia, and wound healing [32]. In mouse the loss of Dicer in vascular smooth muscle cells lead to a dilated, thin-walled blood vessels that also exhibited impaired contractility [33]. Zebrafish maternal-zygotic Dicer mutants (that lack both maternal and zygotic Dicer) display abnormal morphogenesis during gastrulation, brain formation, somitogenesis, and heart development. However, it has been shown that the injection of miRNA-430 rescues early gastrulation defects, enabling the analysis of the loss of microRNAs in tissues that otherwise would not develop [28, 34]. Recent studies on zebrafish *Dicer* mutant embryos demonstrated excessive endocardial cell formation, suggesting a role for miRNA in cardiac development [35]

Fig. 1 Molecular and functional consequences of the loss of Dicer activity in maternal-zygotic dicer mutants embryos (MZdicer KO). In zebrafish mutants lacking dicer enzymatic activity, pre-microRNAs are not processed in mature microRNAs, leading to several developmental defects. The lack of miR-430 miRNAs leads mainly to gastrulation, brain formation, and somitogenesis defects. The loss of other microRNAs affects embryo morphogenesis and organogenesis, as well as heart development and blood circulation

(Fig. 1). These studies indicate that the zebrafish system can be a valuable tool to investigate microRNA's function in vivo, especially for studying cardiovascular development and homeostasis where this model presents some unique features among other vertebrate and mammalian models [36].

The zebrafish model system to study cardiovascular miRNA functions

The zebrafish as a model for cardiovascular studies

The zebrafish (*Danio rerio*) system has emerged in the past years as an ideal vertebrate model organism to study a wide variety of biological processes [37–39]. Some of the advantages of the zebrafish animal model system include high fecundity, external fertilization, rapid development, and optical clarity. Last but not least, the zebrafish has low-cost maintenance compared to other vertebrate model systems. The importance of zebrafish for cardiovascular research relies on some unique characteristics such as the early development of a functional cardiovascular system (beating heart, aorta, cardinal vein, and blood) that is already formed by 24 h post-fertilization (hpf) [20, 40]. Since the zebrafish embryo is relatively small and aquatic, oxygen can diffuse passively through tissues, and thus embryos can live more than 3 days without a functional cardiovascular system [41]. This allows genetic manipulation of cardiovascular development for longer than it would be in mammals, where the absence of a functional cardiovascular system is fatal at its early stage of development. Until 5 days post-fertilization (dpf), the embryos are nearly transparent, allowing in vivo visualization during development (even at

single-cell resolution) of the heart, blood vessels, and other tissues, without instrumentation or manipulation other than the use of a stereomicroscope. Furthermore, the generation of tissue-specific transgenic lines is relatively easy thanks to the availability of hundreds of fertilized eggs with the injection of DNA and RNA constructs [42]. Such a method usually uses a native tissue-specific promoter to drive expression of a fluorescent reporter protein, such as green fluorescent protein (GFP). The following transgenic lines are of great interest for cardiovascular studies: *Fli1:GFP* (which expresses GFP in endothelial cells and some neural crest-derived cells) [43], *kdr1:GFP* (which expresses GFP localized to endothelial cells) [44], *gata1:dsRED* (which expresses dsRED in erythrocytes) [45], *cd41:GFP* (which expresses GFP in thrombocytes) [46], *cmlc2:GFP* (which expresses GFP in cardiomyocytes) [47], and *tgn1:Cherry* (which expresses cherry in smooth muscle cells) [48]. Coupled with impressive optical clarity, these transgenic lines allow observation of in vivo cellular behavior in a way impossible in other models. Transparent zebrafish embryos are also well suitable for in vivo time-lapse imaging. The fast acquisition speed of a spinning disk and two-photon confocal microscopy reduces the recording time significantly when millimeter-sized embryos need to be imaged at high resolution and at short time intervals. Light sheet fluorescence microscopy could also be very useful in zebrafish [49, 50]. These attributes, including the possibility of carrying out gain- and loss-of-function studies have led to the emergence of the zebrafish as an excellent embryological model that provides a unique opportunity to uncover novel insights into the molecular genetics of development and disease.

More recently, the zebrafish has become an attractive model to study the effects of genetic variations identified in

patients with cardiovascular defects by candidate gene or whole-genome-association studies [51, 52]. Moreover, zebrafish are suitable for forward genetics approaches, which allow the unbiased identification of novel and unanticipated cardiovascular genes. Small and large mutagenesis screens have been conducted in zebrafish and have led to a wealth of heart and endothelial mutants, several of which have already been characterized in detail and positionally cloned [41, 53]. Zebrafish mutants with various cardiovascular phenotypes that closely correlate with human disease, such as congenital heart disease, cardiomyopathies, and arrhythmias, have also been isolated [54]. The pool of zebrafish mutants for which the causal gene mutation has been identified is constantly growing. The human orthologs of several of these zebrafish genes have been shown to be involved in the pathogenesis of human cardiovascular diseases (CVD), known to be the most common cause of worldwide deaths [51, 54]. Cardiovascular zebrafish models also provide the opportunity to develop and test novel therapeutic strategies, using innovative technologies such as high-throughput in vivo small molecule screens.

The zebrafish as a model for miRNA studies

Zebrafish is a helpful model system to study miRNA function and characterize miRNA:mRNA target interactions [55–58]. Important information regarding the conservation of miRNA genes in vertebrates has been derived from the identification and characterization of miRNAs in zebrafish. *Danio rerio* have also proven to be a particularly valuable system to study miRNA functions in developmental processes. In the last decade different analyses have led to the identification of 415 zebrafish miRNAs, and based on seed sequence identity, it has been possible to group these into 44 families [59–61]. However, the exact number of zebrafish miRNAs might be higher, since genome sequencing and annotation has not been fully completed. Thus far, most zebrafish miRNAs exhibit tissue-specific patterns of expression [62].

miRNA expression analyses

To understand the function of specific microRNAs, it is important to determine their spatial and temporal expression pattern. Several methods have been developed to evaluate miRNA expression in zebrafish. Due to their relatively small size and optical transparency, zebrafish are useful for whole-mount in situ hybridization (ISH) analyses. However, compared to gene expression analysis, ISH detection of miRNAs can sometimes be challenging because of the microRNA's relatively small size (usually 21–25 nt), which then lowers the probe signals. Therefore,

non-radioactive-labeled synthetic RNA oligonucleotide probes, such as 2-*O*-methyl RNA oligonucleotides, and locked nucleic acids (LNA) are currently used since they can overcome this problem, allowing an overall enhancing of the probe signal [6, 63]. To distinguish between genes that encode the same mature miRNA sequence, Postlethwait and collaborators [64] developed the use of digoxigenin-labeled riboprobes designed to bind to miRNA primary transcripts, allowing the observation of miRNA expression for both intergenic and intronic miRNAs. Alternatively to ISH, zebrafish microRNA expression patterns have also been investigated by qPCR, microarray, RNA-seq, and Northern blot analyses at different developmental stages and in different tissues [55, 59, 65]. As a result, a set of specific miRNAs displays cardiovascular-specific expression patterns, suggesting functional roles in differentiation or homeostasis of these miRNAs during development in zebrafish.

Control of microRNA expression

Modulation of miRNA expression can be achieved in several ways in zebrafish. For gain-of-function studies, the most widely used technique is the injection in one-cell eggs of double-stranded miRNA mimics to promote single miRNA overexpression [66, 67]. It must be taken into consideration that in this way the specific miRNA is overexpressed not only in the tissues where it is endogenously expressed, but also in the entire embryo. However, it could be useful to mimic situations in which a microRNA is spatio-temporally deregulated. To analyze the role of microRNA gain of function in a specific tissue, DNA constructs in which a tissue-specific promoter controls pri-miRNA expression can be used. More sophisticated genetic techniques based on the CreERT2/loxP system have been described in zebrafish, and they can also provide temporal and tissue-specific miRNA expression in zebrafish embryos [68, 69]. These strategies will allow the study of miRNA functions not only in the developing embryos, but also in adult animals and their exclusive tissues of interest.

Inhibition of specific miRNAs can be obtained by the injection of miRNA modulators, among which are cholesterol-modified antagomirs, modifications such as 2' *O*-methoxyethyl phosphorothioate or 2' fluoro substitutions, as well as oligonucleotides stabilized by LNA phosphorothioate chemistries. Although these techniques, especially antagomirs, are successfully used in mammals, they do not work quite efficiently in zebrafish embryos, mainly because their chemistry is not compatible with the lipid-enriched zebrafish yolk sac [70]. Conversely, antisense morpholino phosphorodiamidate oligonucleotides (MOs) have been extensively used to knock down gene functions in zebrafish embryos and larvae [71–73]. To inhibit specific miRNAs,

MOs must be designed to block the processing of pre-miRNA or the activity of the mature miRNA [74, 75]. A recent alternative approach to inhibit miRNA function *in vivo* is to use miRNA ‘sponges’ or ‘decoys.’ They contain several miRNA-binding sites, and, when present in a cell as RNA, act as competitive inhibitors for miRNA binding [76, 77]. Placing a miRNA sponge in the 3′-UTR of a reporter gene (dGFP in most cases), it is also possible to obtain information about the miRNA expression pattern and to down-modulate it at the same time. Interestingly, miRNA sponges have also been used to generate miRNA tissue-specific and stable knockdown [78, 79]. The use of sponges can be useful to overcome morpholinos off-target effects and to obtain a stable or prolonged knockdown not achievable with morpholinos because of their relative short half-life. Recently, targeted gene inactivation via zinc finger nucleases (ZFN) has been established for zebrafish genome [80, 81]. This technique, allowing the introduction of targeted lesions in the zebrafish genome, could, in principle, be used to selective targeted pre-miRNA sequences. The TALEN technology could also be used to induce somatic mutagenesis in zebrafish microRNA genes [82, 83].

Finally, compared to mammalian models the zebrafish system also has the unique feature of being able to perform “transient” knockdown (by morpholino) and rescuing (by mimics) of microRNAs in whole embryos. Such genetic techniques cannot be achieved so easily in rodents in a short period of time [36].

Target prediction and validation tools

Several open-access bioinformatic databases have been developed to facilitate the analysis of miRNAs and their target prediction. However, prediction of miRNA–mRNA interactions still remains challenging, owing to the short length of miRNAs, requirement of only partial homology for binding, redundancy among members of an miRNA family, and the existence of multiple putative miRNA recognition sites [84, 85]. The majority of the computational target prediction programs are based on several features, such as complementarity between the 5′-seed of the miRNA and the 3′-UTR of the target mRNA, thermodynamic stability of the miRNA–mRNA duplex, conservation among species, and the presence of several miRNA target sites. However, in zebrafish different *in silico* target prediction programs exist. They use prediction algorithms based on mammalian counterparts [86]. Such prediction programs must be used carefully in zebrafish studies since (1) the zebrafish genome is not fully sequenced, (2) only a limited number of miRNA targets have been experimentally validated, and (3) there is low conservation among species regarding mature microRNA sequences as well as

the target prediction. Experimental high-throughput studies (such as poly(A)-site mapping followed by RNA-Seq) are needed to allow for generation of zebrafish precise prediction algorithms [27]. To validate predicted miRNA–mRNA interactions, several *in vitro* experimental approaches can be used, such as biochemical methods (luciferase assays, qRT-PCR, Western blot, RNA seq), “omics” approaches (SILAC, LAMP), and RISCome analysis (quantification of mRNAs in the RNA-induced sequencing complex; RISC RNA sequencing, RIP-chip) [87]. The zebrafish system allows the validation of target mRNA *in vivo*. Novel techniques have been recently implemented in the zebrafish model such as fluorescent miRNA sensor and MO target protector. In the first case, validation of the target prediction is tested using a reporter assay based on monitoring GFP/cherry fluorescence in zebrafish embryos microinjected with mRNA encoding the fluorescent reporter fused to the 3′-UTR of the target gene in the presence or absence of a specific miRNA duplex [67, 88]. In this assay, a decrease in fluorescence in the presence of miRNA duplex indicated miRNA-mediated repression and then the confirmation of the target *in vivo*. In the second case, MOs are designed to be complementary to specific miRNA-binding sites in target mRNAs. In this way, MOs have been shown to efficiently protect the target mRNA from translational inhibition or degradation [89, 90]. To study the regulation of a particular target, it is important to first establish that the 3′-UTR is regulated by a particular miRNA. miRNAs may speed degradation or slow translation of their targets, but repression caused by either mechanism can be assessed by measuring the protein output of a reporter. The efficiency of “target protectors” to block the interaction of miRNAs with a particular binding site in a target mRNA can also be investigated by co-injections with reporter mRNAs containing a GFP/cherry coding sequence and a 3′-UTR region with the miRNA-binding site [65, 91]. Subsequently, a tested target protector can be used to study whether protection of a specific target mRNA from silencing has any biological effect also in a tissue-specific manner (by generation of a cell autonomous miRNA sensor) [65, 67].

MicroRNAs involved in heart and vascular development and homeostasis in zebrafish

The functions of individual miRNAs in normal and pathological angiogenesis are just starting to be unveiled. Single miRNAs may regulate cardiovascular functions either cell-autonomously or non-cell autonomously. Recent progress toward understanding the functions of specific miRNAs in cardiovascular development and homeostasis has been done. Here, we described those miRNAs whose

Table 1 Summary of vertebrate miRNAs involved in cardiovascular development and homeostasis in zebrafish

microRNA	Targets	Functions	Technique	Reference
Cardio(vascular)				
23	Has2 Icat Tmem2	Cardiac valve formation	Knockdown (morpholino); overexpression (mimic)	Lagendijk et al. [35]
133	Mps1 Cx43	Cardiac regeneration	Conditional knockdown (transgene); overexpression (transgene)	Yin et al. [79]
138	aldh1a2 cspg2	Cardiac patterning	Knockdown (morpholino)	Morton et al. [70]
143	aldh1a2 rxrab add3	Cardiogenesis	Overexpression (RNA injection)	Miyasaka et al. [100]
218	Robo1	Migration of heart precursors	Knockdown (morpholino)	Deacon et al. [101]
(Cardio)vascular				
24	Pak4 Gata2	Cardiac and patterning angiogenesis	Overexpression (pre-miRNA injection)	Fiedler et al. [110]
31	PROX1	Vascular lineage-specific differentiation and sprouting	Knockdown (morpholino)	Pedrioli et al. [112]
92a	ITGA5	Patterning angiogenesis	Overexpression (pre-miRNA injection)	Bonauer et al. [114]
126	spred1 pi3kr2	Vascular integrity and maturation	Overexpression (pre-miRNA injection)	Fish et al., [119]
221	cdkn1b pi3krl	Endothelial tip cell behaviour	Knockdown (morpholino)	Nicoli et al. [67]
			Overexpression (miRNA duplex)	Nicoli et al. [65]

Fig. 2 Schematic representation of zebrafish heart at 72 hpf, which shows the bulbus arteriosus (BuA), atrium (A), ventricle (V), and the atrio-ventricular valve (AVC, dashed circle). Green indicates the pericardium, purple the myocardium, pink the endocardium, and blue smooth muscle cells. Each of these tissues expresses specific microRNAs that are shown along with their targets. Inhibition of the specific target expression allows for a specific functional role of the different microRNAs in cardiogenesis (see text for details)

functions in the zebrafish cardiovascular system have been addressed (Table 1).

Heart-specific miRNAs

Heart formation and function are precisely controlled by networks of transcription factors closely intertwined with families of miRNAs that modulate multiple aspects of cardiac development, function, and dysfunctions. Here, we report some miRNAs whose function has also been characterized through the use of zebrafish (Fig. 2).

The miRNA 23

Defects in cardiac valves are the most common subtype of cardiovascular malformations, and in adults they remain a major cause of morbidity and mortality [92]. Increasing the molecular understanding of processes that control heart valve development and remodeling could be useful to improve or develop new therapeutic approaches. In this sense it could be relevant that the discovery of one gene involved in heart valve development is miRNA-23 [35]. It has been showed that zebrafish MZDicer mutants have endocardial defects, including excessive endocardial cushion formation, and Legendijk et al. demonstrated that they are caused by the loss of miRNA-23 activity. In this process, miRNA-23 acts to downregulate *has2* (hyaluronic acid synthase 2), an extracellular remodeling enzyme known to be required for endocardial cushion and valve formation [93] and whose upregulation in MZDicer mutants authors showed to be responsible for the observed valve defects. Furthermore, they demonstrated that in mouse endothelial cells, miRNA-23 is able to inhibit a transforming growth factor-beta (TGF- β)-induced endothelial-to-

mesenchymal transition (EMT), a process that normally occurs during heart valve development. So the authors proposed a model in which miRNA-23, *has2*, and its product hyaluronic acid (HA) create a feedback regulatory loop that could respond to the AVC signal, such as TGF- β , and restrict endocardial cushion formation to the AVC region. Data shown in zebrafish and mouse endothelial cells suggest a conserved role of miRNA-23 whose function in mammalian valve development is still unknown but which could be interesting to investigate.

The miRNA 138

A precise regulation in time and space of gene expression and protein activity is fundamental for a proper development of a chambered heart. Morton et al. [70] demonstrated that miRNA-138 is necessary to establish appropriate chamber-specific gene expression patterns during embryonic development. This microRNA is expressed in specific domains of the heart, and its downregulation leads to ectopic expression in the ventricle of genes normally expressed in the atrioventricular valve region (AVC). As a consequence, ventricular cardiomyocyte morphology and cardiac function are affected. The microRNA function is to partially repress *aldh1a2* in the ventricle, a gene that encodes retinoic acid (RA) dehydrogenase, an enzyme involved in RA synthesis. miRNA-138 also mediates ventricular repression of *cspg2*, a gene positively regulated by RA signaling. This study demonstrates that, targeting multiple members of a common pathway, miRNA-138 establishes discrete temporal and spatial domains of gene expression during cardiac morphogenesis, which is necessary for the correct development of the heart. This microRNA is highly conserved across species as well as

transcriptional networks that control heart patterning [94], and so it could be interesting to analyze miRNA-138 functions in mammalian heart in which RA signaling plays a key role during development [95].

The miRNA 143/145 cluster

In mammals, miRNA-143 and miRNA-145 are two smooth muscle cell-specific microRNAs playing a role in vascular SMC phenotypic switch [96–99]. These microRNAs are expressed in early cardiac progenitors and throughout the embryonic heart before becoming restricted to vascular and visceral smooth muscle cells. Despite being expressed at high levels in the developing heart, no cardiac function for the miRNA-143/145 cluster in developing mammalian heart has been found [98, 99]. Although the downregulation of miRNA-145 affects cardiac development [66], its role in this organ has not yet been fully investigated in zebrafish. On the contrary, a role of miRNA-143 in cardiogenesis in zebrafish has been reported. In particular, it has been demonstrated that miRNA-143 is involved in the conversion of mechanical stimuli into biological output (mechano-transduction pathway) in the heart [100], a process that is important during the later phases of cardiac development when hemodynamic forces play an essential role in the morphogenesis and adaptation of the heart to the circulatory demands. The authors showed that miRNA-143 expression in the myocardium and endocardium of outflow tract and ventricle is heartbeat-dependent. Blocking the heartbeat with the injection of *tmt2* (cardiac troponin T2) morpholino or with BDM (2,3-butanedione monoxime) treatment, miRNA-143 cardiac expression is reduced or abolished. They showed that the loss of miRNA-143 leads to pericardial edema and heart defects: the atrium of miRNA-143 morphants is swollen, and the ventricle doesn't inflate, maintaining a tube-like structure; the ventricle doesn't beat, and there are blood flow defects. Downregulation of miRNA-143 disrupts ventricular myocardial and endocardial architectures, probably through the repression of *aldh1a2* and *rxrab*, which encode the retinoid X receptor alpha b, and then through an aberrant activation of RA signaling. This study demonstrates that mechanical forces regulate the expression of miRNA-143 to set up a gradient of RA signaling in the developing heart. Understanding how mechanical stimuli are converted in biological responses in the developing heart can be useful for giving insight into the mechanism of both congenital and acquired heart disease and then how to prevent or combat them.

Another study showed that miRNA-143 is required for heart chamber morphogenesis through a direct repression of *add3* (adducin 3), which encodes an F-actin capping protein, in myocardial cells [101]. Deacon et al. observed

that downregulation of miRNA-143 inhibits ventricular cardiomyocyte F-actin remodeling. As a consequence, cellular growth and elongation are blocked and cause ventricular collapse and decreased contractility.

We would speculate that the fact that a phenotype is evident in zebrafish but not in mouse would reside in a second copy of miRNA-143, which is present in the zebrafish genome but not in the mouse genome. The second copy of the miRNA-143 is identical to the one located in the miRNA-143/145 cluster, but it is in a different genomic region, and its expression could possibly be controlled by a different transcriptional unit (data not shown). Alternatively, it is conceivable that the cardiac phenotype of the miRNA-143 KO mouse model has not been characterized under ischemic conditions, such as I/R [98].

The miRNA 218

MiRNA-218 is a conserved microRNA situated intronically in *Slit2* *Slit3*, genes that encode ligands of Robo receptors. Fish et al. [102] demonstrated that the *Slit*/miRNA-218/*robo* signaling pathway regulates migration of heart precursors in the midline, in part modulating Vegf signaling in the endocardium. The authors observed that downregulation of miRNA-218 causes a delayed heart field migration that results in pericardial edema and abnormal heart looping. The same phenotypes are caused by the overexpression of *robo1*. Demonstrating that *robo1* is a direct target of this microRNA, they found that a ligand-encoded microRNA regulates the expression of its own receptor. They subsequently also observed that Vegf is a mediator of heart field migration in zebrafish: its temporal inhibition leads to delayed heart field fusion. In vivo it interacts with *robo1*, which in turn could facilitate the phosphorylation of Vegfr2 as demonstrated in vitro. This study provides evidence of a *Slit*/miRNA-218/*Robo*/*Vegf* feedback regulatory loop that precisely regulates the migration of heart fields to form the linear heart tube. It has also been demonstrated in mouse that miRNA-218 regulates vascular patterning by modulation of Slit-Robo signaling [103], highlighting a conserved role of miRNA-218 in this pathway.

The miRNA 133

The zebrafish heart is similar to the human heart in many respects. But unlike the human heart, after injury the zebrafish heart closes wounds rapidly and regenerates cardiac tissue to fully restored cardiac muscle function [104]. Specific molecular networks have been associated to these regeneration events [105, 106]. These findings have suggested a potential role of microRNA in the regulation of heart regeneration and offer hope that this event could also

Fig. 3 Schematic representation of zebrafish trunk vessels showing endothelial cells (EC, pink), tip cells (red), and smooth muscle cells (SMC, blue) of the dorsal aorta (DA), posterior cardinal vein (PCV), and intersegmental vessels (Se). The expression of select miRNAs is shown, along with their observed target and functional roles (see text for details)

be induced in mammals. Yin et al. [79], through a microarray analysis of microRNA differentially expressed after the resection of the ventricle apex of the adult zebrafish heart, found that miRNA-133 expression correlates with the regeneration process. In zebrafish, miRNA-133 is downregulated after apex amputation, but progressively returns to be expressed to baseline when regeneration is completed. To study miRNA-133 function in adult heart regeneration, these authors generated two zebrafish transgenic lines through which they modulate miRNA levels in adult animals during regeneration. For overexpression studies, they generated a transgenic line containing 250 bp of genomic sequences flanking the sides of the miRNA-133a1 precursor sequence downstream of the *hsp70* promoter. Adult transgenic animals exposed to a single heat treatment showed a 1.8 increase of mature miR-133. For downregulating miRNA-133, the authors created another transgenic line using an miRNA-133 sponge construct (encoding EGFP cDNA followed by triplicate perfect binding sites for miRNA-133) placed downstream of the *hsp70* promoter. They showed that, remarkably, a single heat shock was sufficient to transiently but efficiently knock down adult cardiac miRNA-133 expression. Using these transgenic lines it was possible to discover that miRNA-133 inhibits myocardium regeneration reducing cardiomyocyte proliferation. Microarray analyses on miR-133-overexpressing/downregulated adult zebrafish identified cell cycle regulators such as monopolar spindle 1 (*mps1*), an essential factor for heart regeneration [107], and cytoskeletal/structural component proteins such as connexin-43 (*cx43*), a gene implicated in human cardiomyopathies [108], as direct targets of miRNA-133. The results of Yin et al. are also consistent with the elevated cardiomyocyte proliferation displayed by miR-133a1/a2 double knockout mice [109]. This work highlights the utility of the zebrafish model for studying the role of microRNAs also in the adult

cardiovascular system in physiopathological contexts. Although we are still far from the possibility of inducing heart regeneration mammals, dissecting this process in zebrafish can provide context for understanding and possibly enhancing the cardiac regenerative capacity in mammals.

In summary, miRNA-133, a conserved miRNA known to have a role in cardiac development and disease in mammals [12], regulates zebrafish heart regeneration [79]. This work further demonstrates the versatility of the zebrafish model for miRNA studies not only during embryonic development, but also in the adult animal.

Vascular-specific miRNAs

Formation of a functional vasculature during vertebrate development is essential for embryonic survival. Physiological aspects of the vascular system have been extensively studied, resulting in its clinical importance. Thus far studies have shown a high degree of conservation of the molecular mechanisms that govern zebrafish vascular development as well as in the anatomical structures that comprise the vascular system in vertebrates. In zebrafish embryos both endothelial cell functions and angiogenesis are critically regulated by microRNAs (Fig. 3). Here are some examples of studies conducted by knockdown or ectopic overexpression of specific microRNAs and their effects on angiogenesis and endothelial cell behavior.

The miRNA 24

The miRNA-23/27/24 gene cluster is enriched in human endothelial cells and in highly vascularized tissues such as the heart. Recently, several reports have demonstrated that the miRNA-23/24/27 cluster is involved in heart and

vascular development and homeostasis in zebrafish [35, 110]. Overexpression of miRNA-24 in zebrafish embryos resulted in pericardial edema as well as blood accumulation. Studies in transgenic zebrafish that express GFP in the vasculature showed that miRNA-24 overexpression impaired intersegmental vessel formation and resulted in impaired blood transportation [110]. Morpholino-mediated silencing of the miRNA-24 targets *gata2* and *pak4* in zebrafish mimicked the effects of miRNA-24 overexpression, showing an involvement and potential therapeutic importance of those targets in angiogenesis [110]. These recent data also propose miRNA-24 as a suitable target for therapeutic intervention in cardiovascular diseases [111].

The miRNA 31

Using a TaqMan-based qRT-PCR profiling platform for signature miRNAs, miRNA-31 showed up as an important endothelial miRNA [112]. In vivo gain-of-function studies with zebrafish embryos established that overexpression of miRNA-31 impaired lymphatic development and reduced endothelial venous sprouting. Taken together, these findings indicate that miRNA-31 plays a pivotal role in regulating lineage-specific differentiation within the developing vasculature of vertebrates.

The miRNA 92

Many microRNAs involved in pathological angiogenesis, such as tumor angiogenesis, modulate tumor growth by controlling the production of angiogenic factors and then neovascularization. The miRNA 17–92 cluster is transcribed as a polycistronic unit and selectively targets anti-angiogenic proteins [113]. Inside this cluster, miRNA-92a seems to be important in the control of cardiovascular system homeostasis in vertebrates. By using antagomir approaches, Dimmeler's group determined that miRNA-92a is highly expressed in endothelial cells and regulates angiogenic functions of endothelial cells as evidenced by gain of function and loss of function experiments in vitro [114]. To demonstrate the conservative role of this microRNA, overexpression of miRNA-92a in zebrafish not only impairs in vitro angiogenesis but also blocks intersomitic vessel growth [114]. One of the mechanisms through which miRNA-92 controls endothelial cell functions seems to be the direct regulation of integrin subunit $\alpha 5$ (*itga5*), a crucial regulator of vessel growth and angiogenesis [115]. MiRNA-92a is expressed also in mammal cardiac fibroblasts and cardiomyocytes, and antagomir-92a treatment significantly reduced apoptosis of cardiomyocytes in vivo, but did not directly affect survival of cardiomyocytes in cell culture assays in vitro, indicating that antagomir-92a may influence cardiomyocyte apoptosis via a paracrine

mechanism. Interestingly, it has been observed that atheroprotective flow patterns decrease the level of miRNA-92a, which in turn increases KLF2 expression to maintain endothelial homeostasis [116].

The miRNA 126

MiRNA-126 is one of the most evolutionary conserved and enriched microRNAs in human and mouse endothelial cells [117, 118] and is also expressed in the zebrafish vascular tree throughout development [67, 119]. In 2008, different groups analyzed and reported vascular defects in miRNA-126-deficient cells, ranging from zebrafish to human and mice. MiRNA-126 has been clearly shown to be a master non coding RNA in vascular functions acting as a regulator of angiogenic integrity and signaling in vitro and in vivo [119–121] and recently also in heart valve development [122]. In zebrafish, the downregulation of miRNA-126 induced collapsed blood vessels and cranial hemorrhages in the developing organism, suggesting its primary role for maintaining vascular structure during development [119]. Sprouty-related EVH1 domain-containing protein 1 (*Spred1*) and a regulatory subunit of PI3K, PIK3R2 (also known as p85 β), two direct targets of miRNA-126, seem to explain these angiogenic defects. MiRNA-126 regulates *Spred1* and *PIK3R2*, which are negative regulators of MAPK [123] and PI3K [124] signaling pathways, respectively. As a consequence, miRNA-126 promotes VEGF and other growth factor signaling in ECs in vivo. As a matter of fact, miRNA-126 may fine-tune angiogenic responses by targeting multiple signaling pathways [119, 125]. Remarkably, deletion of miRNA-126 in mice resulted in the same vascular developmental defects observed in zebrafish, such as delayed angiogenic sprouting, widespread hemorrhaging, and partial embryonic lethality [120, 121], supporting the idea that the zebrafish model can be a simple and easy tool to discovery miRNA functions in vertebrates. Recent studies in zebrafish demonstrate for the first time that miRNA-126 expression and the *egfl7* gene are both under the control of blood flow. Blood flow-dependent Kruppel-like factor 2a (*klf2a*) induces expression of miRNA-126 to modulate Vegf signaling [67].

The miRNA 221

Formation of new vessels in normal and pathological conditions requires coordination of distinct cell behaviors between tip and stalk cells [126]. The Vegf and Notch signaling pathways govern this process. Recently, through deep sequencing and functional screening in zebrafish, a new microRNA has been associated to this process [65]. MiRNA-221 has been found to be essential for angiogenesis since it is required for tip cell proliferation and

migration. miRNA-221 promoted tip cell behavior through repression of two targets: cyclin-dependent kinase inhibitor 1b (*cdkn1b*) and phosphoinositide-3-kinase regulatory subunit 1 (*pik3r1*). Importantly, these results in zebrafish indicate the central role of miRNA-221 as a regulatory node through which tip cell migration and proliferation are controlled during angiogenesis [65, 127]. A main role for miRNA-221/222 in endothelial therapeutic applications in humans has also been proposed [128]. In fact, recent work demonstrate that miRNA-221 is regulated by hyperglycemia in HUVEC cells, and lipid-lowering therapies (LLT) with atorvastatin increases EPC numbers and decreases miRNA-221/222 levels in patients with coronary artery disease [129, 130].

Conclusion

In this review, we pointed out the important role of the vertebrate *Danio rerio*, commonly called “zebrafish,” in helping to understand microRNA functions in heart and vascular cells not only during development but also in pathological conditions.

Recent discoveries have demonstrated that the zebrafish system can be used to study tissue regeneration since adult zebrafish can effectively regenerate injured hearts submitted to partial surgical amputation of the ventricle area in vivo. Limited information is currently available about the molecular mechanisms that regulate this process [131]. In particular, which microRNAs can be involved in this process remains unclear. Identification of microRNAs active during the regenerative potential of vertebrate/zebrafish tissue may implicate the discovery of new therapeutic approaches to supplement or replace conventional pharmacotherapeutic interventions in humans.

Recently, the possibility has also been suggested that microRNAs can be transported between cardiovascular tissues and regulate athero-vascular protection by an extracellular-vesicle-mediated mechanism [132, 133]. Demonstration of a role for circulating microRNAs could be validated in the zebrafish model system by microinjecting exogenous microRNA-containing liposomes in the blood flow of developing embryos to detect a functional role for mature microRNAs in cardiovascular tissues.

Interestingly, the zebrafish model can be used to perform pharmaceutical high-throughput modifier screens. The use of small RNA molecules such as microRNA products through such screens will serve as ideal entry points for novel drug development in the treatment of many vascular dysfunctions. Also, for miRNA-based therapeutic strategies the zebrafish model will come in handy. Antisense oligonucleotide approaches for inhibiting miRNA function and gain-of-function technologies for replacement

of miRNAs are currently being explored as tools for uncovering miRNA biology and as potential therapeutic agents. We think that in the near future significant progress in understanding miRNA biology will be derived from the zebrafish system, and this model will help to advance the technology for therapeutic modulation of miRNA activity in cardiovascular diseases.

Acknowledgments We apologize to the many researchers whose work was not cited in this review because of space limitations. We would like to thank all members of Santoro’s laboratory for support and discussion, Stefania Nicoli for sharing information before publication, Ellen Jane Corcoran for editorial assistance, and our anonymous reviewers for their insightful comments on this manuscript. Work in Massimo Santoro’s laboratory is supported by a HFSP Career Developmental Award, Marie Curie Action (IRG 247852), Telethon (GGP10195), and AIRC (MFAG 8911).

References

- Ambros V (2004) The functions of animal microRNAs. *Nature* 431(7006):350–355. doi:10.1038/nature02871
- Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116(2):281–297
- Bartel DP (2009) MicroRNAs: target recognition and regulatory functions. *Cell* 136(2):215–233. doi:10.1016/j.cell.2009.01.002
- Filipowicz W, Bhattacharyya SN, Sonenberg N (2008) Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 9(2):102–114. doi:10.1038/nrg2290
- Du T, Zamore PD (2005) microPrimer: the biogenesis and function of microRNA. *Development* 132(21):4645–4652. doi:10.1242/dev.02070
- Wienholds E, Kloosterman WP, Miska E, Alvarez-Saavedra E, Berezikov E, de Bruijn E, Horvitz HR, Kauppinen S, Plasterk RH (2005) MicroRNA expression in zebrafish embryonic development. *Science* 309(5732):310–311. doi:10.1126/science.1114519
- Krol J, Loedige I, Filipowicz W (2010) The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* 11(9):597–610. doi:10.1038/nrg2843
- Vasudevan S, Tong Y, Steitz JA (2007) Switching from repression to activation: microRNAs can up-regulate translation. *Science* 318(5858):1931–1934. doi:10.1126/science.1149460
- Seitz H (2009) Redefining microRNA targets. *Curr Biol* 19(10):870–873. doi:10.1016/j.cub.2009.03.059
- Davis BN, Hata A (2009) Regulation of MicroRNA Biogenesis: a miRiad of mechanisms. *Cell Commun Signal* 7:18. doi:10.1186/1478-811X-7-18
- Kai ZS, Pasquinelli AE (2010) MicroRNA assassins: factors that regulate the disappearance of miRNAs. *Nat Struct Mol Biol* 17(1):5–10. doi:10.1038/nsmb.1762
- Liu N, Olson EN (2010) MicroRNA regulatory networks in cardiovascular development. *Dev Cell* 18(4):510–525. doi:10.1016/j.devcel.2010.03.010
- Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, Conlon FL, Wang DZ (2006) The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 38(2):228–233. doi:10.1038/ng1725
- Zhao Z, Boyle TJ, Liu Z, Murray JJ, Wood WB, Waterston RH (2010) A negative regulatory loop between microRNA and Hox gene controls posterior identities in *Caenorhabditis elegans*. *PLoS genetics* 6(9). doi:10.1371/journal.pgen.1001089

15. Didiano D, Hobert O (2008) Molecular architecture of a miRNA-regulated 3' UTR. *RNA* 14(7):1297–1317. doi:[10.1261/rna.1082708](https://doi.org/10.1261/rna.1082708)
16. Plasterk RH (2006) Micro RNAs in animal development. *Cell* 124(5):877–881. doi:[10.1016/j.cell.2006.02.030](https://doi.org/10.1016/j.cell.2006.02.030)
17. Bushati N, Cohen SM (2008) MicroRNAs in neurodegeneration. *Curr Opin Neurobiol* 18(3):292–296. doi:[10.1016/j.conb.2008.07.001](https://doi.org/10.1016/j.conb.2008.07.001)
18. Bruneau BG (2008) The developmental genetics of congenital heart disease. *Nature* 451(7181):943–948. doi:[10.1038/nature06801](https://doi.org/10.1038/nature06801)
19. Olson EN, Schneider MD (2003) Sizing up the heart: development redux in disease. *Genes Dev* 17(16):1937–1956. doi:[10.1101/gad.1110103](https://doi.org/10.1101/gad.1110103)
20. Stainier DY (2001) Zebrafish genetics and vertebrate heart formation. *Nat Rev Genet* 2(1):39–48. doi:[10.1038/35047564](https://doi.org/10.1038/35047564)
21. Carmeliet P (2005) Angiogenesis in life, disease and medicine. *Nature* 438(7070):932–936. doi:[10.1038/nature04478](https://doi.org/10.1038/nature04478)
22. Red-Horse K, Crawford Y, Shojaei F, Ferrara N (2007) Endothelium-microenvironment interactions in the developing embryo and in the adult. *Dev Cell* 12(2):181–194. doi:[10.1016/j.devcel.2007.01.013](https://doi.org/10.1016/j.devcel.2007.01.013)
23. Jain RK (2003) Molecular regulation of vessel maturation. *Nat Med* 9(6):685–693. doi:[10.1038/nm0603-685](https://doi.org/10.1038/nm0603-685)
24. Santoro MM, Pesce G, Stainier DY (2009) Characterization of vascular mural cells during zebrafish development. *Mech Dev* 126(8–9):638–649. doi:[10.1016/j.mod.2009.06.1080](https://doi.org/10.1016/j.mod.2009.06.1080)
25. Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, Hannon GJ (2003) Dicer is essential for mouse development. *Nat Genet* 35(3):215–217. doi:[10.1038/ng1253](https://doi.org/10.1038/ng1253)
26. Murchison EP, Partridge JF, Tam OH, Cheloufi S, Hannon GJ (2005) Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci USA* 102(34):12135–12140. doi:[10.1073/pnas.0505479102](https://doi.org/10.1073/pnas.0505479102)
27. Yang WJ, Yang DD, Na S, Sandusky GE, Zhang Q, Zhao G (2005) Dicer is required for embryonic angiogenesis during mouse development. *J Biol Chem* 280(10):9330–9335. doi:[10.1074/jbc.M413394200](https://doi.org/10.1074/jbc.M413394200)
28. Giraldez AJ, Cinalli RM, Glasner ME, Enright AJ, Thomson JM, Baskerville S, Hammond SM, Bartel DP, Schier AF (2005) MicroRNAs regulate brain morphogenesis in zebrafish. *Science* 308(5723):833–838. doi:[10.1126/science.1109020](https://doi.org/10.1126/science.1109020)
29. Wienholds E, Koudijs MJ, van Eeden FJ, Cuppen E, Plasterk RH (2003) The microRNA-producing enzyme Dicer1 is essential for zebrafish development. *Nat Genet* 35(3):217–218. doi:[10.1038/ng1251](https://doi.org/10.1038/ng1251)
30. Harfe BD, McManus MT, Mansfield JH, Hornstein E, Tabin CJ (2005) The RNaseIII enzyme Dicer is required for morphogenesis but not patterning of the vertebrate limb. *Proc Natl Acad Sci USA* 102(31):10898–10903. doi:[10.1073/pnas.0504834102](https://doi.org/10.1073/pnas.0504834102)
31. Chen JF, Murchison EP, Tang R, Callis TE, Tatsuguchi M, Deng Z, Rojas M, Hammond SM, Schneider MD, Selzman CH, Meissner G, Patterson C, Hannon GJ, Wang DZ (2008) Targeted deletion of Dicer in the heart leads to dilated cardiomyopathy and heart failure. *Proc Natl Acad Sci USA* 105(6):2111–2116. doi:[10.1073/pnas.0710228105](https://doi.org/10.1073/pnas.0710228105)
32. Suarez Y, Fernandez-Hernando C, Yu J, Gerber SA, Harrison KD, Pober JS, Iruela-Arispe ML, Merkenschlager M, Sessa WC (2008) Dicer-dependent endothelial microRNAs are necessary for postnatal angiogenesis. *Proc Natl Acad Sci USA* 105(37):14082–14087. doi:[10.1073/pnas.0804597105](https://doi.org/10.1073/pnas.0804597105)
33. Albinsson S, Suarez Y, Skoura A, Offermanns S, Miano JM, Sessa WC (2010) MicroRNAs are necessary for vascular smooth muscle growth, differentiation, and function. *Arterioscler Thromb Vasc Biol* 30(6):1118–1126. doi:[10.1161/ATVBAHA.109.200873](https://doi.org/10.1161/ATVBAHA.109.200873)
34. Giraldez AJ, Mishima Y, Rihel J, Grocock RJ, Van Dongen S, Inoue K, Enright AJ, Schier AF (2006) Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* 312(5770):75–79. doi:[10.1126/science.1122689](https://doi.org/10.1126/science.1122689)
35. Lagendijk AK, Goumans MJ, Burkhard SB, Bakkers J (2011) MicroRNA-23 restricts cardiac valve formation by inhibiting Has2 and extracellular hyaluronic acid production. *Circ Res* 109(6):649–657. doi:[10.1161/CIRCRESAHA.111.247635](https://doi.org/10.1161/CIRCRESAHA.111.247635)
36. Santoro MM (2011) “Fishing” for endothelial microRNA functions and dysfunction. *Vascul Pharmacol* 55(4):60–68. doi:[10.1016/j.vph.2011.08.224](https://doi.org/10.1016/j.vph.2011.08.224)
37. Thisse C, Zon LI (2002) Organogenesis—heart and blood formation from the zebrafish point of view. *Science* 295(5554):457–462. doi:[10.1126/science.1063654](https://doi.org/10.1126/science.1063654)
38. Lieschke GJ, Currie PD (2007) Animal models of human disease: zebrafish swim into view. *Nat Rev Genet* 8(5):353–367. doi:[10.1038/nrg2091](https://doi.org/10.1038/nrg2091)
39. Skromne I, Prince VE (2008) Current perspectives in zebrafish reverse genetics: moving forward. *Dev Dyn* 237(4):861–882. doi:[10.1002/dvdy.21484](https://doi.org/10.1002/dvdy.21484)
40. Lawson ND, Weinstein BM (2002) Arteries and veins: making a difference with zebrafish. *Nat Rev Genet* 3(9):674–682. doi:[10.1038/nrg888](https://doi.org/10.1038/nrg888)
41. Stainier DY, Fouquet B, Chen JN, Warren KS, Weinstein BM, Meiler SE, Mohideen MA, Neuhauss SC, Solnica-Krezel L, Schier AF, Zwartkruis F, Stemple DL, Malicki J, Driever W, Fishman MC (1996) Mutations affecting the formation and function of the cardiovascular system in the zebrafish embryo. *Development* 123:285–292
42. Kawakami K (2007) Tol2: a versatile gene transfer vector in vertebrates. *Genome Biol* 8(Suppl 1):S7. doi:[10.1186/gb-2007-8-s1-s7](https://doi.org/10.1186/gb-2007-8-s1-s7)
43. Lawson ND, Weinstein BM (2002) In vivo imaging of embryonic vascular development using transgenic zebrafish. *Dev Biol* 248(2):307–318
44. Jin SW, Beis D, Mitchell T, Chen JN, Stainier DY (2005) Cellular and molecular analyses of vascular tube and lumen formation in zebrafish. *Development* 132(23):5199–5209. doi:[10.1242/dev.02087](https://doi.org/10.1242/dev.02087)
45. Traver D, Paw BH, Poss KD, Penberthy WT, Lin S, Zon LI (2003) Transplantation and in vivo imaging of multilineage engraftment in zebrafish bloodless mutants. *Nat Immunol* 4(12):1238–1246. doi:[10.1038/ni1007](https://doi.org/10.1038/ni1007)
46. Lin HF, Traver D, Zhu H, Dooley K, Paw BH, Zon LI, Handin RI (2005) Analysis of thrombocyte development in CD41-GFP transgenic zebrafish. *Blood* 106(12):3803–3810. doi:[10.1182/blood-2005-01-0179](https://doi.org/10.1182/blood-2005-01-0179)
47. Burns CG, Milan DJ, Grande EJ, Rottbauer W, MacRae CA, Fishman MC (2005) High-throughput assay for small molecules that modulate zebrafish embryonic heart rate. *Nat Chem Biol* 1(5):263–264. doi:[10.1038/nchembio732](https://doi.org/10.1038/nchembio732)
48. Seiler C, Abrams J, Pack M (2010) Characterization of zebrafish intestinal smooth muscle development using a novel sm22alpha-b promoter. *Dev Dyn* 239(11):2806–2812. doi:[10.1002/dvdy.22420](https://doi.org/10.1002/dvdy.22420)
49. Scherz PJ, Huisken J, Sahai-Hernandez P, Stainier DY (2008) High-speed imaging of developing heart valves reveals interplay of morphogenesis and function. *Development* 135(6):1179–1187. doi:[10.1242/dev.010694](https://doi.org/10.1242/dev.010694)
50. Keller PJ, Schmidt AD, Wittbrodt J, Stelzer EH (2008) Reconstruction of zebrafish early embryonic development by scanned light sheet microscopy. *Science* 322(5904):1065–1069. doi:[10.1126/science.1162493](https://doi.org/10.1126/science.1162493)

51. Bakkers J (2011) Zebrafish as a model to study cardiac development and human cardiac disease. *Cardiovasc Res* 91(2):279–288. doi:[10.1093/cvr/cvr098](https://doi.org/10.1093/cvr/cvr098)
52. Tu S, Chi NC (2012) Zebrafish models in cardiac development and congenital heart birth defects. *Differentiation*. doi:[10.1016/j.diff.2012.05.005](https://doi.org/10.1016/j.diff.2012.05.005)
53. Jin SW, Herzog W, Santoro MM, Mitchell TS, Frantsve J, Jungblut B, Beis D, Scott IC, D'Amico LA, Ober EA, Verkade H, Field HA, Chi NC, Wehman AM, Baier H, Stainier DY (2007) A transgene-assisted genetic screen identifies essential regulators of vascular development in vertebrate embryos. *Dev Biol* 307(1):29–42. doi:[10.1016/j.ydbio.2007.03.526](https://doi.org/10.1016/j.ydbio.2007.03.526)
54. Dahme T, Katus HA, Rottbauer W (2009) Fishing for the genetic basis of cardiovascular disease. *Dis Models Mech* 2(1–2):18–22. doi:[10.1242/dmm.000687](https://doi.org/10.1242/dmm.000687)
55. Wienholds E, Plasterk RH (2005) MicroRNA function in animal development. *FEBS Lett* 579(26):5911–5922. doi:[10.1016/j.febslet.2005.07.070](https://doi.org/10.1016/j.febslet.2005.07.070)
56. Schier AF, Giraldez AJ (2006) MicroRNA function and mechanism: insights from zebra fish. *Cold Spring Harb Symp Quant Biol* 71:195–203. doi:[10.1101/sqb.2006.71.055](https://doi.org/10.1101/sqb.2006.71.055)
57. Begemann G (2008) MicroRNAs and RNA interference in zebrafish development. *Zebrafish* 5(2):111–119. doi:[10.1089/zeb.2008.0528](https://doi.org/10.1089/zeb.2008.0528)
58. Takacs CM, Giraldez AJ (2010) MicroRNAs as genetic sculptors: fishing for clues. *Semin Cell Dev Biol* 21(7):760–767. doi:[10.1016/j.semcdb.2010.02.003](https://doi.org/10.1016/j.semcdb.2010.02.003)
59. Chen PY, Manninga H, Slanchev K, Chien M, Russo JJ, Ju J, Sheridan R, John B, Marks DS, Gaidatzis D, Sander C, Zavolan M, Tuschl T (2005) The developmental miRNA profiles of zebrafish as determined by small RNA cloning. *Genes Dev* 19(11):1288–1293. doi:[10.1101/gad.1310605](https://doi.org/10.1101/gad.1310605)
60. Kloosterman WP, Steiner FA, Berezikov E, de Bruijn E, van de Belt J, Verheul M, Cuppen E, Plasterk RH (2006) Cloning and expression of new microRNAs from zebrafish. *Nucleic Acids Res* 34(9):2558–2569. doi:[10.1093/nar/gkl278](https://doi.org/10.1093/nar/gkl278)
61. Thatcher EJ, Bond J, Paydar I, Patton JG (2008) Genomic organization of zebrafish microRNAs. *BMC Genom* 9:253. doi:[10.1186/1471-2164-9-253](https://doi.org/10.1186/1471-2164-9-253)
62. Thatcher EJ, Flynt AS, Li N, Patton JR, Patton JG (2007) MiRNA expression analysis during normal zebrafish development and following inhibition of the Hedgehog and Notch signaling pathways. *Dev Dyn* 236(8):2172–2180. doi:[10.1002/dvdy.21211](https://doi.org/10.1002/dvdy.21211)
63. Kloosterman WP, Wienholds E, de Bruijn E, Kauppinen S, Plasterk RH (2006) In situ detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. *Nat Methods* 3(1):27–29. doi:[10.1038/nmeth843](https://doi.org/10.1038/nmeth843)
64. He X, Yan YL, DeLaurier A, Postlethwait JH (2011) Observation of miRNA gene expression in zebrafish embryos by in situ hybridization to microRNA primary transcripts. *Zebrafish* 8(1):1–8. doi:[10.1089/zeb.2010.0680](https://doi.org/10.1089/zeb.2010.0680)
65. Nicoli S, Knyphausen CP, Zhu LJ, Lakshmanan A, Lawson ND (2012) miR-221 is required for endothelial tip cell behaviors during vascular development. *Dev Cell* 22(2):418–429
66. Zeng L, Carter AD, Childs SJ (2009) miR-145 directs intestinal maturation in zebrafish. *Proc Natl Acad Sci USA* 106(42):17793–17798. doi:[10.1073/pnas.0903693106](https://doi.org/10.1073/pnas.0903693106)
67. Nicoli S, Standley C, Walker P, Hurlstone A, Fogarty KE, Lawson ND (2010) MicroRNA-mediated integration of haemodynamics and Vegf signalling during angiogenesis. *Nature* 464(7292):1196–1200. doi:[10.1038/nature08889](https://doi.org/10.1038/nature08889)
68. Hans S, Freudenreich D, Geffarth M, Kaslin J, Machate A, Brand M (2011) Generation of a non-leaky heat shock-inducible Cre line for conditional Cre/lox strategies in zebrafish. *Dev Dyn* 240(1):108–115. doi:[10.1002/dvdy.22497](https://doi.org/10.1002/dvdy.22497)
69. Mosimann C, Kaufman CK, Li P, Pugach EK, Tamplin OJ, Zon LI (2011) Ubiquitous transgene expression and Cre-based recombination driven by the ubiquitin promoter in zebrafish. *Development* 138(1):169–177. doi:[10.1242/dev.059345](https://doi.org/10.1242/dev.059345)
70. Morton SU, Scherz PJ, Cordes KR, Ivey KN, Stainier DY, Srivastava D (2008) microRNA-138 modulates cardiac patterning during embryonic development. *Proc Natl Acad Sci USA* 105(46):17830–17835. doi:[10.1073/pnas.0804673105](https://doi.org/10.1073/pnas.0804673105)
71. Nasevicius A, Ekker SC (2000) Effective targeted gene ‘knockdown’ in zebrafish. *Nat Genet* 26(2):216–220. doi:[10.1038/79951](https://doi.org/10.1038/79951)
72. Chen E, Ekker SC (2004) Zebrafish as a genomics research model. *Curr Pharm Biotechnol* 5(5):409–413
73. Eisen JS, Smith JC (2008) Controlling morpholino experiments: don't stop making antisense. *Development* 135(10):1735–1743. doi:[10.1242/dev.001115](https://doi.org/10.1242/dev.001115)
74. Flynt AS, Li N, Thatcher EJ, Solnica-Krezel L, Patton JG (2007) Zebrafish miR-214 modulates Hedgehog signaling to specify muscle cell fate. *Nat Genet* 39(2):259–263. doi:[10.1038/ng1953](https://doi.org/10.1038/ng1953)
75. Kloosterman WP, Lagendijk AK, Ketting RF, Moulton JD, Plasterk RH (2007) Targeted inhibition of miRNA maturation with morpholinos reveals a role for miR-375 in pancreatic islet development. *PLoS Biol* 5(8):e203. doi:[10.1371/journal.pbio.0050203](https://doi.org/10.1371/journal.pbio.0050203)
76. Brown BD, Naldini L (2009) Exploiting and antagonizing microRNA regulation for therapeutic and experimental applications. *Nat Rev Genet* 10(8):578–585. doi:[10.1038/nrg2628](https://doi.org/10.1038/nrg2628)
77. Ebert MS, Sharp PA (2010) Emerging roles for natural microRNA sponges. *Curr Biol* 20(19):R858–R861. doi:[10.1016/j.cub.2010.08.052](https://doi.org/10.1016/j.cub.2010.08.052)
78. Brown BD, Gentner B, Cantore A, Colleoni S, Amendola M, Zingale A, Baccarini A, Lazzari G, Galli C, Naldini L (2007) Endogenous microRNA can be broadly exploited to regulate transgene expression according to tissue, lineage and differentiation state. *Nat Biotechnol* 25(12):1457–1467. doi:[10.1038/nbt1372](https://doi.org/10.1038/nbt1372)
79. Yin VP, Lepilina A, Smith A, Poss KD (2012) Regulation of zebrafish heart regeneration by miR-133. *Dev Biol* 365(2):319–327. doi:[10.1016/j.ydbio.2012.02.018](https://doi.org/10.1016/j.ydbio.2012.02.018)
80. Zhu C, Smith T, McNulty J, Rayla AL, Lakshmanan A, Siekmann AF, Buffardi M, Meng X, Shin J, Padmanabhan A, Cifuentes D, Giraldez AJ, Look AT, Epstein JA, Lawson ND, Wolfe SA (2011) Evaluation and application of modularly assembled zinc-finger nucleases in zebrafish. *Development* 138(20):4555–4564. doi:[10.1242/dev.066779](https://doi.org/10.1242/dev.066779)
81. Lawson ND, Wolfe SA (2011) Forward and reverse genetic approaches for the analysis of vertebrate development in the zebrafish. *Dev Cell* 21(1):48–64. doi:[10.1016/j.devcel.2011.06.007](https://doi.org/10.1016/j.devcel.2011.06.007)
82. Cade L, Reyon D, Hwang WY, Tsai SQ, Patel S, Khayter C, Joung JK, Sander JD, Peterson RT, Yeh JR (2012) Highly efficient generation of heritable zebrafish gene mutations using homo- and heterodimeric TALENs. *Nucleic Acids Res*. doi:[10.1093/nar/gks518](https://doi.org/10.1093/nar/gks518)
83. Moore FE, Reyon D, Sander JD, Martinez SA, Blackburn JS, Khayter C, Ramirez CL, Joung JK, Langenau DM (2012) Improved somatic mutagenesis in zebrafish using transcription activator-like effector nucleases (TALENs). *PLoS ONE* 7(5):e37877. doi:[10.1371/journal.pone.0037877](https://doi.org/10.1371/journal.pone.0037877)
84. Yoon S, De Micheli G (2006) Computational identification of microRNAs and their targets. *Birth Defects Res Part C Embryo Today Rev* 78(2):118–128. doi:[10.1002/bdrc.20067](https://doi.org/10.1002/bdrc.20067)
85. Witkos TM, Koscińska E, Krzyżosiak WJ (2011) Practical aspects of microRNA target prediction. *Curr Mol Med* 11(2):93–109
86. Min H, Yoon S (2010) Got target? Computational methods for microRNA target prediction and their extension. *Exp Mol Med* 42(4):233–244

87. Dangwal S, Bang C, Thum T (2011) Novel techniques and targets in cardiovascular microRNA research. *Cardiovasc Res*. doi:[10.1093/cvr/cvr297](https://doi.org/10.1093/cvr/cvr297)
88. Pase L, Lieschke GJ (2009) Validating microRNA target transcripts using zebrafish assays. *Methods Mol Biol* 546:227–240. doi:[10.1007/978-1-60327-977-2_14](https://doi.org/10.1007/978-1-60327-977-2_14)
89. Choi J, Dong L, Ahn J, Dao D, Hammerschmidt M, Chen JN (2007) FoxH1 negatively modulates flk1 gene expression and vascular formation in zebrafish. *Dev Biol* 304(2):735–744. doi:[10.1016/j.ydbio.2007.01.023](https://doi.org/10.1016/j.ydbio.2007.01.023)
90. Staton AA, Giraldez AJ (2011) Use of target protector morpholinos to analyze the physiological roles of specific miRNA-mRNA pairs in vivo. *Nat Protoc* 6(12):2035–2049. doi:[10.1038/nprot.2011.423](https://doi.org/10.1038/nprot.2011.423)
91. Staton AA, Knaut H, Giraldez AJ (2011) miRNA regulation of Sdf1 chemokine signaling provides genetic robustness to germ cell migration. *Nat Genet* 43(3):204–211. doi:[10.1038/ng.758](https://doi.org/10.1038/ng.758)
92. Armstrong EJ, Bischoff J (2004) Heart valve development: endothelial cell signaling and differentiation. *Circ Res* 95(5):459–470. doi:[10.1161/01.RES.0000141146.95728.da](https://doi.org/10.1161/01.RES.0000141146.95728.da)
93. Camenisch TD, Spicer AP, Brehm-Gibson T, Biesterfeldt J, Augustine ML, Calabro A Jr, Kubalak S, Klewer SE, McDonald JA (2000) Disruption of hyaluronan synthase-2 abrogates normal cardiac morphogenesis and hyaluronan-mediated transformation of epithelium to mesenchyme. *J Clin Invest* 106(3):349–360. doi:[10.1172/JCI10272](https://doi.org/10.1172/JCI10272)
94. Srivastava D (2006) Making or breaking the heart: from lineage determination to morphogenesis. *Cell* 126(6):1037–1048. doi:[10.1016/j.cell.2006.09.003](https://doi.org/10.1016/j.cell.2006.09.003)
95. Niederreither K, Subbarayan V, Dolle P, Chambon P (1999) Embryonic retinoic acid synthesis is essential for early mouse post-implantation development. *Nat Genet* 21(4):444–448. doi:[10.1038/7788](https://doi.org/10.1038/7788)
96. Cordes KR, Sheehy NT, White MP, Berry EC, Morton SU, Muth AN, Lee TH, Miano JM, Ivey KN, Srivastava D (2009) miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature* 460(7256):705–710. doi:[10.1038/nature08195](https://doi.org/10.1038/nature08195)
97. Elia L, Quintavalle M, Zhang J, Contu R, Cossu L, Latronico MV, Peterson KL, Indolfi C, Catalucci D, Chen J, Courtneidge SA, Condorelli G (2009) The knockout of miR-143 and -145 alters smooth muscle cell maintenance and vascular homeostasis in mice: correlates with human disease. *Cell Death Differ* 16(12):1590–1598. doi:[10.1038/cdd.2009.153](https://doi.org/10.1038/cdd.2009.153)
98. Xin M, Small EM, Sutherland LB, Qi X, McAnally J, Plato CF, Richardson JA, Bassel-Duby R, Olson EN (2009) MicroRNAs miR-143 and miR-145 modulate cytoskeletal dynamics and responsiveness of smooth muscle cells to injury. *Genes Dev* 23(18):2166–2178. doi:[10.1101/gad.1842409](https://doi.org/10.1101/gad.1842409)
99. Boettger T, Beetz N, Kostin S, Schneider J, Kruger M, Hein L, Braun T (2009) Acquisition of the contractile phenotype by murine arterial smooth muscle cells depends on the Mir143/145 gene cluster. *J Clin Invest* 119(9):2634–2647. doi:[10.1172/JCI38864](https://doi.org/10.1172/JCI38864)
100. Miyasaka KY, Kida YS, Banjo T, Ueki Y, Nagayama K, Matsumoto T, Sato M, Ogura T (2011) Heartbeat regulates cardiogenesis by suppressing retinoic acid signaling via expression of miR-143. *Mech Dev* 128(1–2):18–28. doi:[10.1016/j.mod.2010.09.002](https://doi.org/10.1016/j.mod.2010.09.002)
101. Deacon DC, Nevis KR, Cashman TJ, Zhou Y, Zhao L, Washko D, Guner-Ataman B, Burns CG, Burns CE (2010) The miR-143-adducin3 pathway is essential for cardiac chamber morphogenesis. *Development* 137(11):1887–1896. doi:[10.1242/dev.050526](https://doi.org/10.1242/dev.050526)
102. Fish JE, Wythe JD, Xiao T, Bruneau BG, Stainier DY, Srivastava D, Woo S (2011) A Slit/miR-218/Robo regulatory loop is required during heart tube formation in zebrafish. *Development* 138(7):1409–1419. doi:[10.1242/dev.060046](https://doi.org/10.1242/dev.060046)
103. Small EM, Sutherland LB, Rajagopalan KN, Wang S, Olson EN (2010) MicroRNA-218 regulates vascular patterning by modulation of Slit-Robo signaling. *Circ Res* 107(11):1336–1344. doi:[10.1161/CIRCRESAHA.110.227926](https://doi.org/10.1161/CIRCRESAHA.110.227926)
104. Poss KD, Wilson LG, Keating MT (2002) Heart regeneration in zebrafish. *Science* 298(5601):2188–2190. doi:[10.1126/science.1077857](https://doi.org/10.1126/science.1077857)
105. Jopling C, Sleep E, Raya M, Marti M, Raya A, Izpisua Belmonte JC (2010) Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature* 464(7288):606–609. doi:[10.1038/nature08899](https://doi.org/10.1038/nature08899)
106. Kikuchi K, Holdway JE, Werdich AA, Anderson RM, Fang Y, Egnaczyk GF, Evans T, Macrae CA, Stainier DY, Poss KD (2010) Primary contribution to zebrafish heart regeneration by gata4(+) cardiomyocytes. *Nature* 464(7288):601–605. doi:[10.1038/nature08804](https://doi.org/10.1038/nature08804)
107. Poss KD, Nechiporuk A, Hillam AM, Johnson SL, Keating MT (2002) Mps1 defines a proximal blastemal proliferative compartment essential for zebrafish fin regeneration. *Development* 129(22):5141–5149
108. Bruce AF, Rothery S, Dupont E, Severs NJ (2008) Gap junction remodelling in human heart failure is associated with increased interaction of connexin43 with ZO-1. *Cardiovasc Res* 77(4):757–765. doi:[10.1093/cvr/cvm083](https://doi.org/10.1093/cvr/cvm083)
109. Liu N, Bezprozvannaya S, Williams AH, Qi X, Richardson JA, Bassel-Duby R, Olson EN (2008) microRNA-133a regulates cardiomyocyte proliferation and suppresses smooth muscle gene expression in the heart. *Genes Dev* 22(23):3242–3254. doi:[10.1101/gad.1738708](https://doi.org/10.1101/gad.1738708)
110. Fiedler J, Jazbutyte V, Kirchmaier BC, Gupta SK, Lorenzen J, Hartmann D, Galuppo P, Kneitz S, Pena JT, Sohn-Lee C, Loyer X, Soutschek J, Brand T, Tuschl T, Heineke J, Martin U, Schulte-Merker S, Ertl G, Engelhardt S, Bauersachs J, Thum T (2011) MicroRNA-24 regulates vascularity after myocardial infarction. *Circulation* 124(6):720–730. doi:[10.1161/CIRCULATIONAHA.111.039008](https://doi.org/10.1161/CIRCULATIONAHA.111.039008)
111. Bang C, Fiedler J, Thum T (2011) Cardiovascular importance of the microRNA-23/27/24 family. *Microcirculation*. doi:[10.1111/j.1549-8719.2011.00153.x](https://doi.org/10.1111/j.1549-8719.2011.00153.x)
112. Pedrioli DM, Karpanen T, Dabouras V, Jurisic G, van de Hoek G, Shin JW, Marino D, Kalin RE, Leidel S, Cinelli P, Schulte-Merker S, Brandli AW, Detmar M (2010) miR-31 functions as a negative regulator of lymphatic vascular lineage-specific differentiation in vitro and vascular development in vivo. *Mol Cell Biol* 30(14):3620–3634. doi:[10.1128/MCB.00185-10](https://doi.org/10.1128/MCB.00185-10)
113. Dews M, Homayouni A, Yu D, Murphy D, Seignani C, Wentzel E, Furth EE, Lee WM, Enders GH, Mendell JT, Thomas-Tikhonenko A (2006) Augmentation of tumor angiogenesis by a Myc-activated microRNA cluster. *Nat Genet* 38(9):1060–1065. doi:[10.1038/ng1855](https://doi.org/10.1038/ng1855)
114. Bonauer A, Carmona G, Iwasaki M, Mione M, Koyanagi M, Fischer A, Burchfield J, Fox H, Doebele C, Ohtani K, Chavakis E, Potente M, Tjwa M, Urbich C, Zeiher AM, Dimmeler S (2009) MicroRNA-92a controls angiogenesis and functional recovery of ischemic tissues in mice. *Science* 324(5935):1710–1713. doi:[10.1126/science.1174381](https://doi.org/10.1126/science.1174381)
115. Francis SE, Goh KL, Hodivala-Dilke K, Bader BL, Stark M, Davidson D, Hynes RO (2002) Central roles of alpha5beta1 integrin and fibronectin in vascular development in mouse embryos and embryoid bodies. *Arterioscler Thromb Vasc Biol* 22(6):927–933
116. Wu W, Xiao H, Laguna-Fernandez A, Villarreal G Jr, Wang KC, Geary GG, Zhang Y, Wang WC, Huang HD, Zhou J, Li YS, Chien S, Garcia-Cardena G, Shyy JY (2011) Flow-dependent regulation of Kruppel-like factor 2 is mediated by microRNA-92a. *Circulation* 124(5):633–641. doi:[10.1161/CIRCULATIONAHA.110.005108](https://doi.org/10.1161/CIRCULATIONAHA.110.005108)

117. Suarez Y, Fernandez-Hernando C, Pober JS, Sessa WC (2007) Dicer dependent microRNAs regulate gene expression and functions in human endothelial cells. *Circ Res* 100(8):1164–1173. doi:[10.1161/01.RES.0000265065.26744.17](https://doi.org/10.1161/01.RES.0000265065.26744.17)
118. Polisenio L, Tuccoli A, Mariani L, Evangelista M, Citti L, Woods K, Mercatanti A, Hammond S, Rainaldi G (2006) MicroRNAs modulate the angiogenic properties of HUVECs. *Blood* 108(9):3068–3071. doi:[10.1182/blood-2006-01-012369](https://doi.org/10.1182/blood-2006-01-012369)
119. Fish JE, Santoro MM, Morton SU, Yu S, Yeh RF, Wythe JD, Ivey KN, Bruneau BG, Stainier DY, Srivastava D (2008) miR-126 regulates angiogenic signaling and vascular integrity. *Dev Cell* 15(2):272–284. doi:[10.1016/j.devcel.2008.07.008](https://doi.org/10.1016/j.devcel.2008.07.008)
120. Kuhnert F, Mancuso MR, Hampton J, Stankunas K, Asano T, Chen CZ, Kuo CJ (2008) Attribution of vascular phenotypes of the murine *Egfl7* locus to the microRNA miR-126. *Development* 135(24):3989–3993. doi:[10.1242/dev.029736](https://doi.org/10.1242/dev.029736)
121. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, Richardson JA, Bassel-Duby R, Olson EN (2008) The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Dev Cell* 15(2):261–271. doi:[10.1016/j.devcel.2008.07.002](https://doi.org/10.1016/j.devcel.2008.07.002)
122. Stankunas K, Ma GK, Kuhnert FJ, Kuo CJ, Chang CP (2010) VEGF signaling has distinct spatiotemporal roles during heart valve development. *Dev Biol* 347(2):325–336. doi:[10.1016/j.ydbio.2010.08.030](https://doi.org/10.1016/j.ydbio.2010.08.030)
123. Wakioka T, Sasaki A, Kato R, Shouda T, Matsumoto A, Miyoshi K, Tsuneoka M, Komiya S, Baron R, Yoshimura A (2001) Sprd is a sprouty-related suppressor of Ras signalling. *Nature* 412(6847):647–651. doi:[10.1038/35088082](https://doi.org/10.1038/35088082)
124. Ueki K, Fruman DA, Yballe CM, Fasshauer M, Klein J, Asano T, Cantley LC, Kahn CR (2003) Positive and negative roles of p85 alpha and p85 beta regulatory subunits of phosphoinositide 3-kinase in insulin signaling. *J Biol Chem* 278(48):48453–48466. doi:[10.1074/jbc.M305602200](https://doi.org/10.1074/jbc.M305602200)
125. Fish JE, Srivastava D (2009) MicroRNAs: opening a new vein in angiogenesis research. *Science signaling* 2 (52):pe1. doi:[10.1126/scisignal.252pe1](https://doi.org/10.1126/scisignal.252pe1)
126. Potente M, Gerhardt H, Carmeliet P (2011) Basic and therapeutic aspects of angiogenesis. *Cell* 146(6):873–887. doi:[10.1016/j.cell.2011.08.039](https://doi.org/10.1016/j.cell.2011.08.039)
127. Kuehnbacher A, Urbich C, Zeiher AM, Dimmeler S (2007) Role of Dicer and Drosha for endothelial microRNA expression and angiogenesis. *Circ Res* 101(1):59–68. doi:[10.1161/CIRCRES.AHA.107.153916](https://doi.org/10.1161/CIRCRES.AHA.107.153916)
128. Liu X, Cheng Y, Yang J, Xu L, Zhang C (2012) Cell-specific effects of miR-221/222 in vessels: molecular mechanism and therapeutic application. *J Mol Cell Cardiol* 52(1):245–255. doi:[10.1016/j.yjmcc.2011.11.008](https://doi.org/10.1016/j.yjmcc.2011.11.008)
129. Li Y, Song YH, Li F, Yang T, Lu YW, Geng YJ (2009) MicroRNA-221 regulates high glucose-induced endothelial dysfunction. *Biochem Biophys Res Commun* 381(1):81–83. doi:[10.1016/j.bbrc.2009.02.013](https://doi.org/10.1016/j.bbrc.2009.02.013)
130. Minami Y, Satoh M, Maesawa C, Takahashi Y, Tabuchi T, Itoh T, Nakamura M (2009) Effect of atorvastatin on microRNA 221/222 expression in endothelial progenitor cells obtained from patients with coronary artery disease. *Eur J Clin Invest* 39(5):359–367. doi:[10.1111/j.1365-2362.2009.02110.x](https://doi.org/10.1111/j.1365-2362.2009.02110.x)
131. Poss KD (2007) Getting to the heart of regeneration in zebrafish. *Semin Cell Dev Biol* 18(1):36–45. doi:[10.1016/j.semcdb.2006.11.009](https://doi.org/10.1016/j.semcdb.2006.11.009)
132. Zerneck A, Bidzhekov K, Noels H, Shagdarsuren E, Gan L, Denecke B, Hristov M, Koppel T, Jahantigh MN, Lutgens E, Wang S, Olson EN, Schober A, Weber C (2009) Delivery of microRNA-126 by apoptotic bodies induces CXCL12-dependent vascular protection. *Sci Signal* 2(100):ra81. doi:[10.1126/scisignal.2000610](https://doi.org/10.1126/scisignal.2000610)
133. Hergenreider E, Heydt S, Treguer K, Boettger T, Horrevoets AJ, Zeiher AM, Scheffer MP, Frangakis AS, Yin X, Mayr M, Braun T, Urbich C, Boon RA, Dimmeler S (2012) Atheroprotective communication between endothelial cells and smooth muscle cells through miRNAs. *Nat Cell Biol*. doi:[10.1038/ncb2441](https://doi.org/10.1038/ncb2441)