

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cancer transition and priorities for cancer control

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/122502> since

Published version:

DOI:10.1016/S1470-2045(12)70268-1

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Lancet Oncology, 13 (8), 2012, DOI: 10.1016/S1470-2045(12)70268-1]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://www.sciencedirect.com/science/article/pii/S1470204512702681?via=ihub>]

Cancer transition and priorities for cancer control

Milena Maulea, Franco Merletti

Refers To

Freddie Bray, Ahmedin Jemal, Nathan Grey, Jacques Ferlay, David Forman

Global cancer transitions according to the Human Development Index (2008–2030): a population-based study

The Lancet Oncology, Volume 13, Issue 8, August 2012, Pages 790-801

Referred to by

Freddie Bray, Ahmedin Jemal, Nathan Grey, Jacques Ferlay, David Forman

Global cancer transitions according to the Human Development Index (2008–2030): a population-based study

The Lancet Oncology, Volume 13, Issue 8, August 2012, Pages 790-801

In *The Lancet Oncology*, Freddie Bray and colleagues¹ assess worldwide patterns of cancer burden, in terms of both incidence and mortality, and predict future scenarios in relation to different levels of socioeconomic development, which they measure using the Human Development Index (HDI). Their paper, which provides a good explanation of the theory of cancer transition, serves both purposes of research and guidance in setting priorities for cancer control.

Cancer transition can be regarded as an extension or completion of Omran's theory on epidemiological transition.² and ³ In analogy with the third stage of epidemiological transition—a shift from infectious to non-communicable diseases—the theory of cancer transition sees a shift from a predominance of cancers linked to infections to cancers associated with risk factors that are mainly non-infectious and possibly related to a so-called western lifestyle. So, whereas in very high and high HDI areas four cancer types (lung, female breast, colorectum, and prostate) account for almost half the total cancer incidence burden, in middle and low HDI areas, in addition to cancers of the lung, colorectum, and female breast, cancers of the stomach, liver, and cervix account for a substantial burden of cancer.¹ In areas with low HDI, the most frequent cancer types seem to be those of the cervix, female breast, liver, Kaposi's sarcoma, and non-Hodgkin lymphoma.¹

Bray and colleagues' findings¹ show that, in 2008, the largest cancer burden was in very high HDI areas, which bore almost 40% of the global incidence burden despite accounting for only 15% of the world's population; low HDI areas bore only 2% of the burden, with a population accounting for nearly 6% of the global population. However, Bray and colleagues' projections for 2030 predict the largest absolute increase in low HDI areas. And the proportion of the global burden of cancer borne in low HDI areas might be higher if only the population older than 50 years are considered—people in this age group are most susceptible to cancer and are likely to account for a smaller proportion of the national population in low HDI countries. Furthermore, with respect to projected burdens, the socioeconomic level in low HDI countries is expected to improve (as implicit in the demographic projections that assume decreasing population growth), which will in turn increase the incidence of cancer because of both population ageing and the spread of new risk factors. For these reasons, cancer burden in lower HDI areas will probably become a more serious problem in the future than it seems now.

The analysis of cancer burden and trends together with future projections shows where different populations stand along the epidemic curve and might help to identify the most useful interventions.

The knowledge gained in areas where more advanced stages of cancer transition have been reached is crucial to guide implementation of preventive action in those that still lag behind.

In the theory of epidemiological transition,³ the shift from infections to degenerative diseases in Europe was not associated with progress in medical science but rather with improved socioeconomic conditions (that in turn improved nutrition and hygiene).⁴ Medical progress became more important at a later stage in high-income countries, but was seen to have a great effect in low-income countries, causing accelerated epidemiological transitions. Should the analogy with Omran's theory hold, implementation of preventive measures to eradicate known risk factors in low and middle HDI countries could be most effective and could change the shape of the cancer epidemiological curve for most of the world's population (more than 70% putting together middle and low HDI countries). Public health programmes should include campaigns for immunisation,⁵ tobacco control, reduction in alcohol consumption and obesity,⁶ and the implementation of cancer-screening programmes that have been shown to reduce mortality effectively.⁷

In very high HDI areas, there is strong evidence of the simultaneous start of decline of lung cancer incidence,⁸ at least in men, and the rise and increasing relative importance of other cancer types (eg, kidney, pancreas,⁸ testis⁹), often with unknown cause or known risk factors that explain only a small fraction of cases. In such areas, which are at the front edge of the cancer epidemic curve, has cancer transition theory been fully realised or has a new epidemic stage—a shift to other cancer types related to risk factors that are still unknown—become apparent? The answer and the possible strategies to counteract the rise of new diseases depend entirely on our ability to understand the patterns of their emergence and identify their causes.

We declare that we have no conflicts of interest.

References

1

F Bray, A Jemal, N Grey, J Ferlay, D Forman

Global cancer transitions according to the Human Development Index (2008–2030): a population-based study

Lancet Oncol, 13 (2012), pp. 790–801

2

O Gersten, JR Wilmoth

The cancer transition in Japan since 1951

Demographic Research, 7 (2002), pp. 271–306

3

AR Omran

The epidemiologic transition. A theory of the epidemiology of population change

Milbank Mem Fund Q, 49 (1971), pp. 509–538

4

T McKeown, RG Brown

Medical evidence related to English population change in the eighteenth century

Population Studies, 9 (1955), pp. 119–141

5

BS Sylla, CP Wild

A million africans a year dying from cancer by 2030: what can cancer research and control offer to the continent?

Int J Cancer, 130 (2011), pp. 245–250

6

R Beaglehole, R Bonita, R Horton et al.

Priority actions for the non-communicable disease crisis

Lancet, 377 (2011), pp. 1438–1447

7

J Peto, C Gilham, O Fletcher, FE Matthews

The cervical cancer epidemic that screening has prevented in the UK

Lancet, 364 (2004), pp. 249–256

8

BA Kohler, E Ward, BJ McCarthy et al.

Annual report to the nation on the status of cancer, 1975–2007, featuring tumors of the brain and other nervous system

J Natl Cancer Inst, 103 (2011), pp. 714–736

9

VM Chia, SM Quraishi, SS Devesa, MP Purdue, MB Cook, KA McGlynn

International trends in the incidence of testicular cancer, 1973–2002

Cancer Epidemiol Biomarkers Prev, 19 (2010), pp. 1151–1159