



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

An exoproteome approach to monitor safety of a cheese-isolated Lactococcus lactis

This is the author's manuscript							
Original Citation:							
Availability:							
This version is available http://hdl.handle.net/2318/130143 since							
Published version:							
DOI:10.1016/j.foodres.2012.12.017							
Terms of use:							
Open Access							
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.							

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [*Food Research International, 54, 1, November 2013,* http://dx.doi.org/10.1016/j.foodres.2012.12.017].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect*® *platform*]

An exoproteome approach to monitor safety of a cheese-isolated *Lactococcus lactis*

Federica Genovese^{a, c,*}, Jean Daniel Coïsson^b, Avishek Majumder^c, Alessandro Pessione^a, Birte Svensson^c, Susanne Jacobsen^c, Enrica Pessione^a

^a Dipartimento di Biologia Animale e dell'Uomo, Università di Torino, Via Accademia Albertina 13, 10123 Torino, Italy

^b DISCAFF-Novara, Università del Piemonte Orientale A. Avogardo, Via Giovanni Bovio 6, 28100 Novara, Italy

^C Enzyme and Protein Chemistry, Department of Systems Biology, Technical University of Denmark, Søltofts Plads Building 224, DK-2800 Kgs., Lyngby, Denmark

*Corresponding author at: Dipartimento di Biologia Animale e dell'Uomo, Via Accademia Albertina 13, 10123 Torino, Italy. Tel.: + 39 011 6704691; fax: + 39 011 6704508.

Abstract

The safety of the cheese-isolated and potential starter *Lactococcus lactis* 11D was explored by means of an extracellular proteomic study. A preliminary analysis showed good caseification/proteolytic behavior of the strain, absence of production of biogenic amines and good survival at acidic pH. The extracellular proteome map was analyzed to investigate the presence of potential virulence factors. Five moonlighting proteins with adhesive properties (ornithine carbamoyltransferase, fructose bisphosphate aldolase, trigger factor, EF-Tu and GroEL) were identified in the exoproteome, as well as the potential plasminogen binding proteins enolase, GAPDH and phosphoglycerate mutase. Adhesive properties are fundamental features for good starters and commensal strains, although some controversial aspects arise from plasminogen binding proteins as we shall discuss. Noticeably, GroEL, chitinase, and triose phosphate isomerase were abundant in the L. lactis 11D exoproteome. These proteins play a role in bacterial aggregation and in bacteria-fungi interactions, therefore their presence may indicate a good competition potential of the strain against other microorganisms in both food and the gastrointestinal habitat. A DIGE comparative exoproteomic analysis was performed on the L. lactis 11D strain grown on glucose and the disaccharide trehalose, examined here due to its common use as lyophilization stabilizer, respectively. The experiment showed that chitinase biosynthesis was enhanced in presence of trehalose. This is to our knowledge the first extracellular proteomic mapping of L. lactis with relevance for bacterial strain-typing in food safety.

Highlights

▶ Bacterial strain-typing is an important issue in food safety. ▶ The potential starter *L*. *lactis* 11D was analyzed by an exoproteome approach. ▶ The exoproteome showed high presence of moonlighting proteins involved in adhesion. ▶ A DIGE exoproteome comparing growth on glucose and on trehalose was performed. ▶ Chitinase biosynthesis was enhanced in presence of trehalose.

Abbreviations

- CCB, Colloidal Coomassie blue;
- CFU, colony forming units;
- DIGE, difference gel electrophoresis;
- EF-Tu,elongation factor thermo unstable;
- GRAS, Generally regarded as safe;
- IEF, Isoelectric focusing;
- LAB,Lactic acid bacteria;
- PDO, Protected Designation of Origin;
- Pg, Plasminogen;
- SDS-PAGE, Sodium Dodecyl Sulfate PolyAcrylamide Gel Electrophoresis;
- SGF, Simulated gastric fluid

Keywords

- Lactococcus lactis;
- Adhesion;
- Exoproteome;
- Plasminogen activation;
- Chitinase;
- DIGE

1. Introduction

Lactic acid bacteria (LAB) are widely exploited in food industry as spontaneous or starter cultures for several fermentation processes (<u>Hugas, Garriga, & Aymerich, 2003</u>). They are also used as bio-control agents for food safety improvement, due to their ability to counteract spoilage and pathogenic bacteria by means of nutritional competition, acidification, available enzymes and bacteriocin production (<u>De Vuyst & Leroy, 2007</u>).

The search for new starter microorganisms to be employed in food manufacturing for improving texture and for adding particular aromas is also of growing interest to satisfy consumers' demand of novel tastes. New starter microorganisms can be searched in the spontaneous microbiota of PDO (Protected Designation of Origin) products. Strain safety is an essential pre-requisite for their use in industrial food production, thus lack of excessive biogenic amine production and of secreted toxic molecules must be verified.

In the past few years production of biogenic amines by several LAB involved in food fermentation processes was highlighted (wine, beer, and cheese). Thus high contents of histamine and tyramine can cause allergies, hypertension and headache, and may have severe consequences such as brain hemorrhage and death (Millichap & Yee, 2003). The ability to biosynthesize amines is related to a specific strain rather than to a species, therefore even bacteria generally regarded as safe (GRAS) may present risks (Chander, Batish, Babu, & Bhatia, 1988).

Another important feature to be excluded is the secretion by the strain of known toxic molecules including proteins that may be involved in the activation of proteolytic cascades. This process is well known in pathogenic bacteria, nevertheless also non-virulent bacterial species can display this feature (<u>Candela et al., 2007</u> and <u>Hurmalainen et al., 2007</u>). Most of these proteins are intracellular metabolic enzymes or stress proteins that act as plasminogen (Pg) receptors

triggering localized proteolysis when secreted (<u>Carneiro et al., 2004</u>, <u>Kinnby et al., 2008</u>, <u>Pancholi</u> and <u>Fischetti</u>, <u>1992</u> and <u>Sibbald et al., 2006</u>) and thus facilitating pathogenic bacteria invasion (<u>Gohar et al., 2005</u> and <u>Hurmalainen et al., 2007</u>). It is therefore important to confirm the absence of potential proteolytic activators in strains employed in food technology in particular considering that some food bacteria display good resistance to acidic pH allowing them to reach the gut ecological niche. Exoproteome analysis is a valid method to exclude the presence of secreted virulence factors in bacteria of both biotechnological importance (<u>Hansmeier, Chao, Puhler, Tauch, & Kalinowski, 2006</u>) and food application interests (<u>Gagnaire et al., 2004</u>).

The aim of the present investigation was to identify possible starter candidates among the microbiota of Piedmont PDO artisanal-made cheeses, to be employed in cheese manufacturing. Towards this goal a preliminary screen focused on technological properties and biogenic amine production was performed. Subsequently the potential safety for human health was evaluated by exoproteome analysis of the chosen strain *Lactococcus lactis* 11D. Furthermore, a differential exoproteomic analysis of the strain grown on glucose and on trehalose, respectively, was performed by DIGE to exclude occurrence of trehalose induction of virulence factors. This is an important consideration since trehalose is a common starter cultures stabilizer used in the lyophilization process as cryoprotectant (<u>Tymczyszyn, Gomez-Zavaglia, & Disalvo, 2007</u>).

2. Materials and methods

2.1. Bacterial strains selection and their technological evaluation

Three *L. lactis* subsp. *lactis* strains were selected directly from an artisanal made cheese isolated and used to produce miniature cheeses as previously described (<u>Bertolino, Zeppa, Gerbi, & McSweeney, 2008</u>). Proteolysis parameters (total nitrogen and soluble nitrogen) were determined after 60 days of ripening using Official Methods of Analysis (<u>Helrich, 1990</u>). The ripening index was calculated as percentage of water soluble nitrogen of the total nitrogen.

2.2. Biogenic amine determination

Biogenic amine extraction was performed on cheese (20 g) in 95 mL pure HPLC water (Milli-Q, Millipore). Each sample was homogenized (2 min), and thereafter added trichloroacetic acid (5 mL; TCA, Fluka, 100% w/v). After centrifugation at $2500 \times g$ (15 min), the supernatants (50 mL) were extracted three times with ether (each 15 mL). The volume of the aqueous solution was adjusted to 50 mL with HPLC water and filtered (0.22 µm syringe filter; Millipore, Bedford, MA, USA) before HPLC analyses (Arlorio, Coïsson, & Martelli, 1998). Four biogenic amines (tyramine, histamine, 2-phenylethylamine and tryptamine) and their precursor amino acids (tyrosine, histidine, phenylalanine and tryptophan) were quantified using an optimized ion-pair HPLC method (Coïsson, Cerutti, Travaglia, & Arlorio, 2004) and 10 µL of sample injected.

2.3. Strain maintenance

The selected strain was maintained in 50% glycerol aliquots of frozen culture, and transferred to fresh M17 broth before performing proteomic analysis.

2.4. Acid tolerance of L. lactis 11D

Simulated gastric fluid (SGF) was formulated according to the guidelines of the U.S. Pharmacopeia. Briefly, SGF was composed of 3.2 g/L pepsin and 2.0 g/L NaCl, and the pH was adjusted to 1.5, 2.0, 2.5, and 3.0 by the addition of 5 M HCl. Middle exponential phase cultures

(1 mL) of *L. lactis* subsp. *lactis* 11D in M17 broth were added to SGF (19 mL) at 37 °C under mild agitation (200 rpm). After 30 min 1 mL was collected, mixed in sterile M17 broth and plated onto M17 agar after serial 10-fold dilutions. Plates were incubated at 37 °C for 24 h. The average number of CFU from triplicate analyses of every dilution was determined.

2.5. Proteomic experiments

2.5.1. Bacterial growth

For the extracellular and cytosolic maps, *L. lactis* 11D was grown on M17 broth. For the comparative exoproteome analysis the strain was grown on M17 broth deprived of lactose supplemented with 1% (w/v) glucose and 1% (w/v) trehalose for the two different conditions. *L. lactis* 11D was pre-cultured in M17 (10 mL) at 37 °C overnight (to late exponential phase) and inoculated into 45 mL of M17, glucose-M17 or trehalose-M17, respectively. These cultures were incubated at 37 °C until middle exponential phase. Four biological and four technical replicates were prepared at each condition.

2.5.2. Sample preparation for extracellular maps

The middle-exponential phase cultures were centrifuged (9000 ×*g*; 20 min) and the cell free supernatants were filtered (0.2 µm syringe filter, Millipore Millex filter unit); after a cooling step at -80 °C (3 h) cultures were freeze-dried overnight (Scanvac CoolSafeTM instrument, Labo-Gene). The freeze-dried supernatants were dialyzed (MWCO 3500 Da molecular porous membrane, SpectraPor) against 3×5 L 5 mM Tris HCl pH 7.5 (overnight at 4 °C) and precipitated by three volumes 10:90 (v/v) TCA:acetone. After incubation at -80 °C overnight, the samples were centrifuged (25,000 ×*g* for 30 min, at 4 °C) and protein pellets were resuspended in rehydration buffer (7 M urea, 2 M thiourea, 3% CHAPS, 10 mM Tris HCl pH 8.5). Protein concentration was assessed by 2D-Quant kit (GE Healthcare).

2.5.3. Sample preparation for the cytosolic proteome

The cell pellet obtained after centrifugation of the middle-exponential phase culture of *L. lactis* 11D in M17 was washed twice with 0.95% NaCl and transferred to a Fast-Prep tube. One part of glass beads (< 106 µm; Sigma) was added to one part of cells. The cells were added 300 µL saturated phenol (phenol with 10 mM Tris HCl pH 8 kept at 4 °C for 6 h), 300 µL 10 mM Tris HCl pH 7.5 and placed on ice. The cell disruption was carried out (Bio 101 Savant FastPrep FP 120 bead beater, Savant, Farmingdale, USA) for four cycles (45 s, 6 m s⁻¹) with 60 s intermittent cooling on ice between cycles. After centrifugation (15,000 ×*g*; 15 min, at 4 °C) three phases were formed. Three volumes of 96% ethanol (v/v) were added to the isolated middle phase and stored overnight at – 20 °C. After centrifugation as above, the protein pellet was resuspended in rehydration buffer and added 1 µL benzonase nuclease (25 U/µL, Novagen) to avoid DNA streaking. Protein concentrations were assessed by 2D-Quant kit (GE Healthcare).

2.5.4. Isoelectric focusing (IEF)

An equal amount of proteins resuspended in rehydration buffer were added 100 mM DTT, 10% isopropanol, 5% glycerol and 1% IPG buffer pH 4–7 (GE Healthcare) and loaded by anodic cuploading on 11 cm Immobiline DryStrip (GE Healthcare) pH 4–7, which had been previously rehydrated overnight with 200 μ L of rehydration buffer. IEF was carried out at 20 °C,

 $50 \,\mu$ A/strip, following the steps: $30 \,\text{V}$ 70 min, $100 \,\text{V}$ (gradient) 60 min, $100 \,\text{V}$ 8 h, $2000 \,\text{V}$ (gradient) 6 h, $8000 \,\text{V}$ (gradient) 2 h, $8000 \,\text{V}$ until $33,000 \,\text{V}$ h.

2.5.5. IEF for DIGE (Difference Gel Electrophoresis)

Prior to IEF, proteins were labeled with Cy2, Cy3 and Cy5 CyDye DIGE Fluors for Ettan DIGE (GE Healthcare) according to the manufacturer's instructions. Each dye (100 pmol) was added to sample or internal standard (each containing 15 μ g protein) and incubated for 30 min on ice. The reaction was stopped by addition of 1.5 μ L L-lysine (100 mg/mL) on ice for 10 min. Samples labeled with the different dyes were mixed together and added to the rehydration buffer. IPG strips, previously rehydrated overnight with 200 μ L of rehydration buffer, were loaded by anodic cup-loading with the mixture of labeled proteins (45 μ g). The run was performed using an IPG-phor (GE Healthcare) covered from direct light, following the program described in Section 2.5.4.

2.5.6. SDS-PAGE

IPG strips were soaked (15 min) in equilibration buffer (5 mL 6 M urea, 30% glycerol, 50 mM Tris HCl pH 8.8, 0.01% bromophenol blue and 2% SDS) with DTT (10 mg/mL) followed by 15 min in the same buffer (5 mL) containing iodoacetamide (25 mg/mL). The second dimension SDS-PAGE was performed on PROTEAN Plus Precast Gels for 2-D (BioRad) at constant voltage of 200 V until the bromophenol blue reached the bottom of the gel. Molecular weight marker MARK 12TM unstained standard (Invitrogen) was used.

For the comparative exoproteome analysis by DIGE, SDS-PAGE was performed in 24 cm gels using ETTANTM DALT SIX Electrophoresis unit (GE Healthcare) with the following program: 2 W per gel, 600 V, 400 mA for 30 min, then 12 W per gel until the bromophenol blue reached the bottom of the gel. The electrophoresis unit was covered from direct light during the run. All gels were fixed and stained according to the Blue silver method for Colloidal Coomassie Blue (CCB) staining (Candiano et al., 2004) and destained in distilled water overnight.

2.5.7. Image acquisition

Gels were scanned with the ScanMaker 9800 XL scanner (Microtek), both in reflective and in transparent mode. DIGE gels were scanned with the Typhoon 9410 Variable Mode Imager (GE Healthcare). Imaging of DIGE gels was done immediately after the SDS-PAGE at excitation/emission wavelengths of 488/520 nm (Cy2), 532/580 nm (Cy3) and 633/670 nm (Cy5), respectively.

2.5.8. Image analysis

Progenesis SameSpots (version 3.3, nonlinear Dynamics Ltd, Newcastle upon Tyne, UK) was used for spot detection in all experiments. DIGE gel images were also aligned by automated calculation of ten manually assigned alignment landmark vectors. Scanned gels were analyzed by intra-gel (difference in-gel) and inter-gel (biological variance) analysis. A 1.5-fold threshold (spot volume ratio change) and ANOVA $p \le 0.05$ was chosen as criterion in the identification of differentially expressed protein candidates.

2.5.9. Mass spectrometry analysis

All visible spots in the exoproteome map and the selected spots in the DIGE maps were manually excised from the freshly CCB-stained gel with a scalpel and transferred to EppendorfTM tubes. Trypsin digestion was carried out as previously described (Hellman, Wernstedt, Gonez, & Heldin, 1995) and the tryptic digests (2 μ L) were loaded to the AnchorChip target plate (Brucker-Daltonics) with α -cyano-4-hydroxycinnamic acid (CHCA) matrix (Beavis, Chaudhary, & Chait, 1992).

MS and MS/MS spectra were obtained by Ultraflex II MALDI-TOF MS mass spectrometer (Bruker-Daltonics) in auto-mode using Flex Control v3.0 (Bruker-Daltonics) and processed by Flex Analysis v3.0 (Bruker-Daltonics). Peptide mass maps were acquired in reflectron mode with 500 laser shots per spectrum. Spectra were externally calibrated using a tryptic digest of β lactoglobulin (5 pmol/µL); MS/MS data were acquired with stop conditions so that 1000–1600 laser shots were accumulated for each spectrum. The combined peptide mass fingerprint and MS/MS search was performed on **NCBInr** database for bacteria (NCBInr20100323; 10606545 sequences; 3615943919 residues) using the MASCOT 2.0 software (http://www.matrixscience.com) integrated together with BioTools v3.1 (Bruker-Daltonics). The parameters used for the search were: monoisotopic peptide mass accuracy of 80 ppm, fragment mass accuracy to ± 0.7 Da; maximum of one missed cleavage; carbamidomethylation of cysteine as fixed modification and partial oxidation of methionine as variable modification. There were no restrictions with respect to protein Mw and pI. Filtering of peaks was done for known autocatalytic trypsin peaks and keratin peaks; the signal to noise threshold ratio was set to 1:6. The significance threshold was set at p < 0.05, and identification required that each protein contained at least one peptide or an e-value < 0.05.

3. Results and discussion

3.1. Proteolytic potential, caseification properties and biogenic amines production

To select possible candidates to be employed as new starters in cheese manufuring, the microbiota of an Italian artisanal-made cheese was screened and three selected *L. lactis* strains were used to produce miniature-cheeses. To evaluate the technological value of these strains and their impact on human health, the proteolytic potential, the free amino acid and the most common biogenic amine concentrations were measured.

All three strains displayed good technological features as starter for cheese production in terms of pH and ripening index (<u>Table 1</u>). As regards biogenic amines, one strain (*L. lactis* 11D) did not produce any, another strain (*L. lactis* 150) gave $< 20 \text{ mg kg}^{-1}$ for both tyramine and histamine, and the third strain (*L. lactis* 370) showed a tyramine level reaching 73 mg kg⁻¹ (<u>Table 1</u>). The findings demonstrate that *L. lactis* 11D is biogenic amine-free and this strain was therefore chosen for further analyses regarding its safety features.

Table 1.

Technological features (proteins, pH, and ripening index), free amino acids and biogenic amines concentration in cheese samples inoculated with *Lactococcus lactis* 11D, 150 and 370.

L. lactis 11D L. lactis 150 L. lactis 370

	L. lactis 11D	L. lactis 150	L. lactis 370
Proteins (%)	18.35 ± 0.24	21.24 ± 1.19	20.79 ± 0.08
рН	5.53 ± 0.01	4.93 ± 0.02	5.47 ± 0.03
Ripening index	48.53 ± 0.01	29.74 ± 0.01	68.99 ± 0.03
Tyrosine	40.15 ± 0.15	42.27 ± 0.15	14.13 ± 0.31
Histidine	13.12 ± 1.07	15.23 ± 0.31	5.15 ± 0.31
Phenylalanine	170.85 ± 0.16	116.20 ± 0.75	161.92 ± 0.59
Tryptophan	4.53 ± 0.04	18.60 ± 0.09	1.51 ± 0.08
Tyramine	< LOD	10.90 ± 0.65	73.35 ± 0.94
Histamine	< LOD	18.36 ± 0.20	10.66 ± 1.89
2-Phenylethylamine	< LOD	< LOD	< LOD
Tryptamine	< LOD	< LOD	< LOD

Ripening index = $N_{sol} / N_{tot} * 100$, biogenic amine and amino acid concentrations are measured in mg kg⁻¹, LOD = limit of detection.

3.2. Acid tolerance

As shown in Table 2, L. lactis 11D is able to survive under acidic conditions for 30 min. No significant differences between the initial population at 0 min (cell count of $6.77 \times 10^9 \pm 4.9 \times 10^8$) and the population after 30 min at pH 2.5 (cell count of $6.47 \times 10^9 \pm 1.05 \times 10^9$) and pH 3 (cell count of $6.08 \times 10^9 \pm 4.4 \times 10^8$) were observed; by contrast *L. lactis* 11D is not able to grow after 30 min at $pH \le 2.0$. It is thus demonstrated that the strain may be able to overcome the gastric acidic barrier and reach the gut. This promising property motivated the analysis of proteins released to the extracellular environment that may be important both for understanding adhesion/invasion mechanisms and for assessing the strain safety for human consumption.

Table 2.

Identified extracellular proteins of Lactococcus lactis 11D.

Protein	Spot #	Gene ID	Score	E-value	Mw_	pI <u> </u>	Accession no.
Cell wall processing chitinase and su	rface ant	ioen					

ell wall processing, chitinase and surface antigen

Protein	Spot #	Gene ID	Score	E-value	Mw_	pI <u></u>	Accession no.
N-acetylmuramidase	5	acmD	242	3.10E- 18	37,517	4.5	<u>15672523</u>
N-acetylmuramoyl-L-alanine amidase	11	acmB	165	1.60E- 10	49,556	4.82	<u>15673889</u>
Chitinase	58	chiA	160	4.90E- 10	54,487	5.32	<u>15673932</u>
Surface antigen	70–80		186	1.20E- 12	46,212	7.68	<u>116513168</u>
Adhesion proteins							
Fructose-bisphosphate aldolase	9	fbaA	146	1.20E- 08	32,197	5.04	<u>15673891</u>
Ornithine carbamoyltransferase	30	arcB	148	7.80E- 09	39,488	5.44	<u>15674018</u>
Trigger factor	87A	tig	107	9.80E- 05	46,901	4.43	<u>15672533</u>
Triosephosphate isomerase	63	tpiA	263	2.50E- 20	27,060	4.63	<u>15673116</u>
Chaperonin GroEL	2	groEL	363	2.50E- 30	57,166	4.75	<u>15672376</u>
Elongation factor Tu	64–65	tuf	410	4.90E- 35	43,185	4.89	<u>15673843</u>
Plasminogen-binding proteins							
Glyceraldehyde 3-phosphate dehydrogenase	85	gapB	268	7.80E- 21	35,911	5.57	<u>15674228</u>
Phosphoglycerate mutase	69	gpmA	173	2.50E- 11	26,297	5.3	<u>116511195</u>
α-Enolase	55	eno	557	9.80E- 50	46,940	4.68	<u>15672626</u>
Molecular chaperone DnaK	23	dnaK	420	4.90E- 36	64,947	4.62	<u>15672936</u>

Protein	Spot #	Gene ID	Score	E-value	Mw_	pI <u> </u>	Accession no.
Putative moonlighting proteins and carbohydrate metabolism							
Phosphoglucose isomerase	28	pgi	147	9.80E- 09	49,564	4.89	15674150
Elongation factor Ts	40	tsf	118	7.80E- 06	36,647	4.92	<u>15674134</u>
Elongation factor G	66	fusA	157	9.80E- 10	77,907	4.75	<u>15674242</u>
Thioredoxin reductase	8	trxB1	82	0.031	33,987	4.76	<u>15672922</u>
30S Ribosomal protein S1	17	rpsA	350	4.90E- 29	44,683	4.99	<u>15672820</u>
Phosphofructokinase	32	<i>pfk</i>	124	2.00E- 06	35,638	5.56	<u>399697</u>
Phosphotransferase system mannose/fructose-specific component IIA	1	ptsIIB	129	6.20E- 07	35,085	5.23	<u>116512556</u>
Phosphoenolpyruvate-protein phosphotransferase	15	ptsI	97	0.00092	62,635	4.67	<u>15672100</u>
Tagatose 1.6-diphosphate aldolase	83	tbpa	423	2.50E- 36	36,641	4.82	<u>116326612</u>
UDP-Glucose 4-epimerase	60	galE	102	0.00031	36,321	5.18	<u>15673961</u>
Galactose-6-phosphate isomerase subunit LacB	36–45	lacB	93	0.0027	19,085	5.13	<u>116326614</u>
L-Lactate dehydrogenase	24	ldh	121	3.90E- 06	35,086	4.97	<u>15673313</u>
Proteins with other function							
50S Ribosomal protein L10	54	rplJ	101	0.00039	18,076	5.08	<u>15673251</u>
DNA polymerase III subunit beta	46	dnaN	81	0.039	42,238	5.17	<u>15671984</u>
Ubiquitin	16	ub	214	2.00E- 15	3592	4.99	<u>207094432</u>

Protein	Spot #	Gene ID	Score	E-value	Mw_	pI <u> </u>	Accession no.
Dipeptidase PepV	34	pepV	97	0.0011	52,025	4.75	15672824
Cell division protein FtsZ	87E	ftsZ	115	1.60E- 05	43,961	4.54	<u>15673851</u>
Hypothetical protein L22960	10	ymcF	278	7.80E- 22	38,305	4.95	<u>15673185</u>
Hypothetical protein L107870	57	ytcC	107	2.10E- 05	39,963	6.92	<u>15673838</u>
Hypothetical protein L0370	43	typA	103	0.00025	68,258	4.91	<u>15673997</u>
Hypothetical protein L96658	21– 29–42	usp45	173	2.50E- 11	47,036	8.33	<u>15674211</u>
Hypothetical protein L22496	19	yacG	412	3.10E- 35	50,338	8.47	<u>15672000</u>

The values indicated for Mw and pI are the theoretical ones.

3.3. Exoproteome analysis

Proteins related to adhesive/invasive properties are generally distributed in the pH 4–7 range (<u>www.jvirgel.de</u>). The present study therefore focused on the acidic exoproteome analyzed for cells collected during the mid-logarithmic phase, to reduce the number of dead and lysed cells that increase in the late phases of the bacterial growth, when lack of nutrients occurs. Even though a non-specific release of cytoplasmatic components by cell autolysis cannot be excluded, a 2-DE map of the intracellular proteins was performed under the same conditions and displayed a different protein pattern (Fig. 1).



Fig. 1.

Comparison between the cytosolic (left) and the extracellular (right) map of *Lactococcus lactis* 11D. The maps exhibit different patterns, demonstrating no predominant cytosolic contamination in the extracellular sample.

Progenesis SameSpots analysis revealed 91 protein spots in the exoproteome (Fig. 2) representing 37 identified unique proteins (Appendix A Table 1), while the remaining spots represented multiple forms of the same proteins with a different pI or were too faint to be adequately identified by mass spectrometry analysis. It is worth noticing that no known virulence protein was identified in the acidic exoproteome of L. lactis 11D. The identified proteins were divided into main functional (Table 2) according the KEGG five groups to pathways (http://www.genome.jp/kegg/): (i) cell wall processing, chitinase and surface antigen; (ii) adhesion proteins; (iii) plasminogen binding proteins; (iv) putative moonlighting and carbohydrate metabolism; (v) proteins with other functions.



Fig. 2.

Exoproteome map of *Lactococcus lactis* 11D shows 91 spots by analysis with Progenesis SameSpots. Unique proteins (37) were identified with statistical significance, these included several moonlighting proteins

Proteins of the first group are chitinase, N-acetylmuramidase, and N-acetylmuramoyl-L-alanine amidase. Chitinase (spot 58) is of a special interest as it hydrolyzes chitin, the major cell-wall component of fungi and a prominent component of yeast cell walls. Since often bacteria and fungi share the same ecological niche, this enzyme may play a role in competition for nutrients. Carranza et al. (2009) reported evidence of an extracellular chitinase in *Chronobacter turicensis* displaying a fungi-inhibiting action. N-acetylmuramidase (spot 5) and N-acetylmuramoyl-L-alanine amidase (spot 11) are proteins involved in peptidoglycan renewal (maturation and degradation), but also in bacterial interspecies interactions since they act as general cell-wall lysis factors (Salazar & Asenjo, 2007). The presence of these proteins is consistent with the harvesting in mid-exponential phase, when the highest bacterial cells duplication occurs. The most abundant protein (spots 70–80) was identified as a generic surface antigen. Considering its high abundance it is of future interest to elucidate the function of this protein.

Among the proteins identified in the supernatant from *L. lactis* 11D more than 20% are anchorless multifunctional proteins (<u>Jeffery, 2009</u>) involved in adhesion and plasminogen binding mechanisms. These proteins are usually expressed in the intracellular compartment, where they exert metabolic roles, but when released to the extracellular environment (<u>Bergonzelli et al., 2006</u> and <u>Granato et al., 2004</u>) or expressed on the cell surface (<u>Alam et al., 2009</u>, <u>Hussain et al., 1999</u> and <u>Tunio et al., 2010</u>) they display additional functions. Such proteins are generally referred to as moonlighting (<u>Jeffery, 2009</u>).

The finding in the extracellular proteome of several adhesion proteins that should exert their function when surface-bound is a common phenomenon reflecting that a dynamic equilibrium between these two cell districts establishes, especially during the logarithmic growth phase.

Several factors such as pH variations and cell wall renewal may favor the cell wall detachment of several proteins (<u>Antikainen, Kuparinen, Lahteenmaki, & Korhonen, 2007</u>).

Fructose bisphosphate aldolase (FBA) (spot 9) and ornithine carbamovltransferase (OTC) (spot 30), exert their adhesion activity when surface-bound. FBA is involved in the binding to a large trans-membrane receptor of the cadherin superfamily in the pathogenic Gram-negative Neisseria meningitidis and Gram-positive Streptococcus pneumoniae (Tunio et al., 2010). OTC, a nitrogen metabolism enzyme, has been found on the surface of several streptococci, Clostridium perfringens (Alam et al., 2009), and the opportunistic Staphylococcus epidermidis (Hussain et al., 1999), where it acts as adhesin. Trigger factor (spot 87A) and DnaK (spot 23) are also connected with adhesion, since it has been reported that surface adhesin P1 from Streptococcus mutans cannot efficiently work in the absence of DnaK and trigger factor (Crowley et al., 2008). Furthermore, it has been demonstrated in Lactobacillus reuteri NCIB 11951 that a collagen I binding protein, not belonging to the S-layer, shares high sequence homology with Escherichia colitrigger factor (Aleljung et al., 1994). Triose phosphate isomerase (TPI) (spot 63) was also proven as an adhesion factor for fungal mannans in *Staphylococcus aureus* when surface exposed (Furuya & Ikeda, 2009). Besides, both GroEL (spot 2) and EF-Tu (spots 64 and 65) in Lactobacillus johnsonii display adhesive capabilities towards human tissues (Bergonzelli et al., 2006). In L. johnsonii, GroEL induces a strong aggregation of the pathogen Helicobacter *pylori*, thus decreasing the bacterial load and facilitating clearance of the aggregated pathogens by the mucus (Bergonzelli et al., 2006). Its presence has been detected in the spent culture medium throughout the whole logarithmic phase, which is in agreement with the results of the present investigation, where high abundance of both GroEL and EF-Tu is evident in the extracellular environment. GroEL and EF-Tu have also been detected in the extracellular proteome of *Bifidobacterium animalis* subsp. *lactis* as putative moonlighting proteins with adhesion roles that contribute to the probiotic features of the strain (Gilad, Svensson, Viborg, Stuer-Lauridsen, & Jacobsen, 2011).

Three glycolytic enzymes, enolase (spot 55), GAPDH (spot 85) and phosphoglycerate mutase (PGM) (spot 69) as well as the stress protein DnaK (spot 23) behave as plasminogen (Pg) receptors in pathogenic bacteria, like S. aureus and streptococci (Kinnby et al., 2008). Once bound on the bacterial cell-wall, Pg is converted by specific virulence factors produced by both staphylococci (staphylokinases) and streptococci (streptokinases) into plasmin (Lahteenmaki, Kuusela, & Korhonen, 2001) able to activate proteolytic cascades, resulting in the degradation of the extracellular matrix and subsequent migration of bacteria through host tissues (Gladysheva, Turner, Sazonova, Liu, & Reed, 2003).a-enolase, in particular, is able to directly bind to extracellular matrix proteins like laminin (Carneiro et al., 2004), collagen and fibronectin (Candela et al., 2007), thus favoring damage via the proteolytic cascade. Nevertheless GAPDH and α -enolase have both been detected on the surface of certain LAB (Lactobacillus acidophilus, L. amylovorus, L. gallinarum, L. gasseri, L. crispatus and L. *johnsonii*) (Hurmalainen et al., 2007) and bifidobacteria (Bifidobacterium lactis, B. longum, B. bifidum and B. animalis) (Candela et al., 2007 and Gilad et al., 2011). Both of these enzymes allow Pg-binding on the cell-wall, but no homologue genes coding for any of the known Pgactivators (strepto- and staphylokinases) have been found in the genome of the sequenced lactobacilli species. This may indicate that LAB lacks an endogenous potential for Pg activation by conversion to plasmin. Some authors, however, suggest that Pg attachment to the cell-wall receptor can cause a conformational change of the latter, enhancing Pg susceptibility to host Pgactivating factors, namely uPA (urokinase-type Pg activator) and tPA (tissue Pg activator) (Candela et al., 2007 and Crowley et al., 2008). These controversial reports concerning bacteria-Pg interactions and the possibility to activate proteolytic cascades, suggest the need of an accurate typing of both surface and extracellular proteome of starter strains before use, especially for those able to bypass the gastric barrier.

Many ribosomal proteins have been predicted to be extracellular by in silico analysis (CELLO: sub cellular localization predictive system (Yu, Chen, Lu, & Hwang, 2006)) and several identified proteins, such as EF-G (spot 66), EF-Ts (spot 40), thioredoxin reductase (spot 8) and 30S ribosomal protein S1 (spot 17) were previously classified as signal peptide-lacking exoproteins in *S. aureus* (Sibbald et al., 2006) and *Bacillus anthracis* (Gohar et al., 2005) and as surface protein in *L. lactis* NZ9000 (Berlec, Zadravec, Jevnikar, & Strukelj, 2011). All these considerations suggest that some of the detected proteins might play an extracellular role that remains to be identified.

3.4. Comparative exoproteome map by DIGE (Difference Gel Electrophoresis)

In order to ensure the viability and to control safety of the selected strain during and after the freeze-dry process, a culture was set up on the cryoprotectant sugar trehalose. Growth curves kinetic analysis of *L. lactis* 11D highlighted a lower growth rate ($\mu_{trehalose-M17} = 0.58 \text{ h}^{-1}$, $\mu_{glucose-M17} = 0.78 \text{ h}^{-1}$) but a higher final biomass concentration (0.97 g/L in trehalose-M17, 0.69 g/L in glucose-M17) for bacteria grown in presence of trehalose, as compared to the control (Fig. 3). This suggests that trehalose is a suitable carbon source for *L. lactis* 11D, which does not damage physiological functions. Furthermore, the lower growth rate is a good index that the strain cannot consume all the cryoprotectant/antioxidant present.



Fig. 3.

Lactococcus lactis 11D growth curves in glucose-M17 and trehalose-M17: trehalose stimulated growth of the strain (white dots), although the growth rate was lower as compared with the control (glucose, black dots).

Figure options

A comparative proteomic analysis was performed by DIGE to detect different extracellular protein expression between the growth of *L. lactis* 11D on trehalose and the control (glucose). The comparative image analysis of the DIGE gels revealed 72 differentially expressed spots: 35

spots were down-regulated and 37 spots were up-regulated in trehalose-M17 (Fig. 4). A significant number of spots were faint and very difficult to identify by mass spectrometry, successful hits were obtained for 8 spots. Thus five down-regulated and three up-regulated proteins were identified with statistical significance (Appendix A Tables 2–3). In spite of these limits, DIGE proved to be a sensitive method for spot detection, as expected (Marouga, David, & Hawkins, 2005).



Fig. 4.

Down-regulated (left) and up-regulated (right) spots in trehalose-M17. Numbers indicate spots with a different expression ratio among the two tested conditions higher than 1.5 fold and with an ANOVA p < 0.05.

It is worth noticing that even in presence of trehalose no known virulence factor was up-regulated in the acidic exoproteome indicating that as regards the analyzed proteins, the use of this cryoprotectant seems not to enhance risks for human health.

Chitinase proved to be up-regulated in *L. lactis* 11D grown on trehalose suggesting that this carbon source, commonly used in freeze-dry processes, can enhance the antagonistic potential of the bacterial strain against fungi. Moreover, the growth in trehalose seems to attenuate adhesin expression, as demonstrated by GroEL and FBA down-regulation.

4. Conclusions

The selection of bacteria to be directly employed as starters in the artisanal cheese matrix opens new possibilities for improving organoleptic properties, texture and added value of food, provided that a reliable safety control is carried out. Several analyses to verify the safety of a strain for human health must be performed especially in the light of several pathogenic pathways being strain-specific and not species-specific. An example of this is the controversial interaction between bacteria and plasminogen, which can lead to proteolytic cascades activation. This suggests the need of an accurate typing of extracellular proteomes of starter strains, especially for those able to bypass the gastric barrier that can reach and interact with the gut epithelium. In the present study the acidic exoproteome of *L. lactis* 11D was analyzed and shows potential to become a relevant tool for bacterial strain-typing to ensure bacterial safety.

Acknowledgments

This work was performed with the financial support of FEMS Research Fellowship 2010 and C.I.B. (Consorzio Interuniversitario per le Biotecnologie) to Federica Genovese and of the Danish Strategic Research Council's Program Committee on Health, Food and Welfare (FøSu) to Avishek Majumder and the EPC group. Part of the funding was supplied by the project CIPE 2006 (Regione Piemonte). Professor Giuseppe Zeppa (DIVAPRA, Università degli studi di Torino) is thanked for kindly providing the miniature cheeses. Anne Blicher, Bjarne Gregers Schmidt, and Birgit Andersen are thanked for technical assistance. The Typhoon scanner was granted by the Danish Research Council for Natural Science and the Danish Center for Advanced Food Studies contributed to the Bruker Ultraflex II mass spectrometer.

References

- S.I. Alam, S. Bansod, R.B. Kumar, N. Sengupta, L. Singh Differential proteomic analysis of *Clostridium perfringens* ATCC13124: Identification of dominant, surface and structure associated proteins BMC Microbiology, 9 (2009), p. 162
- P. Aleljung, W. Shen, B. Rozalska, U. Hellman, A. Ljungh, T. Wadstrom Purification of collagen-binding proteins of *Lactobacillus reuteri* NCIB 11951 Current Microbiology, 28 (1994), pp. 231–236
- J. Antikainen, V. Kuparinen, K. Lahteenmaki, T.K. Korhonen pH-dependent association of enolase and glyceraldehyde-3-phosphate dehydrogenase of *Lactobacillus crispatus* with the cell wall and lipoteichoic acids Journal of Bacteriology, 189 (2007), pp. 4539–4543
- M. Arlorio, J.D. Coïsson, A. Martelli Ion-pair HPLC determination of biogenic amines and precursor aminoacids. Application of a method based on simultaneous use of heptanesulphonate and octylamine to some foods

Chromatographia (1998), pp. 763–769

5. R.C. Beavis, T. Chaudhary, B.T. Chait α-Cyano-4-hydroxycinnamic acid as a matrix for matrix assisted laser desorption mass spectrometry

Organic Mass Spectrometry, 27 (1992), pp. 156–158

6. G.E. Bergonzelli, D. Granato, R.D. Pridmore, L.F. Marvin-Guy, D. Donnicola, I.E. Corthesy-Theulaz

GroEL of *Lactobacillus johnsonii* La1 (NCC 533) is cell surface associated: Potential role in interactions with the host and the gastric pathogen *Helicobacter pylori* Infection and Immunity, 74 (2006), pp. 425–434

7. A. Berlec, P. Zadravec, Z. Jevnikar, B. Strukelj

Identification of candidate carrier proteins for surface display on *Lactococcus lactis* by **theoretical and experimental analyses of the surface proteome** Applied and Environmental Microbiology, 77 (2011), pp. 1292–1300

8. M. Bertolino, G. Zeppa, V. Gerbi, P. McSweeney Study of proteolysis in miniature Toma Piemontese cheese made using wild bacteria

Italian Journal of Food Science, 20 (2008), pp. 57-73

- M. Candela, S. Bergmann, M. Vici, B. Vitali, S. Turroni, B.J. Eikmanns *et al.* Binding of human plasminogen to *Bifidobacterium* Journal of Bacteriology, 189 (2007), pp. 5929–5936
- 10. G. Candiano, M. Bruschi, L. Musante, L. Santucci, G.M. Ghiggeri, B. Carnemolla *et al.* Blue silver: A very sensitive colloidal Coomassie G-250 staining for proteome analysis

Electrophoresis, 25 (2004), pp. 1327–1333

- 11. C.R. Carneiro, E. Postol, R. Nomizo, L.F. Reis, R.R. Brentani Identification of enolase as a laminin-binding protein on the surface of *Staphylococcus aureus* Microbes and Infection, 6 (2004), pp. 604–608
- P. Carranza, I. Hartmann, A. Lehner, R. Stephan, P. Gehrig, J. Grossmann *et al.* Proteomic profiling of *Cronobacter turicensis* 3032, a food-borne opportunistic pathogen Proteomics, 9 (2009), pp. 3564–3579
- 13. H. Chander, V.K. Batish, S. Babu, K.L. Bhatia Amine production by Streptococcus lactis under different growth conditions Acta Microbiologica Polonica, 37 (1988), pp. 61–64
- 14. J. Coïsson, C. Cerutti, F. Travaglia, M. Arlorio Production of biogenic amines in "Salamini italiani alla cacciatora PDO"

Meat Science, 67 (2004), pp. 343-349

15. P.J. Crowley, T.B. Seifert, R.van. Isoda, M. Tilburg, M.W. Oli, R.A. Robinette *et al.* **Requirements for surface expression and function of adhesin P1 from** *Streptococcus mutans*

Infection and Immunity, 76 (2008), pp. 2456–2468

16. L. De Vuyst, F. Leroy Bacteriocins from lactic acid bacteria: Production, purification, and food applications

Journal of Molecular Microbiology and Biotechnology, 13 (2007), pp. 194-199

17. H. Furuya, R. Ikeda

Interaction of triosephosphate isomerase from the cell surface of *Staphylococcus* aureus and alpha-(1 > 3)-mannooligosaccharides derived from glucuronoxylomannan of *Cryptococcus neoformans*

Microbiology, 155 (2009), pp. 2707-2713

18. V. Gagnaire, M. Piot, B. Camier, J.P. Vissers, G. Jan, J. Leonil Survey of bacterial proteins released in cheese: A proteomic approach

International Journal of Food Microbiology, 94 (2004), pp. 185-201

- 19. O. Gilad, B. Svensson, A.H. Viborg, B. Stuer-Lauridsen, S. Jacobsen The extracellular proteome of *Bifidobacterium animalis* subsp. *lactis* BB-12 reveals proteins with putative roles in probiotic effects Proteomics, 11 (2011), pp. 2503–2514
- 20. I.P. Gladysheva, R.B. Turner, I.Y. Sazonova, L. Liu, G.L. Reed Coevolutionary patterns in plasminogen activation

Proceedings of the National Academy of Sciences of the United States of America, 100 (2003), pp. 9168–9172

21. M. Gohar, N. Gilois, R. Graveline, C. Garreau, V. Sanchis, D. Lereclus

A comparative study of *Bacillus cereus*, *Bacillus thuringiensis* and *Bacillus anthracis*extracellular proteomes Proteomics, 5 (2005), pp. 3696–3711

- 22. D. Granato, G.E. Bergonzelli, R.D. Pridmore, L. Marvin, M. Rouvet, I.E. Corthesy-Theulaz Cell surface-associated elongation factor Tu mediates the attachment of *Lactobacillus johnsonii* NCC533 (La1) to human intestinal cells and mucins Infection and Immunity, 72 (2004), pp. 2160–2169
- 23. N. Hansmeier, T.C. Chao, A. Puhler, A. Tauch, J. Kalinowski The cytosolic, cell surface and extracellular proteomes of the biotechnologically important soil bacterium *Corynebacterium* efficiens YS-314 in comparison to those of*Corynebacterium glutamicum* ATCC 13032 Proteomics, 6 (2006), pp. 233–250
- 24. U. Hellman, C. Wernstedt, J. Gonez, C.H. Heldin Improvement of an "In-Gel" digestion procedure for the micropreparation of internal protein fragments for amino acid sequencing

Analytical Biochemistry, 224 (1995), pp. 451-455

- 25. Helrich, K. (1990). *Official methods of analysis of the Association of Official Analytical Chemists*. (AOAC Virginia ed.).
- 26. M. Hugas, M. Garriga, M.T. Aymerich Functionality of enterococci in meat products

International Journal of Food Microbiology, 88 (2003), pp. 223-23

27. V. Hurmalainen, S. Edelman, J. Antikainen, M. Baumann, K. Lahteenmaki, T.K. Korhonen
 Extracellular proteins of *Lactobacillus crispatus* enhance activation of human plasminogen

Microbiology, 153 (2007), pp. 1112–1122

28. M. Hussain, G. Peters, G.S. Chhatwal, M. Herrmann

A lithium chloride-extracted, broad-spectrum-adhesive 42-kilodalton protein of*Staphylococcus epidermidis* **is ornithine carbamoyltransferase** Infection and Immunity, 67 (1999), pp. 6688–6690

29. C.J. Jeffery

Moonlighting proteins — An update

Molecular BioSystems, 5 (2009), pp. 345–350

30. B. Kinnby, N.A. Booth, G. Svensater

Plasminogen binding by oral streptococci from dental plaque and inflammatory lesions

Microbiology, 154 (2008), pp. 924-931

31. K. Lahteenmaki, P. Kuusela, T.K. Korhonen Bacterial plasminogen activators and receptors

FEMS Microbiology Reviews, 25 (2001), pp. 531-552

32. R. Marouga, S. David, E. Hawkins

The development of the DIGE system: 2D fluorescence difference gel analysis technology

Analytical and Bioanalytical Chemistry, 382 (2005), pp. 669-678

33. J.G. Millichap, M.M. Yee

The diet factor in pediatric and adolescent migraine

Pediatric Neurology, 28 (2003), pp. 9-15

34. V. Pancholi, V.A. Fischetti

A major surface protein on group A streptococci is a glyceraldehyde-3-phosphatedehydrogenase with multiple binding activity

The Journal of Experimental Medicine, 176 (1992), pp. 415-426

35. O. Salazar, J.A. Asenjo Enzymatic lysis of microbial cells

Biotechnology Letters, 29 (2007), pp. 985–994

36. M.J. Sibbald, A.K. Ziebandt, S. Engelmann, M. Hecker, A. de Jong, H.J. Harmsen *et al.* Mapping the pathways to staphylococcal pathogenesis by comparative secretomics

Microbiology and Molecular Biology Reviews, 70 (2006), pp. 755-788

- 37. S.A. Tunio, N.J. Oldfield, A. Berry, D.A. a'Aldeen, K.G. Wooldridge, D.P. Turner The moonlighting protein fructose-1, 6-bisphosphate aldolase of *Neisseria meningitidis*: Surface localization and role in host cell adhesion Molecular Microbiology, 76 (2010), pp. 605–615
- 38. E.E. Tymczyszyn, A. Gomez-Zavaglia, E.A. Disalvo Effect of sugars and growth media on the dehydration of Lactobacillus delbrueckii ssp.bulgaricus Journal of Applied Microbiology, 102 (2007), pp. 845–851
- 39. C.S. Yu, Y.C. Chen, C.H. Lu, J.K. Hwang **Prediction of protein subcellular localization**

Proteins, 64 (2006), pp. 643-651