

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cav1.3 and Cav1.2 channels of adrenal chromaffin cells: Emerging views on cAMP/cGMP-mediated phosphorylation and role in pacemaking

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/132208> since

Published version:

DOI:10.1016/j.bbamem.2012.11.013

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *BIOCHIMICA ET BIOPHYSICA ACTA-BIOMEMBRANES*, 1828, 2013, 10.1016/j.bbamem.2012.11.013.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.bbamem.2012.11.013

The definitive version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S000527361200404X>

**Cav1.3 and Cav1.2 channels of adrenal chromaffin cells:
emerging views on cAMP/cGMP-mediated phosphorylation and role in pacemaking**

D.H.F. Vandael[†], S. Mahapatra[†], C. Calorio, A. Marcantoni, E. Carbone

*Department of Drug Science, Laboratory of Cellular & Molecular Neuroscience, NIS Center,
CNISM, University of Torino, 10125 Torino, Italy*

Correspondence to: Emilio Carbone

[†] These authors contributed equally to this work

Key words: Voltage-gated calcium channels, BK and SK channels, cAMP/PKA and cGMP/PKG modulation, exocytosis, catecholamines release, endocytosis, adrenal medulla

Acknowledgments: The authors thank the Marie Curie Research Training Network “CavNET”, Italian M.I.U.R., Regione Piemonte, Università di Torino, Compagnia di San Paolo di Torino for funding.

Abstract

Voltage-gated Ca^{2+} channels (VGCCs) are voltage sensors that convert membrane depolarizations into Ca^{2+} signals. In the chromaffin cells of the adrenal medulla, the Ca^{2+} signals driven by VGCCs regulate catecholamine secretion, vesicle retrievals, action potential shape and firing frequency. Among the VGCC-types expressed in these cells (N-, L-, P/Q-, R- and T-type), the two L-type isoforms, Cav1.2 and Cav1.3, control key activities due to their particular activation-inactivation gating and high-density of expression in rodents and humans. The two isoforms are also effectively modulated by G protein-coupled receptor pathways delimited in membrane micro-domains and by the cAMP/PKA and NO/cGMP/PKG phosphorylation pathways which induce prominent Ca^{2+} current changes if opposingly regulated.

The two L-type isoforms shape the action potential and directly participate to vesicle exocytosis and endocytosis. The low-threshold of activation and slow rate of inactivation of Cav1.3 confer to this channel the unique property of carrying sufficient inward current at subthreshold potentials able to activate BK and SK channels which set the resting potential, the action potential shape, the cell firing mode and the degree of spike frequency adaptation during spontaneous firing or sustained depolarizations. These properties help chromaffin cells to optimally adapt when switching from normal to stress-mimicking conditions. Here, we will review past and recent findings on cAMP- and cGMP-mediated modulations of Cav1.2 and Cav1.3 and the role that these channels play in the control of chromaffin cell firing.

1. - The voltage-gated L-type channels: a brief overview

L-type calcium channels (LTCCs; Cav1) belong to the family of voltage-gated Ca^{2+} channels (VGCCs) that are permeable to Ca^{2+} and sensitive to membrane voltage. VGCCs include also the Cav2 (N, P/Q, R) and Cav3 types (T) [1] and are often classified as high-voltage (HVA; L, N, P/Q and R-type) and low-voltage activated (T-type) [2-4]. LTCCs are hetero-oligomers consisting of a pore-forming α_1 -subunit of 190–250 kDa in association with auxiliary β -, $\alpha_2\delta$ - and γ -subunits [5]. The α_1 -subunit is composed of four membrane-spanning domains (I–IV) linked together in a single polypeptide chain. Each domain contains six putative transmembrane segments (S1–S6) plus a “P” loop that dips partially into the pore to form the pore lining [5]. Like the other VGCCs, LTCCs open readily during membrane depolarization and allow Ca^{2+} to enter the cell. LTCCs, thus, regulate cell excitability and control a variety of Ca^{2+} -dependent physiological processes, like: excitation-contraction coupling in cardiac, skeletal and smooth muscles, gene expression, synaptic plasticity, brain aging, hormone secretion, vesicle retrieval and pacemaker activity in heart, neurons and neuroendocrine cells [6-13].

Presently, four genes are found to code for the Cav1.1, Cav1.2, Cav1.3 and Cav1.4 subunits [1]. Of these, Cav1.1 and Cav1.4 are mainly expressed in skeletal muscle, pituitary cells and the retina, whereas Cav1.2 and Cav1.3 are widely distributed throughout the central nervous system, sensory and endocrine cells, atrial myocytes and cardiac sino-atrial node cells [12, 14-18]. All four Cav1 isoforms are highly sensitive to 1,4-dihydropyridines blockers (Ca^{2+} antagonists) or activators (Ca^{2+} agonists), which are the most common used compounds for identifying functional LTCCs. The blocking potency of 1,4-DHPs differs depending on holding potential [7, 19-21] but their separation on this basis is complex and rarely used for quantitative assays. This justifies the continuous search for new selective blockers for Cav1.2 and Cav1.3 [16, 22], which are the most widely expressed Cav1 isoforms.

2. - Cav1.2 and Cav1.3 as multitasks channels controlling excitability, secretion and endocytosis in chromaffin cells

Cav1.2 and Cav1.3 channels are highly expressed in chromaffin cells of the adrenal medulla [23-27] where they control excitability, secretion of catecholamines and vesicle retrieval. Excitability is primarily controlled by Cav1.3 which activates at very low voltages and inactivates slowly in a voltage-dependent manner (measured in Ba^{2+}) with respect to Cav1.2 and other high-threshold channels (N, P/Q, R) [7, 28-30]. Depending on the C-terminal regulatory domain [30], Cav1.3 exhibits weak or pronounced Ca^{2+} -dependent inactivation (CDI). Assuming that at the pacemaker potential (-50 to -40 mV) the intracellular level of Ca^{2+} is rather low, it is clear that CDI contributes little to the total inactivation of the channel at these potentials. This enables Cav1.3 to carry most of the pacemaker current that sustains chromaffin cell spontaneous activity [10-13, 18] and to drive sufficient SK current activation that helps adapting the firing rate to a sustainable frequency during prolonged depolarizations (spike-frequency adaptation, SFA) [31]. In addition, Cav1.3 has been shown to be tightly coupled to fast inactivating BK channels, suggesting also a key role in the control of the AP shape [11, 12]. Concerning the second main activity, it is widely accepted that catecholamine secretion is particularly sensitive to L-type currents in rat (RCCs) and mouse chromaffin cells (MCCs) [10, 32-37]. In RCCs L-type channels carry nearly half of the total Ca^{2+} current and are responsible for the corresponding exocytosis [36, 38]. Concerning vesicle retrieval, LTCCs have a direct control of compensatory and excess endocytosis, which are strongly attenuated when LTCCs are blocked [39-41]. At present, however, it is not clear yet if both isoforms have distinct roles in this process.

In addition, Cav1.2 and Cav1.3 are shown to be effectively modulated by G-protein coupled receptors in membrane micro-domains [42-45], or up- and down-regulated by cAMP/PKA and NO/cGMP/PKG pathways, respectively [10, 46-49]. These two mechanisms act opposingly on channel gating (see [50-52] for reviews) and give rise to Ca^{2+} current changes of one order of magnitude under proper stimulation of one pathway and inhibition of the other [49]. All these peculiarities highlight the strategic role that Cav1.3 and Cav1.2 exert on chromaffin cell functioning.

3. - The autocrine modulation of LTCCs: a feedback for regulating catecholamine secretion

Chromaffin cells constantly release catecholamines, opioids and ATP during low and sustained sympathetic stimulation (see [51-53] for reviews). As a consequence, Ca^{2+} channels experience basal and stimulus-induced autocrine modulations, which affect their activation-inactivation gating and probability of channel opening, altering in this way the amount of Ca^{2+} entry into the cell. This autocrine effect is mediated by the adrenergic, opioidergic and purinergic autoreceptors highly expressed in chromaffin cells and coupled to G-proteins. This occurs in isolated chromaffin cells [42, 44, 48, 49] but it is more prominent in cell clusters [54] and in adrenal gland slices [45], where the released neurotransmitters can accumulate more effectively between closely packed chromaffin cells.

3.1 - Direct and remote modulation of LTCCs

For Cav2.1 (P/Q) and Cav2.2 (N) channels the autocrine modulation driven by purinergic, opioidergic and adrenergic receptors is mostly inhibitory and voltage-dependent [42, 44, 45, 55]. Cav2.1 and Cav2.2 channel activation is markedly delayed, but short pre-pulses to positive voltages can fully remove the $G_i\beta\gamma$ -mediated delay and channel activation returns normal during the immediately following depolarization [43, 56-62]. In contrast to this, LTCCs modulation is mainly voltage-independent and can occur through two distinct pathways(see [52] for a review): 1) by quickly inhibiting LTCCs gating via PTX-sensitive G-proteins (*direct mode*); or 2) by slowly up-regulating LTCCs activity through a cAMP/PKA-mediated phosphorylation (*remote mode*), similarly to the up-regulation of cardiac Cav1.2 channels during β_1 -AR-mediated sympathetic stimulation. The direct modulation is fully resolved in membrane micro-patches and derives from the interaction of G-protein subunits with LTCCs within membrane micro-domains where Ca^{2+} channels, membrane autoreceptors and coupled G-proteins coexist [44-46, 58]. The remote modulation is more complex and involves the presence of functional adenylate cyclases, an effective cAMP-mediated PKA up-regulation and LTCCs phosphorylation (see [51] and [52] for a review). The cAMP/PKA mediated up-regulation of LTCCs exists in BCCs ([46]), RCCs ([48]) and MCCs ([10]) and is thus a general property of chromaffin cells, related to the expression of β -ARs in these and other animal species [63, 64].

Although not directly related to the neurotransmitter-mediated modulation described above, LTCCs are also regulated by a remote mechanism which originates from the autocrine NO/cGMP/PKG

pathway which tonically inhibits LTCCs and catecholamine release in BCCs [47,65] and MCCs [49]. Chromaffin cells possess sufficient levels of neuronal-like NO-synthase and guanylyl cyclase to synthesize NO and cGMP [65] which can tonically down-regulate L-type currents and catecholamine release. A marked depression of L-type currents occurs also when NO-donors or permeable forms of cGMP are exogenously applied to isolated BCCs [47] or MCCs [49]. The action is PKG-dependent, develops slowly by reducing the open channel probability and causes no changes to the time course of L-type current activation. The NO/cGMP/PKG-mediated down-regulation of LTCCs exerts an opposite action to the cAMP/PKA-mediated up-regulation, closely resembling the effects that the two pathways exert on cardiac Cav1.2 channels.

As underlined elsewhere [49, 66, 67], the existence of the two modulatory pathways in MCCs, which express equal densities of functional Cav1.2 and Cav1.3 channels [11, 13], raises key questions concerning LTCC modulation:

- *Are Cav1.2 and Cav1.3 equally prone to cAMP/PKA and cGMP/PKG phosphorylation at basal conditions and during stimulation?*
- *Do the PKA- and PKG-mediated phosphorylations act independently on LTCCs?*
- *To what extent the two opposing mechanisms could synergistically affect the size of Cav1.2 and Cav1.3 Ca^{2+} currents?*

The answers to these questions are not obvious given the little amount of information available on cAMP/PKA and cGMP/PKG modulation of native Cav1.3 channels (see ref. [5]). In the following paragraph we briefly summarize the new findings that recently appeared on the sensitivity of Cav1.2 and Cav1.3 to cAMP/PKA and cGMP/PKG-mediated phosphorylation.

3.2 - Cav1.2 and Cav1.3 are equally modulated by cAMP/PKA and cGMP/PKG

Given the lack of selective blockers for Cav1.2 and Cav1.3, a suitable approach to quantify the effects of PKA/PKG activators and inhibitors on the two Cav1 isoforms is to use knock-out (KO) mice strains missing either one of the two isoforms. Comparing the effects on WT and Cav1.3^{-/-} KO mice and normalizing the effects by the size of L-type currents furnishes an immediate estimate of PKA/PKG actions on Cav1.2 in Cav1.3^{-/-} MCCs and Cav1.2 + Cav1.3 in WT MCCs. In MCCs, the estimate is

further simplified since Cav1.2 and Cav1.3 carry the same quantity of Ca^{2+} currents [11, 13]. An equal modulation of the two channels (either positive or negative) would turn out in a double effect on WT with respect to Cav1.3^{-/-} MCCs. Following this approach it has been possible to show that Cav1.2 and Cav1.3 are equally modulated by cAMP and cGMP in an opposing manner [49].

The cAMP/PKA pathway is already active at basal levels where both isoforms are equally down-regulated by the PKA blocker H89 (Fig. 1a) and are fully up-regulated by the adenylyl cyclase activator forskolin which causes a current increase proportional to the L-type current size in both WT and Cav1.3^{-/-} MCCs (Fig. 1b). Similar findings are reported for the LTCCs of RCCs [48], BCCs [46] and are comparable to the up-regulatory action of cAMP/PKA on cardiac Cav1.2 channels [68, 69]. This proves unequivocally that Cav1.3 is effectively up-regulated by the cAMP/PKA pathway, in good agreement with works reporting the existence of: 1) PKA phosphorylation sites in the C-terminal tail of recombinant Cav1.3 channels [70], 2) co-localization of the neuronal Cav1.3 C-terminal with the A-kinase anchoring protein 15 (AKAP15) and β_2 -AR in mouse brain [71] and 3) a PKA and cAMP-mediated up-regulation of Ca^{2+} currents carried by the Cav1.3 long C-terminal splice variant [72] and the recombinant 180 kDA Cav1.3 isoform [73]. These two latter works also suggest that at variance of Cav1.2, AKAPs are not specifically required for reconstituting PKA-mediated up-regulation of Cav1.3 in heterologous expression systems, in apparent contradiction with the observation that AKAP15 co-localizes with neuronal Cav1.3 α_1 subunit [71]. The presence of AKAPs is nevertheless critical for the reconstituted PKA-modulation of Cav1.2 in tsA201 cells. This is due to the existence of *in vivo* C-terminal proteolytic processing which cleave a distal segment of the C-terminal regulatory domain and make the truncated form of the channel insensitive to PKA-modulation [74]. Under these conditions, PKA-modulation of Cav1.2 is re-established only when the truncated Cav1.2 subunit is reconstituted in the presence of the distal C-terminal domain, AKAP15 and the auxiliary $\alpha_2\delta_1$ and β_{2b} subunits [74].

As for cAMP/PKA, also the cGMP/PKG pathway is already active at basal levels, with the main difference that it down-regulates the L-type currents when activated in MCCs [49] and BCCs [47, 75]. Also in this case, comparing the effects of PKG inhibitors (KT 5823) and cGMP analogues (8-pCPT-

cGMP) on WT and Cav1.3^{-/-} MCCs, Cav1.2 and Cav1.3 are shown to be equally down-regulated at rest by basal levels of active PKG (Fig. 1d) and elevations of cGMP cause a proportional down-regulation of L-type currents in WT and Cav1.3^{-/-} MCCs (Fig. 1c). Thus, as for Cav1.2, also Cav1.3 is effectively down-regulated by cGMP/PKG. This agrees with data on cochlear inner hair cells reporting a NO/cGMP-mediated down-regulation of Cav1.3 currents [76].

3.3 - cAMP- and cGMP-mediated phosphorylation act opposingly and independently on LTCCs gating

The existence of two opposing modulations mediated by cAMP and cGMP in neuroendocrine cells, closely resembling those acting on the Cav1.2 isoform of ventricular myocytes, opens interesting questions about the role that these two modulatory pathways are likely to play in chromaffin cell functioning. A number of functional and structural evidence supports the view that the two systems act independently on both channel isoforms (or auxiliary β -subunits). The first set of evidence comes from protein microsequencing and peptide mapping studies which suggest that the two kinases phosphorylate different serine sites of the Cav1.2 α_1 and β subunits. PKA phosphorylates Cav1.2 α_1 -subunit at Ser1700 (cardiac) and Ser1901 (neuronal) sites [70, 74, 77, 78] and Cav1.2 β -subunit at Ser 478/479 and Ser 459 [79-81], while PKG phosphorylates Cav1.2 α_1 -subunit at Ser533 [77, 82-83] and Cav1.2 β subunit at Ser 496 [82]. The same could occur for the phosphorylating action of PKA and PKG on Cav1.3 α_1 -subunit in which PKA is shown to phosphorylate Ser1964, Ser1743 and Ser1816 [84, 85], but nothing is known about the phosphorylation sites of PKG. PKA and PKG phosphorylate also Ser 1928 [74, 83], which is also phosphorylated by PKC [86], but this serine residue is shown not to be functionally linked to Cav1.2 channel gating modification [87]. Apart from this singularity, all the other sites sensitive to PKA and PKG phosphorylation appear to be distinct. It should be noticed, however, that the existence of distinct phosphorylation sites for PKA and PKG do not rule out the possibility that they may mutually interfere on channel gating.

The second and most convincing evidence in favor of an independent action of the two pathways is the synergist effect that cAMP and cGMP can produce when opposingly modulated (one activated and the other inhibited) to give either minimal or maximal values of L-type currents differing by one order of magnitude (18% vs. 180% of control value). Fig. 2 illustrates the effects of the two opposing

synergistic regulations of L-type currents. In one case, the application of the cAMP/PKA activator forskolin followed by the selective PKG inhibitor KT 5823 leads to a nearly doubling of the L-type current (Fig. 2 bottom-right), while in the other case the sequential application of the PKA-blocker H89 and the cGMP analogue, 8-pCPT-cGMP, leads to a reduction of about 80% of the same current (Fig. 2 top-right). The cumulative PKA- and PKG-mediated modulation of L-type currents summarized in Fig. 2-left are among the most impressive changes of neuronal L-type current amplitude that can be driven by intracellular modulatory pathways. They are strikingly similar to the changes observed for the cardiac Cav1.2 currents when separately enhanced by cAMP or down-regulated by cGMP [5, 69, 88, 89], which set the systolic strength and sino-atrial node frequency during sympathetic and parasympathetic regulation of heart beating.

3.4 - Functional relevance of Cav1.2 and Cav1.3 modulation on chromaffin cell functioning

Recent works on MCCs, BCCs and RCCs [46, 48, 49] point to a converging idea that cAMP/PKA and cGMP/PKG pathways are already active at rest due to the basal activity of the two cyclases (AC and GC). AC is mainly activated by PACAP [90], Ca²⁺ entry and G protein subunits that are activated by the basal activity of hormones and neurotransmitters released by sympathetic neurons [91], surrounding capillaries [92, 93] and by the autocrine activity of chromaffin cells [42, 48, 57, 58]. This latter is most probably the main cause of the high basal level of cAMP in culture conditions (2.2 mM) that rises 2- to 3-fold during β_1 -AR stimulation and/or PDE-4 selective inhibition [10]. The soluble GC is activated by the resting NO levels generated by the Ca²⁺-calmodulin-mediated activation of NO synthase (NOS) expressed in most chromaffin cells [65, 94]. Under these conditions, cGMP/PKG appears to work as a 'break' to limit the potentiating effects of cAMP/PKA and helps setting the resting levels of Cav1.2 and Cav1.3 currents.

A *synergistic potentiation* of LTCCs could occur during sustained sympathetic stimulations that releases PACAP and induces massive secretion of adrenaline from chromaffin cells, which would further raise the levels of cAMP/PKA through the autocrine activation of the β_1 -ARs present in RCCs and MCCs [10, 48]. The increased Ca²⁺ entry during high-frequency stimulation could in turn activate the cGMP-specific Ca²⁺-calmodulin-dependent PDE (PDE1) that regulates the resting levels of cGMP

[65, 95]. Thus, activation of a cGMP-specific PDE that lowers cGMP/PKG levels and the parallel increase of PKA during PACAP release and β_1 -AR stimulation could markedly enhance Cav1.2 and Cav1.3 currents. This would sustain the rapid increase of firing activity and catecholamine release that ensure the fast activation of the 'fight-or-flight response' in chromaffin cells.

A reversed action (*synergistic inhibition*) could occur if, as in most mammalian ventricular myocytes, chromaffin cells possess cGMP-activated PDE2 isoform that hydrolyses cAMP [96]. Any robust up-regulation of the NO/cGMP/PKG pathway under these conditions would enhance cGMP and down-regulate cAMP which would rapidly depress Cav1.2 and Cav1.3 channel gating. The existence of other PDEs acting on cAMP beside PDE4, is supported by the findings that in MCCs the non-specific PDE blocker IBMX increases basal cAMP levels more potently than the PDE-4-specific blocker rolipram [10].

The Cav1.3 up- and down-regulation by PKA and PKG described here could have also key physiological significance if extrapolated to the Cav1.3 channels of other tissues where the channel is highly expressed and functional [12]. In cardiac sino-atrial and atrio-ventricular node cells, Cav1.3 contributes to the pacemaker current controlling heart beating [8, 97] and, thus, a β_1 -AR or a NO/cGMP/PKG-driven modulation of its gating could either accelerate or decelerate the heart rate. Effective modulations driven by the cAMP/PKA and cGMP/PKG pathways could occur also to the Cav1.3 channels of cochlear inner hair cells [98] and dopaminergic neurons of substantia nigra pars compacta [99] which control key functions of hearing sensory transduction and motor control.

4. - Cav1.3 as pacemaker channels in chromaffin cells

The first recordings of AP firing in chromaffin cells date back to 1976. Current injection or application of ACh was able to make gerbil chromaffin cells fire when measured in acute isolated slices or primary cultures [100]. Neither the absence of extracellular Ca^{2+} , nor the block of VGCCs by Co^{2+} was able to induce a cessation of firing in their preparation, indicating a key contribution of voltage-gated Na^+ channels to spontaneous firing. Data on RCCs, published in the same year, were also reported by Brandt et al., 1976 [101]. Interestingly, these authors reported that by means of extracellular

recordings, RCCs fired in a slow but spontaneous manner (0.05-0.1 spikes/sec). This was not observed in recordings where the cell was impaled by a sharp micropipette, presumably due to damage of the cell membrane that typically results in a lower input resistance. A later study by Kidokoro and Ritchie [102] strengthened the existing evidence that RCCs possess the prerequisites to spontaneously fire AP's. They measured spontaneously discharged AP's by extracellular recordings in physiological conditions. It has to be mentioned that Brandt et al. [101] measured APs in 10 mM extracellular Ca^{2+} that might elevate tonic Ca^{2+} -activated K^+ currents which could interfere with cell's excitability. An increased divalent cation concentration in the bath is known to lead to a positive shift of the activation threshold of voltage-gated channels [103, 104]. This thus implicates that the cell has to bridge a bigger voltage gap in order to trigger an AP. Under physiological conditions as used by Kidokoro and Ritchie [102], the ability to generate spikes was abolished by Co^{2+} (a non-selective VGCC blocker) and not by TTX, suggesting a key role for VGCCs in spontaneous spike generation in RCCs. Until recently, the general belief was that pacemaker channels were either belonging to the HCN family, the low-threshold persistent Na_v channels or the low-voltage activated (LVA; T-type) Ca^{2+} channels. One of the criteria for a channel to contribute to the pacemaker is in fact that it has to open at low membrane potentials and carry inward cation currents. This was the reason why HVA Ca^{2+} channels were excluded "by definition". Dopaminergic neurons with an L-type dependent pacemaker mechanism were in fact considered rather an exception [105].

4.1 – Nav1 channels of chromaffin cells do not fulfill the requirements of pacemaker channels

The Na^+ currents of chromaffin cells are typically transient in nature and mainly TTX-sensitive [13, 106, 107]. All inward current ascribed to Na^+ can be blocked by TTX at concentrations as low as 100 nM [108]. Klugbauer et al. [109] were the first showing that BCCs express high densities of the $\text{Na}_v1.7$ isoform. It is even believed that the $\text{Na}_v1.7$ channel that originally was called PN1 (peripheral nervous system specific) is the only isoform expressed in chromaffin cells [106, 107]. $\text{Na}_v1.7$ in fact shows a peculiar expression pattern with high levels of expression in chromaffin cells, sympathetic ganglia and $\text{A}\beta$ - as well as C-type dorsal root ganglion (DRG) neurons [107, 110-112].

Na_v1.7 channels produce fast activating and inactivating currents that generally recover slowly from fast inactivation [107]. This slow recovery from inactivation is generally believed not to permit and sustain prolonged high-frequency firing of excitable cells. Na_v1.7 channels, however, respond to slow depolarizing ramps due to the slow onset of closed state inactivation and could trigger boosting depolarizations to bring V_{rest} closer to V_{thresh} [110]. Data from our lab unequivocally indicate that Na_v1 block by 300 nM TTX does not block spontaneous membrane oscillations of MCCs [12, 13]. AP-clamp experiments prove clearly that Na⁺ currents in MCCs are confined to the AP-upstroke phase, showing a rather complete inactivation during the broad Ca²⁺-dependent spike waveforms [31] (Fig. 3). When MCCs are forced into fast firing patterns (e.g. by blocking SK channels) we observed a continuous and steady decrease of the spike amplitude, spike threshold and dV/dt_{max} which are all indicative of a reduced Na_v channel availability [31]. This was not found to interfere with the ability of MCCs to elicit spontaneous APs, indicating that fast inactivating Na⁺ currents fulfill only a minor role in MCCs pacemaking.

4.1 – LTCCs as pacemaker channels in chromaffin cells

In contrast with TTX, nifedipine (3 μM) just like Cd²⁺ is very effective in blocking MCCs spontaneous activity [10-12, 50]. A property that is not limited to chromaffin cells but is shared with several central neurons in which LTCCs are responsible for pacemaking [see [12] for a review]. Given that MCCs and RCCs express only Cav1.2 and Cav1.3 channels [11, 16, 24] the pacemaking of these cells is obviously restricted to either one of the two channels. Cav1.3 in particular exhibits biophysical features that are critical for pacemaking. The channel possesses a low-threshold of activation (-50 mV in 2 mM Ca²⁺), fast activation kinetics and relatively slow inactivation characteristics [6, 113]. The interest in Cav1.3 as a pacemaker channel grew substantially after it was shown that it contributes to the spontaneous firing patterns of substantia nigra (SN) dopaminergic neurons [9, 114]. Given this, Marcantoni et al. [11] investigated the exact nature of the LTCC-dependent pacemaker of MCCs. Since no Cav1 selective DHP's are available yet, this information was provided by experiments conducted on Cav1.3^{-/-} mice. Strikingly, Cav1.3 deficiency leads to a loss of firing in the majority of MCCs. The small portion of cells that maintained their spontaneous activity, fired at elevated

frequencies and were characterized by depolarized resting membrane potentials of -40 to -45 mV. WT-MCCs typically fluctuate around a V_{rest} of -45 to -50 mV that perfectly coincides with the activation threshold of Cav1.3 channels and is about 9 and 25 mV more negative than that of Cav1.2 and Nav1 channels, respectively [13]. Especially at V_{rest} the input resistance is notably elevated (4-5 G Ω) ensuring significant membrane depolarizations, even upon small inwardly directed Ca^{2+} currents (2-3 pA). Voltage- and Ca^{2+} -dependent inactivation of Cav1.3 is moreover minimal around the pacemaker potential, favoring the development of persistent Ca^{2+} currents that cover the interspike intervals as is evident for WT MCCs (Fig. 3). This pacemaker current in fact is strongly attenuated in Cav1.3^{-/-} MCCs, unequivocally proving its direct involvement in MCCs pacemaking. Impeding Ca^{2+} ions to flow in the cell through LTCCs leads to a cell depolarization instead of an expected hyperpolarization. This apparent anomaly can be explained by predicting a specific coupling of LTCCs (Cav1.3 in particular) to Ca^{2+} -activated K^+ channels that fine tune MCCs pacemaking.

4.2 - MCCs pacemaking: a healthy balance between Cav1.3 inward Ca^{2+} currents and Cav1.3-driven outward K^+ currents

As mentioned before, Ca^{2+} ions serve a multitude of roles in cellular physiology, ranging from neural growth and brain plasticity to cell death. Ca^{2+} is furthermore able to affect gene expression and trigger exocytosis of neurotransmitters [115]. The "power" of Ca^{2+} as an effective second messenger partly resides in the huge concentration gradient difference that is maintained across the cell membrane. Ca^{2+} outside the cell is 10^4 times more concentrated as the unbound Ca^{2+} inside. Thus, cells that make use of a Ca^{2+} -dependent pacemaker need enhanced metabolic controls in order to guarantee efficient Ca^{2+} handling [114, 116]. An additional way to reduce the total amount of Ca^{2+} influx is by shortening the AP duration and slowing down the pace of firing by activating Ca^{2+} -activated K^+ channels. For MCCs in fact the "drive" of pacemaking (Cav1.3) is strongly coupled to BK and SK channels that act as a "brake" in these cells [11, 12, 31]. Cav1.3 deficiency results in a strong and almost complete block of a transient outward current that covers the region in between the spike peak and the antipeak. This current was shown to reflect a paxilline-sensitive fast-inactivating BK current, which is characteristic

of MCCs [11, 50], RCCs [117] and BCCs [118]. Loss of this BK current typically results in broader APs and reduced fast after-hyperpolarization phases [11].

Although the acute application of paxilline leads to an increase of the firing frequency (as is the case of a fraction of spontaneously firing Cav1.3^{-/-} MCCs), BK^{-/-} MCCs did exhibit reduced firing frequencies [12]. Similarly, Purkinje neurons of BK^{-/-} mice show reduced firing frequencies as compared to WT's [119]. Upon stimulation we observed furthermore that BK loss reduces the onset of instantaneous firing frequency and fastens the time course of spike frequency adaptation of MCCs (Vandael & Carbone, unpublished data). A similar phenomenon has been observed for hippocampal pyramidal neurons [120]. A possible explanation could be that more Ca²⁺ enters during the broader AP's and triggers more prominent SK currents. Thus even though Cav1.3 deficiency leads to an aberrant activation of BK channels, this phenomenon is not likely the only reason of the observed increased firing frequencies of the spontaneously active Cav1.3^{-/-} MCCs. Moreover, Cav1.3^{-/-} MCCs typically respond to BayK 8644 with a sudden strong rise in the firing frequency followed by a depolarization block, a phenomenon that was not always observed in MCCs lacking the BK channels [12]. All BK^{-/-} MCCs could be forced into depolarization block when BayK 8644 was co-applied together with apamin, a potent SK channel blocker. This thus indicates that Cav1.3^{-/-} MCCs show an aberrant activation of both BK and SK channels.

4.3 - A functional coupling between Cav1.3 (the motor) and SK channels (the brake) drives MCCs firing

The above mentioned issues were the drive for studying the role of Cav1.3 in SK channel activation. SK channels were first identified in BCCs by Marty and Neher [121], and later in RCCs by Neely & Lingle [122] but their role on chromaffin cell excitability remained elusive. We have recently shown that MCCs express SK1-3 channels that contribute to a tonic current that slows-down basal firing frequencies [31]. Interestingly, the resting firing frequency does reflect the amount of SK channels that contribute to MCCs pacemaking. A negative correlation in fact exists between the basal frequency and its percentage increase induced by apamin [31] (Fig. 4). In other words, the more SK channels are

available in a chromaffin cell the lower is its resting firing frequency. Fig. 4 shows two examples of MCCs displaying low and high firing frequency and how apamin affects more markedly the MCC firing slowly. In line with this, SK channels contribute to spike frequency adaptation upon current injection (Fig. 5a,c). In brief, SK channels give rise to a gradual increase in interspike interval duration when MCCs are stimulated. Spike frequency adaptation furthermore enhances the availability of Na_v channels, necessary to maintain stable AP waveforms. As suggested by Engel et al. [123], SK currents might shunt the membrane resistance, meaning that more current is required to induce a certain change in voltage. This also implicates that cells activating a sufficient amount of SK channels can deal with stronger current inputs as compared to cells with little SK activation. In this regard we found that $\text{Cav1.3}^{-/-}$ MCCs show a remarkable lower degree of spike frequency adaptation (Fig. 5 a,c) and an earlier switch into depolarization block as compared to WTs [31]. Both findings suggest that $\text{Cav1.3}^{-/-}$ MCCs have difficulties in triggering SK currents. It is thus tempting to hypothesize that the observed increased firing frequencies of the subgroup of spontaneously active $\text{Cav1.3}^{-/-}$ MCCs can be attributed to a reduced degree of SK activation.

When SK currents are measured during a train of APs it is striking that they build-up during the interspike intervals in WT MCCs and that this phenomenon is strongly attenuated in $\text{Cav1.3}^{-/-}$ MCCs (Fig. 5b,d). Nifedipine was moreover able to induce a complete block of the interspike interval specific SK current, enforcing the idea that LTCCs are critical in triggering SK channels activation in MCCs. Voltage-clamp experiments on slow SK tail currents pointed out although that LTCCs and non-LTCCs contribute equally to SK activation. Indeed, a train of APs is not at all comparable to a square pulse voltage-clamp protocol. Analyzing the total amount of Ca^{2+} charge flowing in the cell during the firing cycle we concluded that Cav1.3 shows a major contribution as compared to the other HVA channels. Already the interspike interval is in charge of 50% of the total Ca^{2+} charge surging into the cell and is largely absent in $\text{Cav1.3}^{-/-}$ MCCs. Thus, Cav1.3 is coupled in a functional rather than a molecular manner to SK channels in MCCs.

Noteworthy is the overlap between the outward Cav1.3-dependent SK current and the inwardly directed Cav1.3 current itself during the pacemaker cycle (Fig. 5b,d). The sum of both has its

implications on the rate of depolarization and consequently on the duration of the interspike interval. MCCs are thus a perfect example of a spontaneously active cell type where the “motor” of the system (Cav1.3) controls the “brake” (SK) in order to prevent over-excitation. These findings furthermore indicate that modulation of the "brake" could have as important consequences on catecholamine release as modulation of the "motor".

5. L-type channels and secretion: any specific role for Cav1.2 and Cav1.3?

In chromaffin cells, the different VGCCs (L, N, P/Q, R, T) contribute to exocytosis proportionally to their density of expression and gating properties [33-36, 124-128]. Secretion is not particularly linked to any specific Ca^{2+} channel type and either deletion or up-regulation of one of them causes a proportional change in secretion. For instance, Cav2.1 deletion causes a loss of the P/Q-type currents with a compensatory increase of L-type currents and secretion [129]. Similarly, when up-regulated by cAMP [130] or chronic hypoxia,[131] T-type channels contribute to low-threshold exocytosis with the same Ca^{2+} -dependence of L-type channels [38].

As shown above, Cav1.2 and Cav1.3 represent the final target of different modulatory pathways mediated by direct activation of G protein-coupled autoreceptors or remote pathways involving PKA and PKG activation. In principle, by acting on the two isoforms, any of these signaling loops can exert also a potent modulatory effect on the exocytotic response. In fact, the L-type current increase induced by cAMP stimulation only accounts for 20% of the total secretory response, suggesting an additional down-stream effect on the secretory machinery [36]. It is also interesting to notice that, due to their slower and less complete time-dependent inactivation, LTCCs (in particular Cav1.3) are favored in triggering exocytosis with respect to other HVA Ca^{2+} channels during sustained stimuli. Nevertheless, the contribution of Cav1.2 and Cav1.3 to exocytosis remains proportional to the quantity of Ca^{2+} ions entering the cell, suggesting that there is no preferential co-localization of LTCCs to secretory granules [51].

Although the critical role of LTCCs in triggering exocytosis is well established [36, 53, 132, 133] there are no clear indications of a possible distinct role of Cav1.2 and Cav1.3 to exocytosis, despite the

different inactivation kinetics and voltage range of activation of the two isoforms [7]. Preliminary observations show that deletion of the Cav1.3 subunit in MCCs lowers the amount of exocytosis at very negative potentials (-50 to -30 mV in 10 mM Ca^{2+}) (Navarro V, Carbone E, Carabelli V, unpublished data), indicating that besides sustaining action potential firing, Cav1.3 preferentially contributes to exocytosis at low membrane potentials [66]. This is in good agreement with what recently reported by Pérez-Alvarez et al. [26], in which secretion of Cav1.3^{-/-} mice is partially reduced at very negative potentials. In this way, Cav1.3 contributes to the low-threshold exocytosis similarly to the T-type Cav3.2 channel when expressed during cAMP/PKA elevations [38, 127] or chronic hypoxia conditions [131].

6. L-type channels and endocytosis: are Cav1.3 and Cav1.2 equally involved?

Another open question to solve concerns the role of Ca^{2+} and Ca^{2+} channels in the retrieval of synaptic vesicles during endocytosis. Neuroendocrine chromaffin cells exhibit different types of endocytosis, according to cell activity and stimulation protocols. Square pulse depolarizations cause exocytosis followed by a decline in membrane capacitance, which can reach the pre-stimulus level (*compensatory* endocytosis) or fall even below (*excess* endocytosis) [134, 135]. Transition between these two modes appears to be regulated by intracellular Ca^{2+} : the retrieval being accelerated and potentiated by increasing Ca^{2+} levels. It is interesting that in chromaffin cells endocytosis is also supported by barium [136] by the activation of kinase/phosphatase-mediated pathways [134, 137] and by additional pathways of vesicle recycling independent of dynamin and calmodulin [40]. In bovine chromaffin cells, the mechanism of action involves sphingosine and originates at the intracellular site [138]. The phenomenon is lost when applying repeated stimuli [139].

Concerning the role of Ca^{2+} channels in sustaining endocytosis, a preferential coupling of LTCCs to endocytosis has been recently proposed: Ca^{2+} -entry through LTCCs in bovine chromaffin cells is more effective in triggering endocytosis than exocytosis [39] (see also [41] for a recent review). Even at the mammalian neuromuscular junction, endocytosis is mainly sustained by LTCCs, while P/Q-type channels trigger exocytosis [140]. Block of LTCCs by DHPs decreases endocytosis and directs newly

formed synaptic vesicles to a slow-release vesicle pool during high-frequency stimulation. According to the proposed model, in the absence of functioning LTCCs, endocytosis cannot be sufficiently fast to balance exocytosis causing vesicular membrane to accumulate at the presynaptic surface.

Given the main role of LTCCs in controlling compensatory and excess endocytosis, the next issue is whether Cav1.2 or Cav1.3 has a preferential control on vesicle retrieval. One argument in favor of Cav1.3 is its slower and less complete time-dependent inactivation with respect to Cav1.2. The delayed inactivation of Cav1.3 could be physiologically relevant for sustaining prolonged Ca^{2+} influxes that support normal endocytosis. This can be certainly clarified comparing the endocytotic responses to Ca^{2+} loads in WT and Cav1.3^{-/-} MCCs and check whether the hypothesis of an equal role of Cav1.2 and Cav1.3 on vesicle retrieval is satisfied. Under these conditions, a nearly half reduction of functional LTCCs, as occurs in Cav1.3^{-/-} MCCs, should cause a proportional decrease of endocytosis. Finally, also the coupling of Cav1.2 and Cav1.3 to calmodulin, the Ca^{2+} sensor of different forms of endocytosis [141] could be a further molecular target of differential regulation of the endocytosis that needs to be tested.

Concluding remarks

The importance of Cav1.2 and Cav1.3 channels in the control of chromaffin cell firing, catecholamine secretion and vesicle retrieval is of growing interest and supported by an increasing number of upcoming papers [12, 18, 41, 67]. Cav1.3 and Cav1.2 contribute to shape the AP waveform and support the pacemaker current. Cav1.3, in particular, activates BK and SK channels that in turn set the frequency of spontaneously firing cells and help adapting the spike frequency during prolonged depolarization. At present, there are no indications of any preferential coupling of either one isoform to the exo- and endocytotic machinery. Since both channels are equally modulated by PKA- and PKG-mediated pathways and may undergo drastic up and down autocrine regulations, it is expected that both isoforms play a critical role in chromaffin cell functioning.

Future works will certainly help to identify how the voltage-dependence of activation-inactivation gating and modulatory pathways acting on these Cav1 channels may fine tune cell firing and Ca^{2+} -

dependent excitation-secretion coupling. New findings on the functional role of these two LTCCs on chromaffin cells activity will be beneficial not only for understanding stress-related pathologies associated to the adrenal gland [67] but also for resolving the origin of diseases of central neurons, sensory and cardiac cells where the two LTCCs are highly expressed and play vital roles [12, 16].

References

- [1] W.A. Catterall, E. Perez-Reyez, T.P. Snutch, J. Striessnig, Nomenclature and structure-function relationships of voltage-gated calcium channels, *Pharmacol. Rev.* 57 (2005) 411-425.
- [2] L. Lacinová, N. Klugbauer, F. Hofmann, Low voltage activated calcium channels: from genes to function, *Gen. Physiol. Biophys.* 19 (2000) 121-136.
- [3] E. Pérez-Reyes, Molecular physiology of low-voltage-activated T-type calcium channels, *Physiol. Rev.* 83 (2003) 117-161.
- [4] E. Carbone, A. Marcantoni, A. Gianniccoli, D. Guido, V. Carabelli, T-type channels-secretion coupling: evidence for a fast low-threshold exocytosis, *Pflügers Arch.* 453 (2006) 373-383.
- [5] W.A. Catterall, Voltage-gated calcium channels, *Cold Spring Harb. Perspect. Biol.* 3 (2011) a003947.
- [6] J. Platzer, J. Engel, A. Schrott-Fischer, K. Stephan, S. Bova, H. Chen, H. Zheng, J. Striessnig, Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type Ca^{2+} channels, *Cell* 102 (2000) 89-97.
- [7] A. Koschak, D. Reimer, I. Huber, M. Grabner, H. Glossmann, J. Engel, J. Striessnig, Alpha 1D (Cav1.3) subunits can form L-type Ca^{2+} channels activating at negative voltages, *J. Biol. Chem.* 276 (2001) 22100-22106.
- [8] M.E. Mangoni, B. Couette, E. Bourinet, J. Platzer, D. Reimer, J. Striessnig, J. Nargeot, Functional role of L-type $Ca_v1.3$ Ca^{2+} channels in cardiac pacemaker activity, *Proc. Natl. Acad. Sci. USA* 100 (2003) 5543-5548.
- [9] C.S. Chan, J.N. Guzman, E. Llijic, J.N. Mercer, C. Rick, T. Tkatch, G.E. Meredith, D.J. Surmeier. 'Rejuvenation' protects neurons in mouse models of Parkinson's disease, *Nature* 447 (2007) 1081-1086.
- [10] A. Marcantoni, V. Carabelli, D.H. Vandael, V. Comunanza, E. Carbone, PDE type-4 inhibition increases L-type Ca^{2+} currents, action potential firing and quantal size of exocytosis in mouse chromaffin cells, *Pflügers Arch.* 457 (2009) 1093-1110.
- [11] A. Marcantoni, D.H.F. Vandael, S. Mahapatra, V. Carabelli, M.J. Sinnegger-Brauns, J. Striessnig, E. Carbone, Loss of Cav1.3 channels reveals the critical role of L-type and BK channel coupling in pacemaking mouse adrenal chromaffin cells, *J. Neurosci.* 30 (2010) 491-504.
- [12] D.H.F. Vandael, A. Marcantoni, S. Mahapatra, A. Caro, P. Ruth, A. Zuccotti, M. Knipper, E. Carbone, Cav1.3 and BK channels for timing and regulating cell firing, *Molec. Neurobiol.* 41 (2010) 185-198.
- [13] S. Mahapatra, A. Marcantoni, D.H.F. Vandael, J. Striessnig, E. Carbone, Are Cav1.3 pacemaker channels in chromaffin cells? Possible bias from resting cell conditions and DHP blocker usage, *Channels (Austin)* 5 (2011) 219-224.
- [14] I. Calin-Jageman, A. Lee, Ca_v1 L-type Ca^{2+} channel signaling complexes in neurons, *J. Neurochem.* 105 (2008) 573-583.
- [15] S.S. Stojilkovic, J. Tabak, R. Bertram, Ion channels and signaling in the pituitary gland, *Endocrine Rev.* 31 (2010) 845-915.
- [16] A. Zuccotti, S. Clementi, T. Reinbothe, A. Torrente, D.H.F. Vandael, A. Pirone, Structural and functional differences between L-type calcium channels: crucial issues for further selective targeting, *Trends Pharmacol. Sci.* 32 (2011) 366-375.
- [17] H.C. Tuckwell, Quantitative aspects of L-type Ca^{2+} currents, *Prog. Neurobiol.* 96 (2012) 1-31.
- [18] S. Mahapatra, C. Calorio, D.H.F. Vandael, A. Marcantoni, V. Carabelli, E. Carbone. Calcium channel types contributing to chromaffin cell excitability, exocytosis and endocytosis, *Cell Calcium* 51 (2012) 321-330.

- [19] B.P. Bean, Nitrendipine block of cardiac calcium channels: high-affinity binding to the inactivated state, *Proc. Natl. Acad. Sci. USA* 81 (1984) 6388-6392.
- [20] A. Welling, A. Ludwig, S. Zimmer, N. Klugbauer, V. Flockerzi, F. Hofmann, Alternatively spliced IS6 segments of the alpha 1C gene determine the tissue-specific dihydropyridine sensitivity of cardiac and vascular smooth muscle L-type Ca²⁺ channels, *Circ. Res.* 81 (1997) 526-532.
- [21] T.D. Helton, W. Xu, D. Lipscombe, Neuronal L-type calcium channels open quickly and are inhibited slowly, *J. Neurosci.* 25 (2005) 10247-10251.
- [22] D.J. Triggle, Calcium channel antagonists: clinical uses--past, present and future, *Biochem. Pharmacol.* 74 (2007) 1-9.
- [23] E. García-Palomero, I. Cuchillo, A.G. García, J. Renart, A. Albillos, C. Montiel, Greater diversity than previously thought of chromaffin cell Ca²⁺ channels, derived from mRNA identification studies, *FEBS Letters* 481 (2000) 235-239.
- [24] P. Baldelli, J.M. Hernández-Guijo, V. Carabelli, M. Novara, T. Cesetti, E. Andrés-Mateos, C. Montiel, E. Carbone, Direct and remote modulation of L-channels in chromaffin cells: distinct actions on α 1C and α 1D subunits?, *Mol. Neurobiol.* 29 (2004) 73-96.
- [25] A. Benavides, S. Calvo, D. Tornero, C. Gonzalez-Garcia, V. Ceña, Adrenal medulla calcium channel population is not conserved in bovine chromaffin cells in culture, *Neuroscience* 128 (2004) 99-109.
- [26] A. Pérez-Alvarez, A. Hernández-Vivanco, J.C. Caba-González, A. Albillos, Different roles of Cav1 channel subtypes in spontaneous action potential firing and fine tuning of exocytosis in mouse chromaffin cells, *J. Neurochem.* 116 (2011) 105-121.
- [27] P. Segura-Chama, C.V. Rivera-Cerecedo, R. González-Ramírez, R. Felix, J.M. Hernández-Guijo, A. Hernández-Cruz, Atypical Ca²⁺ currents in chromaffin cells from SHR and WKY rat strains result from the deficient expression of a splice variant of the α 1D Ca²⁺ channel, *Am. J. Physiol. Heart Circ. Physiol.* 302 (2012) H467-H478.
- [28] W. Xu, D. Lipscombe, Neuronal Cav1.3 α 1 L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines, *J. Neurosci.* 21 (2001) 5944-5951.
- [29] B.L. Brandt, S. Hagiwara, Y. Kidokoro, S. Miyazaki, Action potentials in the rat chromaffin cell and effects of acetylcholine, *J. Physiol. (Lond)* 263 (1976) 417-439.
- [30] A. Lieb, A. Scharinger, S. Sartori, M.J. Sinnegger-Brauns, J. Striessnig, Structural determinants of Cav1.3 L-type calcium channel gating, *Channels (Austin)* 6 (2012) 197-205.
- [31] D.H.F. Vandael, A. Zuccotti, J. Striessnig, E. Carbone, Cav1.3-driven SK channel activation regulates pacemaking and spike frequency adaptation in mouse chromaffin cells, *J. Neurosci.* in press (2012) (DOI:10.1523/jneurosci.3715-12.2012)
- [32] T. Nagayama, T. Matsumoto, F. Kuwakubo, Y. Fukushima, M. Yoshida, M. Suzuki-Kusaba, H. Hisa, T. Kimura, S. Satoh, Role of calcium channels in catecholamine secretion in the rat adrenal gland, *J. Physiol. (Lond)* 520 (1999) 503-512.
- [33] S.J. Kim, W. Lim, J. Kim, Contribution of L- and N-type calcium currents to exocytosis in rat adrenal medullary chromaffin cells, *Brain Res.* 675 (1995) 289-296.
- [34] K.L. Engisch, M.C. Nowycky, Calcium dependence of large dense-cored vesicle exocytosis evoked by calcium influx in bovine adrenal chromaffin cells, *J. Neurosci.* 16 (1996) 1359-1369.
- [35] E.A. Lukyanetz, E. Neher, Different types of calcium channels and secretion from bovine chromaffin cells, *Eur. J. Neurosci.* 11 (1999) 2865-2873.
- [36] V. Carabelli, A. Giancippoli, P. Baldelli, E. Carbone, A.R. Artalejo, Distinct potentiation of L-type currents and secretion by cAMP in rat chromaffin cells, *Biophys. J.* 85 (2003) 1326-1337.

- [37] T. Akiyama, T. Yamazaki, H. Mori, K. Sunagawa, Effects of Ca²⁺ channel antagonists on acetylcholine and catecholamine releases in the in vivo rat adrenal medulla, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287 (1) (2004) R161–R166.
- [38] V. Carabelli, A. Marcantoni, V. Comunanza, E. Carbone, Fast exocytosis mediated by T- and L-type channels in chromaffin cells: distinct voltage-dependence but similar Ca²⁺-dependence, *Europ. Biophys. J.* 36 (2007) 753-762.
- [39] J.M. Rosa, A.M. de Diego, L. Gandía, A.G. García, L-type calcium channels are preferentially coupled to endocytosis in bovine chromaffin cells, *Biochem. Biophys. Res. Commun.* 357 (2007) 834-839.
- [40] J.M. Rosa, L. Gandía, A.G. García, Permissive role of sphingosine on calcium-dependent endocytosis in chromaffin cells, *Pflügers Arch.* 460 (2010) 901-914.
- [41] J.M. Rosa, C. Nanclares, A. Orozco, I. Colmena, R. de Pascual, A.G. García, L. Gandía, Regulation by L-type calcium channels of endocytosis: an overview, *J. Mol. Neurosci.* 48 (2012) 360-367.
- [42] A. Albillos, L. Gandía, P. Michelena, J.A. Gilabert, M. del Valle, E. Carbone, A.G. García, The mechanism of calcium channel facilitation in bovine chromaffin cells, *J. Physiol. (Lond)* 494 (1996) 687–695.
- [43] V. Carabelli, M. Lovallo, V. Magnelli, H. Zucker, E. Carbone, Voltage-dependent modulation of single N-type Ca²⁺ channel kinetics by receptor agonists in IMR32 cells, *Biophys. J.* 70 (1996) 2144-2154.
- [44] J.M. Hernández-Guijo, V. Carabelli, L. Gandía, A.G. García, E. Carbone, Voltage-independent autocrine modulation of L-type channels mediated by ATP, opioids and catecholamines in rat chromaffin cells, *Eur. J. Neurosci.* 11 (1999) 3574-3584.
- [45] A. Hernández, P. Segura-Chama, N. Jiménez, A.G. García, J.M. Hernández-Guijo, A. Hernández-Cruz, Modulation by endogenously released ATP and opioids of chromaffin cell calcium channels in mouse adrenal slices, *Am. J. Physiol. Cell Physiol.* 300 (2011) C610–C623.
- [46] V. Carabelli, J.M. Hernández-Guijo, P. Baldelli, E. Carbone, Direct autocrine inhibition and cAMP-dependent potentiation of single L-type Ca²⁺ channels in bovine chromaffin cells, *J. Physiol. (Lond)* 532 (2001) 73–90.
- [47] V. Carabelli, M. D'Ascenzo, E. Carbone, C. Grassi, Nitric oxide inhibits neuroendocrine Ca(v)1 L-channel gating via cGMP-dependent protein kinase in cell-attached patches of bovine chromaffin cells, *J. Physiol. (Lond)* 541 (2002) 351-366.
- [48] T. Cesetti, J.M. Hernandez-Guijo, P. Baldelli, V. Carabelli, E. Carbone, Opposite action of beta1- and beta2-adrenergic receptors on Ca(V)1 L-channel current in rat adrenal chromaffin cells, *J. Neurosci.* 23 (2003) 73-83.
- [49] S. Mahapatra, A. Marcantoni, A. Zuccotti, V. Carabelli, E. Carbone, Equal sensitivity of Cav1.2 and Cav1.3 channels to the opposing modulations of PKA and PKG in mouse chromaffin cells, *J. Physiol. (Lond)* 590 (2012) 5053–5073.
- [50] A. Marcantoni, P. Baldelli, J.M. Hernández-Guijo, V. Comunanza, V. Carabelli, E. Carbone, L-type calcium channels in adrenal chromaffin cells: role in pace-making and secretion, *Cell Calcium* 42 (2007) 397-408.
- [51] A. Marcantoni, V. Carabelli, V. Comunanza, H. Hoddah, E. Carbone, Calcium channels in chromaffin cells: focus on L- and T-types, *Acta Physiol.* 192 (2008) 233–246.
- [52] E. Carbone, V. Carabelli, T. Cesetti, P. Baldelli, J.M. Hernández-Guijo, M. L. Giusta, G-protein- and cAMP-dependent L-channel gating modulation: a manifold system to control calcium entry in neurosecretory cells, *Pflügers Arch.* 442 (2001) 801–813.

- [53] A.G. Garcia, A.M. Garcia-De-Diego, L. Gandía, R. Borges, J. García-Sancho, Calcium signalling and exocytosis in adrenal chromaffin cells, *Physiol. Rev.* 86 (2006) 1093-1131.
- [54] L. Gandía, I. Mayorgas, P. Michelena, I. Cuchillo, R. de Pascual, F. Abad, J.M. Novalbos, E. Larrañaga, A.G. García, Human adrenal chromaffin cell calcium channels: drastic current facilitation in cell clusters, but not in isolated cells, *Pflugers Arch.* 436 (1998) 696-704.
- [55] L. Gandía, A.G. García, M. Morad, ATP modulation of calcium channels in chromaffin cells, *J. Physiol. (Lond)* 470 (1993) 55-72.
- [56] A. Albillos, E. Carbone, L. Gandía, A.G. García, A. Pollo, Opioid inhibition of Ca²⁺ channel subtypes in bovine chromaffin cells: selectivity of action and voltage-dependence, *Eur. J. Neurosci.* 8 (1996) 1561-1570.
- [57] K.P.M. Currie, A.P. Fox, ATP serves as a negative feedback inhibitor of voltage-gated Ca²⁺ channel currents in cultured bovine chromaffin cells, *Neuron* 16 (1996) 1027-1036.
- [58] V. Carabelli, I. Carra, E. Carbone, Localized secretion of ATP and opioids revealed through single Ca²⁺ channel modulation in bovine chromaffin cells, *Neuron* 20 (1998) 1255-1268.
- [59] K.S. Elmslie, W. Zhou, S.W. Jones, LHRH and GTP- γ -S modify calcium current activation in bullfrog sympathetic, *Neuron* 5 (1990) 75-80.
- [60] L. Boland, B.P. Bean, Modulation of N-type calcium channels in bullfrog sympathetic neurons by luteinizing hormone-releasing hormone: kinetics and voltage-dependence, *J. Neurosci.* 13 (1993) 516-533.
- [61] S.R. Ikeda, Voltage-dependent modulation of N-type calcium channels by G-protein $\beta\gamma$ subunits, *Nature* 380 (1996) 255-258.
- [62] S. Herlitze, G.H. Hockerman, T. Scheuer, W.A. Catterall, Molecular determinants of inactivation and G protein modulation in the intracellular loop connecting domains I and II of the calcium channel α 1A subunit, *Proc. Natl. Acad. Sci. USA* 94 (1997) 1512-1516.
- [63] S. Foucart, R. Nadeau, J. de Champlain, Local modulation of adrenal catecholamines release by beta-2 adrenoceptors in the anaesthetized dog Naunyn Schmiedebergs, *Arch. Pharmacol.* 337 (1988) 29-34.
- [64] V. Cortez, M. Santana, A.P. Marques, A. Mota, J. Rosmaninho-Salgado, C. Cavadas, Regulation of catecholamine release in human adrenal chromaffin cells by β -adrenoceptors, *Neurochem. Int.* 60 (2012) 387-393.
- [65] P.M. Schwarz, F. Rodriguez-Pascual, D. Koesling, M. Torres, U. Forstermann, Functional coupling of nitric oxide synthase and soluble guanylyl cyclase in controlling catecholamine secretion from bovine chromaffin cells, *Neuroscience* 82 (1998) 255-265.
- [66] V. Comunanza, A. Marcantoni, D.H.F. Vandael, S. Mahapatra, D. Gavello, V. Carabelli, E. Carbone, Cav1.3 as pacemaker channels in adrenal chromaffin cells: specific role on exo- and endocytosis?, *Channels (Austin)* 4 (2010) 440-446.
- [67] N.C. Guérineau, M.G. Desarménien, V. Carabelli, E. Carbone, Functional chromaffin cell plasticity in response to stress: focus on nicotinic, gap junction, and voltage-gated Ca²⁺ channels, *J. Mol. Neurosci.* 48 (2012) 368-386.
- [68] H. Reuter, Calcium channel modulation by neurotransmitters, enzymes and drugs, *Nature.* 301 (1983) 569-574.
- [69] H.C. Hartzell, R. Fischmeister, Opposite effects of cyclic GMP and cyclic AMP on Ca²⁺ current in single heart cells, *Nature* 323 (1986) 273-275.
- [70] J. Mitterdorfer, M. Froschmayr, M. Grabner, F.F. Moebius, H. Glossmann, J. Striessnig, Identification of PKA phosphorylation sites in the carboxyl terminus of L-type calcium channel α ₁ subunits, *Biochemistry* 35 (1996) 9400-9406.

- [71] M.R. Marshall, J.P. Clark III, R. Westenbroek, F.H. Yu, T. Scheuer, W.A. Catterall, Functional roles of a C-terminal signaling complex of Ca_v1 channels and A-kinase anchoring protein 15 in brain neurons, *J. Biol. Chem.* 286 (2011) 12627–12639.
- [72] Y. Liang, S.J. Tavalin, Auxiliary β subunits differentially determine PKA utilization of distinct regulatory sites on Cav1.3 L-type Ca^{2+} channels, *Channels (Austin)* 1 (2007) 102–112.
- [73] Y. Qu, G. Baroudi, Y. Yue, N. El-Sherif, M. Boutjdir, Localization and modulation of α_1D (Cav1.3) L-type Ca channel by protein kinase A, *Am. J. Physiol. Heart Circ. Physiol.* 288 (2005) H2123–H2130.
- [74] M.D. Fuller, M.A. Emrick, M. Sadilek, T. Scheuer, W.A. Catterall, Molecular mechanism of calcium channel regulation in the fight-or-flight response, *Sci. Signal* 3 (2010) ra70
- [75] F. Rodriguez-Pascual, M.T. Miras-Portugal, M. Torres, Modulation of the dihydropyridine-insensitive Ca^{2+} influx by 8-bromo-guanosine-3':5'-monophosphate, cyclic (8-Br-cGMP) in bovine adrenal chromaffin cells, *Neurosci. Lett.* 180 (1994) 269-272.
- [76] A. Almanza, F. Navarrete, R. Vega, E. Soto, Modulation of voltage-gated Ca^{2+} current in vestibular hair cells by nitric oxide, *J. Neurophysiol.* 7 (2007) 1188–1195.
- [77] K.S. De Jongh, B.J. Murphy, A.A. Colvin, J.W. Hell, M. Takahashi, W.A. Catterall, Specific phosphorylation of a site in the full-length form of the α_1 subunit of the cardiac L-type calcium channel by adenosine 3',5'-cyclic monophosphate-dependent protein kinase, *Biochemistry* 35 (1996) 10392–10402.
- [78] P. Gui, X. Wu, S.C. Stotz, R.J. Winkfein, E. Wilson, G.E. Davis, A.P. Braun, G.W. Zamponi, M.J. Davis, Integrin receptor activation triggers converging regulation of Cav1.2 calcium channels by c-Src and protein kinase A pathways, *J. Biol. Chem.* 281 (2006) 14015-14025.
- [79] B.L. Gerhardstein, T.S. Puri, A.J. Chien, M.M. Hosey, Identification of the sites phosphorylated by cyclic AMP-dependent protein kinase on the beta 2 subunit of L-type voltage-dependent calcium channels. *Biochemistry*, 38(1999) 10361-10370.
- [80] M. Bünemann, B.L. Gerhardstein, T. Gao, M.M. Hosey, Functional regulation of L-type calcium channels via protein kinase A-mediated phosphorylation of the beta(2) subunit, *Biol. Chem.* 274(1999) 33851-33854.
- [81] H. Haase, S. Bartel, P. Karczewski, I. Morano, E.G. Krause, In-vivo phosphorylation of the cardiac L-type calcium channel beta-subunit in response to catecholamines. *Mol Cell Biochem* (1996) 163-164: 99-106.
- [82] L. Yang, G. Liu, S.I. Zakharov, A.M. Bellinger, M. Mongillo, S.O. Marx, Protein kinase G phosphorylates Cav1.2 α_1C and β_2 subunits, *Circ. Res.* 101 (2007) 465–474.
- [83] L.H. Jiang, D.J. Gawler, N. Hodson, C.J. Milligan, H.A. Pearson, V. Porter, D. Wray, Regulation of cloned cardiac L-type calcium channels by cGMP-dependent protein kinase, *J. Biol. Chem.* 275 (2000) 6135–6143.
- [84] M.A. Emrick, M. Sadilek, K. Konoki, W.A. Catterall, Beta-adrenergic-regulated phosphorylation of the skeletal muscle $Ca(V)1.1$ channel in the fight-or-flight response, *Proc. Natl. Acad. Sci. USA* 107(2010) 18712-18717.
- [85] O. Ramadan, Y. Qu, R. Wadgaonkar, G. Baroudi, E. Karnabi, M. Chahine, M. Boutjdir, Phosphorylation of the consensus sites of protein kinase A on α_1D L-type calcium channel, *J. Biol. Chem.* 284 (2009) 5042-5049.
- [86] L. Yang, G. Liu, S.I. Zakharov, J.P. Morrow, V.O. Rybin, S.F. Steinberg, S.O. Marx, Ser1928 is a common site for Cav1.2 phosphorylation by protein kinase C isoforms, *J. Biol. Chem.* 280 (2005) 207–214.
- [87] T. Lemke, A. Welling, C.J. Christel, A. Blaich, D. Bernhard, P. Lenhardt, F. Hofmann, S.

- Moosmang, Unchanged beta-adrenergic stimulation of cardiac L-type calcium channels in Ca_v 1.2 phosphorylation site S1928A mutant mice, *J. Biol. Chem.* 283 (2008) 34738-34744.
- [88] P.F. Méry, C. Pavoine, L. Belhassen, F. Pecker, R. Fishmeister, Nitric oxide regulates cardiac Ca²⁺ current. Involvement of cGMP-inhibited and cGMP-stimulated phosphodiesterases through guanylyl cyclase activation, *J. Biol. Chem.* 268 (1993) 26286–26295.
- [89] J. Schlossmann, R. Feil, F. Hofmann, Signaling through NO and cGMP-dependent protein kinases, *Ann. Med.* 35 (2003) 21-27.
- [90] D.A. Przywara, X. Guo, M.L. Angelilli, T.D. Wakade, A.R. Wakade, A non-cholinergic transmitter, pituitary adenylate cyclase activating polypeptide, utilizes a novel mechanism to evoke catecholamine secretion in rat adrenal chromaffin cells. *J. Biol. Chem.* 271 (1996) 10545–10550.
- [91] K. Anderson, P.J. Robinson, P.D. Marley, Cholinoceptor regulation of cyclic AMP levels in bovine adrenal medullary cells. *Br. J. Pharmacol.* 106 (1992) 360–366.
- [92] S.P. Wilson, Vasoactive intestinal peptide elevates cyclic AMP levels and potentiates secretion in bovine adrenal chromaffin cells. *Neuropeptides* 11 (1988) 17–21.
- [93] P.D. Marley, Mechanisms in histamine-mediated secretion from adrenal chromaffin cells. *Pharmacol Ther* 98 (2003) 1–34.
- [94] M.J. Oset-Gasque, M. Parramon, S. Hortelano, L. Bosca, M.P. Gonzalez, Nitric oxide implication in the control of neurosecretion by chromaffin cells, *J. Neurochem.* 63 (1994) 1693–1700.
- [95] S. Vicente, M.P. Gonzalez, M.J. Oset-Gasque, Neuronal nitric oxide synthase modulates basal catecholamine secretion in bovine chromaffin cells, *J. Neurosci. Res.* 69 (2002) 327–340.
- [96] C. Lugnier, Cyclic nucleotide phosphodiesterase (PDE) superfamily: a new target for the development of specific therapeutic agents, *Pharmacol. Ther.* 109 (2006) 366–398.
- [97] Q. Zhang, V. Timofeyev, H. Qiu, L. Lu, N. Li, A. Singapuri, C.L. Torado, H.S. Shin, N. Chiamvimonvat, Expression and roles of Cav1.3 (α_1D) L-type Ca²⁺ channel in atrioventricular node automaticity, *J. Mol. Cell. Cardiol.* 50 (2011) 194–202.
- [98] W. Marcotti, S.L. Johnson, A. Rusch, C.J. Kros, Sodium and calcium currents shape action potentials in immature mouse inner hair cells, *J. Physiol. (Lond)* 552 (2003) 743–761.
- [99] J.M. Guzman, J. Sanchez-Padilla, C.S. Chan, D.J. Surmeier, Robust pacemaking in substantia nigra dopaminergic neurons, *J. Neurosci.* 29 (2009) 11011–11019.
- [100] B. Biales, M. Dichter, A. Tischler, Electrical excitability of cultured adrenal chromaffin cells, *J. Physiol. (Lond)* 262 (1976) 743-753
- [101] B.L. Brandt, S. Hagiwara, Y. Kidokoro, S. Miyazaki, Action potentials in the rat chromaffin cell and effects of acetylcholine, *J. Physiol. (Lond)* 263 (1976) 417-439.
- [102] Y. Kidokoro, A.K. Ritchie. Chromaffin cell action potentials and their possible role in adrenaline secretion from rat adrenal medulla, *J. Physiol. (Lond)* 307 (1980) 199-216.
- [103] B. Frankenhaeuser, A.L. Hodgkin, The action of calcium on the electrical properties of squid axons, *J. Physiol. (Lond)* 137 (1957) 218-244.
- [104] B. Hille, Charges and potentials at the nerve surface. Divalent ions and pH, *J. Gen. Physiol.* 51(1968) 221-236
- [105] B.P. Bean, The action potential in mammalian central neurons, *Nat. Rev. Neurosci.* 8 (2007) 451-465.
- [106] H. Kobayashi, S. Shiraishi, T. Yanagia, H. Yokoo, R. Yamamoto, S. Minami, T. Saitoh, A. Wada, Regulation of voltage-dependent sodium channel expression in adrenal chromaffin cells: involvement of multiple calcium signaling pathways, *Ann. NY Acad. Sci.* 971 (2002) 127-134.
- [107] A. Wada, E. Wanke, F. Gullo, E. Schiavon, Voltage-dependent Nav1.7 sodium channels: multiple roles in adrenal chromaffin cells and peripheral nervous system, *Acta Physiol. (Oxf)* 192

- (2008) 221-231.
- [108] B. Hollins, S.R. Ikeda, Inward currents underlying action potentials in rat adrenal chromaffin cells, *J. Neurophysiol.* 76 (1996) 1195-1211.
- [109] N. Klugbauer, L. Lacinova, V. Flockerzi, F. Hofmann, Structure and functional expression of a new member of the tetrodotoxin-sensitive voltage-activated sodium channel family from human neuroendocrine cells, *EMBO J.* 14 (1995) 1084-1090.
- [110] S.D. Dib-Hajj, T.R. Cummins, J.A. Black, S.G. Waxman, From genes to pain: Nav 1.7 and human pain disorders, *Trends Neurosci.* 30 (2007) 555-563.
- [111] L. Djouhri, R. Newton, S.R. Levinson, C.M. Berry, B. Carruthers, S.N. Lawson, Sensory and electrophysiological properties of guinea-pig sensory neurones expressing Nav1.7 (PN1) Na⁺ channel alpha subunit protein, *J. Physiol. (Lond)* 546 (2003) 565-576.
- [112] J.J. Toledo-Aral, B.L. Moss, Z.J. He, A.G. Koszowski, T. Whisenand, S.R. Levinson, J.J. Wolf, I. Silos-Santiago, S. Halegoua, G. Mandel, Identification of PN1, a predominant voltage-dependent sodium channel expressed principally in peripheral neurons, *Proc. Natl. Acad. Sci. USA* 94 (1997) 1527-1532.
- [113] A. Koschak, Impact of gating modulation in Cav1.3 L-type calcium channels, *Channels (Austin)* 4 (2010) 523--525.
- [114] D.J. Surmeier, J.N. Guzman, J. Sanchez, P.T. Schumacker, Physiological phenotype and vulnerability in Parkinson's disease, *Cold Spring Harb. Perspect Med.* 2 (2012) a009290
- [115] G.J. Augustine, T. Santamaria, K. Tanaka, Local calcium signaling in neurons, *Neuron* 40(2) (2003) 331-346.
- [116] C.S. Chan, T.S. Gertler, D.J. Surmeier, A molecular basis for the increased vulnerability of substantia nigra dopamine neurons in aging and Parkinson's disease, *Mov. Disord.* 25 (2010) S63-S70.
- [117] M. Prakriya, C.J. Lingle, BK channel activation by brief depolarizations requires Ca²⁺ influx through L- and Q-type Ca²⁺ channels in rat chromaffin cells, *J. Neurophysiol.* 81 (1999) 2267-2278.
- [118] R.S. Scott, D. Bustillo, L.A. Olivos-ore, I. Cuchillo-Ibanez, M.V. Barahona, E. Carbone, A.R. Artalejo, Contribution of BK channels to action potential repolarisation at minimal cytosolic Ca²⁺ concentration in chromaffin cells, *Pflugers Arch* 462 (2011) 545-557.
- [119] M. Sausbier, H. Hu, C. Arntz, S. Feil, S. Kamm, H. Adelsberger, U. Sausbier, C.A. Sailer, R. Feil, F. Hoffmann, M. Korth, M.J. Shipston, H.G. Knaus, D.P. Wolfer, C.M. Pedroarena, J.F. Storm, P. Ruth, Cerebellar ataxia and Purkinje cell dysfunction caused by Ca²⁺ activated K⁺ channel deficiency, *Proc. Natl. Acad. Sci. USA* 101 (2004) 9474-9478.
- [120] N. Gu, K. Vervaeke, J.F. Storm, BK potassium channels facilitate high-frequency firing and cause spike frequency adaptation in rat CA1 hippocampal pyramidal cells, *J. Physiol. (Lond)* 580 (2007) 859-882
- [121] A. Marty, E. Neher, Potassium channels in cultured bovine adrenal chromaffin cells, *J. Physiol. (Lond)* 367 (1985) 117-141
- [122] A. Neely, C.J. Lingle, Two components of calcium-activated potassium current in rat adrenal chromaffin cells, *J. Physiol.* 453 (1992) 97-131
- [123] J. Engel, H.A. Schultens, D. Schild, Small conductance potassium channels cause an activity-dependent spike frequency adaptation and make the transfer function of neurons logarithmic, *Biophys. J.* 76 (1999) 1310-1319
- [124] J. Klingauf, E. Neher, Modeling buffered Ca²⁺ diffusion near the membrane: implications for secretion in neuroendocrine cells, *Biophys. J.* 72 (1997) 674-690.
- [125] G. Ulate, R.S. Scott, J. González, J.A. Gilabert, A.R. Artalejo, Extracellular ATP regulates

- exocytosis by inhibiting multiple Ca^{2+} channel types in bovine chromaffin cells, *Pflügers Arch.* 439 (2000) 304-314.
- [126] S.A. Chan, L. Polo-Parada, C. Smith, Action potential stimulation reveals an increased role for P/Q-calcium channel-dependent exocytosis in mouse adrenal tissue slices, *Arch. Bioch. Bioph.* 435 (2005) 65–73.
- [127] A. Giacippoli, M. Novara, A. de Luca, P. Baldelli, A. Marcantoni, E. Carbone, V. Carabelli, Low-threshold exocytosis induced by cAMP-recruited Cav3.2 ($\alpha_1\text{H}$) channels in rat chromaffin cells, *Biophys. J.* 90 (2006) 1830-1841.
- [128] L. Polo-Parada, S.A. Chan, C. Smith, An activity dependent increased role for L-type calcium channels in exocytosis regulated by adrenergic signaling in chromaffin cells, *Neuroscience* 143 (2006) 445-459.
- [129] M. Aldea, K. Jun, H.S. Shin, E. Andrés-Mateos, L.M. Solís-Garrido, C. Montiel, A.G. García, A. Albillos, A perforated patch-clamp study of calcium currents and exocytosis in chromaffin cells of wild-type and $\alpha_1\text{A}$ knockout mice, *J. Neurochem.* 81 (2002) 911-921.
- [130] M. Novara, P. Baldelli, D. Cavallari, V. Carabelli, A. Giacippoli, E. Carbone, Exposure to cAMP and β -adrenergic stimulation recruits Ca_v3 T-type channels in rat chromaffin cells through Epac cAMP-receptor proteins, *J. Physiol. (Lond)* 558 (2004) 433-449.
- [131] V. Carabelli, A. Marcantoni, V. Comunanza, A. de Luca, J. Díaz, R. Borges, E. Carbone, Chronic hypoxia up-regulates $\alpha_1\text{H}$ T-type channels and low threshold catecholamine secretion in rat chromaffin cells, *J. Physiol. (Lond)* 584 (2007) 149–165.
- [132] A.G. García, F. Sala, J.A.Reig, S. Viniegra, J. Frías, R. Fontériz, L. Gandía, Dihydropyridine BAY-K-8644 activates chromaffin cell calcium channels, *Nature* 309 (1984) 69-71.
- [133] M.G. López, A. Albillos, M.T. de la Fuente, R. Borges, L. Gandía, E. Carbone, A.G. García, A.R. Artalejo, Localized L-type calcium channels control exocytosis in cat chromaffin cells, *Pflügers Arch.* 427 (1994) 348–354.
- [134] K.L. Engisch, M.C. Nowycky, Compensatory and excess retrieval: two types of endocytosis following single step depolarizations in bovine adrenal chromaffin cells, *J. Physiol. (Lond)* 506 (1998) 591-608.
- [135] S.A. Chan, C. Smith, Physiological stimuli evoke two forms of endocytosis in bovine chromaffin cells, *J. Physiol. (Lond)* 537 (2001) 871-885.
- [136] P.G. Nucifora, A.P. Fox, Barium triggers rapid endocytosis in calf adrenal chromaffin cells, *J. Physiol. (Lond)*. 508 (1998) 483-494.
- [137] C.R. Artalejo, J.R. Henley, M.A. McNiven, H.C. Palfrey, Rapid endocytosis coupled to exocytosis in adrenal chromaffin cells involves Ca^{2+} , GTP and dynamin but not clathrin, *Proc. Natl. Acad. Sci. USA* 92 (1995) 8328-8332.
- [138] F. Darios, C. Wasser, A. Shakirzyanova, A. Giniatullin, K. Goodman, J.L. Munoz-Bravo, J. Raingo, J. Jorgacevski, M. Kreft, R. Zorec, J.M. Rosa, L. Gandia, L.M. Gutiérrez, T. Binz, R. Giniatullin, E.T. Kavalali, B. Davletov, Sphingosine facilitates SNARE complex assembly and activates synaptic vesicle exocytosis, *Neuron* 62 (2009) 683-694.
- [139] R.D. Burgoyne, Fast exocytosis and endocytosis triggered by depolarisation in single adrenal chromaffin cells before rapid Ca^{2+} current run-down, *Pflügers Arch.* 430 (1995) 213-219.
- [140] P.P. Perissinotti, B. Giugovaz Tropper, O.D. Uchitel, L-type calcium channels are involved in fast endocytosis at the mouse neuromuscular junction, *Eur. J. Neurosci.* 27 (2008) 1333-1344.
- [141] M.W. Wu, A. Llobet, L. Lagnado, Loose coupling between calcium channels and sites of exocytosis in chromaffin cells, *J. Physiol. (Lond)* 587 (2009) 5377–5391.

Figure legends

Figure 1 – Ca²⁺ current modulation by cAMP activators, cGMP analogues and PKA- and PKG-blockers in mouse chromaffin cells. *a*, Inhibitory effect of the PKA blocker H-89 (5 μM) on the total Ca²⁺ currents of a WT MCC viewed by plotting the peak Ca²⁺ current amplitude elicited by voltage steps of 20 ms to +10 mV versus time (holding potential -50 mV). The test of H-89 was preceded by an assay of the blocking potency of 3 μM nifedipine to quantify the size of L-type currents. The bars to the right show the percentage of total Ca²⁺ current inhibited by H-89 in WT MCCs (27.7%; n= 8), which is about twice the amount blocked in Cav1.3^{-/-} MCCs (12.8 %; n= 8). *b*, The potentiating action of forskolin (100 μM) on the L-type currents remaining after blocking N-, P/Q- and R-type channels with a mixture of ω-Ctx-MVIIC (10 μM), ω-Ctx-GVIA (3.2 μM) and SNX 482 (0.4 μM) in a WT MCC. The percentage of L-type current potentiation induced in WT MCCs (13.4%; n= 9) is almost twice of that induced in Cav1.3^{-/-} MCCs (7.2 %; n=8). *c*, Inhibitory effect of the cGMP membrane permeable analogue 8-pCPT-cGMP (1 mM). The action of 8-pCPT-cGMP was selective for LTCCs and the percentage of current inhibition was 36.5% for WT MCCs (n= 4) and 19.2% for Cav1.3^{-/-} MCCs (n=6). *d*, The potentiating effects of the PKG-blocker KT 5823 (1 μM) on the Ca²⁺ currents of a WT MCC. On average the percentage of potentiation was 20.9% for WT MMCs (n= 11) and 9.7% for Cav1.3^{-/-} MCCs (n= 7). The four panels were adapted from ref. [49].

Figure 2 - The synergistic effects of cAMP/PKA and cGMP/PKG pathways in WT MCCs. To the left is shown a schematic representation of Cav1 channel α1 subunit with two PKA and PKG phosphorylation sites (P) during *basal conditions*, *synergistic potentiation* and *synergistic inhibition*. Basally, both PKA and PKG P-sites are partially phosphorylated and unphosphorylated. Phosphorylation and dephosphorylation driven by up- and down-regulation of PKA and PKG proceed independently of each other to reach two extreme conditions in which either two PKA P-sites are dephosphorylated and two PKG P-sites are phosphorylated (minimal Cav1 current; top right) or the PKA and PKG P-sites are oppositely phosphorylated-dephosphorylated (maximal Cav1 current; bottom right). The location and the number of P-sites at the intracellular side is arbitrary. The two panels to the right show the time course of synergistic down-regulation of LTCCs induced by the sequential application of H-89 and 8-pCPT-cGMP (top) and the synergistic up-regulation of LTCCs by sequential application of forskolin and KT 5823 (bottom). In both cases the WT MCCs were pretreated with a toxin mixture containing ω-Ctx-MVIIC, ω-Ctx-GVIA and SNX 482 to block N-, P/Q- and R-type channels (see Fig. 1b). The drawing and the experimental data are adapted from ref. [49].

Figure 3 - Contribution of Cav1 and Nav1 channels to MCCs action potentials. *Top*, representative AP of a WT MCC. The inset represents a phase plane plot (dV/dt vs V) of the respective AP waveform. *Bottom*, overlapping of Nav1 (black) and Cav1 currents (red) supporting MCCs pacemaking. The grey regions indicate the currents that flow in between membrane potentials of -40 and -50 mV. Notice that Nav1 currents contribute specifically to the spike upstroke while Cav1 currents support the slow potential rise of the pacemaker phase.

Figure 4 - SK channels slow down the MCC pacemaker cycle. *Top left*, representative current clamp recordings of spontaneously firing MCCs without current injection. Notice that apamin (200 nM) leads to stronger increases of firing rate in cells with low basal firing frequencies as compared to cells with elevated basal firing rates. *Top right*, close ups of the APs indicated by the asterisks (left). Again the effects of apamin on the AP shape are more evident in cells that show low basal firing rates. *Bottom*, graphical representation of the degree of correlation between the basal firing rate and the percentage increase (or decrease) on frequency (left), afterhyperpolarization (middle) and baseline (right) induced by apamin.

Figure 5 - Spike frequency adaptation of MCCs is dependent on Cav1.3-driven SK channels. *a*, representative current clamp traces of a MCC after 15 pA current injection in control (top), after block of BK channels by 1 μ M paxilline (middle) and after block of SK channels by 200 nM apamin (bottom). Notice that apamin and not paxilline reduces the naturally occurring spike frequency adaptation of WT MCCs. *b*, representative AP-clamp traces of a WT MCC. The top trace represents the AP train recorded in response to a 15 pA current stimulus. To the bottom are illustrated the SK currents (black) and the total Ca²⁺ currents (blue) that flow during the spike train. Red traces are recorded in the presence of 3 μ M nifedipine to block the L-type currents. Notice that the L-type Ca²⁺ current perfectly overlaps with the interspike interval specific outward SK current. *c* and *d*, Same as in *a* and *b* but referring to Cav1.3^{-/-} MCCs. Notice that the reduced spike frequency adaptation of Cav1.3^{-/-} MCCs can be ascribed to the inability of these cells to build-up sizeable SK currents (adapted from ref. [31]).





Fig. 1






Fig. 2




Fig. 3







Fig. 4




Fig. 5