

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effect of buffer strips and soil texture on runoff losses of flufenacet and isoxaflutole from maize fields**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/138971> since 2016-07-05T17:41:36Z

*Published version:*

DOI:10.1080/03601234.2013.824239

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

M. Milan, A. Ferrero, M. Letey, F. De Palo, F. Vidotto  
Effect of buffer strips and soil texture on runoff losses of flufenacet and  
isoxaflutole from maize fields  
JOURNAL OF ENVIRONMENTAL SCIENCE AND HEALTH. PART B.  
PESTICIDES, FOOD CONTAMINANTS, AND AGRICULTURAL  
WASTES (2013) 48  
DOI: 10.1080/03601234.2013.824239

The definitive version is available at:

<http://www.tandfonline.com/doi/abs/10.1080/03601234.2013.824239>

1 **Effect of buffer strips and soil texture on runoff losses of flufenacet and isoxaflutole from**  
2 **maize fields**

3  
4 MARCO MILAN, ALDO FERRERO, MARILISA LETEY, FERNANDO DE PALO and  
5 FRANCESCO VIDOTTO

6  
7 *Dipartimento di Scienze Agrarie, Forestali e Alimentari (DISAFA), Sezione agronomia, Università di*  
8 *Torino, Via Leonardo da Vinci 44, 10095 Grugliasco (TO), Italy.*

9  
10 -----  
11 \*Address correspondence to Marco Milan, Dipartimento di Scienze Agrarie, Forestali e Alimentari,  
12 (DISAFA) Università di Torino, Via Leonardo da Vinci 44, 10095 Grugliasco, Torino, Italy. Tel:  
13 (+39) 0116708897; Fax: (+39) 0116708798;  
14 E-mail: marco.milan@unito.it

15

16

17 **Abstract**

18

19 The influence of buffer strips and soil texture on runoff of flufenacet and isoxaflutole was studied  
20 for two years in Northern Italy. The efficacy of buffer strips was evaluated on six plots  
21 characterized by different soil textures; two plots had Riva soil (18.6% sand, 63.1% silt, 18.3%  
22 clay) while the remaining four plots had Tetto Frati (TF) soil (37.1% sand, 57% silt, 5.9% clay).  
23 Additionally, the width of the buffer strips, constituted of spontaneous vegetation grown after crop  
24 sowing, was also compared for their ability to abate runoff waters. Chemical residues in water  
25 following runoff events were investigated, as well as their dissipation in the soil. After the first  
26 runoff events, concentrations of herbicides in water samples collected from Riva plots were as  
27 much as four times lower in waters from TF plots. On average of two growing season, the field  
28 half-life of flufenacet in the upper soil layer (5 cm) ranged between 8.1 and 12.8 days in Riva soil,  
29 8.5 and 9.3 days in TF soil. Isoxaflutole field half-life was less than 1 day. Buffer strip was very  
30 affective by the uniformity of the vegetative cover, particularly, at the beginning of the season. In  
31 TF plots, concentration differences were generally due to the presence or absence of the buffer strip,  
32 regardless of its width.

33

34 **Keywords:** Runoff, buffer strip, flufenacet, isoxaflutole, soil

35

36 **Introduction**

37

38 Diffuse water contamination by pesticides used on croplands has been reported in studies around  
39 the world. <sup>[1-3]</sup> The most frequently detected pollutants in surface and ground water are the most  
40 used in croplands and urban areas. Among the pesticides, herbicides and their metabolites are the  
41 most commonly detected substances in surface waters. In fact, the most recent monitoring campaign  
42 of the Piemonte region (Northwest Italy) conducted by the Regional Environmental Agency

43 Authority (ARPA) found that of the top 20 chemicals detected in surface water, 17 were herbicides  
44 or herbicide metabolites. <sup>[4]</sup> Similar results have been observed nationwide and worldwide. <sup>[5-8]</sup> In  
45 the last decade, protection and prevention of water resource contamination (surface and ground  
46 waters) has become a top priority of European policy as evidenced by the Directive on Sustainable  
47 Use of Pesticides (2009/128/EC) that mandates European Member States do more to reduce water  
48 pollution related to drift, runoff, and leaching of pesticides and other agricultural products.

49 Surface water contamination is mainly due to runoff from croplands and/or spray drift during  
50 pesticide application. The magnitude of the problem is highly related to several factors: rainfall  
51 intensity, pesticide characteristics, soil slope, and soil texture. Vegetative buffer strips (VBSs) are  
52 important tools to prevent runoff from entering the water stream and/or carrying away valuable  
53 sediment, organic materials, nutrients, and chemicals. In most cases, runoff events that occur shortly  
54 after herbicide application account for the largest losses. In general, intense rainfall shortly after  
55 application generally results in herbicide losses usually less than 0.5% of the amount applied, for  
56 most herbicides. <sup>[9]</sup>

57 Several field actions can be adopted to prevent diffuse pesticide pollution and/or nutrient losses via  
58 runoff and drift; however, a catchment scale approach is necessary to optimize efforts. To reduce  
59 pesticide transport via runoff, <sup>[10]</sup> in particular most mitigation efforts involve soil management and  
60 cropping practices, VBS use, retention and dispersion structures, proper pesticide use, and in some  
61 agriculture areas, attention to irrigation management. <sup>[11]</sup> Of course, each of these measures has a  
62 different impact on runoff relative to the local soil and climatic conditions.

63 In the case of VBSs, they are usually set up along streams, ponds, or lakes to prevent water  
64 pollution. VBSs have been a useful tool to reduce runoff and erosion, <sup>[10, 12, 13]</sup> and their efficacy is  
65 generally expressed as a percent reduction in pesticide concentration as compared to a non-buffered  
66 control. According to the literature, VBS effectiveness is generally above 50%. Typically, runoff  
67 volume retention (intended as infiltration) averages 45% (ranging between 0 and 100%) across  
68 different studies under both natural and simulated experimental conditions. <sup>[14]</sup>

69 Of particular interest is the Footprint Project which looked at the efficacy of buffer strips to reduce  
70 pesticide runoff. <sup>[15]</sup> Buffer strips with widths ranging from 2 m to 21 m, the median reduction  
71 observed in the pesticides considered ranged between 65% (2 m buffer strip) and 95% (21 m buffer  
72 strip). The results presented by the FOCUS group working on a dataset from European studies only,  
73 resulted in a reduced mean efficacy of 74% (pesticides in water phase) to 79% (sediment phase) for  
74 buffer strip widths ranging from 1 to 20 m. <sup>[15]</sup> In an accurate review of the mitigation measures  
75 available to prevent runoff and erosion of pesticides, Reichenberger et al. <sup>[10]</sup> found buffer strips  
76 vary in effectiveness at the farm scale (from high to very low, according to the local conditions), but  
77 they generally show a very high efficacy when adopted at the catchment scale. In some European  
78 countries, the adoption of buffer strips between agricultural lands and waterways is already  
79 mandatory (e.g. the introduction of 10m buffer strips along waterways from September 1 in  
80 Denmark), <sup>[16]</sup> or included in EU cross-compliance measures (France and Italy).

81 Complementary to these results, many factors have been shown to influence VBS effectiveness:  
82 slope, micro-topography, soil type, rainfall intensity, infiltration capacity, strip width, and irrigation  
83 volume. Pesticide characteristics (solubility, *K<sub>oc</sub>*, and persistence), as well as soil texture, organic  
84 content, and crop and tillage management also play important roles. <sup>[9, 17, 18, 19]</sup> Finally, buffer strips  
85 filtration activity can vary with the specific PPP used, the sediment amount carried by runoff water  
86 into the strip, the water retention time in the VBS, the soil infiltration rate, the uniformity of water  
87 flow through the VBS, and maintenance of the strip itself. In general, however, the greater the width  
88 of the buffer strip, the higher the runoff retention and infiltration capability, as well as the sediment  
89 transport reduction. In the case of larger buffer strips, both infiltration and dilution of runoff flow  
90 are promoted while the effect on sediment settling is less important. <sup>[20]</sup>

91 Despite the many studies that have investigated buffer strip efficacy to limit pesticide runoff, there  
92 still remains a need for more research in this field because most of the studies have been carried out  
93 on small plots. Furthermore, most studies were conducted under simulated rainfall and in a single  
94 soil condition. The aim of this study was to evaluate the efficacy of buffer strips to mitigate the

95 runoff losses of flufenacet and isoxaflutole under natural rainfall conditions and in two different soil  
96 textures. Moreover, in one of the two soils, different buffer strip widths were compared. The two  
97 herbicides in the study are commonly used in Italy to control both grasses and broad-leaved weeds  
98 in several crops. Specifically, flufenacet is an oxyacetamide herbicide effective in pre- and early  
99 post-emergence against many grasses in corn, wheat, rice, tomato, soybean, potato, and sunflower.  
100 <sup>[21]</sup> Isoxaflutole is a broad spectrum proherbicide of the isoxazole family, used in pre-emergence or  
101 pre-plant mostly in maize and sugar-cane against grass and broad-leaved weed species. <sup>[21]</sup> A  
102 complete frame of the mode of action of isoxaflutole is reported by Pallet et al. <sup>[22]</sup> Both herbicides  
103 are applied on many important crops and in different periods of the year, so both carry a high  
104 potential to contaminate water resources.

105

## 106 **Material and Methods**

107

### 108 *Experimental Site*

109 The study was conducted at the experimental station of the Dipartimento di Scienze Agrarie,  
110 Forestali e Alimentari of Università di Torino, Italy. The station is located in the Po Valley in  
111 Northwest Italy in the municipality of Carmagnola (44° 53' 08.99'' N, 7° 41' 11.33'' E; WGS84) in  
112 an area traditionally cultivated with maize.

### 113 *Experimental Design*

114 The study was carried out on six adjoining large plots cultivated with maize, each measuring 7 x  
115 150m with a 0.5% slope (Fig. 1). Measurements were taken on the same plots during the 2009 and  
116 2010 growing seasons, which are regarded as temporal replications.

117 Four plots (TF plots, from “Tetto Frati”, which is the name of the hamlet where the station is  
118 located) were characterized by silty-loam soil original to the site (37.1% sand, 57% silt, 5.9% clay)  
119 with 1.3% organic matter and a pH=8. The remaining two plots (RIVA plots) were also a silty-loam

120 soil, but it was transferred from Riva municipality twenty years before (18.6% sand, 63.1% silt,  
121 18.3% clay), with of organic matter and a pH=6,2.

122 At the downhill edges of each plot the runoff water was intercepted by a transversal drainage ditch.  
123 Each drainage ditch was connected to an independent automatic sampler. In 2010, the system was  
124 operated by flow metering devices, formed by a series of V-notch weirs fitted with magnetostrictive  
125 water level transmitters. The water level in the weirs was continuously recorded with a datalogger.  
126 On the four TF plots, a control without a buffer strip (TF TEST) was compared to three plots with 2  
127 m (TF2), 4 m (TF4), and 6 m (TF6)-wide vegetated buffer strips. The buffer strips were sowed with  
128 maize as was the rest part of the field; weeds were allowed to grow freely. Buffer strips were  
129 mowed as needed. Weeds grown in the buffer strip were representative of the common maize weeds  
130 of Northern Italian: *Echinochloa crus-galli* (L.) P. Beauv., *Panicum dichotomiflorum* Michx.,  
131 *Chenopodium album* L., *Portulaca oleracea* L., *Trifolium repens* L., *Galinsoga quadriradiata* Cav.,  
132 *Poa pratensis* L. and *Setaria viridis* L. Their density, expressed as percentage of soil coverage,  
133 ranged from 20% (a week post crop sowing) to 100% during the rest of the season. The buffer strips  
134 were mowed at least twice a year, after which the hay was removed. In the RIVA plots, a control  
135 *sans* buffer strip (RIVA TEST) was compared to a plot with a 6-m buffer strip (RIVA6).  
136 Characteristics of the buffer strip were similar to the buffers established in TF plots.

137 The plots were cultivated according to local agronomic practices, and the crop was sown on April 9  
138 and April 20 in 2009 and 2010, respectively.

139 Herbicides were applied during pre-emergence, within three/four days after sowing, using a rear-  
140 mounted boom sprayer. Over the two growing seasons water was supplied as needed to the crop by  
141 a furrow irrigation system. In 2009, two irrigations were carried out on June 16 (37 m<sup>3</sup>, average of  
142 the six plots) and August 3 (35 m<sup>3</sup>), respectively; in 2010, fields were irrigated only once on July 21  
143 (38 m<sup>3</sup>).

144

145 ***Herbicides Studied***



146

147 All plots, except the buffer strips, were treated with flufenacet (4'-fluoro-N-isopropyl-2-[5-  
148 (trifluoromethyl)-1,3,4-thiadiazol-2-yloxy]acetanilide) and isoxaflutole (5-cyclopropyl-1,2-oxazol-  
149 4-yl)( $\alpha,\alpha,\alpha$ -trifluoro-2-mesyl-p-tolyl)methanone) at 240 g a.i. ha<sup>-1</sup> and 50 g a.s. ha<sup>-1</sup>, respectively,  
150 by spray application of 500 g ha<sup>-1</sup> of the commercial herbicide Merlin GP<sup>®</sup> (Bayer CropScience  
151 Italia). To avoid deposition from spray drift, the buffer strip was covered with a plastic film during  
152 herbicide application. Table 1 shows the physical and chemical properties of the studied substances.

153

#### 154 **Soil sampling**

155

156 Three soil samples were collected from the treated areas at various times: before herbicide  
157 application (to assess the residual previous soil contamination,  $t_1$ ), immediately after spraying (to  
158 assess initial herbicide concentrations,  $t_0$ ), and at increasing intervals from herbicide application (at  
159 0, 1, 7, 15, 40 and 90 days after treatment in 2009; and at 0, 3, 7, 17, 31, 42 and 90 days after  
160 treatment in 2010). Samples were taken from the upper 5 cm of the soil surface, with a 50 mm  
161 diameter soil core sampler. At each sampling time, and for each plot, 10 soil cores samples were  
162 randomly collected in the treated areas. After collection, soil samples were stored at -20°C until  
163 analysis.

164

#### 165 ***Water sampling***

166

167 Samples of runoff water were collected after each runoff event by automatic samplers adjusted to  
168 collect a bulk sample made by 500 mL sub-samples gathered at 10-min intervals for the duration of  
169 the event. The bulk samples had volumes ranging from 0.5 L to 25 L, which correlated to runoff  
170 event duration and intensity. Within about two hours from the end of each event, up to three 1 L  
171 subsamples were derived from the bulk sample and stored at -20°C until analysis. Water samples

172 were collected at 4, 5, 11, 12, 13, 62, 68, 77, 82, 93, 110, and 115 days after treatment (DAT) in  
173 2009 and at 10, 11, 17, 44, 53, 54, 57, 89, and 110 DAT in 2010.

174

### 175 ***Herbicide Extraction and Analysis in the Soil***

176

#### 177 *Flufenacet*

178

179 The extraction of flufenacet from the soil was performed on 50 g samples. The samples were  
180 transferred into a 250 mL glass bottle (Duran, Germany) and 100 mL of acetone was added (J.T.  
181 Baker, USA). Thereafter, the solution was sonicated for 30 min in an ultrasonic bath (Sonorex RK  
182 156BH, Germany). The sonicated solution was then passed through a 150 mm diameter Büchner  
183 funnel (Büchner, Germany) connected to a side-arm 250 mL flask (Duran, Germany) using a  
184 neoprene adapter with a tube to a vacuum pump (Supelco). Two filter paper disks (Perfecte 2<sup>®</sup>,  
185 Cartiera di Cordenons, Italy) were placed on the Büchner surface, then covered with a layer of celite  
186 (Celite 545 J.T. Baker). The soil and the glass bottle used for the extraction were washed with 50  
187 mL of acetone and 50 mL of deionized water and the resulting volume was passed through the  
188 Büchner funnel. All the filtrate obtained was transferred into a 250 mL volumetric flask and  
189 adjusted to volume with deionized water. A final volume of 100 mL was then transferred to a 500  
190 mL volumetric flask and adjusted to volume with deionized water. Herbicide extraction from this  
191 solution was carried out using solid phase extraction (SPE) cartridges. The cartridges (SupelcoSil  
192 LC-18, 6 mL, 0.5 g C18 sorbent material) were previously activated with 6 mL of n-hexane (J.T.  
193 Baker, USA) and 6 mL of methanol (J.T. Baker, USA), and finally washed with 6 mL of deionized  
194 water. The entire volume (0.5 L) flowed through the cartridges under vacuum at a rate of 500 mL h<sup>-1</sup>.  
195 <sup>1</sup>. The cartridges were let to dry. The adsorbed herbicide was eluted with n-hexane until a final  
196 volume of 5 mL was reached. The eluted volume of 5 mL was then filtered through a 0.2 µm nylon  
197 filter (Whatman, USA) to eliminate impurities. **The final volume was dried under nitrogen flow by**

198 a nitrogen generator (Claind, Italy) and recovered with 1 mL of n-hexane (Sigma Aldrich,  
199 Steinheim, Germany). Analysis was performed by GC-MS as described in the next paragraphs.

200

201 *Isoxaflutole*

202

203 The extraction of isoxaflutole from the soil was performed on 50 g samples. The samples were  
204 transferred into a 500 mL polyethylene bottle and 150 mL of a solution (80/20 v/v) of acetonitrile  
205 (Sigma Aldrich, Steinheim, Germany) and formic acid (0.8%) (Sigma Aldrich, Steinheim,  
206 Germany) was added. Thereafter, the solution was sonicated for 30 min in an ultrasonic bath  
207 (Sonorex RK 156BH, Germany). The sonicated solution was then transferred in a 250 mL vacuum  
208 flask through a funnel with its hole covered with a cotton stopper. The filtrates obtained were  
209 concentrated and dried in a rotary evaporator, then re-dissolved with 10 mL of a solution of  
210 acetonitrile and not-brought water (50/50 v/v). An aliquot of the eluted volume was transferred in  
211 the vials for the analysis. Analysis was performed by HPLC as indicated in the next paragraphs.

212

213 ***Herbicide Extraction and Analysis in the Water***

214

215 Herbicides extraction from the water samples was carried out using solid phase extraction (SPE)  
216 cartridges. The cartridges (SupelcoSil LC-18, 6 mL, 0.5 g C18 sorbent material) were previously  
217 activated with 6 mL of acetonitrile (Sigma Aldrich, Steinheim, Germany) and then washed with  
218 20 mL of distilled water. The entire volume (1 L) of the water sample flowed through the cartridges  
219 under vacuum at a flow of 500 mL h<sup>-1</sup>. The cartridges were let to dry. The adsorbed herbicides were  
220 eluted with acetonitrile until a final volume of 5 mL was reached. The eluted volume of 5 mL was  
221 then filtered through a 0.20 µm nylon filter (Whatman, USA) to eliminate impurities. The final  
222 volume was dried under nitrogen flow by a nitrogen generator (Claind, Italy) and recovered with 1  
223 mL of n-hexane (Sigma Aldrich, Steinheim, Germany). Analysis was performed by GC-MS.

224 ***High-performance liquid chromatography analysis***

225

226 Analysis of isoxaflutole in soil was done by high-performance liquid chromatography (HPLC)  
227 using a Spectraphysics P2000 equipped with a C18 Varian Pursuit column (150 mm × 4.6 mm i.d.,  
228 5µm particle size), an ultra-violet (UV) detector at 270 nm for isoxaflutole, and a mobile phase  
229 composed by brought water with pH=2, and acetonitrile (50/50 v/v) with the flow rate set to 1 mL  
230 min<sup>-1</sup>. Analytical-grade isoxaflutole supplied by Sigma Aldrich, Germany was used as the  
231 analytical standard.

232

233 ***Gas Chromatography Analysis***

234

235 An Agilent 6890N GC and Agilent 5975 MS single-quadrupole, equipped with an MS detector, an  
236 autosampler (Agilent) and split-splitless injector, connected to an Agilent Chemstation was used.  
237 The Supelco Equity5 TM column (30 m x 0.25 mm i.d.) contained 5% diphenyl and 95% dimethyl  
238 siloxane. The MS source temperature was 270°C and the gas carrier was helium. Analytical-grade  
239 flufenacet and isoxaflutole, supplied by Sigma Aldrich, Germany, was used as the analytical  
240 standards. Retention times for flufenacet in the soil samples were 17.6 min and 16.3 min in 2009  
241 and 2010, respectively. Retention times for flufenacet in the water were 22 min and 25.6 min in  
242 2009 and 2010, respectively. Retention times for isoxaflutole in the water were 26.1 min and 17.7  
243 min in 2009 and 2010, respectively. Retention times for isoxaflutole in the soil samples were 9.2  
244 min and 9.8 min in 2009 and 2010, respectively.

245

246 ***Recovery and detection limits***

247

248 The recovery tests for the extraction of the herbicides in the water were conducted both with tap  
249 water and surface water not contaminated. Three samples (500 mL) of not treated water were

250 contaminated with 1 mL of a stock solution 100 mg/L of flufenacet and isoxaflutole. The initial  
251 concentration was 0.2 mg/L. Extraction was carried out using SPE cartridges previously activated  
252 with 6mL of acetonitrile and then washed with 20 mL of distilled water. The same procedure was  
253 repeated contaminating other water samples with 1 mL of a stock solution 10 mg/L and 1 mL of a  
254 stock solution 1 mg/L of flufenacet and isoxaflutole.

255 The recovery tests for the extraction of the herbicides in the soil were conducted with two accession  
256 of the soil used in the experiment not contaminated. For each soil, three samples of soil (50 g) were  
257 contaminated with 100 µL of the stock solution 100 mg/L, with an initial concentration of 0.2 mg/L.  
258 The same procedure was carried out for flufenacet and isoxaflutole, but using the methods of  
259 extraction for each herbicide indicated in the previous paragraphs.

260 The mean recoveries of flufenacet and isoxaflutole in water were 98% and 87%, respectively; those  
261 in soil were 70% and 82 % for flufenacet and isoxaflutole, respectively. The limit of quantifications  
262 of the instrument (LOQ<sub>i</sub>) achieved in the water samples were 0.08 µg L<sup>-1</sup> for flufenacet and 0.1 µg  
263 L<sup>-1</sup> isoxaflutole while the limit of quantification of the method (LOQ<sub>m</sub>) were 5 µg kg<sup>-1</sup> for both for  
264 flufenacet and isoxaflutole.

265

### 266 *Statistical Analysis*

267

268 A statistical analysis was employed to determine the significance of differences among the  
269 concentrations observed in the waters collected from the check and buffered fields at the different  
270 sampling times. The values presented are the means of the three data values. SPSS, version 17.00,  
271 (SPSS, IBM Corporation, 2008), was used for the statistical analysis. A Ryan-Eynot-Gabriel-  
272 Welsch-F test (\*P <0.05) was employed to determine the statistical significance of differences  
273 among the concentrations observed in the waters collected from the check field and the buffered  
274 field at the different sampling time. Soil data were subjected to ANOVA to test effect of the year,

275 soil, days elapsed from the treatment (DAT) and the interaction between them. Flufenacet  
276 dissipation in the soil was fitted to a 2-parameter exponential decay model:

$$277 \quad C_t = C_0 e^{-kt} \quad [1]$$

278 where  $C_t$  is the concentration at time  $t$ ,  $C_0$  is the initial concentration,  $t$  is time, and  $k$  is the rate  
279 constant. Soil half-lives ( $T_{1/2}$ ) values for flufenacet were calculated from the following equation:

$$280 \quad T_{1/2} = \ln(2/k) \quad [2]$$

281 where  $k$  is the rate constant.

282 Model fitting was performed using the function *drm* of the add-on package *drc* of the R software.

283 Data from 2009 and 2010 were first analyzed separately and then pooled to fit into a single model.

284 The *anova* function of the R software was used to test if the pooled dataset was significantly better  
285 explained by single curves data separately (fitting both years and both soils) than by a single model  
286 fitting all data.

287

### 288 ***Rainfall Distribution and Runoff Events***

289

290 Total rainfall measured during the crop growing seasons were 577 mm and 545 mm in 2009 and  
291 2010, respectively. Weather data were collected daily from the meteorological station located near  
292 the experimental fields. In both years, the periods close to herbicide application were characterized  
293 by rainfall events that directly affected runoff losses and herbicide dissipation. The spring of 2009  
294 was particularly rainy; in fact, a 282 mm rainfall was recorded in April. In the 2009 season, 13  
295 runoff events were recorded while in 2010, 9 events were recorded.

296

## 297 **Result and discussion**

298

### 299 ***Herbicide Dissipation in the Soil***

300 Flufenacet and isoxaflutole dissipation was studied in the soil of the treated areas. Persistence of  
301 both herbicides in the soil varied slightly between years, according to the different climatic  
302 conditions.

303 Flufenacet showed a rapid decay in both seasons (Figs. 2a and 2b). In 2009 the field dissipation  
304 half-life ( $DT_{50}$ ) was 8.1 days in RIVA soil and 9.3 days in TF. During 2010 the persistence of  
305 flufenacet in the upper soil layers did not change significantly relative to the previous year, resulting  
306 in a soil half-life of 12.8 days in RIVA and 8.5 days on TF soil. In soil samples collected before the  
307 2010 herbicide application, residues of flufenacet were below the detection limit. Also, the different  
308 rainfall pattern recorded in the two years during the first days after herbicide application did not  
309 significantly influence the flufenacet dissipation trend (Figs. 2a and 2b).

310 The statistical analysis conducted did not show a significant effect of the year and of the type of soil  
311 on the dissipation dynamics of flufenacet. The only significant factor was the time elapsed from the  
312 treatment (DAT). The data of the two years pooled in a single model revealed a  $DT_{50}$  of 10.2 days  
313 with a confidence interval ( $\alpha=0.05$ ) comprised between 6.4 to 14 days. As reported in the literature,  
314 flufenacet dissipation follows a first order kinetics. <sup>[23]</sup> In both years, three months after herbicide  
315 application on TF soil, flufenacet was below the limit of quantification. The only exception was in  
316 2009 in RIVA soil when 90 days after treatment the concentration of the herbicide was still in the  
317 detectable range, but no higher than  $29 \mu\text{g kg}^{-1}$ .

318 The rapid flufenacet field dissipation can be attributed partly to the sampling procedure adopted, in  
319 which only the superficial soil layer was sampled. This result agrees with Rouchaud et al., <sup>[24]</sup> who  
320 found no flufenacet residues after the wheat harvest in summer and after the corn harvest during the  
321 fall in the 0-20 cm soil layer with a similar LOQ. In top soil, the dissipation dynamics are generally  
322 faster compared to that of deeper soil layers. Furthermore, the microbial degradation which is the  
323 principal means of dissipation of flufenacet in soil must be considered. Since microbial activity is  
324 enhanced during the spring, a shorter half-life could be expected at that time. Soil half-lives for  
325 aerobic microbial degradation have ranged from 10 to 34 days in varying soil types at

326 approximately 1.0 ppm at 20-21°C. <sup>[25]</sup> In a study conducted by Rouchaud et al., <sup>[26]</sup> the half-life of  
327 flufenacet in different soil ranged between 66±3.9 days and 44±2.2 days. However, their study was  
328 conducted on soils characterized by a history of organic fertilization and thus with a highest organic  
329 matter content. <sup>[26]</sup> Persistence was also affected by the time of herbicide application, with high  
330 persistence after fall applications. According to Gupta and Gajbhiye <sup>[23]</sup> flufenacet half-life ranged  
331 from 10.1 to 30.1 days on three different Indian soils. Dissipation studies conducted in Italy and  
332 France have reported soil half-lives of 13-16 days when applied during early spring and of 15-53  
333 days during spring application. Autumn applications are generally characterized by longer  
334 persistence periods. <sup>[27, 28]</sup> Soil moisture content and pH affected flufenacet dissipation less.  
335 Conversely, the type of soil, its adsorption capacity, and the rate of application can have a  
336 significant effect on dissipation behavior. Gupta and Gajbhiye <sup>[23]</sup> observed that dissipation of  
337 flufenacet is slower in soil with high adsorption capacity and less desorption.

338 The isoxaflutole soil half-life observed over the two years in the treated areas of the two soils  
339 studied was short, less than 1 day, and soil dissipation resulted faster in TF soils compared to Riva  
340 soil. In general isoxaflutole dissipation follows a first order kinetic. <sup>[29]</sup> The field dissipation half-  
341 lives in this study are similar to those reported by other studies conducted worldwide: from 0.5 to 4  
342 days, <sup>[27]</sup> from 1.4 to 3 days, <sup>[30]</sup> and from 0.5 to 2.4 days. <sup>[28]</sup> Other documents indicate the field  
343 half-life was less than 2 days <sup>[29]</sup> while Papiernik et al., <sup>[31]</sup> reported a soil half-life for the sum of  
344 isoxaflutole+diketonitrile as within 8 to 14 days in the top 1 m of three different soils. Our result  
345 agreed with other studies <sup>[32]</sup> and is explained by the abiotically-governed transformation of  
346 isoxaflutole into the active form diketonitrile, which is the key step in the dissipation pathway of the  
347 herbicide. <sup>[33]</sup>

348 In this study, the observed rapid dissipation can be partially attributed to the sampling procedure;  
349 only the superficial (5 cm) soil layer was sampled.

350 Furthermore, during 2009, a cumulative amount of 46.2 mm of rain was recorded during the week  
351 preceding herbicide application (Fig. 2a). The higher water content of the soil observed just before



352 the treatment had likely facilitated the conversion of isoxaflutole into proherbicide diketonitrile. In  
353 2010, the soil surface was very dry at spraying due the absence of rainfall in the previous weeks.  
354 According to Taylor et al.,<sup>[34]</sup> under dry conditions, isoxaflutole is very stable and unavailable, and  
355 it persists more at the surface.<sup>[35]</sup> However, a succeeding rainfall might promote the rapid  
356 transformation of isoxaflutole into its active form. Indeed, this condition was verified during 2010,  
357 when the soil was dry before treatment, but just 6 hours after treatment, a light rain occurred and  
358 caused transformation of the parent compound. Pallet et al.<sup>[35]</sup> observed that the shorter half-life  
359 recorded for isoxaflutole under increased moisture content might relate to the need of isoxaflutole to  
360 be in solution in order to be transformed into diketonitrile.

361 Conversion of isoxaflutole to diketonitrile is rapid and become faster with higher temperatures,  
362 higher soil moisture levels, and at basic pH.<sup>[33]</sup> Since the higher clay content of RIVA soil did not  
363 affect the sorption of the molecule,<sup>[36]</sup> the diverse and low persistence might instead be associated  
364 with the differing pH of the two soils. Mitra et al.<sup>[36]</sup> and Rouchaud et al.<sup>[37]</sup> observed faster  
365 dissipation of isoxaflutole at basic pHs. In the present study, TF soil has a sub-alkaline reaction (pH  
366 8.2) while the pH of RIVA soil is sub-acid (6.2) (Table 1). Hence, the slightly faster dissipation  
367 observed in TF soil over the two years might result from the combined effects of soil reaction and  
368 different soil moisture at the time of herbicide application.

369 As previously indicated, the two soils studied had similar low organic matter content, but they  
370 differed in pH and clay content. These two parameters may affect the soil dissipation of many  
371 pesticides, including isoxaflutole. A study conducted by Mitra et al.,<sup>[36]</sup> pointed out that sorption of  
372 isoxaflutole was not influenced by clay content.<sup>[38]</sup> On the contrary, it was highly related to organic  
373 matter content<sup>[39]</sup> and to the soil pH.<sup>[39, 40]</sup> Specifically, sorption of isoxaflutole increases with  
374 increasing organic matter content, and sorption increases at decreasing pH.<sup>[36]</sup> Beltrán et al.<sup>[41]</sup> has  
375 discussed the influence of soil pH on the dissipation reaction rate of isoxaflutole, and they found  
376 that the isomerization of IFT into DKN is rapid, depends strongly on pH, and is governed by a  
377 chemical process.

378 *Dissolved Flufenacet in Runoff Water*

379 The runoff of flufenacet was studied during the growing seasons in relation to rainfall and irrigation  
380 occurrences. In Table 2 are listed all the runoff events that occurred and the subsequent  
381 concentrations of flufenacet detected. The different rainfall distributions affected the transport of  
382 flufenacet and isoxaflutole, and thus runoff losses. In particular, several rainfall events occurred  
383 early after herbicide application causing relevant runoff outflows.

384 In both years, the highest flufenacet concentrations in runoff waters were measured during the  
385 runoff events that occurred in the first two weeks after application. This is consistent with several  
386 studies <sup>[18,42]</sup> that showed that major losses occur during runoff events close to herbicide application.  
387 [2, 9, 43-47]

388 Flufenacet was found in all samples collected with significant differences in the plots containing  
389 buffer strips and related to soil texture differences. The first runoff event occurred only four days  
390 after herbicide application. As shown in Table 2, during 2009 the presence of the buffer strip did  
391 not affect the amount of flufenacet transported much. The highest concentrations detected in TF  
392 plots ranged between 9.2 µg L<sup>-1</sup> (TF2) and 14.9 µg L<sup>-1</sup> (TF4) with no significant difference among  
393 plots. These concentrations were four times lower than those found in the RIVA runoff waters  
394 (Table 2). In spite of this, the presence of the buffer did not significantly reduce the amount of  
395 flufenacet transported. The lack of a clear effect of the buffer to reduce the transported dissolved  
396 flufenacet was likely due to the low weed coverage (around 20%) at that time.

397 Thereafter, a continuous rainfall occurred between 11<sup>th</sup> and 13<sup>th</sup> day after treatment, which resulted  
398 in a runoff event that endured during the entire rain period. At the beginning of this event, the  
399 flufenacet concentrations were lower than in the first event, but by the end of it (at 13 DAT), a  
400 concentration increase was observed, particularly in runoff waters that flowed from RIVA plots  
401 (Table 2). One explanation for this observation might be that after three days of rain, all soil  
402 macropores were saturated by water, and that runoff overcame the infiltration rate. Then, in the  
403 second week of June (at 62 DAT), fields were watered by furrow irrigation. The flufenacet

404 concentrations in runoff waters from the RIVA fields ranged between  $15 \mu\text{g L}^{-1}$  (RIVA TEST) and  
405  $5.9 \mu\text{g L}^{-1}$  (RIVA6) while runoff water flows from TF plots was no higher than  $0.5 \mu\text{g L}^{-1}$  (Table 2).  
406 Beginning with 68 DAT, the presence of a buffer strip generally resulted in reduced losses of  
407 flufenacet via runoff from TF plots, except for runoff events at 77 DAT (heavy thunderstorm) and  
408 110 DAT (second irrigation). On the other hand, flufenacet was always detected in runoff waters  
409 that flowed from RIVA TEST and RIVA6; those detected in RIVA6 were always lower than in the  
410 test plot. Four months after herbicide application, residues of flufenacet were found only in runoff  
411 waters from TF TEST ( $0.2 \mu\text{g L}^{-1}$ ), RIVA TEST ( $0.3 \mu\text{g L}^{-1}$ ), and RIVA6 ( $0.2 \mu\text{g L}^{-1}$ ).  
412 In 2010, the first runoff event occurred at 11 DAT. As shown in Table 2, the highest concentrations  
413 were observed in runoff waters from TF TEST and RIVA TEST. These concentrations were  
414 remarkably lower than those observed at the first runoff event in the previous year. This difference  
415 is probably due to higher weed coverage of the buffer (about 60%) compared to that of 2009  
416 (indicate the percentage here for comparison). In addition, a rainfall of 10.2 mm occurred just 12  
417 hours after herbicide application, caused no field runoff, but favored the chemical movement  
418 through the soil profile. Two weeks after spraying, flufenacet residues in runoff waters from RIVA  
419 plots ranged between  $2.7 \mu\text{g L}^{-1}$  (RIVA TEST) and  $1.6 \mu\text{g L}^{-1}$  (RIVA6) as opposed to values for  
420 runoff waters from the TF plots between  $0.6 \mu\text{g L}^{-1}$  and  $0.9 \mu\text{g L}^{-1}$  (Table 2).  
421 Flufenacet concentrations in runoff water decreased gradually over the next runoff events and at 53  
422 DAT, no flufenacet residues were found in the waters that flowed from TF plots. The herbicide was  
423 still present in runoff waters from RIVA plots up to a month after spraying, with no significant  
424 differences among plots. Next, a storm of 69.6 mm at 110 DAT caused the complete flooding of the  
425 structures where the sampling devices were located, and made it impossible to collect any runoff  
426 samples. No residue of flufenacet was found in the samples collected following a rainfall that  
427 occurred few days later (113 DAT).  
428 One way to predict the fate of pesticides in the environment is to analyze key parameters, such as  
429  $K_{oc}$ , solubility, persistence, and pH stability. As pesticides bind differently to clay particles and to

430 organic matter, studying their  $K_{oc}$  (K of organic carbon) is an effective measure of adsorption to  
431 organic matter or to soil carbon that may help to explain the behavior of a specific pesticide in a  
432 defined environmental compartment. In general, pesticides with higher  $K_{oc}$  values are more bound  
433 to the soil coefficient contrary to those with lower  $K_{oc}$ . The latter tend to be transported more with  
434 water than on sediment. <sup>[13]</sup> As indicated in the review of flufenacet performed by the European  
435 Commission, the mean  $K_{oc}$  for flufenacet is 202 for OC content > 0.23%. <sup>[27]</sup> Consistent with this  
436 information, we found flufenacet to move off fields more easily with water than when attached to  
437 sediment.

438 In this study, concentration differences measured in the water of the two soils clearly highlighted  
439 the effect of soil texture on the amount of flufenacet transported. Soil texture affects infiltration  
440 rates and runoff is generally more pronounced in fine-textured soil. <sup>[45]</sup> RIVA soil has more silt and  
441 more clay compared to TF soil (Section). Silty soils are very vulnerable to surface runoff due to the  
442 changeable behavior of their particles during seasonal changes, <sup>[10]</sup> and their tendency to develop a  
443 superficial crust. Crusting and compaction influence the infiltration rate, favor runoff, and increase  
444 the initial concentrations of pesticides. <sup>[45]</sup> The higher concentrations recorded throughout the  
445 season in runoff waters from RIVA plots might be related to these considerations.

446

#### 447 ***Dissolved Isoxaflutole in Runoff Water***

448

449 The presence of isoxaflutole in runoff waters was assessed in the same temporal interval as that of  
450 flufenacet. As for flufenacet, the highest isoxaflutole concentrations were detected during the first  
451 runoff event in waters from the plots without buffer. In 2009, at the first event (4 DAT), the highest  
452 isoxaflutole concentrations were detected in RIVA plots, where they ranged between  $5 \mu\text{g L}^{-1}$   
453 (RIVA TEST) and  $2.90 \mu\text{g L}^{-1}$  (RIVA6). In TF plot runoff, they did not exceed  $0.16 \mu\text{g L}^{-1}$  (TF  
454 TEST). In the runoff events that occurred later in the season, isoxaflutole was present in runoff  
455 waters from TF4 only at 6 DAT ( $0.13 \mu\text{g L}^{-1}$ ) and TF6 waters were always below the LOQ. In water

456 samples collected from TF2 and TF TEST, isoxaflutole was generally below the LOQ with the  
457 exception of some samples (Table 3). In contrast, during all of 2010, isoxaflutole was found only in  
458 runoff waters from RIVA TEST ( $0.10 \mu\text{g L}^{-1}$ ) collected at the first runoff event (10 DAT).

459 As isoxaflutole is rapidly converted into diketonitrile, its presence in the dissolved phase of runoff  
460 waters appears to be unlikely. The frequency of the detection of isoxaflutole, atrazine, and their  
461 respective metabolites in 10 Iowa rivers that drain important croplands, Meyer et al. <sup>[48]</sup> found  
462 isoxaflutole in only 4 samples out of 75 collected, and only in the period post planting.  
463 Furthermore, the study found diketonitrile and benzoic acid (both isoxaflutole metabolites) in 56  
464 and 43 samples out of 75 collected, respectively which confirmed rapid transformation of the parent  
465 compound. <sup>[48]</sup> Our results showed that if a runoff event occurs in the first weeks after herbicide  
466 application, significant amounts of this herbicide can be transported via runoff waters, despite its  
467 low water solubility ( $6.2 \text{ mg L}^{-1}$ ). <sup>[49]</sup> The differences in the concentrations of isoxaflutole in runoff  
468 waters, observed over the two years, are likely to be related to the different rainfall pattern occurred.  
469 As discussed in the Section 3.1, under dry conditions, isoxaflutole is very stable and unavailable,  
470 and it persists more at the surface. <sup>[34, 35]</sup> Thus, in 2009 the driest condition of the soil had probably  
471 delayed the conversion of isoxaflutole into diketonitrile. In 2010, the soil was dry too at the time of  
472 spraying, but a rainfall occurred few hours later without causing runoff, promoted the conversion of  
473 the herbicide in the metabolite. However, a succeeding rainfall might promote the rapid  
474 transformation of isoxaflutole into its active form. In synthesis, isoxaflutole, due to its low  
475 application rate and likely for its rapid conversion to metabolite, was always found at lower  
476 concentrations and mostly in runoff waters collected at early runoff events after application. Our  
477 results evidenced also that soil texture did greatly affect the amount of isoxaflutole transported by  
478 water.

479

480 *Efficiency of the Vegetative Buffer Strip*

481

482 In general the runoff mitigation effect of a buffer strip decreases as the ratio between the field area  
483 and the buffer area increases. <sup>[50]</sup> Larger buffer strips mitigate sediment transport; for more soluble  
484 pesticides, the effect might be limited. Also, buffer strips are most effective against nutrients and  
485 pesticides bound to sediments and less effectiveness on predominantly-dissolved chemicals. <sup>[51, 52]</sup>  
486 Among the various conditions of our study, we found that with a 6 m buffer strip, the most  
487 favorable ratio (25:1) we could attain was with isoxaflutole on RIVA soil.

488 The Table 4 reports the runoff events and the corresponding measured runoff volumes during 2010.  
489 The higher runoff volumes were generally observed in plots lacking buffer strips, which  
490 demonstrated the positive effect of the buffer. This was particularly true for RIVA plots; it was less  
491 evident in TF plots. In these plots, runoff volumes measured in buffered plots were, at times, higher  
492 than in the control plot. If operating on a field scale, then some modification must be made for the  
493 soil unevenness that characterizes large plots. Weed spots might also have affected the runoff flow  
494 behavior through the field. Overall, the maximum runoff flows were always measured on RIVA  
495 plots, which indicated the high bent of that soil type to surface runoff.

496 Buffer strip efficacy was evaluated in absolute terms by considering both the observed  
497 concentrations in water samples, and by calculating the total losses in relation to the runoff volumes  
498 recorded during 2010. Total losses were calculated for flufenacet only, as isoxaflutole was always  
499 below the quantification limit. Flufenacet is transported both in the water phase and in the solid  
500 phase and adsorbed to particles eroded from the soil surface. For our purposes, only the amounts of  
501 herbicides dissolved in runoff waters were considered for the calculation. Total losses were  
502 calculated for each event by multiplying the volume of runoff by the mean concentration of  
503 dissolved herbicide (Runoff Volume [ $\text{m}^3$ ] x Concentration [ $\mu\text{g L}^{-1}$ ]). It was expressed as a  
504 percentage of the total amount of herbicide applied. In the case of concentrations below LOQ, and  
505 even considering actual concentrations as equal to LOQ, total losses showed only a negligible  
506 difference in the adopted calculation.

507 As expected, our results indicated that RIVA plots accounted for the highest losses in 2010; they  
508 were 0.40% in RIVA TEST and 0.23% in RIVA 6m. Alternatively, TF plot total losses were largely  
509 lower, ranging between 0.07% (TF TEST) and 0.01% (TF2 and TF6). According to Whauchope,<sup>[9]</sup>  
510 runoff losses of these magnitudes can be considered “*intermediate*.” Overall, most of the losses  
511 were due to the first runoff events both in the buffered and non-buffered plots. The total losses were  
512 not calculated during 2009; however, given the high concentrations observed in the first runoff  
513 events, it is reasonable to assume that flufenacet and isoxaflutole had larger losses during this  
514 season.

515 As indicated in the Material and Methods section, the buffer strip was not specifically sowed, but it  
516 was represented by spontaneous vegetation grown after crop sowing. In both seasons, the first  
517 runoff event occurred early after herbicide spraying. Being that the buffer strip vegetation was  
518 comprised only of spontaneous weeds, its coverage was insufficient to fully counter runoff flows,  
519 particularly during 2009. A buffer strip acts by reducing flow velocity and increasing infiltration;  
520 thereby, it reduces the total pesticides transported.<sup>[53]</sup> However, in 2009, during the first runoff  
521 events, the presence of the buffer strip in all plots did not significantly affect the amount of  
522 pesticides transported, especially for flufenacet. Presence of a buffer strip showed a certain effect  
523 late in the season (Table 2), when the vegetation cover of the buffer became more dense and  
524 uniform. This behavior was observed especially in RIVA plots. During 2010, the presence of the  
525 buffer strip significantly affected the amount of herbicide transport, both on RIVA and TF plots.

526 The effect of the different buffer strip widths in reducing runoff in TF plots can be extrapolated  
527 from Table 3. During both years, the width of the buffer seems unrelated to improved buffer  
528 performance, as detected concentrations did not differ greatly between the compared plots. Buffer  
529 strip efficiency was found to be greatly affected by the uniformity of the vegetative cover, in  
530 particular at the beginning of the season. On TF plots, differences in the concentrations were  
531 generally due the presence/absence of buffer strips, regardless of its width. Similar behavior was  
532 observed by Tingle et al.<sup>[53]</sup>

533

534 **Conclusions**

535

536 Flufenacet and isoxaflutole can be easily transported with runoff waters. In both years, the highest  
537 concentrations were found in water samples collected after the first runoff events. Flufenacet was  
538 always found in runoff waters at concentrations higher than isoxaflutole. In general, flufenacet  
539 losses were larger and extended further into the season. The study evidenced the strong effect of  
540 soil texture on the amount of flufenacet and isoxaflutole transported. Both soils were silty-loam  
541 textured, but they differed in their soil properties affecting the amount of each herbicide available  
542 for surface runoff. Flufenacet did not persist longer in the top soil surface. On TF plots,  
543 concentration differences were generally due the presence/absence of buffer strips, regardless of  
544 buffer strip width. It was also observed that buffer strip efficiency was greatly affected by the  
545 degree of development of the spontaneous vegetative cover, particularly at the beginning of the  
546 growing season. This problem could be avoided by sowing a mix of grasses (such as *Festuca* spp.  
547 and *Lolium* spp.) on the buffer strip surface early in the season to ensure better coverage.

548

549 **Acknowledgements**

550

551 This study was supported financially by the Agricultural Department of the Regione Piemonte,  
552 Italy. The authors thank the experimental station and lab technicians who made valuable  
553 contributions to field operations. Appreciation is further extended to group members of Sustainable  
554 Weed Management of the Department of Agronomy, Forest and Land Management of the  
555 University of Turin, who helped during the study. The paper is attributed equally to the authors.

556

557 **References**

558



- 559 [1] Parris, K. Impact of agriculture on water pollution in OECD countries: Recent trends and future  
560 prospects. *Intern. J. Water Resour. Dev.* **2011**, *27* (1), 33-52.
- 561 [2] Riise, G.; Lundekvam, H.; Wu, Q.L.; Haugen, L.E.; Mulder, J. Loss of pesticides from  
562 agricultural fields in SE Norway - Runoff through surface and drainage water. *Envir. Geochem.*  
563 *Health* **2004**, *26* (2), 269-276.
- 564 [3] Legrand, M.P.; Costentin, E.; Bruchet, A. Occurrence of 38 pesticides in various French surface  
565 and ground waters. *Environ. Technol.* **1991**, *12* (11), 985-996.
- 566 [4] Raviola, M.; Ferrero, T.; Fiorenza, A. *Le risorse idriche superficiali-Corsi d'acqua. Rapporto*  
567 *sullo stato dell'ambiente in Piemonte nel 2009*, Rapporto sullo Stato dell'Ambiente in Piemonte;  
568 ARPA, 2010; 63-72.
- 569 [5] Cerejeira, M.J.; Viana, P.; Batista S.; Pereira, T.; Silva, E.; Valerio, M.J. ; Silva, A.; Ferreira,  
570 M.; Silva-Fernandes, A.M. Pesticides in Portuguese surface and ground waters. *Water Res.* **2003**,  
571 *37*(5), 1055-1063.
- 572 [6] Konstantinou, I.K.; Hela, D.G.; Albanis, T.A. The status of pesticide pollution in surface waters  
573 (rivers and lakes) of Greece. Part I. Review on occurrence and levels. *Environ. Poll.* **2006**, *141* (3),  
574 555-570.
- 575 [7] Ryberg, K.R.; Vecchia, A.V.; Martin, J.D.; Gilliom, R.J. *Trends in pesticide concentrations in*  
576 *urban streams in the United States, 1992-2008*, National Water-Quality Assessment Program  
577 Investigations Report 2010-5139; U.S. Geological Survey Scientific, 2010; 101.
- 578 [8] Paris, P.; De Santis, T.; Esposito, D.; Pace, E.; Romoli, D.; Ursino S. *Monitoraggio nazionale*  
579 *dei pesticidi nelle acque - ISPRA. Dati 2007-2008*, Rapporto di sintesi; ISPRA, 2010; 1-79.
- 580 [9] Wauchope, R.D. The pesticide content of surface water draining from agricultural fields - A  
581 review. *J. Environ. Qual.* **1978**, *7* (4), 459-472.
- 582 [10] Reichenberger, S.; Bach, M.; Skitschak, A.; Frede, H.G. Mitigation strategies to reduce  
583 pesticide inputs into ground- and surface water and their effectiveness; A review. *Sci. Total*  
584 *Environ.* **2007**, *384* (1-3), 1-35.

- 585 [11] Tang, X.; Zhu, B.; Katou, H. A review of rapid transport of pesticides from sloping farmland to  
586 surface waters: Processes and mitigation strategies. *J. Environ. Sci.* **2012**, *24* (3), 351-361.
- 587 [12] CORPEN. Les fonction environnementales des zone tampons. Les bases scientifiques et  
588 techniques des fonctions de protection des eaux, 2007. Available at <http://www.developpement->  
589 [durable.gouv.fr/IMG/pdf/DGALN\\_fonctions\\_environn\\_zones\\_temp\\_chapitre1.pdf](http://www.developpement-durable.gouv.fr/IMG/pdf/DGALN_fonctions_environn_zones_temp_chapitre1.pdf) (accessed May  
590 2011).
- 591 [13] USDA. Conservation buffer to reduce pesticides losses. USDA Natural Resources  
592 Conservation Service, 2000. Available at <http://www.in.nrcs.usda.gov/technical/agronomy/>  
593 [newconbuf.pdf](http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf) (accessed June 2011).
- 594 [14] Arora, K.; Mickelson, S.K.; Helmers, M.J.; Baker, J.L. Review of Pesticide retention processes  
595 occurring in buffer strips receiving agricultural runoff. *J. Am. Water Resour. Assoc.* **2010**, *46* (3),  
596 618-647.
- 597 [15] Zanin, G.; Otto, S.; Masin, R.; Ferrero, A.; Milan, M.; Vidotto, F. Protezione delle acque  
598 superficiali da ruscellamento e deriva: efficacia delle fasce tampone e di altre misure di mitigazione.  
599 Atti del convegno Protezione dei corpi idrici superficiali dall'inquinamento da agrofarmaci,  
600 Bologna, Italy, 19 maggio 2009; Vidotto, F., Ferrero, A., Eds.; SIRFI (Società Italiana per la  
601 Ricerca sulla Flora Infestante): Bologna, 2009; 221.
- 602 [16] Stanners P. Farmers threaten to sue government over waterways laws. *The Copenhagen Post*,  
603 Copenhagen, 2012. Available at <http://cphpost.dk/news/national/farmers-threaten-sue-government->  
604 [over-waterways-laws](http://cphpost.dk/news/national/farmers-threaten-sue-government-over-waterways-laws) (accessed March 2012).
- 605 [17] Campanini, L.; Pisa Rossi, P.; Catizone, P. La presenza di erbicidi nelle acque di ruscellamento  
606 superficiale e nel terreno eroso da zone declivi (Herbicide residues in runoff and sediment from  
607 sloping arable fields). Atti Controllo delle piante infestanti, Bologna, Italy, 21-22 gennaio 1992;  
608 Bologna, 1992; 189-214.
- 609 [18] Müller, K.; Trollove, M.; James, T.K.; Rahman, A. Herbicide loss in runoff: Effects of  
610 herbicide properties, slope, and rainfall intensity. *Austral. J. Soil Res.* **2004**, *42* (1), 17-27.

- 611 [19] Patakioutas, G.I.; Albanis, T.A. Runoff of herbicides from cropped and uncropped plots with  
612 different slopes. *Intern. J. Environ. Analyt. Chem.* **2004**, *84* (1-3), 103-121.
- 613 [20] Schmitt, T.J.; Dosskey, M.G.; Hoagland, K.D. Filter strip performance and processes for  
614 different vegetation, widths, and contaminants. *J. Environ. Qual.* **1999**, *28* (5), 1479-1489.
- 615 [21] BCPC. *The pesticide manual*, 11th Ed.; BCPC (The British Crop Protection Council), Eds.;  
616 Page Bros.: Norwich, 1997; 1606.
- 617 [22] Pallett, K.E.; Little, J.P.; Sheekey, M.; Veerasekaran, P. The mode of action of isoxaflutole: I.  
618 Physiological effects, metabolism, and selectivity. *Pest. Biochem. Physiol.* **1998**, *62* (2), 113-124.
- 619 [23] Gupta, S.; Gajbhiye, V.T. Effect of concentration, moisture and soil type on the dissipation of  
620 flufenacet from soil. *Chemosphere* **2002**, *47*(9), 901-906.
- 621 [24] Rouchaud, J.; Neus, O.; Cools, K.; Bulcke, R. Flufenacet soil persistence and mobility in corn  
622 and wheat crops. *Bull. Environ. Contam. Toxic.* **1999**, *63* (4), 460-466.
- 623 [25] Toxnet. Toxicology Data Network-Flufenacet, National Library of Medicine HSDB Database,  
624 2012. Available at [http://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+](http://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+7011)  
625 7011 (accessed Sept 2012).
- 626 [26] Rouchaud, J.; Neus, O.; Eelen, H.; Bulcke, R. Persistence, mobility, and adsorption of the  
627 herbicide flufenacet in the soil of winter wheat crops. *Bull. Environ. Contam. Toxic.* **2001**, *67* (4),  
628 609-616.
- 629 [27] European-Commission. Review report for the active substance flufenacet, Health & Consumer  
630 Protection Directorate-General, European Commission, 2003. Available at  
631 [http://ec.europa.eu/food/plant/protection/evaluation/newactive/list1\\_flufenacet\\_en.pdf](http://ec.europa.eu/food/plant/protection/evaluation/newactive/list1_flufenacet_en.pdf)  
632 (accessed June 2011).
- 633 [28] PPDB, The Pesticide Properties Database (PPDB) developed by the Agriculture &  
634 Environment Research Unit (AERU), University of Hertfordshire, funded by UK national sources  
635 and the EU-funded FOOTPRINT project (FP6-SSP-022704), 2009. Available at  
636 <http://sitem.herts.ac.uk/aeru/footprint/it/index.htm> (accessed Oct 2010).

- 637 [29] Alletto, L.; Coquet, Y.; Bergheaud, V., Benoit, P. Water pressure head and temperature impact  
638 on isoxaflutole degradation in crop residues and loamy surface soil under conventional and  
639 conservation tillage management. *Chemosphere* **2012**, *88* (9), 1043-1050.
- 640 [30] EPA, Pesticide Fact Sheet - Isoxaflutole, United States Environmental Protection Agency.  
641 Office of Prevention, Pesticides and Toxic Substances, 1998, 15. Available at  
642 [www.epa.gov/.../fs\\_PC-123000\\_15-Sep-98.pdf](http://www.epa.gov/.../fs_PC-123000_15-Sep-98.pdf) (accessed Feb 2011).
- 643 [31] Papiernik, S.K.; Yates, S.R.; Koskinen, W.C.; Barber, B. Processes Affecting the Dissipation  
644 of the Herbicide Isoxaflutole and Its Diketonitrile Metabolite in Agricultural Soils under Field  
645 Conditions. *J. Agric. Food Chem.* **2007**, *55* (21), 8630-8639.
- 646 [32] Mitra, S.; Bhowmik, P.C.; Xing, B. Physical and chemical properties of soil influence the  
647 sorption of the diketonitrile metabolite of RPA 201772. *Weed Sci.* **2001**, *49* (3), 423-430.
- 648 [33] Taylor-Lovell, S.; Sims, G.K.; Wax, L.M. Effects of moisture, temperature, and biological  
649 activity on the degradation of isoxaflutole in soil. *J. Agric. Food Chem.* **2002**, *50* (20), 5626-5633.
- 650 [33] Taylor-Lovell, S.; Sims, G.K.; Wax, L.M.; Hassett, J.J. Hydrolysis and soil adsorption of the  
651 labile herbicide isoxaflutole. *Environ. Sci. Technol.* **2000**, *34* (15), 3186-3190.
- 652 [35] Pallett, K.E.; Cramp, S.M.; Little, J.P.; Veerasekaran, P.; Crudace, A.J.; Slater, A.E.  
653 Isoxaflutole: The background to its discovery and the basis of its herbicidal properties. *Pest Manag.*  
654 *Sci.* **2001**, *57*, 133-142.
- 655 [36] Mitra, S.; Bhowmik, P.C.; Xing, B. Sorption of isoxaflutole by five different soils varying in  
656 physical and chemical properties. *Pest. Sci.* **1999**, *55* (2), 935-942.
- 657 [37] Rouchaud, J.; Neus, O.; Callens, D.; Bulcke, R. Isoxaflutole, herbicide soil persistence and  
658 mobility in summer corn and winter wheat crops. *Bull. Environ. Contam. Toxicol.* **1998**, *60* (4),  
659 577-584.
- 660 [38] Oliveira Jr, R.S.; Marchiori Jr, O.; Constantin, J.; Inoue, M.H. Influence of drought periods on  
661 the residual activity of isoxaflutole in soil. *Planta dan.* **2006**, *24* (4), 733-740.

662 [39] Alletto, L.; Benoit, P.; Bergheaud, V.; Coquet, Y. Variability of retention process of  
663 isoxaflutole and its diketonitrile metabolite in soil under conventional and conservation tillage. *Pest*  
664 *Manag. Sci.* **2012**, *68* (4), 610-617.

665 [40] Rice, P.J.; Koskinen, W.C.; Carrizosa, M.J. Effect of soil properties on the degradation of  
666 isoxaflutole and the sorption-desorption of isoxaflutole and its diketonitrile degradate. *J. Agric.*  
667 *Food Chem.* **2004**, *52* (25), 7621-7627.

668 [41] Beltrán, E.; Fenet, H.; Cooper, J.F.; Coste, C.M. Fate of isoxaflutole in soil under controlled  
669 conditions. *J. Agric. Food Chem.* **2003**, *51* (1), 146-151.

670 [42] Milan, M.; Vidotto, F.; Piano, S.; Negre, M.; Ferrero, A. Buffer strip effect on terbuthylazine,  
671 desethyl-terbuthylazine and S-metolachlor runoff from maize fields in Northern Italy. *Environ.*  
672 *Technol.* **2013**, *34* (1), 71-80.

673 [43] Müller, K.; Trolove, M.; James, T.K.; Rahman, A. Herbicide loss in runoff: Effects of  
674 herbicide properties, slope, and rainfall intensity. *Aust. J. Soil Res.* **2004**, *42* (1), 17-27.

675 [44] Gaynor, J.D.; MacTavish, D.C.; Findlay, W.I. Atrazine and metolachlor loss in surface and  
676 subsurface runoff from three tillage treatments in corn. *J. Environ. Qual.* **1995**, *24* (2), 246-256.

677 [45] Ng, H.Y.F.; Clegg, S.B. Atrazine and metolachlor losses in runoff events from an agricultural  
678 watershed: The importance of runoff components. *Sci. Total Environ.* **1997**, *193* (3), 215-228.

679 [46] Leonard, R.A. Movement of pesticides into surface waters. In *Pesticides in the soil -*  
680 *Environment: processes, impacts and modeling*; Soil Science Society of America, Inc: Madison,  
681 Wisconsin, USA, 1990; 303-342.

682 [47] Kim, J.H.; Feagley, S.E. Runoff of trifluralin, metolachlor and metribuzim from a clay-loam  
683 soil of Louisiana. *J. Environ. Sci. Health, Part B* **2002**, *37* (5), 405-415.

684 [48] Meyer, M.T.; Scribner, E.A.; Kalkhoff, S.J. Comparison of Fate and Transport of Isoxaflutole  
685 to Atrazine and Metolachlor in 10 Iowa Rivers. *Environ. Sci. Technol.* **2007**, *41* (20), 6933-6939

686 [49] BCPC. *The pesticide manual*, 11th Ed.; BCPC (The British Crop Protection Council), Eds.;  
687 Page Bros.: Norwich, 1997; 1606.

- 688 [50] Krutz, L.J.; Senseman, S.A.; Zablotowicz, R.M.; Matocha, M.A. Reducing herbicide runoff  
689 from agricultural fields with vegetative filter strips: A review. *Weed Sci.* **2005**, *53* (3), 353-367.
- 690 [51] Smith, A.M. Vegetative filter strip for improved water quality, Iowa State University,  
691 University Extension, 1999, 4. Available at [http://www.extension.iastate.edu/publications/pm](http://www.extension.iastate.edu/publications/pm1507.pdf)  
692 [1507.pdf](http://www.extension.iastate.edu/publications/pm1507.pdf) (accessed April 2011).
- 693 [52] Abu-Zreig, M.; Rudra, R.P.; Lalonde, M.N.; Whiteley, H.R.; Kaushik, N.K. Experimental  
694 investigation of runoff reduction and sediment removal by vegetated filter strips. *Hydrolog.*  
695 *Process.* **2004**, *18* (11), 2029-2037.
- 696 [53] Tingle, C.H.; Shaw, D.R.; Boyette, M.; Murphy, G.P. Metolachlor and metribuzin losses in  
697 runoff as affected by width of vegetative filter strips. *Weed Sci.* **1998**, *46* (4), 475-479.
- 698

699

700

701

## Figure captions

702 **Figure 1:** Experimental layout adopted. A: sampling devices.

703 **Figure 2.** Flufenacet concentrations ( $\mu\text{g kg}^{-1}$ ) in soil of treated areas in 2009 (a) and 2010 (b).

704 Arithmetic mean of three bulk replications  $\pm$ SD.

705

706

707

708

709

710

711

712

713

714

715

716

717

718


719

720

721

722

723



724

725 **Fig. 1**

726

727

728


729

730


731

732





**a**



**b**

733 **Fig. 2**

734

735

736 **Table 1.** Physico-chemical properties of flufenacet and isoxaflutole. Source: PPDB, The Pesticide  
737 Properties Database, AERU, University of Hertfordshire, 2009.

738

<b>Herbicides</b>	<b>Water solubility</b> <b>(mg L<sup>-1</sup>)</b>	<b>Koc</b> <b>(mL g<sup>-1</sup>)</b>	<b>DT50 in field</b> <b>(days)</b>	<b>GUS</b>
<b>Flufenacet</b>	56	401	40	2.4
<b>Isoxaflutole</b>	6.2	145	1.3	0.6

739

*Note: GUS = Ground water ubiquity score*

740 **Table 2.** Concentration of flufenacet detected in water samples collected after each runoff event in 2009 and 2010. Values are expressed in  $\mu\text{g L}^{-1}$ .

741 Arithmetic mean of three replications  $\pm$  SE. Same-letter values are not significantly different [REGWF test (\*P <0.05)].

DAT	Precipitation mm	Temperature $^{\circ}\text{C}$	Flufenacet concentration in the runoff water ( $\mu\text{g L}^{-1}$ )					
			TFTEST	TF2	TF4	TF6	RIVATEST	RIVA6
----- 2009 -----								
4	23.6	9.3	13.7 $\pm$ 5.5 b	9.2 $\pm$ 3.1 b	14.9 $\pm$ 3.6 b	13.5 $\pm$ 8 b	66.3 $\pm$ 10.2 a	57.2 $\pm$ 3.8 a
5	23.2	10.7	13.7 $\pm$ 3.1 c	6.4 $\pm$ 0.8 d	12.3 $\pm$ 1.2 c	6.3 $\pm$ 1.5 d	67 $\pm$ 2.6 b	91.7 $\pm$ 10.2 a
11*	33.2	10.1	1.5	1.5	6.1	4.6	41.4	42.5
12	63.4	9.5	2.4 $\pm$ 0.2 d	2.1 $\pm$ 0.6 d	3.9 $\pm$ 0.9 c	1.3 $\pm$ 0.2 d	19.5 $\pm$ 0.8 a	16 b
12			3.5 $\pm$ 1.2 b	3 $\pm$ 0.2 b	3.8 $\pm$ 0.8 b	4.4 $\pm$ 0.7 b	10.8 $\pm$ 1.1 a	14.3 $\pm$ 3.2 a
13	9.2	11.8	5.4 $\pm$ 0.8 c	2.7 $\pm$ 0.4 c	9.1 $\pm$ 0.2 c	4.9 $\pm$ 1.8 c	64.7 $\pm$ 3.6 a	57.5 $\pm$ 4.3 b
62 (I)	-	24.5	0.4 $\pm$ 0.3 a	0.3 $\pm$ 0.3 a	0.3 $\pm$ 0.1 a	0.5 $\pm$ 0.4 c	15 $\pm$ 3.5 a	5.8 $\pm$ 1.9 b
68	14	18.2	0.8 $\pm$ 0.3 b	< LOQ	< LOQ	< LOQ	1.5 $\pm$ 0.1 a	1.0 $\pm$ 0.0 b
77	46.8	23.2	0.4 $\pm$ 0.10 b	0.1 $\pm$ 0.3 b	0.3 $\pm$ 0.1 b	0.5 $\pm$ 0.1 b	1.9 $\pm$ 0.4 a	1.7 $\pm$ 0.4 a
82	28.6	23.5	0.1 $\pm$ 0.0 c	< LOQ	< LOQ	0.1 $\pm$ 0.0	0.5 $\pm$ 0.0 a	0.3 $\pm$ 0.1 b
93	39	22.4	0.2 $\pm$ 0.1 bc	< LOQ	< LOQ	< LOQ	0.4 $\pm$ 0.01 c	0.3 $\pm$ 0.0 bc
110 (I)	-	25.1	0.2 $\pm$ 0.1 bc	0.2 $\pm$ 0.1	0.2 $\pm$ 0.2 a	0.5 $\pm$ 0.6 a	0.9 $\pm$ 0.9 c	0.7 $\pm$ 0.3 bc
115	30	22.7	0.2 $\pm$ 0.0 b	< LOQ	< LOQ	< LOQ	0.3 $\pm$ 0.0 a	0.2 $\pm$ 0.0 b
----- 2010 -----								
10	25.4	13.8	13.0 $\pm$ 7.5 a	0.6 $\pm$ 0.4 b	2.7 $\pm$ 1.6 b	0.2 $\pm$ 0.1 b	10.4 $\pm$ 5.9 a	0.7 $\pm$ 0.4 b
11	42.6	11.4	1.0 $\pm$ 0.7 c	0.7 $\pm$ 0.3 d	0.8 $\pm$ 0.5 c	0.3 $\pm$ 0.2 d	6.2 $\pm$ 3.6 a	4.2 $\pm$ 2.4 b
17	11.2	15.4	0.9 $\pm$ 0.5 c	0.7 $\pm$ 0.4 c	0.9 $\pm$ 0.5 c	0.6 $\pm$ 0.3 c	2.7 $\pm$ 1.5 a	1.6 $\pm$ 0.9 a
44	37.6	22.0	0.3 $\pm$ 0.2 c	0.2 $\pm$ 0.1 c	0.2 $\pm$ 0.1 c	0.2 $\pm$ 0.1 c	0.5 $\pm$ 0.2 a	0.4 $\pm$ 0.3
53	72.8	17.1	0.1 $\pm$ 0.1 a	< LOQ	0.1 $\pm$ 0.1 a	0.1 $\pm$ 0.1 a	0.2 $\pm$ 0.2 a	0.1 $\pm$ 0.2 a
54	23.2	18.9	< LOQ	< LOQ	< LOQ	< LOQ	0.1 $\pm$ a	0.1 $\pm$ a
57	12.8	17.7	< LOQ	< LOQ	< LOQ	< LOQ	0.1 $\pm$ a	0.1 $\pm$ a
89 (I)	I	23.9	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
110	69.6	17.3	nc	nc	nc	nc	nc	nc

742

743 Note 1: DAT (days after herbicide treatment); I (Irrigation); LOQ (Limit of quantification) =0.05  $\mu\text{g L}^{-1}$  for flufenacet. nc: not collected \* Arithmetic mean of two data.

744

745

746 **Table 3.** Concentration of isoxaflutole detected in water samples collected after each runoff event in 2009 and 2010. Values are expressed in  $\mu\text{g L}^{-1}$ .747 Arithmetic mean of three replications  $\pm$  SE. Same-letter values are not significantly different [REGWF test (\*P <0.05)].

DAT	Precipitations mm	Temperature °C	Isoxaflutole concentration in the runoff water ( $\mu\text{g L}^{-1}$ )					
			TFTEST	TF2	TF4	TF6	RIVATEST	RIVA6
-----2009-----								
4	23.6	9.3	0.16 (0.01) c	< LOQ	0.08 (0.06) c	< LOQ	5 (3.30) a	2.87 (0.45) b
5	23.2	10.7	0.20 (0.03) c	< LOQ	0.13 (0.07) c	< LOQ	4.17 (0.61) a	0.99 (0.18) b
11*	33.2	10.1	< LOQ	< LOQ	< LOQ	< LOQ	0.75 (0.31) a	0.92 (0.32) a
12			< LOQ	< LOQ	< LOQ	< LOQ	0.34 (0.02) a	0.11 (0.09) b
12	63.4	9.5	< LOQ	< LOQ	< LOQ	< LOQ	0.42 (0.03) a	0.35 (0.26) a
13	9.2	11.8	0.13 (0.01) b	< LOQ	< LOQ	< LOQ	1.14 (0.23) a	1.24 (0.15) a
62 (I)	-	24.5	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
68	14	18.2	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
77	46.8	23.2	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
82	28.6	23.5	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
93	39	22.4	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
110 (I)	-	25.1	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
115	30	22.7	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
-----2010-----								
10	25.4	13.8	< LOQ	< LOQ	< LOQ	< LOQ	0.10 (0.05)	< LOQ
11	42.6	11.4	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
17	11.2	15.4	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
44	37.6	22.0	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
53	72.8	17.1	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
54	23.2	18.9	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
57	12.8	17.7	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
89 (I)	-	23.9	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
110	69.6	17.3	nc	nc	nc	nc	nc	nc

748

749 Note 2: DAT (days after treatment); I (Irrigation); LOQ (Limit of quantification) =  $0.02 \mu\text{g L}^{-1}$ . nc: not collected \* Arithmetic mean of two data.

750

751

**Table 4.** Rainfall events and corresponding measured runoff volumes in 2010.

DAT	Rainfall (mm)	Runoff (m <sup>3</sup> )					
		TFTEST	TF2	TF4	TF6	RIVATEST	RIVA6
10	25.4	1.0	0.9	1.0	0.8	1.2	0.8
11	42.6	1.5	1.3	2.5	2.6	11	9.6
17	11.2	0.9	0.5	1.5	0.8	2.8	1.6
44	37.6	7.2	8.0	7.0	6.5	7.8	7.7
53	72.8						
54	23.2	53.7	50.6	53.0	56.3	87.0	78.5
57	12.8	1.8	1.9	1.8	0.8	3.4	1.6
89 (I)	I*	16.7	16.6	14.5	11.9	21.5	17.5
110	69.6	nm	nm	nm	nm	nm	nm

752

Note 3: DAT (days after treatment); NR: no runoff; I: Irrigation; nm: not measured; \* Irrigation volumes were: RIVATEST: 38.7 m<sup>3</sup>;

753

RIVA6:44.0 m<sup>3</sup>; TFTEST: 34.6 m<sup>3</sup> TF2: 39.4 m<sup>3</sup> TF4: 37.6; TF6 35.3 m<sup>3</sup> TF

754

755

756