

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Welding and lung cancer in a pooled analysis of case-control studies.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/142084> since 2016-07-15T11:06:07Z

*Published version:*

DOI:10.1093/aje/kwt201

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## **Welding and Lung Cancer in a Pooled Analysis of Case- Control Studies**

Journal:	<i>American Journal of Epidemiology</i>
Manuscript ID:	AJE-00289-2013
Manuscript Type:	Systematic Reviews, Meta- and Pooled Analyses
Key Words:	case-control studies, lung cancer, occupational exposure, welding

SCHOLARONE™  
Manuscripts

Review

## Welding and Lung Cancer in a Pooled Analysis of Case-Control Studies

### ABSTRACT

Several epidemiologic studies indicated an increased risk of lung cancer among welders. We took advantage of a large collection of lung cancer cases and controls with occupational and smoking histories from the SYNERGY project to assess welding as a risk factor for developing lung cancer. The database comprises 15,483 male lung cancer cases and 18,388 controls from 16 studies in Europe, Canada, China, and New Zealand, conducted between 1985 and 2010. Odds ratios (OR) and 95% confidence intervals (95%CI) for lung cancer were estimated, adjusted for smoking and other confounders. Overall, 568 cases and 427 controls had ever worked as a welder, and 1,994 cases and 1,930 controls had ever worked in occupations with occasional welding. Welding was associated with an elevated OR for lung cancer (welders: 1.44, 95%CI: 1.25, 1.67; occasional welding: 1.19, 95%CI: 1.10, 1.28) and increased with duration of employment. Regular welding showed a stronger association with lung cancer in never and light smokers (0-<10 packyears: 1.96, 95%CI: 1.37, 2.79) and with squamous cell cancer and small cell lung cancer than with adenocarcinoma. Our findings contribute to the increasing evidence that welding is associated with an increased risk of lung cancer.

1  
2  
3 Word count abstract: 194  
4

5 Word count text: 3,194  
6  
7  
8  
9

10 Abbreviations: AdCa, adenocarcinoma; CI, confidence interval; IARC, International Agency  
11 for Research on Cancer; ISCO-68, International Standard Classification of Occupations,  
12 Revision 1968; ISIC, International Standard Industrial Classification; OR, odds ratio; SCLC,  
13 small cell lung cancer; SqCC, squamous cell carcinoma;  
14  
15  
16  
17  
18  
19

20  
21 Running head: Welding and lung cancer  
22  
23

24  
25  
26 Keywords: case-control studies, lung cancer, occupational exposure, welding  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Worldwide several millions of workers are exposed to welding fumes while working as  
4 welders or in occupations or workplaces in which joining of metal parts are commonly  
5 performed. In 1990, the International Agency for Research on Cancer (IARC) classified  
6 welding fumes as a possible human carcinogen (Group 2B) (1). New evidence for an excess  
7 lung cancer risk in welding has been found in several studies, including a recent meta-analysis  
8 and a large record-linkage study (2, 3). Research gaps and recommendations have been  
9 discussed by Ward *et al.* (4).  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 Welding fume is a complex mixture of particles containing metals and gases that are formed  
21 during the burning of electrodes and heating of the base metal. The particle mass  
22 concentration is strongly correlated with iron and manganese as major constituents of steel  
23 (5), but moderately with chromium and nickel, which are added to the production of stainless  
24 steel (6). Ship-building is a prominent example for welding of large parts of mild steel with  
25 gas metal arc welding and is associated with high exposure to particles and lower exposure to  
26 chromium and nickel as compared to settings using stainless steel (6, 7).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Although the association of welding with lung cancer has been repeatedly investigated, a  
40 causal association between welding and lung cancer has not yet been firmly established,  
41 except for stainless steel welders. Further questions remain, for example the contribution that  
42 welding makes to the etiology of the different subtypes of lung cancer and how welding  
43 interacts with smoking in lung cancer development. Smoking is more closely associated with  
44 squamous cell cancer (SqCC) and small cell cancer of the lung (SCLC) than with  
45 adenocarcinoma (AdCa) (8).  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 We took advantage of a large collection of lung cancer cases and controls with occupational  
4 and smoking histories from the SYNERGY project to explore welding as a risk factor for  
5 developing lung cancer. More information about SYNERGY has been previously published  
6  
7  
8  
9  
10 (8, 9) and is available at <http://synergy.iarc.fr>.

## 11 12 13 14 15 **MATERIALS AND METHODS**

### 16 17 18 **Study population**

19  
20 The pooled dataset for this analysis comprised 15,483 male cases and 18,388 male controls  
21 that were enrolled in 16 studies between 1985 and 2010 (supplementary Table S1). We  
22 excluded women due to the low prevalence of welders among them; only 32 women had ever  
23 been employed in welding occupations. The majority of controls were recruited from the  
24 general population (81.6%). Occupational and smoking histories were mainly assessed in  
25 face-to-face interviews (80.6%). The majority of subjects were alive at the time of interview  
26 (87.9%). The Ethics Committees of the individual studies approved the realization of the  
27 study, as did the IARC Institutional Review Board.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 40 41 42 **Assessment of welding activities**

43 Occupations were coded according to the International Standard Classification of  
44 Occupations, Rev. 1968 (ISCO-68) (10). Industries were classified according to the  
45 International Standard Industrial Classification (ISIC) Rev. 2 (11). Using the 5-digit ISCO-68  
46 codes (supplementary Table S2), two classifications of welding-related occupations were  
47 considered: men who ever worked for at least one year as a welder, and men ever working in  
48 occupations with potential welding activities (further referred to as ‘occasional welding’).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Occasional welding was assessed by expert rating of job titles (B.K., B.P., H.K., R.V., and  
4  
5 S.P.) and excluded men who ever worked as a regular welder. We further stratified both  
6  
7 groups by industry where welding is commonly applied (ship building and repair,  
8  
9 construction, manufacture of machines and related equipment, manufacture of motor vehicles  
10  
11 and repair of transport equipment, others) (supplementary Table S3).  
12  
13

### 14 15 16 **Statistical analysis**

17  
18 The odds ratio (OR) for working as a welder or in occasional welding and 95% confidence  
19  
20 intervals (95%CI) were estimated by unconditional logistic regression. OR1 was adjusted for  
21  
22 age (log-transformed) and study centers (22 centers), OR2 was additionally adjusted for  
23  
24 smoking (log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever  
25  
26 other types of tobacco only, stopping smoking 2-7, 8-15, 16-25, 26+ years before  
27  
28 interview/diagnosis, never smokers)). The full adjustment (OR3) additionally considered  
29  
30 employment in occupations at-risk for lung cancer excluding welding-related occupations  
31  
32 (“List A” jobs (12, 13)). Subjects, who had never worked in welding-related occupations,  
33  
34 comprised the reference group. We defined light smokers as subjects who smoked less than  
35  
36 10 packyears. Tests for linear trend were performed for the OR as a function of duration of  
37  
38 employment and time since last employment as a welder. We estimated the lung cancer risk  
39  
40 for regular or occasional welding as ever held and longest-held occupation, stratified by  
41  
42 smoking and subtype of lung cancer, in subjects who never worked in a “List A” job and in  
43  
44 workers ever employed in a ‘blue-collar’ occupation (with 7, 8, or 9 as first digit of their job-  
45  
46 title code according to ISCO-68). Sensitivity analyses were performed by type of controls and  
47  
48 for workers who started welding after 1979, when measures for the improvement of  
49  
50 occupational hygiene became implemented. Meta-regression models with random effects  
51  
52  
53  
54  
55

1  
2  
3 were applied to estimate ORs for the combined and individual studies, and heterogeneity  
4  
5 between studies was assessed by  $I^2$  statistics (Comprehensive Meta-Analysis Version 2.2.027,  
6  
7 Biostat, Englewood, NJ). All other analyses were performed with SAS software, version 9.2  
8  
9 (SAS Institute Inc., Cary, NC).  
10

## 11 12 13 **RESULTS**

### 14 15 **Study population**

16  
17 Table 1 characterizes the study groups. A total of 568 male lung cancer cases (3.7%) and 427  
18  
19 controls (2.3%) ever worked as a welder. A large fraction of 40.1% of welders among the  
20  
21 controls reported this occupation as longest-held job, with a median of 27 years (interquartile  
22  
23 range 18-35 years). Additional 1994 cases (12.9%) and 1930 controls (10.5%) were identified  
24  
25 as having worked in occupations with occasional welding. A similar fraction of 36.1% had  
26  
27 occupation as longest-held job (median of 31 years, interquartile range 23-38 years). More  
28  
29 welders ever smoked or worked in other at-risk occupations as compared to men who had  
30  
31 never worked in welding-related occupations.  
32  
33  
34  
35  
36  
37  
38  
39

### 40 41 **Lung cancer risk among welders and in occasional welding**

42  
43 If not otherwise stated, we report the fully adjusted risk estimates. Figure 1 shows a meta-OR  
44  
45 of 1.42 (95%CI: 1.23, 1.66) for ever working as a welder with low heterogeneity between  
46  
47 individual studies ( $I^2=29.6%$ ,  $P=0.10$ ) and of 1.17 (95%CI: 1.08, 1.27) for occasional welding  
48  
49 ( $I^2=20.9%$ ,  $P=0.19$ ). Table 2 presents the risk estimates from the pooled analysis. Working as  
50  
51 a welder was associated with an increased lung cancer risk (ever welder: 1.44, 95%CI: 1.25,  
52  
53 1.67, longest held job: 1.50 (95%CI: 1.20, 1.88). Occasional welding was also associated with  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 an elevated lung cancer risk (ever: 1.19, 95%CI: 1.10, 1.28). The risk estimates remained  
4  
5 increased when restricted to men never working in a “List A” job or to blue-collar workers  
6  
7 (welder as longest held occupation: 1.63, 95%CI: 1.26, 2.11, and 1.39; 95%CI: 1.11, 1.73,  
8  
9 respectively). ORs were 1.53 (95%CI: 1.29, 1.82) when restricted to population controls, 1.10  
10  
11 (95%CI: 0.84, 1.44) in studies with hospital controls, and 1.31 (95%CI: 0.68, 2.58) for having  
12  
13 started working after 1979 (supplementary Table S4).  
14  
15

### 16 17 18 **Lung cancer risk by industry** 19

20  
21 Welding in the construction industry showed an increased lung cancer risk (ever welder: 1.47,  
22  
23 95%CI: 1.22, 1.78; ever occasional welding: 1.21, 95%CI: 1.09, 1.33). We estimated an  
24  
25 elevated risk for ever working as welder in ship building and repair (1.53, 95%CI: 1.06, 2.21),  
26  
27 but not for occasional welding (0.90, 95%CI: 0.68, 1.20). The estimates of the lung cancer  
28  
29 risk in the motor vehicle and motor bike production were comparatively low, for example  
30  
31 0.62 (95%CI: 0.28, 1.36) for welding as the longest held job. More results are shown in Table  
32  
33 3.  
34  
35  
36  
37

### 38 **Lung cancer risk by duration of employment in regular or occasional welding** 39

40  
41 Short-term exposure (<3 years of welding) was associated with a relative lung cancer risk of  
42  
43 1.14 (95%CI: 0.80, 1.61) in regular welders and of 1.13 (95%CI: 0.94, 1.34) in occasional  
44  
45 welding (Table 4). The risks increased with longer duration of welding. Long-term exposure  
46  
47 (>25 years of welding) was associated with an OR of 1.77 (95%CI: 1.31, 2.39) in welders and  
48  
49 of 1.40 (95%CI: 1.21, 1.62) in occasional welders. We did not observe a risk reduction with  
50  
51 increasing time since last welding (data not shown).  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Lung cancer risk by histological subtype and smoking

Table 5 shows the risk estimates of ever working as a welder for the major subtypes of lung cancer (AdCa 1.23, 95%CI: 0.99, 1.53, SqCC 1.58, 95%CI: 1.32, 1.89, SCLC 1.41, 95%CI: 1.09, 1.82). Among never or light smoking welders, the estimates of the risk for SqCC and SCLC were increased more strongly, but were based on small numbers (never smokers: AdCa 1.89, 95%CI: 0.79, 4.52, SqCC 3.01, 95%CI: 1.07, 8.49, SCLC 4.45, 95%CI: 1.03, 19.18).

This pattern was not observed among never-smoking occasional welders.

### DISCUSSION

Analyses of a large database of occupational and smoking histories from about 33,900 men within the framework of the SYNERGY project, we demonstrated that welding is associated with an increased lung cancer risk. A meta-analysis of 66 epidemiological studies revealed a 26% excess of lung cancer risk among welders (2). Two studies of which were also part of our analysis (14, 15). A large record-linkage study in the Nordic countries estimated a 33% increased lung cancer incidence in welders, although this study did not adjust for smoking (3). The adjustment for smoking habits suggests that confounding by tobacco smoking may explain about 20% of the increase in lung cancer risk in welders.

Welding is a common task in many occupations, where joining of metal parts is occasionally performed. In order to provide better evidence for lung cancer risks in welding, we investigated welders and workers in occupations with potential welding activities separately. In the pooled analysis of 16 case-control studies, we observed a 44% smoking-adjusted increase in risk for ever working as a welder and a 19% increase for occasional welding, but the risks were lower in studies with hospital controls. This estimate showed only minor

1  
2  
3 changes when we restricted the analysis to blue-collar workers or subjects not working in  
4  
5 other at-risk occupations. The risk increased up to 77% for welders worked more than 25  
6  
7 years.  
8  
9

10  
11 There is a variety of welding techniques with a wide range of emissions of particles and  
12  
13 metals (7). The welding technique and the metal content of the electrodes are important  
14  
15 determinants of the composition of welding fumes (6). Flux-cored arc welding is frequently  
16  
17 used for joining large metal parts of mild steel, e.g. in shipyards, leading to high level of  
18  
19 particle exposure and comparatively low concentrations of chromium and nickel. We found a  
20  
21 significantly increased risk for regular welders in ship building but not for occasional  
22  
23 welding. The finding that lung cancer risk is also increased among welders in shipyards where  
24  
25 predominantly mild steel is used and exposure to particulates is high indicates that the risk is  
26  
27 not restricted to stainless steel welding. Overall, we did not observed a significantly increased  
28  
29 lung cancer risk of welding in the manufacture of motor vehicles. Welding of car parts has  
30  
31 been subject to technological improvements, such as automated laser welding in enclosed  
32  
33 spaces, whereas the joining of large ship parts is still performed with high-emission  
34  
35 technologies (2).  
36  
37  
38  
39  
40  
41  
42

43 A limitation of this pooled analysis is that information on the welding process or on the  
44  
45 specific nature of workplace exposures was not available in this job title-based analysis. Still,  
46  
47 the job title of “welder” is one that carries a strong likelihood of exposure to some form of  
48  
49 welding fumes. There is certainly an opportunity for workers with other job titles to engage in  
50  
51 or be in proximity to welding operations. We attempted to identify the main occupations in  
52  
53 our listing of tasks with occasional welding activities. These workers comprise a larger group  
54  
55

1  
2  
3 compared to subjects that worked as a regular welder (16). We cannot fully exclude  
4  
5 misclassification of exposure, especially of occasional welders, where several workers might  
6  
7 not have performed welding. On the other hand, it is possible that some workers in our  
8  
9 “unexposed” category also had been exposed to welding fumes. However, this would only  
10  
11 represent a very small fraction of the reference group and would tend to attenuate the risk  
12  
13 estimates toward unity.  
14  
15

16  
17  
18 When analysing the risk in blue-collar workers only, we observed somewhat lower but still  
19  
20 elevated lung cancer risks in welding. Welders worked more frequently in other occupations  
21  
22 known to entail an increased lung cancer risk. However, the risk estimates did not change  
23  
24 when we adjusted for working in “List A” jobs, or when we restricted the analysis to men  
25  
26 who never worked in other at-risk occupations. The U.S. National Institute of Occupational  
27  
28 Safety and Health concluded in 1988 that there is an elevated risk of lung cancer among  
29  
30 welders that cannot be completely accounted for by smoking or asbestos exposure (17).  
31  
32 Exposure to asbestos can occur during welding, especially in shipyards (1). The risk estimate  
33  
34 was lower for welders who started working after 1979, when concerns about the health risks  
35  
36 of asbestos initiated the implementation of protective measures and a decline in asbestos  
37  
38 consumption (18). Several studies adjusted for exposure to asbestos, including four studies  
39  
40 that were part of our pooled analysis (14, 15, 19, 20). However, the inclusion of life-time  
41  
42 exposure to asbestos into the model did not influence the estimate of the relative lung cancer  
43  
44 risk among welders in the INCO study (19). Exposure to asbestos was usually assessed with  
45  
46 categorical variables based on expert rating. Personal measurements of asbestos fibres in the  
47  
48 breathing zone of welders are rarely available. We adjusted for other at-risk occupations  
49  
50 except welding and considered welding as a complex exposure circumstance, where exposure  
51  
52  
53  
54  
55

56  
57  
58  
59  
60

1  
2  
3 to asbestos could be an integral part. Asbestos could be used as filler of cylinders with  
4 acetylene gas or being part of certain covered rod electrodes. Asbestos fibres are not stable at  
5 the high temperatures during welding (21). Asbestos-containing materials were used for heat  
6 protection, and, for example, to cover the weld in order to delay the cooling process.  
7  
8 Exposure to asbestos is further possible when repairing metal parts with asbestos insulation or  
9 as bystander. Elevated mesothelioma risks among welders have been reported (22). The  
10 asbestos burden in the lungs of welders with mesothelioma was lower than in cases from the  
11 asbestos manufacturing industry or insulation sector (23). Taking this together, welding fume  
12 may exert a lung cancer risk that cannot be sufficiently explained by asbestos. Future research  
13 should consider types of welding and assessment of specific exposures by different welding  
14 scenarios.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Welders were more frequently smokers, as compared to the reference group and adjustment  
30 for smoking attenuated the risk estimates. Residual confounding by smoking can be a  
31 methodological problem. The detailed SYNERGY data allowed a detailed adjustment for  
32 smoking, and the large size of SYNERGY allowed a stratified analysis by smoking status.  
33  
34 The elevated odds ratios observed in never and light smokers are in line with a former report  
35 from the MONTREAL study (20) and support a direct effect of welding on lung cancer. The  
36 higher risk in never or light smokers might reflect the limited capacity of the lung to deal with  
37 exposure to pulmonary carcinogens.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 We found a stronger association of welding with SqCC and SCLC than with AdCa. This  
50 different association was also observed in other studies (24, 25). Also individual SYNERGY  
51 studies (14, 15, 19, 24, 26, 27) have reported on histological subtypes of lung cancer but were  
52  
53  
54  
55

1  
2  
3 based on smaller numbers (20, 28, 29). The same pattern was reported for exposure to  
4  
5 pulmonary carcinogens in uranium miners (30) and was even more pronounced in the  
6  
7 association of smoking with lung cancer (8).  
8  
9

10  
11 Summarizing, our findings contribute to the increasing evidence that welding is associated  
12  
13 with an increased lung cancer risk. The lung cancer risk of regular welders was higher than  
14  
15 for men ever worked in an occupation with occasional welding. We observed a slightly  
16  
17 stronger risk in never and light smokers and a trend with duration of employment in welding-  
18  
19 related jobs, as well as a stronger association of welding with SqCC and SCLC than with  
20  
21 AdCa. The findings from this investigation support the need for additional research to identify  
22  
23 the agent(s) responsible for possible lung cancer risks.  
24  
25  
26  
27  
28  
29  
30

## 31 REFERENCES

- 32  
33  
34 1 IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Chromium,  
35 Nickel and Welding. *IARC Monogr Eval Carcinog Risks to Hum.* 1990;49:1-36.
- 36  
37 2 Ambroise D, Wild P, Moulin JJ. Update of a meta-analysis on lung cancer and welding.  
38 *Scand J Work Environ Health.* 2006;32(1):22-31.
- 39  
40 3 Pukkala E, Martinsen JJ, Lynge E, et al. Occupation and cancer - follow-up of 15  
41 million people in five Nordic countries. *Acta Oncol.* 2009;48:646-790.
- 42  
43 4 Ward EM, Schulte PA, Straif K, et al. Research Recommendations for Selected IARC-  
44 Classified Agents. *Environ Health Perspect.* 2010;118(10):1355-62.
- 45  
46 5 Pesch B, Weiss T, Kendzia B, et al. Levels and predictors of airborne and internal  
47 exposure to manganese and iron among welders. *J Expo Sci Environ Epidemiol.*  
48 2012;22(3):291-8.
- 49  
50 6 Weiss T, Pesch B, Lotz A, et al. Levels and predictors of airborne and internal exposure  
51 to chromium and nickel among welders-Results of the WELDOX study. *Int J Hyg*  
52 *Environ Health.* 2013;216(2):175-83.  
53  
54  
55  
56  
57  
58  
59  
60

- 1
- 2
- 3 7 Lehnert M, Pesch B, Lotz A, et al. Exposure to inhalable, respirable, and ultrafine
- 4 particles in welding fume. *Ann Occup Hyg.* 2012;56(5):557-67.
- 5
- 6 8 Pesch B, Kendzia B, Gustavsson P, et al. Cigarette smoking and lung cancer-relative
- 7 risk estimates for the major histological types from a pooled analysis of case-control
- 8 studies. *Int J Cancer.* 2012;131(5):1210-9.
- 9
- 10 9 Olsson AC, Gustavsson P, Kromhout H, et al. Exposure to diesel motor exhaust and
- 11 lung cancer risk in a pooled analysis from case-control studies in Europe and Canada.
- 12 *Am J Respir Crit Care Med.* 2011;183(7):941-8.
- 13
- 14 10 International Labour Office. *International Standard Classification of Occupations.*
- 15 Geneva, Switzerland: ILO; 1968.
- 16
- 17 11 United Nations.Statistical Office. *International Standard Industrial Classification of all*
- 18 *Economic Activities.* New York: United Nations; 1968.
- 19
- 20 12 Ahrens W, Merletti F. A standard tool for the analysis of occupational lung cancer in
- 21 epidemiologic studies. *Int J Occup Environ Health.* 1998;4(4):236-40.
- 22
- 23 13 Mirabelli D, Chiusolo M, Calisti R, et al. Database of occupations and industrial
- 24 activities that involve the risk of pulmonary tumors. *Epidemiol Prev.* 2001;25(4-5):215-
- 25 21.
- 26
- 27 14 Jöckel KH, Ahrens W, Pohlabeln H, et al. Lung cancer risk and welding: Results from a
- 28 case-control study in Germany. *Am J Ind Med.* 1998;33(4):313-20.
- 29
- 30 15 Gustavsson P, Jakobsson R, Nyberg F, et al. Occupational exposure and lung cancer
- 31 risk: a population-based case-referent study in Sweden. *Am J Epidemiol.*
- 32 2000;152(1):32-40.
- 33
- 34 16 Lillienberg L, Zock JP, Kromhout H, et al. A population-based study on welding
- 35 exposures at work and respiratory symptoms. *Ann Occup Hyg.* 2008;52(2):107-15.
- 36
- 37 17 National Institute for Occupational Safety and Health. *Criteria for a recommended*
- 38 *standard - Welding, Brazing, and Thermal Cutting.* Cincinnati, OH: US Department of
- 39 Health and Humas Service, Centers for Disease Control and Prevention, National
- 40 Institute for Occupational Safety and Health; 1988.
- 41
- 42 18 Virta RL. *Worldwide Asbestos Supply and Consumption Trends from 1900 through*
- 43 *2003.* Reston, VA: U.S. Geological Survey; 2006. Report No.: 03-083, Circular 1298.
- 44
- 45 19 't Mannetje A., Brennan P, Zaridze D, et al. Welding and lung cancer in Central and
- 46 Eastern Europe and the United Kingdom. *Am J Epidemiol.* 2012;175(7):706-14.
- 47
- 48 20 Vallieres E, Pintos J, Lavoue J, et al. Exposure to welding fumes increases lung cancer
- 49 risk among light smokers but not among heavy smokers: evidence from two case-
- 50 control studies in Montreal. *Cancer Med.* 2012;1(1):47-58.
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
- 21 Poeschel E, Köhling A. *Umweltforschungsplan des Bundesministers des Innern - Luftreinhaltung - Asbestersatzstoff-Katalog, Erhebung über im Handel verfügbare Substitute für Asbest und asbesthaltige Produkte, Band 5: Elektroisolation*. Sankt Augustin, Germany: Hauptverband der gewerblichen Berufsgenossenschaften e.V. (HVBG); 1985. Report No.: 104 08 311.
- 22 Rolland P, Gramond C, Lacourt A, et al. Occupations and industries in France at high risk for pleural mesothelioma: A population-based case-control study (1998-2002). *Am J Ind Med*. 2010;53(12):1207-19.
- 23 Neumann V, Gunthe S, Mülle KM, et al. Malignant mesothelioma-German mesothelioma register 1987-1999. *Int Arch Occup Environ Health*. 2001;74(6):383-95.
- 24 Calvert GM, Luckhaupt S, Lee SJ, et al. Lung cancer risk among construction workers in California, 1988-2007. *Am J Ind Med*. 2012;55(5):412-22.
- 25 Siew SS, Kauppinen T, Kyyronen P, et al. Exposure to iron and welding fumes and the risk of lung cancer. *Scand J Work Environ Health*. 2008;34(6):444-50.
- 26 Corbin M, McLean D, 't Mannetje A, et al. Lung Cancer and Occupation: A New Zealand Cancer Registry-Based Case-Control Study. *Am J Ind Med*. 2011;54(2):89-101.
- 27 Richiardi L, Boffetta P, Simonato L, et al. Occupational risk factors for lung cancer in men and women: a population-based case-control study in Italy. *Cancer Causes Control*. 2004;15(3):285-94.
- 28 Guida F, Papadopoulos A, Menvielle G, et al. Risk of lung cancer and occupational history: results of a French population-based case-control study, the ICARE study. *J Occup Environ Med*. 2011;53(9):1068-77.
- 29 Tse LA, Yu IT, Qiu H, et al. Occupational risks and lung cancer burden for Chinese men: a population-based case-referent study. *Cancer Causes Control*. 2012;23(1):121-31.
- 30 Taeger D, Fritsch A, Wiethage T, et al. Role of exposure to radon and silicosis on the cell type of lung carcinoma in German uranium miners. *Cancer*. 2006;106(4):881-9.



**Table 1.** Distribution of various characteristics of the study population according to case-control and welding exposure status, SYNERGY, 1985-2010

Characteristics	Welders				Occasional welding tasks				Never worked in welding-related occupations			
	Controls		Cases		Controls		Cases		Controls		Cases	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
No. of participants	427	2.3	568	3.7	1,930	10.5	1,994	12.9	16,031	87.2	12,921	83.5
Age [years] <sup>a</sup>	62 (55-68)		61 (55-68)		62 (55-68)		62 (55-68)		64 (56-70)		64 (57-70)	
Packyears in smokers <sup>a</sup>	25 (14-40)		39 (28-54)		24 (11-38)		37 (26-50)		24 (11-40)		40 (27-56)	
Duration of employment in welding <sup>a</sup>	10 (3-22)		12 (4-26)		10 (3-25)		10 (3-27)		0		0	
Smoking status of cigarettes												
Never smokers	78	18.3	15	2.6	408	21.1	48	2.4	4,391	27.4	439	3.4
Former smokers	194	45.4	185	32.6	901	46.7	704	35.3	6,995	43.6	4,533	35.1
Current smokers	153	35.8	366	64.4	577	29.9	1,228	61.6	4,268	26.6	7,805	60.4
Other types of tobacco only	2	0.5	2	0.4	44	2.3	14	0.7	377	2.4	144	1.1
“List A” occupations except welding tasks												
Never	314	73.5	394	69.4	1,630	84.5	1,599	80.2	14,765	92.1	11,324	87.6
Ever	113	26.5	174	30.6	300	15.5	395	19.8	1,266	7.9	1,597	12.4
Histological subtype												
Adenocarcinoma	-		132	23.2	-		510	25.6	-		3,339	25.8
Squamous cell cancer	-		264	46.5	-		812	40.7	-		5,294	41.0
Small cell lung cancer	-		92	16.2	-		314	15.7	-		2,005	15.5
Other or mixed	-		80	14.1	-		358	18.0	-		2,283	17.7

<sup>a</sup> Values are expressed as median (interquartile range).

**Table 2.** Lung cancer risk among workers in welding-related occupations, SYNERGY, 1985-2010

Occupations	No. Controls	No. Cases	OR1 <sup>a</sup>	OR2 <sup>b</sup>	CI	OR3 <sup>c</sup>	CI
Reference group <sup>d</sup>	16,031	12,921	1.00	1.00		1.00	
Welders							
Ever	427	568	1.69	1.45	1.25, 1.68	1.44	1.25, 1.67
Longest held occupation	172	246	1.78	1.48	1.19, 1.86	1.50	1.20, 1.88
Occasional welding tasks							
Ever	1,930	1,994	1.27	1.18	1.10, 1.28	1.19	1.10, 1.28
Longest held occupation	697	746	1.37	1.31	1.16, 1.48	1.32	1.17, 1.49
Never employed in a “List A” job							
Never worked in welding-related occupations	14,765	11,323	1.00	1.00			
Welders							
Ever	314	394	1.70	1.46	1.23, 1.74		
Longest held occupation	129	186	1.92	1.63	1.26, 2.11		
Occasional welding tasks							
Ever	1,630	1,599	1.28	1.18	1.09, 1.29		
Longest held occupation	623	627	1.37	1.27	1.11, 1.45		
Ever employed in a “blue collar” job							
Never worked in welding-related occupations	10,289	9,796	1.00	1.00		1.00	
Welders							
Ever	427	568	1.45	1.32	1.14, 1.53	1.33	1.15, 1.54
Longest held occupation	172	246	1.55	1.36	1.09, 1.71	1.39	1.11, 1.73
Occasional welding tasks							
Ever	1,930	1,994	1.08	1.06	0.98, 1.15	1.07	0.99, 1.16
Longest held occupation	697	746	1.16	1.18	1.04, 1.33	1.20	1.06, 1.36

Abbreviations: CI, 95% confidence interval; OR, odds ratio.

<sup>a</sup> OR1 is adjusted for log(age) and study centers.

<sup>b</sup> OR2 is additionally adjusted for log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever other types of tobacco only,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

stopping smoking 2-7, 8-15, 16-25, 26+ years before interview/diagnosis, never smokers).  
<sup>c</sup> OR3 is additionally adjusted for ever working in a “List A” job except welding.  
<sup>d</sup> Defined as subjects, who had never worked in welding-related occupations.

For Peer Review

**Table 3.** Lung cancer risk among workers in welding-related occupations by industry, SYNERGY, 1985-2010

Industries	Ever-held job						Longest held job					
	No. Controls	No. Cases	OR1 <sup>a</sup>	OR2 <sup>b</sup>	OR3 <sup>c</sup>	CI	No. Controls	No. Cases	OR1 <sup>a</sup>	OR2 <sup>b</sup>	OR3 <sup>c</sup>	CI
Reference group <sup>d</sup>	16,031	12,921	1.00	1.00	1.00		16,031	12,921	1.00	1.00	1.00	
Welders												
Ship building and repair	59	93	1.99	1.57	1.53	1.06, 2.21	15	33	2.50	1.73	1.53	0.89, 3.41
Construction and related building services	240	336	1.78	1.50	1.47	1.22, 1.78	32	46	1.67	1.31	1.33	0.81, 2.20
Manufacture of machines, equipment, appliances	271	352	1.65	1.40	1.40	1.17, 1.68	57	104	2.38	2.08	2.11	1.45, 3.08
Manufacture of motor vehicles and motor bikes	93	102	1.56	1.33	1.30	0.94, 1.80	23	12	0.75	0.62	0.62	0.28, 1.36
Repair of transport equipments	101	136	1.70	1.51	1.51	1.12, 2.03	12	16	1.49	1.14	1.10	0.49, 2.46
Others	13	22	1.98	2.27	2.31	0.99, 5.39	33	35	1.29	1.25	1.27	0.74, 2.20
Occasional welding tasks												
Ship building and repair	132	132	1.26	1.17	0.90	0.68, 1.20	47	45	1.77	1.73	1.32	0.78, 2.23
Construction and related building services	1,152	1,238	1.36	1.24	1.21	1.09, 1.33	216	244	1.53	1.44	1.43	1.21, 1.67
Manufacture of machines, equipment,	739	734	1.20	1.17	1.14	1.01, 1.28	132	136	1.33	1.30	1.28	1.02, 1.60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

---

appliances													
Manufacture of motor vehicles and motor bikes	264	228	1.11	0.98	0.95	0.77, 1.16	69	60	1.28	1.04	1.03	0.71, 1.49	
Repair of transport equipments	843	835	1.22	1.14	1.11	0.99, 1.24	170	206	1.23	1.20	1.18	0.98, 1.44	
Others	112	101	1.05	0.89	0.91	0.67, 1.24	63	55	1.08	0.99	1.01	0.66, 1.54	

---

Abbreviations: CI, 95% confidence interval; OR, odds ratio.

<sup>a</sup> OR1 is adjusted for log(age) and study centers.

<sup>b</sup> OR2 is additionally adjusted for log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever other types of tobacco only, stopping smoking 2-7, 8-15, 16-25, 26+ years before interview/diagnosis, never smokers).

<sup>c</sup> OR3 is additionally adjusted for ever working in a “List A” job except welding.

<sup>d</sup> Defined as subjects, who had never worked in welding-related occupations.

Peer Review

**Table 4.** Lung cancer risk of welding-related occupations by histological subtype and duration of employment, SYNERGY, 1985-2010

Duration	No. Controls	All cases			AdCa			SqCC			SCLC		
		No.	OR <sup>3a</sup>	CI	No.	OR <sup>3a</sup>	CI	No.	OR <sup>3a</sup>	CI	No.	OR <sup>3a</sup>	CI
Reference group <sup>c</sup>	16,031	12,921	1.00		3,313	1.00		5,226	1.00		1,979	1.00	
Welders													
	427	568	1.44	1.25, 1.67	132	1.23	0.99, 1.53	264	1.58	1.32, 1.89	92	1.41	1.09, 1.82
1-<3	84	82	1.14	0.80, 1.61	18	0.84	0.49, 1.45	41	1.38	0.90, 2.11	14	1.25	0.67, 2.35
3-<10	124	171	1.46	1.26, 1.91	39	1.14	0.77, 1.68	77	1.62	1.16, 2.25	32	1.49	0.96, 2.32
10-≤25	129	167	1.38	1.06, 1.79	41	1.26	0.85, 1.87	76	1.34	0.97, 1.85	28	1.30	0.82, 2.07
> 25	90	148	1.77	1.31, 2.39	34	1.31	0.85, 2.02	70	1.71	1.19, 2.46	18	1.20	0.69, 2.11
<i>P</i> <sup>b</sup>				<0.0001			0.1041			0.0002			0.1311
Occasional welding tasks													
	1,930	1,994	1.19	1.10, 1.28	510	1.22	1.09, 1.37	812	1.14	1.03, 1.25	314	1.09	0.94, 1.25
1-<3	323	333	1.13	0.94, 1.34	93	1.26	0.98, 1.62	141	1.17	0.93, 1.48	49	0.98	0.70, 1.37
3-<10	642	638	1.11	1.00, 1.24	153	1.11	0.91, 1.35	241	0.90	0.76, 1.06	117	1.05	0.84, 1.31
10-≤25	485	487	1.16	1.00, 1.34	131	1.18	0.95, 1.46	207	1.14	0.94, 1.37	75	1.05	0.80, 1.38
> 25	480	536	1.40	1.21, 1.62	133	1.28	1.03, 1.58	223	1.37	1.13, 1.65	73	1.15	0.87, 1.51
<i>P</i> <sup>b</sup>				<0.0001			0.0193			0.0011			0.5687

Abbreviations: AdCa, Adenocarcinoma; CI, 95% confidence interval; OR, odds ratio; SCLC, Small cell lung cancer; SqCC, Squamous cell cancer.

<sup>a</sup> OR<sup>3</sup> is adjusted for log(age), study centers, log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever other types of tobacco only, stopping smoking 2-7, 8-15, 16-25, 26+ years before interview/diagnosis, never smokers), and for ever working in a "List A" job.

<sup>b</sup> Trend-test P values were computed by entering the continuous variable duration of employment into the model.

<sup>c</sup> Defined as subjects, who had never worked in welding-related occupations.

**Table 5.** Lung cancer risk of welding-related occupations by histological subtype and smoking, SYNERGY, 1985-2010

Occupations	No. Controls	No.	All cases		No.	AdCa		No.	SqCC		No.	SCLC	
			OR3 <sup>a</sup>	CI		OR3 <sup>a</sup>	CI		OR3 <sup>a</sup>	CI		OR3 <sup>a</sup>	CI
Reference group <sup>b</sup>	16,031	12,921	1.00		3,313	1.00		5,226	1.00		1,979	1.00	
Welders	427	568	1.44	1.25, 1.67	132	1.23	0.99, 1.53	264	1.58	1.32, 1.89	92	1.41	1.09, 1.82
Smoking status													
Never	78	15	2.34	1.31, 4.17	6	1.89	0.79, 4.52	4	3.01	1.07, 8.49	2	4.45	1.03, 19.2
Ever	349	553	1.33	1.14, 1.54	126	1.12	0.90, 1.41	260	1.49	1.24, 1.78	90	1.31	1.01, 1.70
Packyears													
0-<10	146	47	1.96	1.37, 2.79	12	1.37	0.74, 2.52	18	2.25	1.30, 3.91	7	2.28	0.99, 5.29
10-35	174	189	1.19	0.95, 1.49	45	1.07	0.75, 1.52	86	1.35	1.01, 1.79	35	1.36	0.91, 2.04
≥35	107	332	1.34	1.06, 1.69	75	1.15	0.84, 1.59	160	1.50	1.16, 1.96	50	1.12	0.78, 1.62
Occasional welding tasks	1,930	1,994	1.19	1.10, 1.28	510	1.22	1.09, 1.37	812	1.14	1.03, 1.25	314	1.09	0.94, 1.25
Smoking status													
Never	408	48	1.31	0.95, 1.81	25	1.16	0.76, 1.78	8	1.13	0.54, 2.37	2	0.76	0.18, 3.19
Ever	1,522	1,946	1.15	1.07, 1.25	485	1.19	1.06, 1.34	804	1.11	1.00, 1.23	312	1.06	0.92, 1.23
Packyears													
0-<10	784	166	1.20	0.99, 1.45	62	1.31	0.98-1.74	53	1.02	0.74, 1.41	22	1.14	0.70, 1.85
10-35	699	767	1.20	1.07, 1.35	190	1.16	0.97, 1.39	313	1.20	1.03, 1.40	122	1.09	0.87, 1.36
≥35	447	1,061	1.09	0.96, 1.24	258	1.16	0.96, 1.39	446	1.06	0.91, 1.23	170	1.01	0.82, 1.24

Abbreviations: AdCa, Adenocarcinoma; CI, 95% confidence interval; OR, odds ratio; SCLC, Small cell lung cancer; SqCC, Squamous cell cancer.

<sup>a</sup> OR3 is adjusted for log(age), study centers, log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever other types of tobacco only, stopping smoking 2-7, 8-15, 16-25, 26+ years before interview/diagnosis, never smokers), and for ever working in a "List A" job.

Smoking variables were omitted when analyzing never smokers.

<sup>b</sup> Defined as subjects, who had never worked in welding-related occupations.

1  
2  
3  
4  
5  
6  
7  
8  
9 **Figure 1.** Study-specific odds ratios (OR3s) with 95% confidence interval (CI) for ever working as a welder (A) or in occasional welding  
10 (B) compared with men who never worked in welding-related occupations adjusted for log(age), study center, log(packyears+1), time-since-  
11 quitting smoking cigarettes, and for ever working in a “List A” occupation except welding  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49



## WEB MATERIAL

**Table S1.** Characteristics of the studies selected for the pooled analysis of lung cancer risk among male welders, SYNERGY, 1985-2010

Study	Country	Recruitment period	Job periods	Cases		Controls		Source of controls	Type of interview
				No. <sup>c</sup>	RR % <sup>b</sup>	No. <sup>c</sup>	RR % <sup>b</sup>		
AUT	Germany	1990–1995	1931–1995	2,659	76	2,702	41	P	Face-to-face
HdA	Germany	1988–1993	1926–1993	839	69	838	68	P	Face-to-face
EAGLE	Italy	2002–2005	1931–2005	1,536	87	1,617	72	P	Face-to-face
TURIN/VENETO	Italy	1990–1994	1922–1994	955	79	1,249	80	P	Face-to-face
ROME	Italy	1993–1996	1926–1996	294	74	264	63	H	Face-to-face
LUCA	France	1989–1992	1926–1992	307	98	302	98	H	Face-to-face
PARIS	France	1988–1992	1923–1992	161	95	219	95	H	Face-to-face
ICARE	France	2001–2007	1937–2007	2,243	80	2,749	76	P	Face-to-face <sup>a</sup>
CAPUA	Spain	2000–2010	1926–2010	774	91	712	96	H	Face-to-face
MORGEN	NL	1993–1997	1945–1997	41	N/A	111	N/A	P	Self-administered questionnaire
INCO	CZ	1999–2002	1935–2002	235	94	294	80	H	Face-to-face
INCO	Hungary	1998–2001	1931–2001	313	90	250	100	H	Face-to-face
INCO	Poland	1998–2002	1936–2002	551	88	574	88	H/P	Face-to-face
INCO	Slovakia	1998–2002	1935–2002	288	90	236	84	H	Face-to-face
INCO	Romania	1998–2002	1936–2002	141	90	152	99	H	Face-to-face
INCO	Russia	1998–2001	1931–2001	521	96	503	90	H	Face-to-face
INCO (LLP)	UK	1998–2005	1932–2005	281	78	573	84	P	Face-to-face
LUCAS	Sweden	1985–1990	1923–1990	1,018	87	2,309	85	P	Mail, telephone <sup>a</sup>
OCANZ	NZ	2003–2009	1941–2009	212	53	417	48	P	Face-to-face, telephone <sup>a</sup>
MONTREAL	Canada	1996–2002	1933–2002	714	85	895	69	P	Face-to-face <sup>a</sup>
TORONTO	Canada	1997–2002	1929–2002	198	62	371	84/60	H/P	Face-to-face
HONG KONG	China	2003-2007	1931–2007	1,202	96	1,051	48	P	Face-to-face, telephone <sup>a</sup>
Overall		1985–2010	1922–2010	15,483	82	18,388	74		

1  
2  
3  
4  
5 Abbreviations: CZ, Czech Republic; H, hospital controls; N/A, not applicable; NL, Netherlands; NZ, New Zealand; P, population controls; RR, response rate.

6 <sup>a</sup> Interviews also with a next of kin.

7 <sup>b</sup> Response rate was reported for both genders.

8 <sup>c</sup> Men with complete occupational and smoking history.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

For Peer Review

**Table S2.** Classification of welders and occupations with potential welding activities according to the International Standard Classification of Occupations (ISCO) 1968, SYNERGY, 1985-2010

ISCO Code	Description
<b>Welders</b>	
87200	Welders
87210	Gas and electric welders (general)
87215	Gas welders
87220	Electric arc welders (hand)
87225	Electric arc welders (machine)
87230	Thermite arc welders
87235	Resistance welders
<b>Occupations with potential welding activities</b>	
87105	Plumbers
87110	Pipe fitters
87120	Gas pipe fitters
87130	Marine pipe fitters
87140	Aircraft pipe and tube fitters
87190	Other plumbers and pipe fitters
87310	Sheet-metal workers
87350	Boilersmith
87360	Ornamental sheet-metal workers
87370	Vehicle sheet-metal workers
87380	Aircraft sheet-metal workers
87390	Other sheet-metal workers
87400	Structural metal preparers and erectors
87430	Structural steel workers (workshop)
87440	Constructional steel erectors
87450	Metal shipwright
87490	Other structural metal preparers and erectors
95350	Metal roofers
84185	Airframe fitter-assemblers
84190	Other machinery fitters and machine assemblers
84910	Machinery mechanics (general)
84920	Diesel engine mechanics (except motor vehicle)
84955	Agricultural machinery mechanics
84300	Motor-vehicle mechanics
84320	Automobile mechanics
84330	Motor-truck mechanics
84340	Motor-cycle mechanics
84390	Other motor-vehicle mechanics
84400	Aircraft engine mechanics
84410	Aircraft engine mechanics (general)
84420	Aircraft engine service mechanics

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

84490	Other aircraft engine mechanics
87290	Other welders and flame-cutters
72700	Metal drawers and extruders
83500	Metal grinders, polishers and tool sharpeners
85320	Electrical equipment assemblers

---

For Peer Review

**Table S3.** Classification of welders in industries according to the International Standard Industrial Classification (ISIC) revision 2, SYNERGY, 1985-2010

ISIC Code	Description
<b>Construction, plumbers and other building services</b>	
5000	Construction
9100	Public administration and defence
4101	Education services
9310	Electric light and power
4102	Gas manufacture and distribution
4103	Steam and hot water supply
6320	Hotels, rooming houses, camps and other lodging places
8324	Engineering, architectural and technical services
8310	Real estate
<b>Manufacture of machines, equipment, appliances</b>	
3819	Manufacture of fabricated metal products except machinery and equipment
3813	Manufacture of structural metal products
3829	Machinery and equipment except electrical not elsewhere classified
3811	Manufacture of cutlery, hand tools and general hardware
3824	Manufacture of special industrial machinery and equipment except metal and wood working machinery
3800	Manufacture of fabricated Metal Products, Machinery and Equipment
3821	Manufacture of engines and turbines
3822	Manufacture of agricultural machinery and equipment
3833	Manufacture of electrical appliances and housewares
3831	Manufacture of electrical industrial machinery and apparatus
3812	Manufacture of furniture and fixtures primarily of metal
3823	Manufacture of metal and wood working machinery
3851	Manufacture of professional and scientific, and measuring and controlling equipment
3830	Manufacture of electrical machinery apparatus, appliances and supplies
3832	Manufacture of radio, television and communication equipment and apparatus
3839	Manufacture of electrical apparatus and supplies not elsewhere classified
3810	Manufacture of fabricated metal products, except machinery and equipment
3820	Manufacture of machinery except electrical
3825	Manufacture of office, computing and accounting machinery
<b>Manufacture of motor vehicles, motor bikes, bikes</b>	
3843	Manufacture of motor vehicles
3844	Manufacture of motorcycles and bicycles
<b>Ship building and repairing</b>	
3841	Ship building and repairing
<b>Repair of transport equipment</b>	
3842	Manufacture of railroad equipment
3845	Manufacture of aircraft
6100	Wholesale trade
7111	Railway transport
9513	Repair of motor vehicles and motorcycles

---

1		
2		
3	7112	Urban, suburban and inter-urban highway passenger transport
4	7110	Land transport
5	3849	Manufacture of transport equipment not elsewhere classified
6	7100	Transport and storage
7	7131	Air transport carriers
8		
9	<b>Others</b>	
10	3710	Iron and steel basic industries
11	3699	Manufacture of non-metallic mineral products not elsewhere classified
12	1110	Agriculture and livestock production
13	3511	Manufacture of basic industrial chemicals except fertilizers
14	3530	Petroleum refineries
15	2100	Coal mining
16	3320	Manufacture of furniture and fixtures, except primarily of metal
17		
18	3513	Manufacture of synthetic resins, plastic materials and man-made fibres except glass
19		
20	3529	Manufacture of chemical products not elsewhere classified
21	3909	Manufacturing industries not elsewhere classified
22	2901	Stone quarrying, clay and sand pits
23	3720	Non-ferrous metal basic industries

---

**Table S4.** Sensitivity analyses for the association between welding and lung cancer in men, SYNERGY, 1985-2010

Occupations	Studies with population controls				Studies with hospital controls				Started working after 1979			
	No. Controls	No. Cases	OR3 <sup>a</sup>	CI	No. Controls	No. Cases	OR3 <sup>a</sup>	CI	No. Controls	No. Cases	OR3 <sup>a</sup>	CI
<b>Reference group<sup>b</sup></b>	13,100	10,366	1.00		2,931	3,196	1.00		551	229	1.00	
<b>Welders</b>	315	432	1.53	1.29, 1.82	112	162	1.10	0.84, 1.44	61	63	1.31	0.68, 2.58
<b>Occasional welding tasks</b>	1,584	1,651	1.17	1.08, 1.28	346	425	1.10	0.93, 1.30	115	101	1.51	0.94, 2.44

Abbreviations: CI, 95% confidence interval; OR, odds ratio.



<sup>a</sup> OR3 is adjusted for log(age), study centers, log(packyears+1), time-since-quitting smoking cigarettes (current smokers, ever other types of tobacco only, stopping smoking 2-7, 8-15, 16-25, 26+ years before interview/diagnosis, never smokers), and for ever working in a "List A" job.

<sup>b</sup> Defined as subjects, who had never worked in welding-related occupations.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

For Peer Review





54

55

56

57

58

59

60