

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

More than 20 years of registration of type 1 diabetes in Sardinian children: temporal variations of incidence with age, period of diagnosis, and year of birth.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/142146> since 2016-07-12T13:05:24Z

Published version:

DOI:10.2337/db12-1771

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Bruno G;Maule M;Biggeri A;Ledda A;Mannu C;Merletti F;Songini M;Sardinian Group for Diabetes Epidemiology. More than 20 years of registration of type 1 diabetes in Sardinian children: temporal variations of incidence with age, period of diagnosis, and year of birth.. DIABETES. 62 (10) pp: 3542-3546.
DOI: 10.2337/db12-1771

The publisher's version is available at:

<http://diabetes.diabetesjournals.org/cgi/doi/10.2337/db12-1771>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/142146>

More than twenty years of registration of type 1 diabetes in Sardinian children: temporal variations of incidence with age, period of diagnosis and year of birth

Running title: incidence time trend of type 1 diabetes in Sardinia

Graziella Bruno¹, Milena Maule^{1,2}, Annibale Biggeri³, Alessia Ledda⁴, Carla Mannu⁴, Franco Merletti^{1,2}, Marco Songini⁴ and Sardinian Group for Diabetes Epidemiology

¹University of Turin, Department of Medical Sciences

²Cancer Epidemiology Unit, CeRMS and CPO-Piemonte, Hospital Città della Salute e della Scienza, Turin, Italy

³Department of Statistics G. Parenti, University of Florence, Italy.

⁴Diabetes Unit, Cagliari, Italy.

Corresponding author and author to receive reprint request:

prof. Graziella Bruno, University of Turin, Department of Medical Sciences.

corso Dogliotti 14, I-10126 Turin, Italy

Tel: +39 11 633 6709

Fax: +39 11 6634 751

Email: graziella.bruno@unito.it

Word count: 2017

ABSTRACT

We analysed Sardinian Registry data to assess time trends in incidence rates of Type 1 diabetes, period 1989-2009 (2371 cases aged 0-14 years). Poisson regression models were used to estimate the effects of sex, age, period of diagnosis and birth cohorts. Incidence rate was 44.8 cases per 100,000 person-years (95% CI 43.1-46.7). The annual increase was 2.12% (1.45-2.80, test for linear trend: $p < 0.001$). For boys, the increasing trend was evident up to 5 years of age, and for girls up to 8 years. Compared with the 1989-94 birth cohort, the relative risk increased from 0.78 (0.61-1.10) in 1974-79 to 1.62 (1.18-2.23) in 2004-09. The increase over period was less striking, with a tendency to regress in more recent years. The best fitting model for boys included age and a linear time trend, for girls age and non-linear effects of calendar period and birth cohort. In conclusion, incidence increased over time, the increase tended to level off in more recent years by calendar period but not by birth cohort, with some evidence of a stronger increase among girls than boys. Should the increase be attributable to the effects of some perinatal environmental factor, this would mean that such a factor has started affecting females before males.

The most striking feature of type 1 diabetes epidemiology is the evidence of large geographical variation in disease risk (1-3). Sardinia has the second highest incidence rates in the world after Finland, and this excess likely started around 1950, as shown by the analysis of incidence in males conscript cohorts in the period 1936-1979 (4).

An increasing trend of type 1 diabetes, ranging around 2-3% has been shown in almost all areas covered by population-based registries (3). The increase involved both children and young adults in Turin, Northern Italy (5), whereas it was limited to children in Sweden, and interpreted as the anticipation of age at disease onset (6). Updated incidence data of registries included in the EURODIAB study have provided the interesting finding that the lowest risk areas of Eastern Europe underwent the highest yearly increase up to year 2004 while, in the highest risk areas of Northern Europe, the increasing trend was lower (7).

Aims of this report from the high risk Sardinia Region were 1) to update the analysis of incidence rates of type 1 diabetes up to 2009 in children aged 0-14 years; 2) to assess temporal trends over the twenty-one-year period 1989-2009; 3) to assess whether it was possible to disentangle the effects of age, calendar period of diagnosis and birth cohort and if such effects were heterogeneous by age and sex.

RESEARCH DESIGN AND METHODS

The study base of this report were all children aged 0 to 14 years resident in Sardinia in the period 1989-2009. As previously described (8), the registration of incident cases was performed according to the EURODIAB criteria. Type 1 diabetes was defined on the basis of clinical diagnosis of insulin-dependent diabetes by a doctor, excluding cases occurring secondary to other conditions. The date of onset was taken as the date of first insulin injection. Diabetes clinics for children and adults and departments of internal medicine and endocrinology were the first source of ascertainment. The second independent data source

was a membership of the local patient associations and the personal National Health System identification cards needed by each patient to obtain medications and devices free of charge. The estimated completeness of ascertainment was 92% in period 1989-97, 90% in period 1998-2003 and 91% in period 2004-09, with no differences among genders and age-groups over the study period .

Population data by sex and year of birth were provided by the Italian National Institute of Statistics (ISTAT, URL: <http://www.demo.istat.it/>). Age- and sex- specific incidence rates/100,000 person-years were calculated for calendar year of diagnosis and birth cohort (9). The chi-square test for trend in incidence rates used is the Mantel extension of the Armitage-Cochran trend test. The trend test is either stratified by sex or calculated separately for the two genders. Poisson regression models were used to estimate the effects of sex, age (five three-year age-groups: 0-2, 3-5, 6-8, 9-11, 12-14), calendar time (seven three-year periods: 1989-91, 1992-94, 1995-97, 1998-2000, 2001-03, 2004-06, 2007-09) and birth cohorts (eleven six-year birth cohorts: 1974-79, 1977-82, 1980-85, 1983-88, 1986-91, 1989-94, 1992-97, 1995-2000, 1998-2003, 2001-06, 2004-09). Six models were fitted to data: sex; sex, age; sex, age, linear time trend (drift); sex, age, cohort; sex, age, period; sex, age, period, cohort (online Figure 1). The latter five models were then fitted to data separately for the two genders. The term drift denotes a linear temporal variation of rates which does not distinguish between the influences of two of the three temporal variables involved in the analysis. The hierarchically ordered models were compared through the likelihood ratio test. Further Poisson regression models were fitted to data including sex (or separate models were fitted to the two genders), age, linear time trend (drift) and quadratic terms for period and cohort, after centering the 2 variables, in order to check if temporal variation was non linear. Statistical analyses were performed using STATA/SE, release 11.0.

RESULTS

In the period 1989-2009, 2371 incident cases of type 1 diabetes were identified among children aged 0-14 years resident in Sardinia. Incidence rates (IR) by sex, age group and calendar period are shown in Table 1. The IR over the study period was 44.8 cases per 100,000 person-years (95% confidence interval [CI] 43.1-46.7), with significantly higher risk in boys (50.6, 48.0-53.4) than in girls (38.7, 36.4-41.2). In the regression analysis, after controlling for age, the rate ratio (RR) for boys with respect to girls was 1.31 (1.21-1.42). IRs in the age groups 0-4, 5-9, 10-14 years were 37.8 (34.9-41.0), 47.0 (43.9-50.3) and 48.4 (45.4-51.5), respectively; corresponding RRs (boys vs. girls) were 1.22 (1.04-1.44), 1.18 (1.03-1.36) and 1.49 (1.31-1.69).

The IR steadily increased over the study period, from 35.8 (32.2-39.7) in 1989-91 to 51.0 (45.7-56.9) in 2007-09 (Table 1). Controlling for age and sex, the annual increase in the period 1989-2009 was 2.12% (1.45-2.80; test for linear trend: $p < 0.001$). Controlling for age, such an increase was higher in girls (2.67%, 1.63-3.73) than in boys (1.73%, 0.85-2.62), although not significantly.

Table 2 shows age-specific IRs by birth cohort and calendar period (diagonals), for all children and for boys and girls separately. The table shows how IRs increased with each subsequent birth cohort in each age group. The yearly increase, adjusted for age and sex, was highest in the age group 0-2 years (4.3%, $p < 0.001$). Time trends were heterogeneous between sexes: in boys, large statistically significant increasing linear trends were evident in the age group 0-2 years (4.4%, $p < 0.001$) and, to a smaller extent, in 3-5 years old (2.4%, $p = 0.034$), whereas in girls, a yearly increase of similar magnitude was evident not only in the age groups 0-2 years (4.2%, $p = 0.006$) and 3-5 (3.0%, $p = 0.013$) but also in the age group 6-8 (4.3%, $p < 0.001$).

A different pattern of increase in time by birth cohorts and by calendar periods of diagnosis is evident analyzing age-adjusted RRs, which are shown in Table 3. With respect to the 1989-94 birth cohort, the RR increased from 0.78 (0.61-1.10) for children born in 1974-79 to 1.62 (1.18-2.23) for those born in 2004-09. In contrast, the pattern of increase of RRs by calendar period of diagnosis was less striking. Indeed, with respect to children diagnosed in period 1998-2000, the RR increased from 0.73 (0.63-0.85) for children diagnosed in 1989-91 to 1.06 (0.91-1.23) for those diagnosed in 2007-09.

In order to try to disentangle the effect of age, calendar period and birth cohort on the overall temporal trend of the disease in period 1989-2009, we fitted hierarchically ordered Poisson regression models to the data, for all incident cases, and separately by gender, since there was evidence of interaction (sex*cohort, $p=0.018$; sex*period, $p=0.143$) (Table 4). For both boys and girls, using the Akaike information criteria, the best model (model 2) was the one with age and a linear time trend (drift). This finding means that the variation over time has a linear component which cannot be ascribed to either the calendar period or the cohort. However, for girls as well as for all children analyzed together, nonlinear effects of both birth cohort (model 4) and period of diagnosis (model 5) are apparent. Fitting models including quadratic terms for period and cohort showed that the temporal variation was non linear. Indeed, this model fitted to the total (boys+girls) data resulted in a slightly improvement of the Akaike information criteria vs model 2. Opposite signs for the coefficients of the quadratic period and cohort terms indicated that, whereas there was a steady increase with birth cohort, in the most recent periods incidence started to show a tendency to decrease. This result was already apparent in Table 2, showing that incidence rates in children diagnosed in 2007-09 (last diagonal from top right to bottom left, incidence rates: 43.6 in 0-2 years, 48.6 in 3-5 years, ...) were lower than those diagnosed in 2004-06 in almost all age groups (next diagonal on the left: 51.2 in 0-2 years, 54.0 in 3-5 years, ...).

CONCLUSIONS

Our analysis of incidence data from the Sardinia Registry of Type 1 diabetes in the twenty-one years period 1989-2009 provides evidence of a strong increase of incidence rates among Sardinian children. Results of our age-period-cohort analysis of time trends are intriguing. Indeed, we provide evidence that the incidence of type 1 diabetes has increased non linearly with the birth cohort. Such a birth cohort effect seems to have started some 5-10 years earlier among girls, starting from those born in 1980-85, than among boys, starting from those born in 1986-91. This finding allows us to explain why annual percent increases are higher and statistically significant in girls up to 8 years of age but affect boys to a lesser extent and up to 5 years of age only. Should the increase be attributable to the effects of some perinatal environmental factor, this would mean that such a factor has started affecting females before males. In Sardinia, both a period and a cohort effect were evident, with heterogeneities among sexes. Indeed, the nonlinear temporal effects were more evident among girls, with larger increases with time than among boys. In more recent years a tendency of time trends to level off was also evident, particularly in the youngest age groups, which could merely reflect year-to-year variability in incidence rates and need to be confirmed over a longer time period.

Recently Patterson and colleagues (10) have pointed out that the potential of age-period-cohort analysis in disentangling the effects of time trends on different age groups at different times is often hampered by the paucity of age classes and the limited length of time series. Indeed, incidence rates up to 14 years of age do not allow us to capture the total effect of time variations in the whole population. However, despite the small number of age groups and the limited sample size, our analysis, taking advantage of more than 20 years of registration,

shows that incidence has increased more than linearly with birth cohort, whereas the increase with calendar period has shown signs of regression in the most recent periods. Previous age-period-cohort analyses have found different and discordant results, including linear and nonlinear period effects (5, 11-13) as well as cohort effects (14-15), often interpreted as a shift towards earlier age at onset (7, 16-18).

Studies have proposed that the decreasing early life exposure to infectious diseases could be involved in the increasing temporal trends of immune-mediate disorders (19). Lower socioeconomic level and higher crowding in early life, proxy measures of early childhood infection, increase the risk of type 1 diabetes up to 3 years and decrease the risk in the following age groups (20). Experimental models have shown a relationship between intestinal microbioma and type 1 diabetes (21). As regards Sardinia, the frequencies of elmintiasis, which was endemic up to the 1980s, has now almost completely disappeared. The role of mycobacterium avium paratuberculosis infection has also been suggested to play a role, and an homology between its epitopes and beta cell antigen ZnT8 has been identified in Sardinian patients with type 1 diabetes (22).

Another intriguing and still unsolved issue in the epidemiology of type 1 diabetes is the heterogeneity of risk among sexes. Whereas in most of countries incidence is higher in males than in females among young adults only, Sardinian boys have a statistically significant 31% higher risk than girls of the same age, which is not chromosoma Y-linked (24), but until now no other hypothesis has been tested. Moreover, our data show heterogeneities among sexes in time trend patterns, with girls appearing to show the potential effects of hypothetical environmental factor exposures at least 5-10 years before boys. This finding is intriguing and further experimental and epidemiological studies should explore sex differences in the interaction between environment and genetic in the pathogenesis of type 1 diabetes.

In conclusion, our study showed a strong increase of incidence rates of type I diabetes among Sardinian children in the period 1989-2009, which was non linear with the birth cohort, whereas the increase with calendar period showed signs of regression in the most recent periods. The birth cohort effect seems to have started earlier among girls than among boys. Should the increase be attributable to the effects of some perinatal environmental factor, this would mean that such a factor has started affecting females before males.

Authors contribution: GB researched data and wrote the manuscript, MM researched data and wrote the manuscript, AB researched data and contributed to the discussion, AL researched data, CM researched data, FM contributed to the discussion and reviewed/edited manuscript, MS researched data, contributed to the discussion and reviewed/edited manuscript.

Acknowledgements: This study has been presented at the EASD meeting in September 2012, in Berlin.

The Authors declare to have no conflict of interest with respect to the study.

Guarantors of the data: Graziella Bruno and Marco Songini are the guarantor of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis."

References

1. Pundziute-Lyckå A, Dahlquist G, Urbonaite B, Zalinkevicius R; Swedish Childhood Diabetes Study Group; Lithuanian Childhood Diabetes Study Group. Time trend of childhood type 1 diabetes incidence in Lithuania and Sweden, 1983-2000. *Acta Paediatr* 2004;93:1519-24
2. Bruno G, Maule M, Merletti F, Novelli G, Falorni A, Iannilli A, Iughetti L, Altobelli E, D'Annunzio G, Piffer S, Pozzilli P, Iafusco D, Songini M, MD, Roncarolo F, Toni S, Cherubini V, Carle F; RIDI Study Group. Age-period-cohort analysis of 1990-2003 incidence time trends of childhood diabetes in Italy: the RIDI study. *Diabetes* 2010; 59:2281-2287
3. Soltész G, Patterson CC, Dahlquist G; EURODIAB Study Group. Worldwide childhood type 1 diabetes incidence. What can we learn from epidemiology? *Pediatr Diabetes* 2007;8(Suppl 6):6-14
4. Casu A, Pascutto C, Bernardinelli L, Songini M; Sardinian Conscript Type 1 Diabetes Registry. Bayesian approach to study the temporal trend and the geographical variation in the risk of type 1 diabetes. The Sardinian Conscript Type 1 Diabetes Registry. *Pediatr Diabetes* 2004;5:32-8
5. Bruno G, Novelli G, Panero F, Perotto M, Monasterolo F, Bona G, Perino A, Rabbone I, Cavallo-Perin P, Cerutti F, and Piedmont Study Group for Diabetes Epidemiology. Incidence of type 1 diabetes is increasing both in children and young adults in Northern Italy: 1984-2004 temporal trend. *Diabetologia* 2009;52:2531-2535
6. Dahlquist GG, Nyström L, Patterson CC; Swedish Childhood Diabetes Study Group; Diabetes Incidence in Sweden Study Group. Incidence of type 1 diabetes in Sweden among individuals aged 0-34 years, 1983-2007: an analysis of time trends. *Diabetes Care* 2011;34:1754-9
7. Patterson CC, Dahlquist GG, Gyürüs E, Green A, Soltész G; EURODIAB Study Group. Incidence trends for childhood type 1 diabetes in Europe during 1989-2003 and predicted new cases 2005-20: a multicentre prospective registration study. *Lancet* 2009;373:2027-33
8. Casu A, Pascutto C, Bernardinelli L, Songini M. Type 1 diabetes among Sardinian children is increasing: the Sardinian diabetes register for children aged 0-14 years (1989-1999). *Diabetes Care* 2004;27:1623-9
9. Carstensen B. Age-period-cohort models for the Lexis diagram. *Stat Med* 2007; 6:3018-45
10. Patterson CC, Gyürüs E, Rosenbauer J, Cinek O, Neu A, Schober E, Parslow RC, Joner G, Svensson J, Castell C, Bingley PJ, Schoenle E, Jarosz-Chobot P, Urbonaitė B, Rothe U, Krzisnik C, Ionescu-Tirgoviste C, Weets I, Kocova M, Stipancic G, Samardzic M, de Beaufort CE, Green A, Dahlquist GG, Soltész G. Trends in childhood type 1 diabetes incidence in Europe during 1989-2008: evidence of non-uniformity over time in rates of increase. *Diabetologia* 2012; 55:2142-7
11. Nyström L, Dahlquist G, Rewers M, Wall S. The Swedish Childhood Diabetes Study. An analysis of the temporal variation in diabetes incidence 1978-1987. *Int J Epidemiol* 1990;19:141-146
12. Tuomilehto J, Rewers M, Reunanen A, Lounamaa P, Lounamaa R, Tuomilehto-Wolf E, Akerblom HK. Increasing trend in Type 1 (insulin-dependent) diabetes mellitus in childhood in Finland. Analysis of age, calendar time and birth cohort effects during 1965 to 1984. *Diabetologia* 1991;34:282-287

13. Aamodt G, Stene LC, Njølstad PR, Søvik O, Joner G; The Norwegian Childhood Diabetes Study Group. Spatiotemporal trends and age-period-cohort modeling of the incidence of type 1 diabetes among children aged <15 years in Norway 1973-1982 and 1989-2003. *Diabetes Care* 2007;30:884-889
14. Svensson J, Carstensen B, Mølbak A, Christau B, Mortensen HB, Nerup J, Borch-Johnsen K. Increased risk of childhood type 1 diabetes in children born after 1985. *Diabetes Care* 2002;25:2197-2201
15. Berhan Y, Waernbaum I, Lind T, Möllsten A, Dahlquist G; Swedish Childhood Diabetes Study Group. Thirty years of prospective nationwide incidence of childhood type 1 diabetes: the accelerating increase by time tends to level off in Sweden. *Diabetes* 2011;60:577-81
16. Feltbower RG, McKinney PA, Parslow RC, Stephenson CR, Bodansky HJ. Type 1 diabetes in Yorkshire, UK: time trends in 0-14 and 15-29-year-olds, age at onset and age-period-cohort modelling. *Diabet Med* 2003;20:437-441
17. Dahlquist G, Mustonen L. Analysis of 20 years of prospective registration of childhood onset diabetes time trends and birth cohort effects. Swedish Childhood Diabetes Study Group. *Acta Paediatr* 2000;89:1231-1237
18. Pundziute-Lyckå A, Dahlquist G, Nyström L, Arnqvist H, Björk E, Blohmé G, Bolinder J, Eriksson JW, Sundkvist G, Ostman J; Swedish Childhood Diabetes Study Group. The incidence of Type I diabetes has not increased but shifted to a younger age at diagnosis in the 0-34 years group in Sweden 1983-1998. *Diabetologia* 2002;45:783-791
19. Rook GA. Review series on helminths, immune modulation and the hygiene hypothesis: the broader implications of the hygiene hypothesis. *Immunology* 2009;126:3-11
20. Bruno G, Spadea T, Picariello R, Gruden G, Barutta F, Cerutti F, Cavallo-Perin p, Costa G, Gnani R, Piedmont Study Group for Diabetes Epidemiology. Early life socioeconomic indicators and the risk of type 1 diabetes in children and young adults. *J Pediatrics* 2013; 162:600-605
21. Vaarala O. Is the origin of type 1 diabetes in the gut? *Immunol Cell Biol* 2012 90:271-6
22. Masala S, Paccagnini D, Cossu D, Brezar V, Pacifico A, Ahmed N, Mallone R, Sechi LA. Antibodies recognizing *Mycobacterium avium* paratuberculosis epitopes cross-react with the beta-cell antigen ZnT8 in Sardinian type 1 diabetic patients. *PLoS One*. 2011;6:e26931
23. Contu D, Morelli L, Zavattari P, Lampis R, Angius E, Frongia P, Murru D, Maioli M, Francalacci P, Todd JA, Cucca F. Sex-related bias and exclusion mapping of the nonrecombinant portion of chromosome Y in human type 1 diabetes in the isolated founder population of Sardinia. *Diabetes* 2002;51:3573-6

Table 1: Incidence rates of type 1 diabetes among Sardinian children 0 to 14 years old in the years 1989-2009 by sex, age group, calendar period.

		Cases N.	Person-years	incidence rate (per 100,000 person-years)	95% CI	
total		2371	5,286,212	44.8	43.1	46.7
sex	boys	1380	2,725,601	50.6	48.0	53.4
	girls	991	2,560,611	38.7	36.4	41.2
age	0-2	320	910,143	35.2	31.5	39.2
	3-5	396	961,497	41.2	37.3	45.5
	6-8	482	1,034,124	46.6	42.6	51.0
	9-11	613	1,127,595	54.4	50.2	58.8
	12-14	560	1,252,853	44.7	41.2	48.6
period	1989-91	346	967,116	35.8	32.2	39.8
	1992-94	334	857,969	38.9	35.0	43.3
	1995-97	362	785,852	46.1	41.6	51.1
	1998-00	351	723,217	48.5	43.7	53.9
	2001-03	311	678,325	45.8	41.0	51.2
	2004-06	348	648,162	53.7	48.3	59.6
	2007-09	319	625,571	51.0	46.0	56.9

Table 2 Age-specific incidence rates (per 100,000 person-years) of type 1 diabetes among Sardinian children 0 to 14 years old in the years 1989-2009, by birth cohorts (1974-79, ..., 2004-09) and by calendar periods (diagonals: 1989-91, ..., 2007-09). Number of cases are in parentheses. Percent yearly increases and test for age specific time trends are shown in the last columns (adjusted by sex, or separately for the 2 genders).

	Age	Birth cohort											% yearly increase	p
		1974-79	1977-82	1980-85	1983-88	1986-91	1989-94	1992-97	1995-00	1998-03	2001-06	2004-09		
total	0-2					22.8 (35)	24.5 (35)	32.4 (43)	36.8 (45)	40.8 (49)	51.2 (61)	43.6 (52)	4.3	<0.001
	3-5				30.9 (52)	34.8 (53)	40.2 (58)	44.0 (58)	41.2 (50)	54.0 (66)	48.6 (59)		2.7	0.001
	6-8			33.8 (64)	45.3 (75)	49.5 (76)	45.1 (66)	42.6 (56)	63.2 (78)	53.9 (67)			2.3	0.002
	9-11		50.6 (107)	44.8 (84)	54.0 (90)	65.9 (102)	48.5 (72)	60.0 (80)	62.3 (78)				1.3	0.052
	12-14	36.1 (88)	41.4 (87)	50.4 (95)	47.6 (80)	53.6 (84)	42.0 (63)	46.6 (63)					1.2	0.093
boys	0-2					25.3 (20)	28.5 (21)	42.2 (29)	47.1 (30)	38.8 (24)	60.1 (37)	55.1 (34)	4.4	<0.001
	3-5				31.2 (27)	42.1 (33)	44.4 (33)	33.7 (23)	34.8 (22)	60.3 (38)	51.1 (32)		2.4	0.034
	6-8			43.0 (42)	54.0 (46)	53.2 (42)	46.6 (35)	51.4 (35)	55.8 (36)	53.1 (34)			0.8	0.423
	9-11		51.5 (56)	50.7 (49)	59.4 (51)	69.2 (55)	49.9 (38)	70.9 (49)	58.1 (38)				1.1	0.214
	12-14	47.9 (60)	54.6 (59)	69.0 (67)	56.7 (49)	55.9 (45)	54.6 (42)	70.0 (49)					1.2	0.180
girls	0-2					20.1 (15)	20.2 (14)	21.9 (14)	25.6 (15)	42.9 (25)	41.6 (24)	31.2 (18)	4.2	0.006
	3-5				30.7 (25)	27.1 (20)	35.7 (25)	55.1 (35)	48.1 (28)	47.3 (28)	45.9 (27)		3.0	0.013
	6-8			23.9 (22)	36.2 (29)	45.6 (34)	43.5 (31)	33.1 (21)	71.2 (42)	54.8 (33)			4.3	<0.001
	9-11		49.7 (51)	38.5 (35)	48.2 (39)	62.4 (47)	47.1 (34)	48.2 (31)	66.9 (40)				1.5	0.127
	12-14	23.6 (28)	27.4 (28)	30.6 (28)	37.9 (31)	51.1 (39)	28.8 (21)	21.5 (14)					1.2	0.314

Table 3: Rate ratios (RR) for each birth cohort taking those born on 1989-1994 as a reference and each calendar period of diagnosis taking those diagnosed in 1998-2000 as a reference. RRs are adjusted for age and for sex when both genders are analysed together (total). RRs for birth cohort do not incorporate period effect and RRs for period do not incorporate birth cohort period effect

Birth cohort	total		boys		Girls	
	RR	95% CI	RR	95% CI	RR	95% CI
1974-79	0.78	0.61-1.01	0.84	0.61-1.14	0.70	0.45-1.08
1977-82	0.94	0.78-1.13	0.95	0.74-1.21	0.93	0.69-1.24
1980-85	0.92	0.78-1.10	1.04	0.83-1.29	0.77	0.58-1.02
1983-88	1.03	0.87-1.21	1.05	0.85-1.30	1	0.78-1.28
1986-91	1.13	0.97-1.32	1.10	0.89-1.35	1.18	0.93-1.49
1989-94	1	-	1	-	1	-
1992-97	1.12	0.96-1.32	1.2	0.97-1.48	1.02	0.80-1.32
1995-00	1.32	1.11-1.56	1.18	0.93-1.48	1.50	1.17-1.93
1998-03	1.45	1.20-1.76	1.36	1.05-1.76	1.57	1.18-2.09
2001-06	1.64	1.31-2.06	1.66	1.23-2.24	1.62	1.15-2.28
2004-09	1.62	1.18-2.23	1.72	1.16-2.57	1.45	0.85-2.45
Period of diagnosis						
1989-91	0.73	0.63-0.85	0.79	0.65-0.97	0.66	0.53-0.83
1992-94	0.80	0.69-0.93	0.91	0.75-1.11	0.67	0.53-0.84
1995-97	0.95	0.82-1.10	1.06	0.88-1.29	0.81	0.65-1.02
1998-00	1	-	1	-	1	-
2001-03	0.95	0.81-1.10	0.91	0.74-1.12	0.99	0.79-1.24
2004-06	1.11	0.96-1.29	1.18	0.97-1.43	1.04	0.83-1.30
2007-09	1.06	0.91-1.23	1.13	0.93-1.38	0.97	0.77-1.22

Table 4: Summary of testing different age-period-cohort effects to incidence rates of type 1 diabetes among Sardinian children 0 to 14 years old in the years 1989-2009. See online supplemental Figure for notation referring to hierarchical models tested. For convention we also use the Variable | Model notation, i.e. the notation Cohort|Age+Drift denotes the effect of Cohort added to the Age+Drift model.

	Residual deviance (df; p)		d.f	LR χ^2	p
TOTAL					
0. Sex	181.51 (68; <0.001)	A. Sex			
1. Sex+Age	135.88 (64; <0.001)	B. Age Sex	4	45.63	<0.0001
2. Sex+Age+Drift	97.46 (63; 0.004)	C. Drift Sex+Age	1	38.41	<0.0001
3. Sex+Age+Cohort	86.34 (54; 0.003)	D. Cohort Sex+Age+Drift	9	11.12	0.268
4. Sex+Age+Period+Cohort	71.08 (49; 0.021)	E. Period Sex+Age+Cohort	5	15.27	0.009
5. Sex+Age+Period	89.52 (58; 0.005)	F. Cohort Sex+Age+Period	9	18.44	0.030
		G. Period Sex+Age+Drift	5	7.95	0.159
BOYS					
1. Age	41.72 (30) (0.088)	B. Age			
2. Age+Drift	26.80 (29) (0.649)	C. Drift +Age	1	14.92	0.0001
3. Age+Cohort	19.76 (20) (0.557)	D. Cohort Age+Drift	9	7.03	0.634
4. Age+Period+Cohort	9.98 (15) (0.869)	E. Period Age+Cohort	5	9.78	0.082
5. Age+Period	18.42 (24) (0.828)	F. Cohort Age+Period	9	8.45	0.490
		G. Period Age+Drift	5	8.37	0.137
GIRLS					
1. Age	68.31 (30) (<0.001)	B. Age			
2. Age+Drift	42.98 (29) (0.051)	C. Drift +Age	1	25.33	<0.0001
3. Age+Cohort	28.79 (20) (0.113)	D. Cohort Age+Drift	9	14.19	0.116
4. Age+Period+Cohort	17.53 (15) (0.369)	E. Period Age+Cohort	5	11.26	0.047
5. Age+Period	35.85 (24) (0.065)	F. Cohort Age+Period	9	18.32	0.032
		G. Period Age+Drift	5	7.13	0.211

