

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## The neglected navigating web of incomprehensibly emerging and re-emerging Sarcoptes mite

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/142746> since

*Published version:*

DOI:10.1016/j.meegid.2013.04.018

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*Infection, Genetics and Evolution*, 17, 2013, 10.1016/j.meegid.2013.04.018]

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

[ISSN: 1567-1348, ESSN: 1567-7257]

Review

## The neglected navigating web of the incomprehensibly emerging and re-emerging *Sarcoptes* mite

Samer Alasaad, Luca Rossid, Jorg Heukelbach, Jesús M. Pérez, Omar Hamarsheh, Moses Otiende, Xing-Quan Zhu

Show more

<http://dx.doi.org/10.1016/j.meegid.2013.04.018>

Get rights and content

Abstract

Parasite presence in any ecosystem generates complex navigating webs (Parasite-NW) within the system, through which parasites move from one to another host. The appropriate assimilation of parasite navigating web is pivotal for a better understanding of pathogen flow in the ecosystem, with implications for disease control. *Sarcoptes* mite has been approached from medical, veterinary, entomological, physiological and, recently, molecular sides, to understand its epidemiological navigating web between isolates from different hosts and geographical regions. The obtained conclusions are still a matter of debate. *Sarcoptes* navigating web (*Sarcoptes*-NW) is intricate and uncertain, with unexplainable pathogenic flow. In this review we summarize by which routes, under what conditions and at what levels the *Sarcoptes* mite moves among its hosts.

Keywords

*Sarcoptes scabiei*; Scabies; *Sarcoptes* mite; Parasite navigating web (Parasite-NW); Parasite flow; Emerging and re-emerging disease

### 1. Introduction

The neglected parasite *Sarcoptes scabiei* affects humans and a wide range of mammalian hosts worldwide (Bornstein et al., 2001, Pence and Ueckermann, 2002, Walton et al., 2004b and Alasaad et al., 2011a), entailing significant mortality in both wild and domestic animals, with considerable economic losses (Bornstein, 1995, Pence and Ueckermann, 2002, Heukelbach and Feldmeier, 2006 and Dagleish et al., 2007), and ravages in human populations (Walton et al., 2004b and Hay et al., 2012).

There are no accurate estimates of the prevalence of *Sarcoptes* mite in the many animal populations worldwide, particularly in sylvatic animals. In humans, estimations indicate that hundreds of million people are infested with scabies worldwide (WHO, 2009). The prevalence of scabies in African children can be as high as 40–80%, and in remote indigenous communities in northern Australia, up to 50% of children and 25% of adults were found to be infested (Kristensen, 1991, Carapetis et al., 1997 and Terry et al., 2001). Considering increasing resistance of topical chemotherapy, there is an urgent need to develop new control strategies (Currie et al., 2004, Bradberry et al., 2005 and Sanderson et al., 2007), and the increasing need

of sensitive and reliable diagnostic tests for humans and many domestic and wild animals ( Haas et al., 2005 and Heukelbach and Feldmeier, 2006).

Sarcoptes mite has been approached from morphological ( Fain, 1978), medical ( Carapetis et al., 1997 and Feldmeier et al., 2009), veterinary ( Bornstein, 1995), entomological ( Fain, 1978), physiological ( Arlian, 1989, Arlian et al., 1996 and Haas et al., 2005) and, recently, molecular points of view ( Zahler et al., 1999, Walton et al., 1997, Walton et al., 1999, Walton et al., 2004a, Alasaad et al., 2008b, Alasaad et al., 2009a, Alasaad et al., 2009b, Alasaad et al., 2011b, Alasaad et al., 2012c, Rasero et al., 2010 and Gakuya et al., 2011) aiming to understand its epidemiology between isolates from different hosts and geographical regions, but this issue is still a matter of vivid debate.

The objective of the present review is to summarize available evidence on Sarcoptes movement among hosts, in order to further elucidate transmission dynamics and provide an evidence-based rationale for sustainable control.

## 2. Emerging and re-emerging *Sarcoptes scabiei*

Parasite navigating webs (Parasite-NW) are the complex webs through which zoonotic parasites move from one host to another within the ecosystem (Polley, 2005). Pivotal to an appreciation of the function of parasite webs is an understanding of parasite flow: by which routes, under what conditions and at what levels the parasite flows among its various hosts, of the same or different species (including vectors where applicable), and between the hosts and the environment (Daszak et al., 2000).

Based on the definition of emergence and re-emergence of diseases by Lederberg et al. (1992), *Sarcoptes* mite emergence could be the result of the spread of *Sarcoptes* mites in one human/animal population from an infectious origin, or simply the realization that an infection has been present in a population but undetected for several reasons. The re-emergence of *Sarcoptes* is defined as the reappearance of *Sarcoptes* after a decline in incidence. Accordingly:

(i) A *Sarcoptes* mite mange outbreak can be 'genuine' emergence of an infestation, which is new to that particular and naive animal/human population. In this case, other infested hosts (humans, domestic and/or wild animals) sharing space with the non-infected population are suspected to be the reservoir and source of the mites through cross-infestation. In ruminants, the possibility that mites, adapted to a "main" reservoir host, may infest other sympatric species has been documented in mange foci in Europe ( León-Vizcaíno et al., 1999, Rossi et al., 2007 and Oleaga et al., 2008), and field evidence has been supported by the results of experimental infestations ( Meneguz and Rossi, 1995 and Lavín et al., 2000). Migration of human populations may also be a driving force for genuine emergence of scabies.

(ii) A *Sarcoptes* mite outbreak can be 'apparent' emergence/re-emergence, where *Sarcoptes* infestation is pre-existing, and the newly recognition is a result of increased detection opportunities ( Kutz et al., 2003 and Oleaga et al., 2008) and/or the modification of a pre-existing host-parasite equilibrium ( Pence et al., 1983 and Lloyd, 1995). In this second case, destabilizing modifications may have occurred on the host side, e.g. a diminished herd resistance, and/or the parasite side, e.g. the selection of a more pathogenic strain or drug resistance ( Pence and Windberg, 1994 and Leung and Grenfell, 2003).

Further epidemiological studies should consider a third category, which is a mixture of genuine and apparent emergence of *Sarcoptes* mite outbreaks. Such mixture category can be revealed using both epidemiological and population genetic approaches.

### 3. *Sarcoptes* transmissions

It is thought that *Sarcoptes* mite originated from a human ancestor and then spread to domestic and then free-living animals ( Fain, 1968 and Walton et al., 2004b). *Sarcoptes* mites lack free-living stages, and individual hosts, depending on their susceptibility and behaviour, are essentially ephemeral habitats providing patchy environments that hamper random mating ( Price, 1980 and Criscione et al., 2005). All mites on an individual host may in fact form an 'infrapopulation' ( Bush et al., 1997 and Alasaad et al., 2008b) that has a number of recurrent generations. The number of generations is influenced by the short generation interval of this parasite (about three weeks), as well as by the infected host's life expectancy and susceptibility ( Bornstein et al., 2001).

Transmission of *S. scabiei* may occur by direct or indirect contact. Larvae and nymphs of *S. scabiei* frequently leave their burrows and wander on the skin ( Arlian and Vyszynski-Moher, 1988), which may, in the case of crusted scabies, harbour hundreds to several thousands of mites/cm<sup>2</sup> ( Zeh, 1974, Arlian et al., 1988 and Pérez et al., 2011). Some may become dislodged from the host and fall off ( Arlian and Vyszynski-Moher, 1988). In controlled environments, mites may survive up until a few weeks if conditions (microclimate) are optimal ( Arlian, 1989). Cooler temperatures and high humidity prolong off-host survival of *S. scabiei*, presumably because under these conditions mite metabolism is reduced ( Davis and Moon, 1987 and Arlian et al., 1989).

However, *Sarcoptes* navigating web (*Sarcoptes*-NW) is intricate and uncertain with fragility of parasite flow, and consequently transmission dynamics are not clear. For example, the mode of scabies transmission in humans is still discussed-whereas some authors define scabies as a sexually-transmitted disease, others suggest close body contact between children and their mothers to be the main route of transmission in endemic communities ( Jackson et al., 2007). Clearly, these aspects depend on many interacting factors and differ from setting to setting.

The time needed for *S. scabiei* var. *hominis*, immediately transferred from one host to another, to initiate penetration into the stratum corneum was ~10 min ( Arlian et al., 1984), and it took the mites ~35 min to

become completely submerged into the epidermis. The time required for complete penetration into the stratum corneum increased as a function of the time the mites had been off their host. Experiments indicate that mites remain infective at least one-half to two-thirds of their survival time when dislodged from their host ( Arlian, 1989). In historical but classical studies, Mellanby (1944) believed that the stage responsible for transmission was the young, newly fertilised adult female, which wandered around on the skin surface before burrowing. His studies on transmission of scabies determined that patients with a high parasite load in excess of 100 adult female mites are more likely to spread the disease than those harbouring a lower number of parasites.

Additional studies on *Sarcoptes* transmission and off-host survival under different host species, climatic conditions and fomite materials are still needed ( Arlian, 1989).

From the other side, the co-infection between *Sarcoptes* mite and other ecto- and endo-parasites still not well-understood and need further studies ( Alasaad et al., 2008a, Alasaad et al., 2008b and Ryser-Degiorgis et al., 2002). The better understanding of mixed infection, involving *S. scabiei*, is of pivotal interest for the better understanding of *Sarcoptes* epidemiology, diagnosis and treatment

### 3.1. Intra-mammalian groups transmission

#### 3.1.1. Intra-humans transmission

Epidemics of human scabies have been hypothesised to occur on a worldwide basis in 15–25 year cycles (Falk, 1982, Arlian, 1989 and Lassa et al., 2011). Nonetheless, this possible “fluctuations” of disease occurrence is not corroborated by appropriate data (Orkin and Maibach, 1978).

Estimations indicate that 300 million people, especially children, are affected (WHO, 2009). Prevalence of scabies in African children can be as high as 40–80% (Kristensen, 1991 and Terry et al., 2001). In low and middle income countries, scabies is a significant public health problem because it is highly prevalent and complications are frequent (Heukelbach et al., 2005, Hay et al., 2012 and Heukelbach et al., 2012). Presence and severity of scabies are associated with young age, presence of many children in the household, illiteracy, low family income, poor housing, sharing clothes and towels, and irregular use of shower (Feldmeier et al., 2009). The high association with sharing clothes and towels points to a substantial contribution of fomites in the transmission of mites between humans, at least in certain settings (Zeh, 1974). However, in practice (and in endemic communities) it is not possible to quantify the importance of each single factor in epidemiological studies. Crusted scabies (also called Norwegian scabies), which is a scabies mite hyper-infection with hundreds of thousands of parasites on a single host, is rarely diagnosed in healthy individuals, but sometimes is seen in elderly, immune-suppressed patients with chronic disease, or patients suffering from malnutrition. These high transmitter individuals, who because of behaviours or mite load or some other factor, may provide a much higher risk of transmitting infestations (de Almeida Barbosa et al., 1996).

In an urban slum in Bangladesh, the incidence of scabies in children younger than 5 years was 952/1000 per year, which means that almost all children experienced at least one infestation per year (Stanton et al., 1987). This is in contrast to the situation in industrialized countries, where the disease occurs sporadically in all age groups, particularly in sexually active adults, or causes epidemics in institutions and nursing homes (Andersen et al., 2000, Heukelbach et al., 2005 and Ariza et al., 2012).

### 3.1.2. Intra-domestic animals transmission

In Africa, mites from sheep were successfully transferred to goats (Abu-Samra et al., 1984), and transmission of mites from goats to sheep was also achieved (Ibrahim and Abu-Samra, 1987). In both cases, infested goats and sheep developed severe lesions, particularly in moistened areas, and observation of skin scrapings revealed large numbers of mites belonging to all developmental stages, which demonstrated active reproduction of transferred mites in the new host. Similar results were obtained with experimental cross-infection between camels and goats (Nayel and Abu-Samra, 1986). Conversely, other attempts to transfer *Sarcoptes* mites between domestic species failed, like those involving dogs, sheep, goats, cattle and cats (Arlian et al., 1988) (Table 1).

Table 1.

Experimental cross-transmission of *Sarcoptes scabiei*, host species and outcomes.

Host of origin	Receiver host	Outcome	Citation
Man	Macacus rhesus	Not successful	Ruch (1959)
Man	Domestic sheep	Not successful	Pirilä et al. (1967)
Man	Cattle	Not successful	Pirilä et al. (1967)
Domestic goat	Northern chamois	Successful	Fiebiger (1917) and Kerschagl (1955)
Domestic goat	Camel	Successful	Nayel and Abu-Samra (1986)
Domestic goat	Domestic sheep	Successful	Ibrahim and Abu-Samra (1987)
Domestic goat	Southern chamois	Successful	Lavín et al. (2000)
Domestic goat	Donkey	Successful	Abu-Samra et al. (1984)
Northern chamois	Domestic goat	Successful	Meneguz and Rossi (1995)
Southern chamois	Cattle	Not successful	Lavín et al. (2000)
Southern chamois	Domestic sheep	Not successful	Lavín et al. (2000)
One humped camel	Domestic goat	Successful	Nayel and Abu-Samra (1986)
Dog	Domestic sheep	Not successful	Arlian et al. (1988)

Dog	Domestic goat	Not successful	Arlian et al. (1988)
Dog	Cattle	Not successful	Arlian et al. (1988)
Dog	Cat	Not successful	Arlian et al. (1988)
Dog	Rabbit	Successful	Arlian et al. (1984)
Dog	Guinea pig	Not successful	Arlian et al. (1984)
Dog	House mouse	Not successful	Arlian et al. (1984)
Dog	Pig	Not successful	Arlian et al. (1984)
Dog	Rat	Not successful	Arlian et al. (1984)
Dog	Man	Not successful	Kutzer and Grunberg (1970)
Coyote Dog		Not successful	Samuel (1981)
Coyote x dog hybrid	Red fox	Successful	Stone et al. (1972)
Wolf Dog		Not successful	Samuel (1981)
Red fox Dog		Not successful	Samuel (1981)
Red fox Dog		Successful	Stone et al. (1972) and Bornstein (1991)
Red fox Coyote x dog hybrid		Successful	Stone et al. (1972)
Red fox Coyote		Successful	Samuel (1981)
Red fox Man		Not successful	Kutzer and Grunberg (1970)
Tapir Man		Not successful	Kutzer and Grunberg (1970)

#### Table options

##### 3.1.3. Intra-wild animals transmission

Mange, caused by *S. scabiei*, has been reported from many species of wild mammals worldwide. Some of the most remarkable hosts include (1) canids in North America (Pence et al., 1983 and Little et al., 1998); (2) red foxes and other canids in Europe (Mörner, 1992, Ippen et al., 1995 and Gortazar et al., 1998); (3) red foxes and dingoes in Australia (McCarthy, 1960); (4) chamois, ibex and a variety of other ungulates in Europe (Ippen et al., 1995, Fernández Morán et al., 1997 and Rossi et al., 2007); (5) felids in Europe (Mörner, 1992) and Africa (Young, 1975); (6) wild boar in Europe (Ippen et al., 1995); (7) wombats in Australia (Skerratt et al., 1998); and (8) a range of ungulates, primates, and canids in Africa (Young, 1975, Kalema et al., 1998, Munang'andu et al., 2010 and Gakuya et al., 2012b). New outbreaks in wildlife are being continually reported worldwide (Alasaad et al., 2012a).

The molecular analyses of sympatric wild animals in Europe show unambiguously that there has been a lack of gene flow or recent admixture between carnivore-, herbivore-, and omnivore-derived *Sarcoptes*



populations ( Rasero et al., 2010). Host-taxon-derived effect seems stronger driver of intra-specific differentiation than geographical separation, and it seems temporally stable ( Alasaad et al., 2011b). The existence of host-taxon-derived Sarcoptes mites could explain why mange-free populations can live in sympatry with mangy animals, as is the case of the mange-free *Capra ibex* and *Rupicapra rupicapra* of the western Italian Alps that live in close proximity with the endemically mangy population of red fox *Vulpes vulpes*, and wild boar *Sus scrofa* ( Rasero et al., 2010). Vice versa, this effect could be the immediate reason why cross-transmission easily occurs, in Europe, between foxes, dogs and other canids, as well as felids including domestic European cats, when these hosts are exposed to the same vulpine-derived strain ( Bornstein 1995) ( Table 1).

The studied wild animals in Europe (from which the host-taxon was coined) lacked of clear predator–prey interaction and putative inter-specific transmission models. Recent epidemiological and molecular studies revealed potential prey-to-predator gene flow between wild animals from Africa. Cheetahs were infected from both wildebeest and Thomson’s gazelle, while lions were only infected from wildebeest. In a predator/prey ecosystem, like Masai Mara in Kenya, it seems that Sarcoptes infestation is from prey-to-predator, in relation to the predator’s “favourite prey”, which could be attributed to the direct skin-to-skin prolonged contact between the predator and the prey during hunting process ( Gakuya et al., 2011 and Gakuya et al., 2012a).

### 3.2. Inter-mammalian groups transmission

#### 3.2.1. Humans-wild/domestic animals transmission

Humans are occasionally infected with *S. scabiei* of animal origin ( Arlian, 1989). This includes human infection from camel, cat, chamois, coyote, dog, ferret, fox, wombat, gazelle, goat, horse, llama, pig, sheep, water buffalo and cattle ( Kutzer, 1970, Fain, 1978, Chakrabarti et al., 1981, Samuel, 1981, Folz, 1984, Chakrabarti, 1990, Mitra et al., 1993, Morsy et al., 1995, Skerratt and Beveridge, 1999, Menzano et al., 2004 and Bazargani et al., 2007). Those infected include farmers, personnel working in slaughterhouses researchers, veterinarians, wildlife biologist, and pet owners.

Most human infections from animal sources involve only topographically circumscribed body regions, are short-lived and self-limiting, lasting from a few days to several weeks (Arlian, 1989). As infestation is less severe and clinical features are different from infestation with *Sarcoptes scabiei* var. *hominis*, this condition is usually referred in the medical literature as “pseudoscabies”. Lesions are frequently seen on the trunk, arms, and abdomen and rarely on the fingerwebs and genitalia. In addition, the incubation period is markedly shorter, and mite burrows are not regularly seen ( Orkin and Maibach, 1991). Pruritus may be as intensive as in classical scabies, but symptoms usually wane within a maximum of 2–4 weeks. The human host is not a source of infection to other humans. Reportedly, nonhuman mites most frequently originate from dogs infested with *S. scabiei* var. *canis* (30–50% of cases) ( Thomsett, 1968, Folz, 1984 and Aydingöz and Mansur, 2011).

The short-life and self-limiting infection of *Sarcoptes* mites of animal origin when transmitted to humans was confirmed by molecular analysis. Walton et al., 1999 and Walton et al., 2004a used multi-locus genotyping applied to microsatellite markers to substantiate previous findings that gene flow between human- and dog-derived mite populations is extremely rare in scabies-endemic Aboriginal communities in Australia.

### 3.2.2. Domestic animals-wildlife transmission

Considering the increasing deforestation and changing ecosystems worldwide, there has been more and more contact of domestic animals with sylvatic animals. In fact, the introduction of infected domestic animals and the success of the *Sarcoptes* mite in adapting to new highly susceptible wild hosts have been proposed as the origin of *Sarcoptes* mite epizootics in previously mange-free wildlife populations ( Arlian, 1989). This scenario was also proposed for a *Sarcoptes* mite outbreak in Iberian ibex in Sierra Nevada mountain range, Spain in the 1970s, possibly from domestic infected goats ( Pérez et al., 1997).

Molecular studies have been carried out to understand *Sarcoptes* transmission between humans and animals, and between wild host species, but never between domestic and wild animals, and hence effort should be made in this direction. We know very little of interspecific transmission between (putative) domestic reservoirs and wildlife, while we know much more on wildlife to wildlife transmission.

In the Sierras de Cazorla, Segura y Las Villas Natural Park (southern Spain) mange has become an endemic disease. At the end of the 1980s, it firstly affected Iberian ibex (*Capra pyrenaica*) causing devastating mortality rates with a reduction of the population by 95%. Then, the disease reached other sympatric wild ungulate species: European mouflon (*Ovis aries*), red deer (*Cervus elaphus*) and fallow deer (*Dama dama*). The origin of this epizooty is attributed to infested domestic goat herds (see, for instance, León-Vizcaíno et al., 1999). Regarding this situation, an attempt to experimentally infect chamois (*Rupicapra rupicapra*) with *Sarcoptes* mites derived from naturally-infested domestic goats was successful ( Lavín et al., 2000). The opposite route of transmission was also previously demonstrated under experimental conditions ( Meneguz and Rossi, 1995). A recent outbreak of *Sarcoptes* mite in domestic goats from northern Italy has been attributed to presumable contact with wild animals under natural conditions ( Menzano et al., 2007).

Domestic-wild animals transmission usually produce high mortality and morbidity rates, probably because of insufficient management actions.

## 4. Predictions of emergence and re-emergence

*Sarcoptes* emergence and re-emergence could be predicted by different approaches, such as (i) endemic cycles, (ii) increasing agglomeration of people, wild and domestic animals, and increasing contact between human-wildlife-livestock. Notwithstanding, the prediction of *Sarcoptes* emergence and re-emergence could be affected by (a) the emerging of resistant against some acaricides ( Currie et al., 2004), (b) the lack of

effective diagnosis method for many host species especially the wild ones ( Alasaad et al., 2012b), and (c) the totally neglected role of fomites in *Sarcoptes* transmission.

Better prediction of scabies and mange emergence and re-emergence could be achieved through better understanding of inter-specific transmission patterns, which requires further studies including the effect of (i) travelling, sexual and familiar contact, biomedical manipulation and migration in intra-humans transmission; (ii) agricultural intensification and animal translocation in intra-domestic animals transmission; (iii) translocation and animal stress in intra-wild animals transmission; (iv) technology and industry manipulation in humans-domestic animals transmission; (v) human encroachment, tourism, and ecological manipulation in humans-wild animals transmission; and (vi) animal encroachment and animal introduction in domestic animals-wildlife transmission (Fig. 1).

Full-size image (80 K)

Fig. 1.

The neglected *Sarcoptes* navigating web (*Sarcoptes*-NW), including multiple putative routes of parasite flow between mammalian hosts, following authors' criteria. Dotted lines: flow through fomites. Solid lines: flow through direct contact (host-to-host). Circular arrow: flow within animal group. Question mark: questionable parasite flow. Curved lines: flow between groups.

Figure options

## 5. Public awareness

A better understanding of *Sarcoptes* navigating web is of pivotal interest for the public health. Unfortunately many facets of *Sarcoptes* navigating web regarding humans are not yet clear, including human-to-human transmission, especially the sexually transmission and family care, and wild and domestic animals to humans transmission. Integral programs should be carried out including epidemiological and genetic studies of *Sarcoptes* navigating web between humans and wild/domestic animals, and review of knowledge and practices regarding *Sarcoptes* infection in humans ( Gakuya et al., 2012b). Effective and operator-friendly methods for direct and indirect diagnosis of scabies are still missing and the resistance of *Sarcoptes* to drugs is now becoming of major concern.

## Acknowledgements

Project support was provided by Proyecto de Excelencia RNM 06400 (Junta de Andalucía, Spain) and Juan de la Cierva Grant (Ministerio Innovación y ciencia, Spain). XQZ is supported by the International Science & Technology Cooperation Program of China (Grant No. 2013DFA31840), the Science Fund for Creative Research Groups of Gansu Province (Grant No. 1210RJA006) and the Special Fund for Agro-scientific Research in the Public Interest (Grant No. 201303037).

## References

Abu-Samra et al., 1984

M.T. Abu-Samra, K.E.E. Ibrahim, M. Abdel Aziz

Experimental infection of goats with *Sarcoptes scabiei* var. *ovis*

Ann. Trop. Med. Parasitol., 78 (1984), pp. 55–61

[View Record in Scopus](#) | [Cited By in Scopus \(4\)](#)

Alasaad et al., 2008a

S. Alasaad, J.E. Granados, F.J. Cano-Manuel, A. Meana, X.Q. Zhu, J.M. Pérez

Epidemiology of fasciolosis affecting Iberian ibex (*Capra pyrenaica*) in southern Spain

Parasitol. Res., 102 (2008), pp. 751–755

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(9\)](#)

Alasaad et al., 2008b

S. Alasaad, D. Soglia, M. Sarasa, R.C. Soriguer, J.M. Pérez, J.E. Granados, R. Rasero, X.Q. Zhu, L. Rossi

Skin-scale genetic structure of *Sarcoptes scabiei* populations from individual hosts: empirical evidence from Iberian ibex-derived mites

Parasitol. Res., 104 (2008), pp. 101–105

[View Record in Scopus](#) | [Cited By in Scopus \(18\)](#)

Alasaad et al., 2009a

S. Alasaad, L. Rossi, R.C. Soriguer, L. Rambozzi, D. Soglia, J.M. Pérez, X.Q. Zhu

*Sarcoptes* mite from collection to DNA extraction: the lost realm of the neglected parasite

Parasitol. Res., 104 (2009), pp. 723–732

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(16\)](#)

Alasaad et al., 2009b

S. Alasaad, D. Soglia, V. Spalenza, S. Maione, R.C. Soriguer, J.M. Pérez, R. Rasero, M.P. Ryser Degiorgis, H. Nimmervoll, X.Q. Zhu, L. Rossi

Is ITS-2 rDNA suitable marker for genetic characterization of Sarcoptes mites from different wild animals in different geographic areas?

Vet. Parasitol., 159 (2009), pp. 181–185

Article | PDF (492 K) | View Record in Scopus | Cited By in Scopus (17)

Alasaad et al., 2011a

S. Alasaad, D. Holt, S. Walton, L. Rossi, S. Bornstein, M. Abu-Madi, R.C. Soriguer, S. Fitzgerald, X.Q. Zhu, W. Zimmermann, U.S. Ugbomoiko, J. Heukelbach

Sarcoptes-World Molecular Network (Sarcoptes-WMN): integrating research on scabies

Int. J. Infect. Dis., 15 (2011), pp. 294–297

Alasaad et al., 2011b

S. Alasaad, A. Oleaga, R. Casais, L. Rossi, A. Molinar Min, R. Soriguer, C. Gortazar

Temporal stability in the genetic structure of *Sarcoptes scabiei* under the host-taxon law: empirical evidences from wildlife-derived *Sarcoptes* mite in Asturias

Spain. Parasit. Vectors, 4 (2011), p. e151

Alasaad et al., 2012a

S. Alasaad, D. Ndeereh, L. Rossi, S. Bornstein, R. Permuanian, R.C. Soriguer, F. Gakuya

The opportunistic *Sarcoptes scabiei*: a new episode from giraffe in the drought-suffering Kenya

Vet. Parasitol., 185 (2012), pp. 359–363

Article | PDF (818 K) | View Record in Scopus | Cited By in Scopus (6)

Alasaad et al., 2012b

S. Alasaad, R. Permuanian, F. Gakuya, M. Mutinda, R.C. Soriguer, L. Rossi

Sarcoptic-mange detector dogs used to identify infected animals during outbreaks in wildlife

BMC Vet. Res., 8 (2012), p. e110

Alasaad et al., 2012c

S. Alasaad, R.K. Schuster, F. Gakuya, M. Theneyan, M.J. Jowers, S. Maione, A. Molinar Min, R.C. Soriguer, L. Rossi

Applicability of molecular markers to determine parasitic infection origins in the animal trade: a case study from *Sarcoptes* mites in wildebeest

Forensic Sci. Med. Pathol., 8 (2012), pp. 280–284

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(6\)](#)

Andersen et al., 2000

B.M. Andersen, H. Haugen, M. Rasch, A.H. Haugen, A. Tageson

Outbreak of scabies in Norwegian nursing homes and home care patients: control and prevention

J. Hosp. Infect., 45 (2000), pp. 160–164

[Article](#) | [PDF \(79 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(51\)](#)

Ariza et al., 2012

L. Ariza, B. Walter, C. Worth, S. Brockmann, ML. Weber, H. Feldmeier

Investigation of a scabies outbreak in a kindergarten in Constance, Germany

Eur. J. Clin. Microbiol. Infect. Dis. [Epub ahead of print] (2012)

Arlian et al., 1984

L.G. Arlian, R.A. Runyan, S.A. Estes

Cross infectivity of *Sarcoptes scabiei*

J. Am Acad. Dermatol., 10 (1984), pp. 979–986

[Article](#) | [PDF \(2869 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(33\)](#)

Arlian and Vyszynski-Moher, 1988

L.G. Arlian, D.L. Vyszynski-Moher

Life cycle of *Sarcoptes scabiei* var. *canis*

J. Parasitol., 74 (1988), pp. 427–430

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (18)

Arlan et al., 1988

L.G. Arlian, M. Ahmed, D.L. Vyszensky-Moher

Effects of *Sarcoptes scabiei* var. *canis* (Acari: Sarcoptidae) in blood indices of parasitized rabbits

J. Med. Entomol., 25 (1988), pp. 360–369

[View Record in Scopus](#) | [Cited By in Scopus](#) (12)

Arlan, 1989

L.G. Arlian

Biology, host relations, and epidemiology of *Sarcoptes scabiei*

Ann. Rev. Entomol., 34 (1989), pp. 139–161

[View Record in Scopus](#) | [Cited By in Scopus](#) (99)

Arlan et al., 1989

L.G. Arlian, D.L. Vyszenski-Moher, M.J. Pole

Survival of adults and developmental stages of *Sarcoptes scabiei* var. *canis* when off the host

Exp. Appl. Acarol., 6 (1989), pp. 181–187

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (31)

Arlan et al., 1996

L.G. Arlian, M.S. Morgan, L.L. Arends

Immunologic cross-reactivity among various strains of *Sarcoptes scabiei*

J. Parasitol., 82 (1996), pp. 66–72

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (31)

Aydingöz and Mansur, 2011

I.E. Aydingöz, A.T. Mansur

Canine scabies in humans: a case report and review of the literature

Dermatology, 223 (2011), pp. 104–106

[View Record in Scopus](#) | [Cited By in Scopus \(1\)](#)

Bazargani et al., 2007

T.T. Bazargani, J.A. Hallan, S. Nabian, S. Rahbari

Sarcoptic mange of gazelle (*Gazella subgutturosa*) and its medical importance in Iran

Parasitol. Res., 101 (2007), pp. 1517–1520

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(2\)](#)

Bradberry et al., 2005

S.M. Bradberry, S.A. Cage, A.T. Proudfoot, J.A. Vale

Poisoning due to pyrethroids

Toxicol. Rev., 24 (2005), pp. 3–106

Bornstein, 1991

S. Bornstein

Experimental infection of dogs with *Sarcoptes scabiei* derived from naturally infected wild red foxes (*Vulpes vulpes*): clinical observations

Vet. Dermatol., 2 (1991), pp. 151–159

[Full Text via CrossRef](#)

Bornstein et al., 2001

S. Bornstein, T. Mörner, W.M. Samuel

*Sarcoptes scabiei* and sarcoptic mange

W.M. Samuel, M.J. Pybus, A.A. Kocan (Eds.), *Parasitic Diseases of Wild Mammals* (second ed.), Iowa State University Press, Ames (2001), pp. 107–119

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(1\)](#)

Bornstein, 1995



Bornstein, S., 1995. *Sarcoptes scabiei* infections of the domestic dog, red fox and pig. PhD thesis. Department of Veterinary Microbiology, Section of Parasitology, Swedish University of Agricultural Sciences and National Veterinary Institute, Uppsala, Sweden.

Bush et al., 1997

A.O. Bush, K.D. Lafferty, J.M. Lotz, A.W. Shostak

Parasitology meets ecology on its own terms: Margolis et al. revisited

J. Parasitol., 83 (1997), pp. 575–583

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(2373\)](#)

Carapetis et al., 1997

J.R. Carapetis, C. Connors, D. Yarmirr, V. Krause, B.J. Currie

Success of a scabies control program in an Australian aboriginal community

Pediatr. Infect. Dis. J., 16 (1997), pp. 494–499

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(77\)](#)

Chakrabarti, 1990

A. Chakrabarti

Pig handler's itch

Int. J. Dermatol., 29 (1990), pp. 205–206

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(3\)](#)

Chakrabarti et al., 1981

A.N. Chakrabarti, A.A. Chatterjee, K. Chakrabarti, D.N. Sengupta

Human scabies from contact with water buffaloes infested with *Sarcoptes scabiei* var. *bubalis*

Ann. Trop. Med. Parasitol., 75 (1981), pp. 353–357

[View Record in Scopus](#) | [Cited By in Scopus \(9\)](#)

Criscione et al., 2005

C.D. Criscione, R. Poulin, S. Blouin

Molecular ecology of parasites: elucidating ecological and microevolutionary processes

Mol. Ecol., 14 (2005), pp. 2247–2257

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(167\)](#)

Currie et al., 2004

B.J. Currie, P. Harumal, M. McKinnon, S.F. Walton

First documentation of in vivo and in vitro ivermectin resistance in *Sarcoptes scabiei*

Clin. Infect. Dis., 39 (2004), pp. 8–12

Dagleish et al., 2007

M.P. Dagleish, Q. Ali, R.K. Powell, D. Butz, M.H. Woodford

Fatal *Sarcoptes scabiei* infection of blue sheep (*Pseudois nayaur*) in Pakistan

J. Wildl. Dis., 43 (2007), pp. 512–517

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(10\)](#)

Daszak et al., 2000

P. Daszak, A.A. Cunningham, A.D. Hyatt

Emerging infectious diseases of wildlife – threats to biodiversity and human health

Sci., 287 (2000), pp. 443–449

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(1173\)](#)

Davis and Moon, 1987

D.P. Davis, R.D. Moon

Survival of *Sarcoptes scabiei* (De Geer) stored in three media at three temperatures

J. Parasitol., 73 (1987), pp. 661–662

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(1\)](#)

de Almeida Barbosa et al., 1996

A. de Almeida Barbosa, T.M. Correia Silva, M.I. Ribeiro Santos, M.F. Gomez Garrido, B. Niljay Patel, M. Cavalcanti, M.C. Oliveira Riccio

Coexistence of an unusual form of scabies and lepromatous leprosy

A case report. *Pathol. Res. Pract.*, 192 (1996), pp. 88–90

[View Record in Scopus](#) | [Cited By in Scopus \(7\)](#)

Fain, 1968

A. Fain

Étude de la variabilité de *Sarcoptes scabiei* avec une revision des Sarcoptidae

*Acta Zool. Pathol. Antverp.*, 47 (1968), pp. 1–196

[View Record in Scopus](#) | [Cited By in Scopus \(47\)](#)

Fain, 1978

A. Fain

Epidemiological problems of scabies

*Int. J. Dermatol.*, 17 (1978), pp. 20–30

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(63\)](#)

Falk, 1982

Falk, E.S., 1982. Scabies: some aspects of its relationship to the immune mechanisms. Ph.D. Thesis. Tromso University, Tromso.

Feldmeier et al., 2009

H. Feldmeier, A. Jackson, L. Ariza, C.M. Lins-Calheiros, V.L. Soares, F.A. Oliveira

The epidemiology of scabies in an impoverished community in rural Brazil: presence and severity of disease are associated with poor living conditions and illiteracy

*J. Am. Acad. Dermatol.*, 60 (2009), pp. 436–443

[Article](#) | [PDF \(154 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(10\)](#)

Fernández Morán et al., 1997

J. Fernández Morán, S. Gómez, F. Ballesteros, P. Quirós, J.L. Benito, C. Feliu, J.M. Nieto

Epizootiology of sarcoptic mange in a population of Cantabrian chamois (*Rupicapra pyrenaica parva*) in northwestern Spain

Vet. Parasitol., 73 (1997), pp. 163–171

Article | PDF (499 K) | View Record in Scopus | Cited By in Scopus (54)

Fiebiger, 1917

J. Fiebiger

Neue Untersuchungen über Genssenräude. Übertragung der Ziegenräude auf Genssen

Wien. tierärztl. Mschr., 4 (1917), pp. 433–450

Folz, 1984

S.D. Folz

Canine scabies (*Sarcoptes scabiei*) infestation

Compend. Contin. Educ. Vet., 6 (1984), pp. 176–180

Gakuya et al., 2011

F. Gakuya, L. Rossi, J. Ombui, N. Maingi, G. Muchemi, W. Ogara, R.C. Soriguer, S. Alasaad

The curse of the prey: *Sarcoptes* mite molecular analysis reveals potential prey-to-predator parasitic infestation in wild animals from Masai Mara

Kenya. Parasit. Vectors, 4 (2011), p. e193

Gakuya et al., 2012a

F. Gakuya, J. Ombui, N. Maingi, G. Muchemi, W. Ogara, R.C. Soriguer, S. Alasaad

Sarcoptic mange and cheetah conservation in Masai Mara (Kenya): epidemiological study in a wildlife/livestock system

Parasitology, 139 (2012), pp. 1587–1595

View Record in Scopus | Full Text via CrossRef | Cited By in Scopus (7)

Gakuya et al., 2012b

F. Gakuya, J. Ombui, J. Heukelbach, N. Maingi, G. Muchemi, W. Ogara, D. Mijele, S. Alasaad

Knowledge of mange among Masai pastoralists in Kenya

PLoS ONE, 7 (2012), p. e43342

[Full Text via CrossRef](#)

Gortazar et al., 1998

C. Gortazar, R. Villafuerte, J.C. Blanco, D. Fernandez-de-Luco

Enzootic sarcoptic mange in red foxes in Spain

Z. Jagdwiss., 44 (1998), pp. 251–256

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (16)

Haas et al., 2005

N. Haas, B. Wagemann, B. Hermes, B.M. Henz, C. Heile, E. Schein

Crossreacting IgG antibodies against fox mite antigens in human scabies

Arch. Dermatol. Res., 296 (2005), pp. 327–331

[View Record in Scopus](#) | [Cited By in Scopus](#) (14)

Hay et al., 2012

R.J. Hay, A.C. Steer, D. Engelman, S. Walton

Scabies in the developing world-its prevalence, complications, and management

Clin. Microbiol. Infect., 18 (2012), pp. 313–323

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (15)

Heukelbach et al., 2005

J. Heukelbach, T. Wilcke, B. Winter, H. Feldmeier

Epidemiology and morbidity of scabies and pediculosis capitis in resource-poor communities in Brazil

Br. J. Dermatol., 153 (2005), pp. 150–156

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(76\)](#)

Heukelbach and Feldmeier, 2006

J. Heukelbach, H. Feldmeier

Scabies

Lancet, 367 (2006), pp. 1767–1774

[Article](#) | [PDF \(106 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(101\)](#)

Heukelbach et al., 2012

J. Heukelbach, H.D. Mazigo, S. Ugbomoiko

Impact of scabies in resource-poor communities

Clin. Microbiol. Infect., 18 (2012), pp. 313–323

[View Record in Scopus](#) | [Cited By in Scopus \(1\)](#)

Ibrahim and Abu-Samra, 1987

K.E.E. Ibrahim, M.T. Abu-Samra

Experimental transmission of a goat strain of *Sarcoptes scabiei* to desert sheep and its treatment with ivermectin

Vet. Parasitol., 26 (1987), pp. 157–164

[Article](#) | [PDF \(584 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(15\)](#)

Ippen et al., 1995

R. Ippen, S. Nickel, H.D. Schröder

Krankheiten des Jagdbaren Wildes

Deutscher Land- wirtschaftsverlag, Berlin (1995) pp. 189–195

Jackson et al., 2007

A. Jackson, J. Heukelbach, H. Feldmeier

Transmission of scabies in a rural community

Braz. J. Infect. Dis., 11 (2007), pp. 386–387

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (10)

Kalema et al., 1998

Kalema, G., Koch, R.A., Macfie, E., 1998. An outbreak of sarcoptic mange in free-ranging mountain gorillas (*Gorilla gorilla berengei*) in Bwindi Impenetrable National Park, South Western Uganda. In: Proceedings AAZV and AAWV Joint Conference, Omaha, Nebraska, USA, p. 438.

Kerschagl, 1955

W. Kerschagl

Übertragbarkeit der Gamsräude auf andere Wildarten

Anblick, 10 (1955), pp. 295–296

[View Record in Scopus](#) | [Cited By in Scopus](#) (1)

Kristensen, 1991

J.K. Kristensen

Scabies and pyoderma in Lilongwe. Malawi. Prevalence and seasonal fluctuation

Int. J. Dermatol., 30 (1991), pp. 699–702

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (43)

Kutz et al., 2003

S.J. Kutz, E.P. Hoberg, J. Nagy, L. Polley, B. Elkin

Emerging parasitic infections in arctic ungulates

Integrat. Comp. Biol., 44 (2003), pp. 109–118

Kutzer, 1970

E. Kutzer

Sarcoptes-Milben und Sarcoptes räude der Haustiere. Merkblätter Über angewandte Parasitenkunde und Schädlingsbekämpfung

Angew. Parasitol., 11 (1970), pp. 1–22

Kutzer and Grunberg, 1970

E. Kutzer, W. Grunberg

Transmission of sarcoptic mange from animals to man

Vet. Bull., 40 (1970), p. 4

[View Record in Scopus](#) | [Cited By in Scopus \(1\)](#)

Lavín et al., 2000

S. Lavín, M. Ruiz-Bascaran, I. Marco, M.D. Fondevila, A.J. Ramis

Experimental infection of chamois (*Rupicapra pyrenaica parva*) with *Sarcoptes scabiei* derived from naturally infected goats

J. Vet. Med., 47 (2000), pp. 693–769

Lassa et al., 2011

S. Lassa, M.J. Campbell, C.E. Bennett

Epidemiology of scabies prevalence in the U.K. from general practice records

Br. J. Dermatol., 164 (2011), pp. 1329–1334

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(8\)](#)

Lederberg et al., 1992

, in: J. Lederberg, R.E. Shope, S.C. Oaks (Eds.), *Emerging Infections: Microbial Threats to Health in the United States* National Academies Press, Washington, DC (1992)

León-Vizcaíno et al., 1999

L. León-Vizcaíno, M.R. Ruíz de Ybáñez, M.J. Cubero

Sarcoptic mange in Spanish ibex from Spain

J. Wildl. Dis., 35 (1999), pp. 647–659



[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(59\)](#)

Leung and Grenfell, 2003

B. Leung, B.T. Grenfell

A spatial stochastic model simulating a scabies epidemic and coyote population dynamics

*Ecol. Model.*, 166 (2003), pp. 41–52

[Article](#) | [PDF \(159 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(11\)](#)

Little et al., 1998

S.E. Little, W.R. Davidson, E.W. Howerth, P.M. Rakich, V.F. Nettles

Diseases diagnosed in red foxes from Southeastern United States

*J. Wildl. Dis.*, 34 (1998), pp. 620–624

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(16\)](#)

Lloyd, 1995

S. Lloyd

Environmental influences on host immunity

B.T. Grenfell, A.P. Dobson (Eds.), *Ecology of Infectious Diseases in Natural Populations*, Cambridge University Press, Cambridge, UK (1995), pp. 327–361

[Full Text via CrossRef](#)

McCarthy, 1960

P.H. McCarthy

The presence of sarcoptic mange in the wild fox (*Vulpes vulpes*) in Central Queensland

*Aust. Vet. J.*, 36 (1960), pp. 359–360

[Full Text via CrossRef](#)

Mellanby, 1944

K. Mellanby

The development of symptoms, parasitic infection and immunity in human scabies

Parasitology, 35 (1944), pp. 197–206

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (63)

Meneguz and Rossi, 1995

P.G. Meneguz, L. Rossi

Experimental transmission of *Sarcoptes scabiei* from chamois to domestic goat

Proc. Soc. It. Sci. Vet., 49 (1995), pp. 755–756

[View Record in Scopus](#) | [Cited By in Scopus](#) (2)

Menzano et al., 2004

A. Menzano, L. Rambozi, L. Rossi

Outbreak of scabies in human beings, acquired from chamois (*Rupicapra rupicapra*)

Vet. Rec., 155 (2004), p. 568

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (10)

Menzano et al., 2007

A. Menzano, L. Rambozzi, L. Rossi

A severe episode of wildlife derived scabies in domestic goats in Italy

Small Rumin. Res., 70 (2007), pp. 154–158

[Article](#) | [PDF \(272 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus](#) (6)

Mitra et al., 1993

M. Mitra, S.K. Mahanta, S. Sen, C. Ghosh, A.K. Hati

*Sarcoptes scabiei* in animals spreading to man

Trop. Geogr. Med., 45 (1993), pp. 142–143

[View Record in Scopus](#) | [Cited By in Scopus](#) (6)

Morsy et al., 1995

T.A. Morsy, M.E. Bakr, M.M. Ahmed, M.M. Kotb

Human scabies acquired from a pet puppy

J. Egypt. Soc. Parasitol., 24 (1995), pp. 305–308

Mörner, 1992

T. Mörner

Sarcoptic mange in Swedish wildlife

Rev. Sci. Tech., 11 (1992), pp. 115–121

Munang'andu et al., 2010

H.M. Munang'andu, V.M. Siamudaala, W. Matandiko, M. Munyeme, M. Chembensofu, E. Mwase

Sarcoptes mite epidemiology and treatment in African buffalo (*Syncerus caffer*) calves captured for translocation from the Kafue game management area to game ranches

BMC Vet. Res., 6 (2010), p. 29

[Full Text via CrossRef](#)

Nayel and Abu-Samra, 1986

N.M. Nayel, M.T. Abu-Samra

Experimental infection of the one-humped camel (*Camelus dromedarius*) with *Sarcoptes scabiei* var. *cameli* and *S. scabiei* var. *ovis*

Ann. Trop. Med. Parasitol., 80 (1986), pp. 553–561

[View Record in Scopus](#) | [Cited By in Scopus](#) (5)

Oleaga et al., 2008

A. Oleaga, R. Casais, P. González-Quirós, M. Prieto, C. Gortazar

Sarcoptic mange in red deer from Spain: improved surveillance or disease emergence?

Vet. Parasitol., 154 (2008), pp. 103–113

[Article](#) | [PDF \(1112 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(13\)](#)

Orkin and Maibach, 1978

M. Orkin, H.I. Maibach

Scabies in children. Symposium on Pediatric Dermatology

*Pediatr. Clin. North Am.*, 25 (1978), pp. 371–386

[View Record in Scopus](#) | [Cited By in Scopus \(7\)](#)

Orkin and Maibach, 1991

Orkin, M., Maibach, H.I., 1991. Ectoparasitic diseases. In: Orkin, M., Maibach, H., Dahl Norwalk, M.V. (Eds.), *Dermatology*, vol. 3. Appleton and Lange, VA, pp. 205–214.

Pence et al., 1983

D.B. Pence, L.A. Windberg, B.C. Pence, R. Sprowls

The epizootiology and pathology of sarcoptic mange in coyotes, *Canis latrans*, from South Texas

*J. Parasitol.*, 69 (1983), pp. 1100–1115

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(39\)](#)

Pence and Windberg, 1994

D.B. Pence, L.A. Windberg

Impact of a sarcoptic mange epizootic on a coyote population

*J. Wildl. Mgmt.*, 58 (1994), pp. 624–633

[Full Text via CrossRef](#)

Pence and Ueckermann, 2002

D.B. Pence, E. Ueckermann

Sarcoptic mange in wildlife

*Rev. Sci. Tech.*, 21 (2002), pp. 385–398

[View Record in Scopus](#) | [Cited By in Scopus \(85\)](#)

Pérez et al., 1997

J.M. Pérez, I. Ruiz-Martínez, J.E. Granados, R.C. Soriguer, P. Fandos

The dynamics of sarcoptic mange in the ibex population of Sierra Nevada in Spain – influence of climatic factors

J. Wildl. Res., 2 (1997), pp. 86–89

[View Record in Scopus](#) | [Cited By in Scopus \(32\)](#)

Pérez et al., 2011

J.M. Pérez, J.E. Granados, M. Sarasa, E. Serrano

Usefulness of estimated surface area of damaged skin as a proxy of mite load in the monitoring of sarcoptic mange in free-ranging populations of Iberian wild goat, *Capra pyrenaica*

Vet. Parasitol., 176 (2011), pp. 258–264

[Article](#) | [PDF \(525 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(9\)](#)

Pirilä et al., 1967

V.V. Pirilä, O.P. Salo, R. Kiistala

Scabies Norvegica

Acta Derm. Venereol., 47 (1967), pp. 267–268

[View Record in Scopus](#) | [Cited By in Scopus \(1\)](#)

Polley, 2005

L. Polley

Navigating parasite webs and parasite flow: emerging and re-emerging parasitic zoonoses of wildlife origin

Int. J. Parasitol., 35 (2005), pp. 1279–1294

[Article](#) | [PDF \(280 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(37\)](#)

Price, 1980

P.W. Price

Evolutionary Biology of Parasites

Princeton University Press, Princeton, NJ (1980)

Rasero et al., 2010

R. Rasero, L. Rossi, D. Soglia, S. Maione, P. Sacchi, L. Rambozzi, S. Sartore, R.C. Soriguer, V. Spalenza, S. Alasaad

Host taxon-derived Sarcoptes mite in European wild animals revealed by microsatellite markers

Biol. Conserv., 143 (2010), pp. 1269–1277

[Article](#) | [PDF \(784 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(14\)](#)

Rossi et al., 2007

L. Rossi, C. Fraquelli, U. Vesco, R. Permunian, G.M. Somnavilla, G. Carmignola, M. Da Pozzo, P.G. Meneguz

Descriptive epidemiology of a scabies epidemic in chamois in the Dolomite Alps

Italy. Eur. J. Wildl. Res., 53 (2007), pp. 131–141

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(28\)](#)

Ruch, 1959

T.C. Ruch

Diseases of laboratory primates

W.B. Saunders Company, Philadelphia (1959)

Ryser-Degiorgis et al., 2002

M.P. Ryser-Degiorgis, A. Ryser, L.N. Bacciarini, C. Angst, B. Gottstein, M. Janovsky, U. Breitenmoser

Notoedric and sarcoptic mange in free-ranging lynx from Switzerland

J. Wildl. Dis., 2002 (38) (2002), pp. 228–232

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(21\)](#)

Samuel, 1981

W.M. Samuel

Attempted experimental transfer of sarcoptic mange (*Sarcoptes scabiei*, Acarina: Sarcoptidae) among red fox, coyote, wolf and dog

J. Wildl. Dis., 17 (1981), pp. 343–347

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (17)

Sanderson et al., 2007

H. Sanderson, B. Laird, L. Pope, R. Brain, C. Wilson, D. Johnson

Assessment of the environmental fate and effects of ivermectin in aquatic mesocosms

Aquat. Toxicol., 85 (2007), pp. 229–240

[Article](#) | [PDF \(1186 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus](#) (38)

Skerratt et al., 1998

L.F. Skerratt, R.W. Martin, K.A. Handasyde

Sarcoptic mange in wombats

Aust. Vet. J., 76 (1998), pp. 408–410

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (21)

Skerratt and Beveridge, 1999

L.F. Skerratt, I. Beveridge

Human scabies of wombat origin

Aust. Vet. J., 9 (1999), p. 607

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus](#) (10)

Stanton et al., 1987

B. Stanton, S. Khanam, H. Nazrul, S. Nurani, T. Khair

Scabies in urban Bangladesh

J. Trop. Med. Hyg., 90 (1987), pp. 219–226

[View Record in Scopus](#) | [Cited By in Scopus \(19\)](#)

Stone et al., 1972

W.B. Stone, E. Parks, B.L. Weber, F.J. Parks

Experimental transfer of sarcoptic mange from red foxes and wild canids to captive wildlife and domestic animals. N.Y

Fish Game J., 19 (1972), pp. 1–11

Terry et al., 2001

B.C. Terry, F. Kanjah, F. Sahr, S. Kortequee, I. Dukulay, A.A. Gbakima

Sarcoptes scabiei infestation among children in a displacement camp in Sierra Leone

Public Health, 115 (2001), pp. 208–211

[Article](#) | [PDF \(60 K\)](#) | [View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(23\)](#)

Thomsett, 1968

L.R. Thomsett

Mite infestation of man contracted from dogs and cats

Br. Med. J., 3 (1968), pp. 93–95

[View Record in Scopus](#) | [Full Text via CrossRef](#) | [Cited By in Scopus \(16\)](#)

Walton et al., 1997

S.F. Walton, B.J. Currie, D.J. Kemp

A DNA fingerprinting system for the ectoparasite *Sarcoptes scabiei*

Mol. Biochem. Parasitol., 85 (1997), pp. 187–196

[Article](#) | [PDF \(561 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(40\)](#)

Walton et al., 1999

S.F. Walton, J.L. Choy, A. Bonson, A. Valle, J. McBroom, D. Taplin, L. Arlian, J.D. Mathews, B. Currie, D.J. Kemp



Genetically distinct dog-derived and human-derived *Sarcoptes scabiei* in scabies-endemic communities in northern Australia

Am. J. Trop. Med. Hyg., 61 (1999), pp. 542–547

[View Record in Scopus](#) | [Cited By in Scopus \(54\)](#)

Walton et al., 2004a

S.F. Walton, A. Dougall, S. Pizzutto, D. Holt, D. Taplin, L.G. Arlian, M. Morgan, B.J. Currie, D.J. Kemp

Genetic epidemiology of *Sarcoptes scabiei* (Acari: Sarcoptidae) in northern Australia

Int. J. Parasitol., 34 (2004), pp. 839–849

[Article](#) | [PDF \(214 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(47\)](#)

Walton et al., 2004b

S.F. Walton, D.C. Holt, B.J. Currie, D.J. Kemp

Scabies: new future for a neglected disease

Adv. Parasitol., 57 (2004), pp. 309–376

[Article](#) | [PDF \(706 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(66\)](#)

WHO, 2009

WHO (World Health Organisation), 2009. Available from: <http://www.who.int/en/>.

Young, 1975

E. Young

Some important parasitic and other diseases of lion, *Panthera leo*, in the Kruger National Park

J. S. Afr. Vet. Assoc., 46 (1975), pp. 181–183

[View Record in Scopus](#) | [Cited By in Scopus \(10\)](#)

Zahler et al., 1999

M. Zahler, A. Essig, R. Gothe, H. Rinder

Molecular analyses suggest monospecificity of the genus *Sarcoptes* (Acari: Sarcoptidae)

Int. J. Parasitol., 29 (1999), pp. 759–766

[Article](#) | [PDF \(143 K\)](#) | [View Record in Scopus](#) | [Cited By in Scopus \(49\)](#)

Zeh, 1974

J.B. Zeh

Infestation of sarcoptic mange on red fox in New York

New York Fish Game J., 21 (1974), pp. 182–183

[View Record in Scopus](#) | [Cited By in Scopus \(1\)](#)

Corresponding author contact information

Corresponding authors. Address: Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC), Avda. Américo Vespucio s/n 41092 Sevilla, Spain. Tel.: +41 44 6354914; fax: +41 44 6355711 (S. Alasaad), tel./fax: +86 931 8342837 (X.-Q. Zhu).

# Mammals

