

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Accuracy of Plateau Pressure and Stress Index to Identify Injurious Ventilation in Patients with Acute Respiratory Distress Syndrome.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/143250> since

*Published version:*

DOI:10.1097/ALN.0b013e3182a05bb8

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[Anesthesiology, 119(4), 2013, doi: 10.1097/ALN.0b013e3182a05bb8]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*[ <http://journals.lww.com/anesthesiology/toc/2013/10000> ]*

# ACCURACY OF PLATEAU PRESSURE AND STRESS INDEX TO IDENTIFY INJURIOUS VENTILATION IN PATIENTS WITH ACUTE RESPIRATORY DISTRESS SYNDROME<sup>1</sup>

Pier Paolo Terragni MD <sup>2,5</sup>, Claudia Filippini PhD <sup>2,6</sup>, Arthur S. Slutsky MD, <sup>3,7</sup>, Alberto Birocco MD, <sup>2,8</sup>,  
Tommaso Tenaglia MD, <sup>2,8</sup>, Salvatore Grasso MD <sup>4,5</sup>, Tania Stripoli MD <sup>4,6</sup>, Daniela Pasero MD <sup>2,5</sup>,  
Rosario Urbino MD <sup>2,5</sup>, Vito Fanelli MD PhD <sup>2,5</sup>, Chiara Faggiano MD <sup>2,8</sup>,  
Luciana Mascia MD PhD <sup>2,9</sup>, and V. Marco Ranieri MD <sup>2,7</sup>

**Word count:** *ABSTRACT: 242; INTRODUCTION: 282; DISCUSSION: 902*

**Abbreviated title:** Accuracy of *PPLAT* and stress index

**Conflict of Interests statement:** Dr. Ranieri consults for Maquet Critical Care (Solna, Sweden) that, as far as we know, is the only company that implements the stress index in a commercially available ventilator, is the principal investigator of clinical trials on Extra-Corporeal Membrane Oxygenation conducted by Maquet Cardio-Vascular (Rastatt, Germany), Hemodec (Salerno, Italy) and is member of the steering committee of a clinical trial conducted by Faron (Turku, Finland) on the use of a pharmacological agent in patients with Acute Respiratory Distress Syndrome. Dr. Slutsky consults for Maquet Critical Care (Solna, Sweden) that is, as far as we know, is the only company that implements the stress index in a commercially available ventilator, consults for Novalung (Heilbronn, Germany), and Gambro (Lund, Sweden) that produce Extra-Corporeal Membrane Oxygenation equipment and is the principal investigator of clinical trial on Extra-Corporeal Membrane Oxygenation devices that can be used in patients with Acute Respiratory Distress Syndrome; St. Michael's Hospital (Toronto, Canada) receives royalties from Maquet Critical Care (Solna, Sweden) for patents related to Neurally Adjusted Ventilatory Assist (not for stress index). All the others do not have any conflict of interests to declare.

Corresponding author and institution to which the work should be attributed:

V. Marco Ranieri, MD

Università di Torino

Dipartimento di Anestesiologia e Rianimazione, Ospedale S. Giovanni Battista-Molinette

Corso Dogliotti 14, 10126 Torino

Tel: 39-011-633-4001 Fax: 39-011-696-0448

e-mail: marco.ranieri@unito.it

1. Supported by Ministero Università e Ricerca (grant # PR60ANRA04, Rome Italy) and Regione Piemonte (grant # 2ZBT-04, Turin, Italy)
2. Dipartimento di Anestesiologia e Rianimazione, Università di Torino, Ospedale S. Giovanni Battista-Molinette, Torino, Italy.
3. Keenan Research Center at the Li Ka Shing Knowledge Institute of St. Michael's Hospital; Department of Medicine and Interdepartmental Division of Critical Care Medicine, University of Toronto
4. Sezione di Anestesiologia e Rianimazione, Dipartimento dell'Emergenza e dei Trapianti d'Organo, Università di Bari, Italy
5. Assistant Professor
6. Statistician
7. Professor
8. Resident
9. Associate Professor

The authors wish to thank: **Manuela Ceccarelli**, PhD (Statistician at the Epidemiology of Cancer Unit, Città della Salute e della Scienza and CPO Piemonte, Turin, Italy) for her help in the statistical analysis; **Ottavio Davini**, MD, Chief Radiologist and **Giovanni Gandini**, MD, Professor (both at the Dipartimento di Radiologia, Università di Torino, Ospedale S. Giovanni Battista-Molinette, Torino, Italy) for performing all CT-scans; **Eleonora Corno**, MD, **Eleonora Menaldo**, MD, and **Giulio Rosboch**, MD (all residents at Dipartimento di

Anestesiologia e Rianimazione, Università di Torino, Ospedale S. Giovanni Battista-Molinette, Torino, Italy)  
for their help in data collection.

## Abstract

**Background:** Guidelines suggest plateau pressure ( $P_{PLAT}$ )  $\leq 30$  cmH<sub>2</sub>O for patients with acute respiratory distress syndrome but ventilation may still be injurious despite adhering to this guideline. The shape of the airway pressure vs. time curve (*STRESS INDEX*) may identify injurious ventilation. We assessed accuracy of  $P_{PLAT}$  and *STRESS INDEX* to identify morphological indexes of injurious ventilation.

**Methods:** Indexes of lung aeration (computerized-tomography) associated with injurious ventilation were used as a “*reference standard*”. Threshold values of  $P_{PLAT}$  and *STRESS INDEX* were determined assessing the receiver-operating characteristics (“*training set*”  $N=30$ ). Accuracy of these values was assessed in a second group of patients (“*validation set*”  $N=20$ ).  $P_{PLAT}$  and *STRESS INDEX* were partitioned between respiratory system ( $P_{PLAT,RS}$  and *STRESS INDEX,RS*) and lung ( $P_{PLAT,L}$  and *STRESS INDEX,L*) (esophageal pressure; “*physiological set*”  $N=50$ ).

**Results:** Sensitivity and specificity of  $P_{PLAT}>30$  cmH<sub>2</sub>O were 0.06 (95% CI: [0.002 to 0.30]) and 1.0 (95% CI: [0.87 to 1.00]).  $P_{PLAT}>25$ cmH<sub>2</sub>O and a *STRESS INDEX* $>1.05$  best identified morphological indexes associated with injurious ventilation. Sensitivity and specificity of these values were 0.75 (95% CI:[0.35 to 0.97]) and 0.75 (95% CI:[0.43 to 0.95]) for  $P_{PLAT}>25$ cmH<sub>2</sub>O vs. 0.88 (95% CI:[0.47 to 1.00]) and 0.50 (95% CI:[0.21 to 0.79]) for *STRESS INDEX* $>1.05$ .  $P_{PLAT,RS}$  did not correlate with  $P_{PLAT,L}$  ( $R^2=0.0099$ ); *STRESS INDEX,RS* and *STRESS INDEX,L* were correlated ( $R^2=0.762$ ).

**Conclusions:** The best threshold values for discriminating morphological indexes associated with injurious ventilation were  $P_{PLAT,RS}>25$  cmH<sub>2</sub>O and *STRESS INDEX,RS* $>1.05$ . While a substantial discrepancy between  $P_{PLAT,RS}$  and  $P_{PLAT,L}$  occurs, *STRESS INDEX,RS* reflects *STRESS*

*INDEX, L.*

## Introduction

The acute respiratory distress syndrome (ARDS) is a type of pulmonary inflammatory response to various inciting events characterized by hypoxemia and bilateral radiographic opacities<sup>1</sup> with non-aerated regions in the dependent lung and relatively normally aerated regions in the non-dependent lung<sup>2-4</sup>. Inappropriate ventilatory settings may over-distend the normally aerated lung and/or continuously open and close the non-aerated regions causing ventilator-induced lung injury<sup>5</sup>.

Current guidelines recommend keeping end-inspiratory plateau airway pressure ( $P_{PLAT}$ )  $\leq 30$  cmH<sub>2</sub>O<sup>6,7</sup>, based on a randomized clinical trial demonstrated that limiting tidal volume ( $V_T$ ) to 6 ml/kg predicted body weight and  $P_{PLAT}$  to 30 cmH<sub>2</sub>O decreased absolute mortality by 9%<sup>8</sup>. However, these recommendations are challenged by results of recent studies showing that (a) ARDS patients may be exposed to forces which can induce injurious ventilation despite values of  $P_{PLAT} \leq 30$  cmH<sub>2</sub>O<sup>9-11</sup>; (b) impairment of chest wall mechanics compromises the ability of  $P_{PLAT}$  to reflect over-distension<sup>12-14</sup>. Another approach to assess the propensity for injurious ventilation is to assess the *STRESS INDEX* based on the shape of the airway pressure vs. time curve during constant flow<sup>15-18</sup>. Although used in clinical studies<sup>19-22</sup> and implemented in a commercially available ventilator<sup>23</sup>, the accuracy of the *STRESS INDEX* to assess the propensity for injurious ventilation has not been tested in humans, and has been questioned in the context of impairment in chest wall mechanics<sup>24,25</sup>.

In the current study we assessed the diagnostic accuracy of  $P_{PLAT}$  and *STRESS INDEX* to identify ventilator settings likely to produce injurious ventilation. As standards we used indexes of lung aeration (computerized-tomography) associated with injurious ventilation<sup>9,10</sup>. In a separate patient cohort, we examined the impact of chest wall mechanics on the utility of  $P_{PLAT}$  and *STRESS INDEX* to identify propensity for injurious ventilation.



## Material and Methods

The institutional review board (Comitato Etico Interaziendale AUO S. Giovanni Battista e CTO di Torino, Italy) approved the study. As the patients were incompetent, patients were included into the study and consent was delayed. The family was informed of the study (although not required). Written permission for using collected data was hence obtained from the patient (if competent) or from the family (in case of death or if the patient remained incompetent)<sup>2</sup>.

Patients admitted from January 2007 to February 2012 to the intensive care units of the Molinette (Turin) and Policlinico (Bari) hospitals were considered for enrollment when the following criteria were met: age  $\geq 18$  years; diagnosis of ARDS<sup>26</sup>. Exclusion criteria were:  $>3$  days elapsed since ARDS criteria were met and mechanical ventilation was initiated; history of ventricular fibrillation or tachyarrhythmia, unstable angina or myocardial infarction within preceding month; chest tube with persistent air leak; pre-existing chronic obstructive pulmonary disease; pregnancy; known intracranial abnormality. Measurements were interrupted and patients withdrawn from the study if any of the following *a priori* defined conditions occurred: (a) presence of inspiratory efforts during measurement of respiratory mechanics despite infusion of sedatives and respiratory muscle paralytics; (b) drop in arterial oxygen saturation  $<80\%$ ; (c) decrease in mean arterial pressure  $>10\%$  of baseline despite 500 ml intravenous bolus.

### MEASUREMENTS

All patients were ventilated (SERVOi, Maquet, Lund, Sweden) according to the "ARDSNet" protective ventilatory strategy<sup>8</sup>. Measurements were performed during absence of spontaneous respiratory muscle effort obtained by increasing doses of midazolam (up to

10mg/h) and/or propofol (150mg/h increments every 10 min)<sup>27,28</sup> or use of neuromuscular blockade (cis-atracurium besylate 2-8 gamma/kg/min). Absence of spontaneous effort was confirmed by inspecting flow and airway pressure traces during an end-expiratory pause of 4-5 seconds<sup>27,28</sup>.

## **RESPIRATORY MECHANICS**

Flow (heated pneumotachograph Fleisch No. 2; Fleisch, Lausanne, Switzerland and differential pressure transducer; Diff-Cap, Special Instruments, Nordlingen, Germany), volume and airway opening pressure ( $P_{aw}$ ) (Special Instruments Digima-Clic  $\pm$  100 cmH<sub>2</sub>O; Nordlingen, Germany) were measured as previously described<sup>16,18,29</sup>. In a subset of 50 patients, intra-thoracic pressure was evaluated by assessment of esophageal pressure ( $P_{ES}$ )<sup>30</sup> using a thin latex balloon-tipped catheter system (Microteck Medical B.V., Zutphen, Netherlands) connected by a polyethylene catheter to a pressure transducer (Special Instruments Digima-Clic  $\pm$  100 cmH<sub>2</sub>O; Nordlingen, Germany)<sup>31</sup>.

All the described variables were displayed and collected for 5 minutes on a personal computer through a 12-bit analog-to-digital converter board (National Instrument DAQCard 700; Austin, TX, USA) at a 200 Hz sample rate of (KleisTEK Engineering; Bari, Italy). End-expiratory and end-inspiratory occlusions were performed. Signals were averaged and smoothed by a filter that averaged the signal over a 120 ms time window<sup>18</sup>.

## ***END-INSPIRATORY PLATEAU PRESSURE***

$P_{PLAT}$  of the respiratory system ( $P_{PLATRS}$ ) was the value of  $P_{aw}$  after an end-inspiratory occlusion. In the subset of patients in whom  $P_{ES}$  was measured, end-inspiratory chest wall plateau pressure ( $P_{PLATCW}$ ) was measured as the variation in  $P_{ES}$  between end-expiratory and

end-inspiratory occlusions; end-inspiratory plateau pressure of the lung ( $P_{PLATL}$ ) were estimated as  $P_{PLATRS} - P_{PLATCW}$ <sup>12-14</sup>.

### ***STRESS INDEX***

The software identified the beginning and the end of each recorded breath by means of a threshold value (0.1 liter/second) on the flow signal<sup>17-19</sup>. Trans-pulmonary pressure ( $P_L$ ) was calculated as  $P_{AW} - P_{ES}$ <sup>12-14</sup>. Individual flow,  $P_{aw}$  and  $P_L$  signals were averaged and smoothed by a filter that averaged the signal over a 120 ms time window. On the resulting mean flow, the software first identified the steady flow level and then the largest portion of flow signal that was considered to be steady flow  $\pm 3\%$ <sup>17-19</sup>. The beginning and the end of this constant portion were marked by cursors. To eliminate on and off flow transient, the constant flow portion was further narrowed by adding a 50 ms offsets after the beginning (**time 0**) and before the end (**time 1**) of the constant flow portion. The portion of the mean  $P_{aw}$ -time and  $P_L$ -time curves encompassed in the time interval **time 0-time 1** was fitted to the equations:

$$P_{aw} = a_{aw} \cdot (\mathbf{time\ 0-time\ 1})^{b_{aw} + c_{aw}}$$

$$P_L = a_L \cdot (\mathbf{time\ 0-time\ 1})^{b_L + c_L}$$

using the *Levenberg-Marquardt* algorithm<sup>17-19</sup>. Values of  $R^2$  were computed and displayed. The coefficients  $b_{aw}$  and  $b_L$  (*STRESS INDEX<sub>RS</sub>* and *STRESS INDEX<sub>L</sub>*) are dimensionless number that describe the shape of the  $P_{aw}$ -time and of the  $P_L$ -time curves. Values of coefficient  $b < 1$  indicate that elastance decreases with time, whereas elastance increases with time for values of coefficient  $b > 1$ . Finally,  $b = 1$  indicates a constant elastance during tidal inflation (**Figure 1**)<sup>17-19</sup>.

### ***COMPUTERIZED TOMOGRAPHY ASSESSMENT OF “VENTILATOR INDUCED LUNG INJURY”***

As soon as targets of the ventilatory protocols were reached and respiratory and hemodynamic parameters (measured at 20-30-min intervals) were stable, patients were transferred to the computerized tomography (CT) scan facility. During the transport and the exam, the ventilator and the ventilator settings were the one used for the clinical management; particular attention was paid to avoid ventilator disconnection<sup>9,10</sup>. Lung scanning was performed from the apex to the base using a Light Speed Qx/i (General Electric Medical System, Milwaukee, WI, USA) at the end of end-expiratory and end-inspiratory occlusions<sup>9,10</sup>. The ventilator settings were identical to those previously set. The CT scanner was set as previously described<sup>9,10</sup>. Each section of the right and left lung was chosen by manually drawing the outer boundary along the inside of the ribs and the inner boundary along the mediastinal organs. Pleural effusions were excluded. *Non-aerated* (density between +100 and -100 Hounsfield units), *poorly aerated* (density between -101 and -500 Hounsfield units), *normally aerated* (density between -501 and -900 Hounsfield units), and *hyperinflated* (density between -901 and -1000 Hounsfield units) lung compartments were identified as previously described<sup>3,4,32</sup>.

Volume of the entire lungs (i.e. the sum of gas plus tissue volume) and of each compartment at end-expiration and end-inspiration was measured for each slice as: [(size of the pixel)<sup>2</sup> multiplied by the number of pixels in each compartment] multiplied by the thickness of the CT lung slice<sup>3,4,32</sup>. “***Tidal hyperinflation***” was defined as the volume of the *hyperinflated* compartment at end-inspiration minus the volume of the *hyperinflated* compartment at end-expiration<sup>9,10</sup>. “***Tidal recruitment***” was defined as the volume of the *non-aerated* and of the *poorly aerated* compartments at end-expiration minus the volume at end-inspiration<sup>9,10</sup>. “***Protected tidal inflation***” was the volume of the *normally aerated* compartment at end-inspiration minus the volume of the *normally aerated* compartment at

end-expiration<sup>9,10</sup>. All were expressed as % of total tidal inflation-related changes in CT lung volume.

### ***PULMONARY INFLAMMATORY RESPONSE***

Five-ten min after CT and respiratory mechanics measurements, a broncho-alveolar lavage was performed and stored at -80 °C. as previously described<sup>9,10,33</sup>. Assay for tumor necrosis factor- $\alpha$  soluble receptors, interleukin 6, interleukin 8 and interleukin 1 $\beta$  and interleukin-1 receptor antagonist were carried out using a solid-phase enzyme-linked immunoabsorbent assay (ELISA) method (Diacclone, Milan, Bender Med Systems, Milan, ITALY and BioSource International Inc., Camarillo, USA)<sup>9,10,33</sup>.

### **STUDY DESIGN**

In *Phase 1*, we evaluated the diagnostic accuracy of  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  to identify the propensity for injurious ventilation using CT criteria to assess the degree of overdistension<sup>9,10</sup>. Accuracy of  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  was determined in a first group of patients (“*training set*”) to select the threshold values that discriminated best between patients with and without the condition of interest; the accuracy of these values was prospectively assessed in a second group of patients (“*validation set*”)<sup>34</sup>. In *Phase 2*, we addressed the question of how chest wall mechanics affects interpretation of  $P_{PLAT,RS}$  and of the  $STRESS\ INDEX_{RS}$  (“*physiological set*”)<sup>12-14</sup>. In this phase, we did not use CT scan or pulmonary concentration of inflammatory cytokines. Patients were assigned to the different data sets depending on the chronological order in which they entered the study.

The maximal degrees of association between CT scan evidence of “*protected tidal inflation*”, “*tidal hyperinflation*” or “*tidal recruitment*” was identified using cluster analysis with cubic clustering criteria<sup>9</sup>. Cluster analysis entails grouping similar objects into distinct,

mutually exclusive subsets referred to as *clusters*; elements within a cluster share a high degree of “natural association,” whereas the clusters are relatively distinct from one another<sup>35</sup>.

Values of  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  that best differentiated the patients who were ventilated with CT scan evidence of “*protected tidal inflation*” from those in whom tidal volume and pressure limitation caused CT scan evidence of “*tidal hyperinflation*” or “*tidal recruitment*” were determined assessing the receiver-operating-characteristics curve<sup>36</sup>.

The area under the receiver-operating characteristics curve for  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  was calculated and confidence intervals (CI) reported. The selected threshold values were those that minimized false negative classifications (i.e. patients who were thought to be protected when in fact they were not) with a specificity value not lower than 0.5. This decision was based on the assumption that from a clinical perspective, a false negative result is worse than a false positive.

The predictive power of the previously selected values of  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  was estimated using the previously selected cut-off values that best discriminated patients with CT scan evidence of “*protected tidal inflation*” from “*tidal hyperinflation*” or “*tidal recruitment*”.

Values of  $STRESS\ INDEX_{RS}$  and  $P_{PLAT,RS}$ , and values of  $STRESS\ INDEX_{L}$  and  $P_{PLAT,L}$  were compared in a third set of patients ventilated according to the “*ARDSNet*” protective ventilator strategy<sup>8</sup>.

## STATISTICAL ANALYSIS

Results are expressed as mean±SD;  $P<0.05$  was considered significant. Comparisons of continuous and categorical data between groups were performed using unpaired t-tests and chi-squared tests. Regression was performed using least-squares. Since values of cytokine

concentrations were not normally distributed,  $\log_{10}$  transformation was performed prior to applying parametric statistics.

A true-positive was defined when  $P_{PLAT}$  and  $STRESS\ INDEX$  predicted “*tidal hyperinflation*” or “*tidal recruitment*” and CT scan analysis was confirmatory. A true-negative was defined when  $P_{PLAT}$  and  $STRESS\ INDEX$  predicted absence of “*tidal hyperinflation*” or “*tidal recruitment*” and CT scan analysis was confirmatory. A false-positive was defined when  $P_{PLAT}$  and  $STRESS\ INDEX$  value predicted presence of “*tidal hyperinflation*” or “*tidal recruitment*” and CT scan was not confirmatory. A false-negative was defined when  $P_{PLAT}$  and  $STRESS\ INDEX$  value predicted absence of “*tidal hyperinflation*” or “*tidal recruitment*” and CT scan analysis was not confirmatory.

Standard formulae were used to calculate sensitivity, specificity, and positive and negative predictive values. Positive and negative likelihood ratios were calculated (SAS software, version 9.1.3; SAS Institute, Cary, North Carolina, USA; MedCalc version 11.1.1, MedCalc software BVBA, Ostend, Belgium).

## Results

Some of the results reported in the present investigation include data obtained from patients enrolled in previously published studies<sup>1,9,10,20</sup>.

Of the 110 patients enrolled, 10 were excluded for the following reasons: >3 days since mechanical ventilation initiation (N=5); chest tube with persistent air leak (N=2); unilateral lung disease (N=2); spontaneous respiratory effort during physiological measurements (N=1). Of the remaining patients 100 patients, 50 patients were included in **PHASE 1** (30 in the “training set” and 20 in the “validation set”), and 50 patients were included in **PHASE 2** (**Table 1**).

The volume of “*protected ventilation*” and of “*tidal hyperinflation*” identified 2 clusters of patients. In a cluster of 28 patients (16 in the “training set” and 12 in the “validation set”), “*tidal hyperinflation*” was  $8.36 \pm 5.51$  (“training set”) and  $9.91 \pm 4.31\%$  (“validation set”), while “*protected ventilation*” was  $71.20 \pm 8.05$  (“training set”) and  $75.68 \pm 8.01\%$  (“validation set”) of the total tidal-inflation associated change in CT lung compartments. These patients were considered relatively protected from injurious ventilation (“*PROTECTED*”). In a cluster of 22 patients (14 in the “training set” and 8 in the “validation set”), “*tidal hyperinflation*” was  $53.76 \pm 7.92$  (“training set”) and  $50.91 \pm 21.78\%$  (“validation set”), and “*protected ventilation*” was  $28.55 \pm 16.33$  (“training set”) and  $25.91 \pm 14.71\%$  (“validation set”) of the total tidal-inflation associated change in CT lung compartments. These patients were considered relatively not protected from injurious ventilation (“*NON PROTECTED*”). “*Tidal recruitment*” was  $20.44 \pm 7.03$  (“training set”) and  $14.81 \pm 7.73\%$  (“validation set”) in “*PROTECTED*” and  $17.69 \pm 8.15$  (“training set”) and  $23.18 \pm 9.32\%$  (“validation set”) in the “*NON-PROTECTED*”. As such, “*tidal recruitment*” could not be identified in the cluster analysis as a distinct entity that could define a “protected” vs. a “non-



protected” ventilator setting and therefore could not be used as additional criterion to define a non-protected tidal inflation.

Ventilator settings, and biological variables in the “*PROTECTED*” and “*NON PROTECTED*” clusters are shown in **Table 2**. Pulmonary concentrations of the inflammatory cytokines were lower in *PROTECTED* than in *NON-PROTECTED* ( $P < 0.05$ ).

The areas under the receiver-operating characteristics curves for  $P_{PLAT,RS}$  and  $STRESS INDEX_{RS}$  (0.833; 95% CI:[0.621 to 0.954] and 0.917; 95%CI:[0.724 to 0.990], respectively) (**Figure 2**) were both significantly ( $P=0.001$ ) larger than that of an arbitrary test that would be expected *a priori* to have no discriminatory value. Sensitivity and specificity of the value of  $P_{PLAT,RS}$  currently suggested by guidelines ( $>30$  cmH<sub>2</sub>O) were 0.06 (95% CI: [0.002 to 0.30]) and 1.0 (95% CI: [0.87 to 1.00]), respectively. The threshold value of  $P_{PLAT,RS}$  that best identified *NON-PROTECTED* patients was 25 cmH<sub>2</sub>O; sensitivity and specificity were 0.82 (95% CI: [0.48 to 0.98]) and 0.67 (95% CI: [0.35 to 0.90]), respectively. A threshold value of  $STRESS INDEX_{RS} > 1.05$  best identified *NON-PROTECTED* patients; sensitivity and specificity were 0.82 (95% CI: [0.48 to 0.98]) and 0.83 (95% CI: [0.52 to 0.98]), respectively.

Sensitivity and specificity of  $P_{PLAT,RS} > 25$  cmH<sub>2</sub>O to identify “*NON-PROTECTED*” patients were 0.75 (95% CI: [0.35 to 0.97]) and 0.75 (95% CI: [0.43 to 0.95]), respectively. Sensitivity and specificity of a  $STRESS INDEX_{RS} > 1.05$  to identify “*NON-PROTECTED*” patients were 0.88 (95% CI: [0.47 to 1.00]) and 0.50 (95% CI: [0.21 to 0.79]) (**Table 3**).

The correlation coefficients relating  $STRESS INDEX_{RS}$  vs.  $STRESS INDEX_{L}$  and of  $P_{PLAT,RS}$  vs.  $P_{PLAT,L}$  were 0.762 and 0.0099, respectively (**Figure 3**).

## Discussion

The main findings of the present investigation are: (a) the value of  $P_{PLAT,RS}$  recommend by current guidelines ( $\leq 30$  cmH<sub>2</sub>O) does not accurately discriminate patients with CT-scan indexes of tidal hyperinflation; (b) the discriminating threshold values that identify associate  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  to CT pattern of lung aeration related to tidal hyperinflation are  $<25$  cmH<sub>2</sub>O and not greater than 1.05, respectively; (c) while  $STRESS\ INDEX_{RS}$  represents a reasonable reflection of  $STRESS\ INDEX_{L}$ , there is substantial discrepancy when using  $P_{PLAT,RS}$  vs.  $P_{PLAT,L}$ .

Rigorous statistical methods have been developed to evaluate the degree of agreement between a test and the best available method for establishing the presence/absence of the condition of interest<sup>37</sup>. Accordingly to these methods, (a) we included a study population representative of ARDS patients<sup>26</sup> excluding only those patients that were mechanically ventilated for  $>72$  hours or in whom measurements of respiratory mechanics could not be performed; (b) we used previously established methods to measure physiological variables<sup>31</sup>; (c) we developed threshold values of  $P_{PLAT,RS}$  and  $STRESS\ INDEX_{RS}$  through the analysis of the receiver-operating characteristics curves obtained in a training set (30 patients) and then evaluated their accuracy in a validation set (20 patients); (c) we selected threshold values giving priority to those that optimized sensitivity minimizing false negative classifications (i.e. patients who were thought to be protected when in fact they were not). This decision was based on the assumption that from a clinical perspective, a false negative result is less acceptable than a false positive. However our use of CT scan indexes of lung aeration as a “*reference standard*” for injurious ventilation has some weaknesses. We used CT scan evidence of “*protected tidal inflation*” and “*tidal hyperinflation*” to select a cluster characterized by a predominant “*protected tidal inflation*” and a cluster characterized by

predominant “*tidal hyperinflation*”. These two clusters may represent different ranges of a continuum<sup>4,38</sup>) and, since tidal recruitment could not be addressed by cluster analysis, the terms “*PROTECTED*” and “*NON PROTECTED*” should be referred only to tidal hyperinflation. Moreover, although we found a concentration of inflammatory mediators higher in patients included in the *NON-PROTECTED* cluster than in patients included in the *PROTECTED* cluster<sup>9,10</sup>, this may be a marker of severity of ARDS and not solely reflect the degree of hyperinflation.

A clinical trial<sup>8</sup> and observational studies<sup>39,40</sup> have demonstrated that limiting  $V_T$  to 6 ml/kg predicted body weight and  $P_{PLAT,RS}$  to 30 cmH<sub>2</sub>O improves survival. Our data show that the threshold value that best identified *NON-PROTECTED* patients was not  $P_{PLAT,RS} > 30$  cmH<sub>2</sub>O but  $P_{PLAT,RS} > 25$  cmH<sub>2</sub>O. These data are in accord with previous studies demonstrating that *tidal hyperinflation* may occur despite limiting  $V_T$  to 6 ml/kg predicted body weight and  $P_{PLAT,RS}$  to 30 cmH<sub>2</sub>O<sup>9-11</sup>.

Previous studies proposed analyzing the  $P_{AW}$ -t curve during constant flow to assess the mechanical properties of the respiratory system of ARDS patients<sup>15,16</sup>. This approach is based on the concept that at constant flow, the rate of change of  $P_{AW}$  with time corresponds to the rate of change of elastance of the respiratory system during tidal inflation<sup>15</sup>, and can be described by a power equation (**pressure** =  $a \cdot \mathbf{time}^{b+c}$ )<sup>16</sup>. A coefficient  $b=1.0$  indicates a linear  $P_{aw}$ -t curve and an unchanging elastance during inflation; coefficient  $b<1.0$  indicates decreasing elastance during inflation; coefficient  $b>1.0$  indicates a increasing elastance. Experimental studies demonstrated that markers of injurious ventilation were minimized using ventilator settings associated with  $0.9 < b < 1.1$  and therefore concluded that the coefficient  $b$  (called *STRESS INDEX*) could be used to detect tidal hyperinflation or tidal recruitment during mechanical ventilation<sup>17,18,41-43</sup>. Although subsequent experimental studies challenged these findings<sup>24,44</sup>, the use of *STRESS INDEX* to detect injurious ventilation has been tested in clinical studies<sup>19-22</sup> and implemented in a commercially available ventilator<sup>23</sup>. We found that a  $STRESS INDEX_{RS} > 1.05$  best identified patients *NON PROTECTED* from injurious

ventilation. The area under the receiver-operating characteristics curve for  $STRESS\ INDEX_{RS}$  and  $P_{PLAT,RS}$  were not statistically different. However, in the validation set, sensitivity of  $P_{PLAT,RS} > 25\text{ cmH}_2\text{O}$  was slightly lower than sensitivity of  $STRESS\ INDEX_{RS} > 1.05$  (0.75 (95% CI:[0.35-0.97]) vs. 0.88 (95% CI: [0.47-1.00])) (**Table 3**).

Our data demonstrate that while alterations of chest wall mechanics may substantially impair the ability of  $P_{PLAT,RS}$  to estimate  $P_{PLAT,L}$ ,  $STRESS\ INDEX_{RS}$  closely reflects  $STRESS\ INDEX_{L}$ . These results may be explained by partitioning the volume-pressure relationship of the respiratory system between the lung and the chest wall<sup>45-47</sup>.  $P_{PLAT,RS}$  as a measure of  $P_{PLAT,L}$  is directly related to the stiffness of the chest wall at the end of an inspiration, which may be substantial in patients with ARDS<sup>12-14</sup>. The  $STRESS\ INDEX$  reflects the *changes* with volume of the elastance of the respiratory system ( $P_{AW}$  vs. time) or of the lung ( $P_L$  vs. time). In the range of changes of lung volume associated with a low tidal volume strategy<sup>8</sup>, the volume-pressure relationship of the chest wall is linear<sup>12,45,48</sup>, and hence the  $STRESS\ INDEX$  should largely reflect the mechanical properties of the lung.

In conclusion, the present study demonstrates that the value of  $P_{PLAT,RS}$  currently recommend by guidelines ( $\leq 30\text{ cmH}_2\text{O}$ ) does not accurately discriminate patients with CT-scan indexes of tidal hyperinflation. The threshold values of  $P_{PLAT,RS}$  and of  $STRESS\ INDEX_{RS}$  that correspond to CT-scan indexes of tidal hyperinflation are  $< 25\text{ cmH}_2\text{O}$  and  $< 1.05$ . While a substantial discrepancy between  $P_{PLAT,RS}$  and  $P_{PLAT,L}$  occurs,  $STRESS\ INDEX_{RS}$  reflects  $STRESS\ INDEX_{L}$  with reasonable accuracy. Clinical trials are required to test whether ventilator settings targeting  $P_{PLAT,RS} \leq 25\text{ cmH}_2\text{O}$  and/or a  $STRESS\ INDEX_{RS} \leq 1.05$  will improve clinical outcomes.

## References

1. Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, Camporota L, Slutsky AS: Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012; 307: 2526-33
2. Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri VM, Quintel M, Russo S, Patroniti N, Cornejo R, Bugeo G: Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med* 2006; 354: 1775-86
3. Puybasset L, Cluzel P, Chao N, Slutsky AS, Coriat P, Rouby JJ: A computed tomography scan assessment of regional lung volume in acute lung injury. The CT Scan ARDS Study Group. *Am J Respir Crit Care Med* 1998; 158: 1644-55
4. Malbouisson LM, Muller JC, Constantin JM, Lu Q, Puybasset L, Rouby JJ: Computed tomography assessment of positive end-expiratory pressure-induced alveolar recruitment in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 163: 1444-50
5. Tremblay LN, Slutsky AS: Ventilator-induced lung injury: from the bench to the bedside. *Intensive Care Med* 2006; 32: 24-33
6. Malhotra A: Low-tidal-volume ventilation in the acute respiratory distress syndrome. *N Engl J Med* 2007; 357: 1113-20
7. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, Reinhart K, Angus DC, Brun-Buisson C, Beale R, Calandra T, Dhainaut JF, Gerlach H, Harvey M, Marini JJ, Marshall J, Ranieri M, Ramsay G, Sevransky J, Thompson BT, Townsend S, Vender JS, Zimmerman JL, Vincent JL: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 2008; 36: 296-327

8. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 342: 1301-8
9. Terragni PP, Rosboch G, Tealdi A, Corno E, Menaldo E, Davini O, Gandini G, Herrmann P, Mascia L, Quintel M, Slutsky AS, Gattinoni L, Ranieri VM: Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2007; 175: 160-6
10. Terragni PP, Del Sorbo L, Mascia L, Urbino R, Martin EL, Birocco A, Faggiano C, Quintel M, Gattinoni L, Ranieri VM: Tidal volume lower than 6 ml/kg enhances lung protection: role of extracorporeal carbon dioxide removal. *Anesthesiology* 2009; 111: 826-35
11. Bellani G, Guerra L, Musch G, Zanella A, Patroniti N, Mauri T, Messa C, Pesenti A: Lung regional metabolic activity and gas volume changes induced by tidal ventilation in patients with acute lung injury. *Am J Respir Crit Care Med* 2011; 183: 1193-9
12. Ranieri VM, Brienza N, Santostasi S, Puntillo F, Mascia L, Vitale N, Giuliani R, Memeo V, Bruno F, Fiore T, Brienza A, Slutsky AS: Impairment of lung and chest wall mechanics in patients with acute respiratory distress syndrome: role of abdominal distension. *Am J Respir Crit Care Med* 1997; 156: 1082-91
13. Gattinoni L, Pelosi P, Suter PM, Pedoto A, Vercesi P, Lissoni A: Acute respiratory distress syndrome caused by pulmonary and extrapulmonary disease. Different syndromes? *Am J Respir Crit Care Med* 1998; 158: 3-11
14. Chiumello D, Carlesso E, Cadringer P, Caironi P, Valenza F, Polli F, Tallarini F, Cozzi P, Cressoni M, Colombo A, Marini JJ, Gattinoni L: Lung stress and strain during mechanical ventilation for acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2008; 178: 346-55

15. Milic-Emili J, Ploysongsang Y: Respiratory mechanics in the adult respiratory distress syndrome. *Crit Care Clin* 1986; 2: 573-84
16. Ranieri VM, Giuliani R, Fiore T, Dambrosio M, Milic-Emili J: Volume-pressure curve of the respiratory system predicts effects of PEEP in ARDS: "occlusion" versus "constant flow" technique. *Am J Respir Crit Care Med* 1994; 149: 19-27
17. Ranieri VM, Zhang H, Mascia L, Aubin M, Lin CY, Mullen JB, Grasso S, Binnie M, Volgyesi GA, Eng P, Slutsky AS: Pressure-time curve predicts minimally injurious ventilatory strategy in an isolated rat lung model. *Anesthesiology* 2000; 93: 1320-8
18. Grasso S, Terragni P, Mascia L, Fanelli V, Quintel M, Herrmann P, Hedenstierna G, Slutsky AS, Ranieri VM: Airway pressure-time curve profile (stress index) detects tidal recruitment/hyperinflation in experimental acute lung injury. *Crit Care Med* 2004; 32: 1018-27
19. Grasso S, Stripoli T, De Michele M, Bruno F, Moschetta M, Angelelli G, Munno I, Ruggiero V, Anaclerio R, Cafarelli A, Driessen B, Fiore T: ARDSnet ventilatory protocol and alveolar hyperinflation: role of positive end-expiratory pressure. *Am J Respir Crit Care Med* 2007; 176: 761-7
20. Grasso S, Terragni P, Birocco A, Urbino R, Del Sorbo L, Filippini C, Mascia L, Pesenti A, Zangrillo A, Gattinoni L, Ranieri VM: ECMO criteria for influenza A (H1N1)-associated ARDS: role of transpulmonary pressure. *Intensive Care Med* 2012; 38: 395-403
21. Guo FM, Ding JJ, Su X, Xu HY, Shi Y: Effects of different levels of end-expiratory positive pressure on lung recruitment and protection in patients with acute respiratory distress syndrome. *Chin Med J (Engl)* 2008; 121: 2218-23
22. Neve V, de la Roque ED, Leclerc F, Leteurtre S, Dorkenoo A, Sadik A, Cremer R, Logier R: Ventilator-induced overdistension in children: dynamic versus low-flow inflation volume-pressure curves. *Am J Respir Crit Care Med* 2000; 162: 139-47

23. Terragni P, Mascia L, Birocco A, Faggiano C, Tenaglia T, Maiolo G, Pernechele J, Degiovanni E, Viscomi E, Ranieri VM: Agreement of two methods for assessing pressure/time curve profile (stress index) in ARDS. *Intensive Care Med* 2011; 37: S195
24. Formenti P, Graf J, Santos A, Gard KE, Faltesek K, Adams AB, Dries DJ, Marini JJ: Non-pulmonary factors strongly influence the stress index. *Intensive Care Med* 2011; 37: 594-600
25. Chiumello D, Gattinoni L: Stress index in presence of pleural effusion: does it have any meaning? *Intensive Care Med* 2011; 37: 561-3
26. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, Lamy M, Legall JR, Morris A, Spragg R: The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994; 149: 818-24
27. Conti G, Vilardi V, Rocco M, DeBlasi RA, Lappa A, Bufi M, Antonelli M, Gasparetto A: Paralysis has no effect on chest wall and respiratory system mechanics of mechanically ventilated, sedated patients. *Intensive Care Med* 1995; 21: 808-12
28. Decailliot F, Demoule A, Maggiore SM, Jonson B, Duvaldestin P, Brochard L: Pressure-volume curves with and without muscle paralysis in acute respiratory distress syndrome. *Intensive Care Med* 2006; 32: 1322-8
29. Grasso S, Mascia L, Del Turco M, Malacarne P, Giunta F, Brochard L, Slutsky AS, Marco Ranieri V: Effects of recruiting maneuvers in patients with acute respiratory distress syndrome ventilated with protective ventilatory strategy. *Anesthesiology* 2002; 96: 795-802
30. Milic-Emili J, Mead J, Turner JM, Glauser EM: Improved Technique for Estimating Pleural Pressure from Esophageal Balloons. *J Appl Physiol* 1964; 19: 207-11



31. Brochard L, Martin GS, Blanch L, Pelosi P, Belda FJ, Jubran A, Gattinoni L, Mancebo J, Ranieri VM, Richard JC, Gommers D, Vieillard-Baron A, Pesenti A, Jaber S, Stenqvist O, Vincent JL: Clinical review: Respiratory monitoring in the ICU - a consensus of 16. *Crit Care* 2012; 16: 219
32. Gattinoni L, Caironi P, Pelosi P, Goodman LR: What has computed tomography taught us about the acute respiratory distress syndrome? *Am J Respir Crit Care Med* 2001; 164: 1701-11
33. Ranieri VM, Suter PM, Tortorella C, De Tullio R, Dayer JM, Brienza A, Bruno F, Slutsky AS: Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: a randomized controlled trial. *JAMA* 1999; 282: 54-61
34. Rennie D: Improving reports of studies of diagnostic tests: the STARD initiative. *JAMA* 2003; 289: 89-90
35. Jones DK, Dardis R, Ervine M, Horsfield MA, Jeffree M, Simmons A, Jarosz J, Strong AJ: Cluster analysis of diffusion tensor magnetic resonance images in human head injury. *Neurosurgery* 2000; 47: 306-13
36. Yang KL, Tobin MJ: A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991; 324: 1445-50
37. Farcomeni A, Ventura L: An overview of robust methods in medical research. *Stat Methods Med Res* 2012; 21: 111-33
38. Rouby JJ, Constantin JM, Roberto De AGC, Zhang M, Lu Q: Mechanical ventilation in patients with acute respiratory distress syndrome. *Anesthesiology* 2004; 101: 228-34

39. Kallet RH, Jasmer RM, Pittet JF, Tang JF, Campbell AR, Dicker R, Hemphill C, Luce JM: Clinical implementation of the ARDS network protocol is associated with reduced hospital mortality compared with historical controls. *Crit Care Med* 2005; 33: 925-9
40. Sakr Y, Vincent JL, Reinhart K, Groeneveld J, Michalopoulos A, Sprung CL, Artigas A, Ranieri VM: High tidal volume and positive fluid balance are associated with worse outcome in acute lung injury. *Chest* 2005; 128: 3098-108
41. de Perrot M, Imai Y, Volgyesi GA, Waddell TK, Liu M, Mullen JB, McRae K, Zhang H, Slutsky AS, Ranieri VM, Keshavjee S: Effect of ventilator-induced lung injury on the development of reperfusion injury in a rat lung transplant model. *J Thorac Cardiovasc Surg* 2002; 124: 1137-44
42. Gama de Abreu M, Heintz M, Heller A, Szechenyi R, Albrecht DM, Koch T: One-lung ventilation with high tidal volumes and zero positive end-expiratory pressure is injurious in the isolated rabbit lung model. *Anesth Analg* 2003; 96: 220-8
43. Fanelli V, Mascia L, Puntorieri V, Assenzio B, Elia V, Fornaro G, Martin EL, Bosco M, Delsedime L, Fiore T, Grasso S, Ranieri VM: Pulmonary atelectasis during low stretch ventilation: "open lung" versus "lung rest" strategy. *Crit Care Med* 2009; 37: 1046-53
44. Henzler D, Hochhausen N, Dembinski R, Orfao S, Rossaint R, Kuhlen R: Parameters derived from the pulmonary pressure volume curve, but not the pressure time curve, indicate recruitment in experimental lung injury. *Anesth Analg* 2007; 105: 1072-8
45. Rahn H, Otis AB, Fenn WO: The pressure-volume diagram of the thorax and lung. *Am J Physiol* 1946; 146: 161-78
46. Agostoni E, Mead J: Statics of the respiratory system. *Handbook of Physiology, Section 3. Respiration. Vol 1.* Edited by W.O. Fenn and H. Rahn. Washington DC, American Physiological Society 1964: 387-409

47. Agostoni E, Hyatt RE: Static behavior of the respiratory system. Handbook of Physiology, Section 3. Respiration. Vol III. Mechanics of Breathing, Part 1. Edited by Fisherman AP, Macbkle PT, Mead J, Geiger SR, American Physiological Society Bethesda, Maryland 1986: 113-130
  
48. Mergoni M, Martelli A, Volpi A, Primavera S, Zuccoli P, Rossi A: Impact of positive end-expiratory pressure on chest wall and lung pressure-volume curve in acute respiratory failure. Am J Respir Crit Care Med 1997; 156: 846-54