

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Fish consumption as a source of human exposure to perfluorinated alkyl substances in Italy:  
Analysis of two edible fish from Lake Maggiore**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/148041> since 2024-11-28T15:40:29Z

*Published version:*

DOI:10.1016/j.chemosphere.2014.04.085

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [Chemosphere, Volume 114, November 2014, Pages 181–186, doi:10.1016/j.chemosphere.2014.04.085].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform*]

# 1 **Fish consumption as a source of human exposure to perfluorinated alkyl** 2 **substances in Italy: Analysis of two edible fish from Lake Maggiore**

3  
4 S. Squadrone\*, V. Ciccotelli, L. Favaro, T. Scanzio, M. Prearo, M.C. Abete

5  
6 Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle D'Aosta, via Bologna 148, 10154 Torino, ITALY.

7 \*Corresponding author. Tel.: +39 011 2686238; fax: 39 011 2686228. E-mail address: stefania.squadrone@izsto.it

## 8 9 **ABSTRACT**

10 Extensive screening analyses of perfluorooctane sulfonate and related perfluorinated compounds (PFCs) in biota  
11 samples from all over the world have identified PFCs as global pollutants and have shown their bioaccumulation into  
12 higher trophic levels in the food chain.

13 Perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) are environmental contaminants belonging to a  
14 chemical group known as perfluorinated compounds. PFOS and PFOA are very persistent in the environment and  
15 bioaccumulate in humans. They are potential reproductive and developmental toxicants and are considered to be  
16 emerging endocrine disruptors.

17 The United States Environmental Protection Agency (US EPA) considers both compounds to be carcinogenic and the  
18 European Food Safety Authority (EFSA) recently pointed out that they are associated with adverse health effects. Diet  
19 is considered the main source of exposure to PFCs, which have been found more frequently in fish and other seafood,  
20 compared to other food groups. In fact, aquatic ecosystems represent the final reservoir for PFCs due to their great  
21 affinity for sedimentary and living organic matter. In these systems, measured levels of persistent organic pollutants  
22 (POPs) could increase along the trophic web, ultimately affecting humans that consume aquatic species.

23 In this study, PFOS and PFOA was detected by LC-MS/MS in muscle samples of *Coregonus lavaretus* (European  
24 whitefish) and *Perca fluviatilis* (European perch) collected from Lake Maggiore, a large lake located on the south side  
25 of the Italian Alps. PFOA was not found in any of the investigated samples above the limit of quantitation of 0.50 ng  
26 g<sup>-1</sup> fresh weight (fw), whereas PFOS was detected in all 90 samples with concentrations of up to 46.0 ng g<sup>-1</sup> fw. Mean  
27 concentrations were 22.2 ng g<sup>-1</sup> fw in *Perca fluviatilis* and 20.0 ng g<sup>-1</sup> fw in *Coregonus lavaretus*.

28 Comparison of our results with literature data on PFOS intake suggested that fish from Lake Maggiore may be a  
29 significant source of dietary PFOS exposure, even if the reported values were lower than the Total Daily Intake (TDI)  
30 proposed by EFSA.

31  
32 *Keywords:* PFOS; fish, Lake Maggiore, Liquid chromatography–tandem mass spectrometry (LC–MS/MS)

## 33 34 35 36 **1. Introduction**

37 Perfluorinated (fully fluorinated) organic compounds, such as perfluorooctane sulfonate (PFOS), represent a class of  
38 compounds showing high thermal, chemical and biological inertness. They can be widely found in the environment,  
39 primarily resulting from anthropogenic sources.

40 PFOS and a number of related perfluorinated organic sulfonates have been found to be present in fish, birds and  
41 mammals. In contrast to other fluorinated compounds that are partially degradable, PFCs are very persistent both to  
42 abiotic and biotic degradation processes. PFCs tend to bioaccumulate, and PFOS is also a suspected carcinogen and a  
43 candidate POP under the Stockholm Convention (EFSA, 2008; Paul *et al.*, 2009).

44 PFOS meets the criteria for being characterized as a persistent organic pollutant (POP), and restrictions for PFOS and  
45 other perfluoroalkylsulfonate (PFAS) commercialization were therefore fixed from 2001 to 2004 by the US EPA  
46 (2003), and in 2006 by the 2006/122/CE Directive.

47 In addition, in 2008, the EFSA Scientific Panel on Contaminants in the Food Chain (CONTAM Panel) acknowledged  
48 the limitations of the dietary exposure assessment, and recommended that more occurrence data of PFASs in different  
49 foodstuffs, along with biomonitoring data from humans should be collected, particularly with respect to monitoring  
50 trends in exposure.

51 PFOS has been shown to bioaccumulate in fish and in experimental animals, with exposure to PFOS resulting in  
52 hepatotoxicity and increased mortality.

53 Toxicology studies show that PFOS and PFOA are readily absorbed after oral exposure and accumulate primarily in the  
54 serum, kidney, and liver. No further metabolism is expected (EFSA 2008). PFOS and PFOA have a long half-life of  
55 about 4 years in humans. This continued exposure could increase body burdens to levels that would result in adverse  
56 outcomes (ATSDR 2009).

57 Acute- and intermediate- duration oral studies in rodents have raised concerns about potential developmental,  
58 reproductive, and other systemic effects of PFOS and PFOA (ATSDR 2009) while the ingestion of PFOA-contaminated  
59 water was found to cause adverse effects on mammary gland development in mice (Post et al. 2012). Both compounds  
60 are suspected endocrine disrupter, affecting fertility and development and are likely to have an analogous mode of  
61 actions (EFSA, 2008). PFOS and PFOA have a high affinity for binding to B-lipoproteins and liver fatty acid-binding  
62 protein. Several studies have shown that these compounds can interfere with fatty acid metabolism and may deregulate  
63 metabolism of lipids and lipoproteins (EFSA 2008). Some toxicological and epidemiological evidence indicate a link  
64 between PFOS/PFOA exposure and reduced fertility (Governini et al., 2011).

65 Preliminary information indicate increasing levels of these substances in the environment within the food chain as well  
66 as in the general human population, there is therefore a crucial need to improve the database to assess the potential risks  
67 associated with human exposure to PFCs (EFSA 2008).

68 Most biota monitoring studies have so far focused either on liver, as a target organ for PFC accumulation, or on the  
69 blood or whole body homogenates (Houde *et al.*, 2006).

70 Fish are environmental bioindicators as they are essential components of various ecosystems (i.e. sea, rivers, lakes, etc.)  
71 as well as representing an important food source for humans.

72 Reports on PFC levels in edible fish muscle (Kannan *et al.*, 2005; Gulkowska *et al.*, 2006; Gruber *et al.*, 2007; Ericson  
73 *et al.*, 2008) have shown that perfluorooctane sulfonate (PFOS) is usually the prevailing PFAS, with concentrations in  
74 the range of <1 ng g<sup>-1</sup> fresh weight (fw) to >100 ng g<sup>-1</sup> fw, depending on fish species and sampling location (Berger *et*  
75 *al.*, 2009). Very little is known about PFAS levels in commercial fish species in Italy, and in particular on freshwater  
76 fish.

77 Lakes play a vital role in the landscape and ecosystems, as well as being central to recreational activities such as  
78 swimming and fishing. Given the high population density in areas surrounding lakes, and their year-round accessibility,

79 these water bodies are generally more vulnerable to contamination, especially those with the nearby presence of  
80 intensive industrial and commercial activities. Although natural processes often provide an effective means of removing  
81 many micro-contaminants, some effluent-derived contaminants seem to be resistant to biodegradation. Discharge of  
82 wastewater effluents into surface waters and drinking water aquifers, and subsequent re-use of the water, is becoming  
83 more common throughout the world, because the continuing growth of the human population creates a corresponding  
84 increase in the demand for fresh water supply.

85 Investigations on the PFOS/PFOA levels in fish muscle in Lake Maggiore, a North West Italian Lake, have not  
86 previously been carried out.

87 This lake, the second largest in Italy, has suffered from mercury and DDT contamination in the past and the quality of  
88 its water decreased substantially from the 1950s to the 1970s because of the effects of domestic and industrial  
89 contamination. Although nutrients and some non-polar contaminants have attracted widespread attention, polar  
90 substances have been less investigated. The region of Lake Maggiore can be characterized as a mountainous and tourist-  
91 affected area with little agricultural land use; there are many developed smaller urban centers, and some industrial  
92 impact on the west side (in Piedmont). Lake Maggiore receives municipal, agricultural, and industrial discharges, both  
93 directly and via its tributary rivers. The lake water is used in some areas for production of drinking water. It is,  
94 therefore, necessary to identify and quantify the chemicals likely to be present in recycled water, evaluate their potential  
95 effects on humans and aquatic ecosystems, and assess approaches for minimizing their release (Loos 2007).

96 The objective of this study was to investigate the occurrence of emerging polar anthropogenic contaminants (PFOS,  
97 PFOA) in two species of commercial fish, *Coregonus lavaretus* and *Perca fluviatilis*, from Lake Maggiore, in order to  
98 establish the contamination levels and to assess human dietary exposure through fish consumption.

## 99 **2. Materials and methods**

### 100 *2.1 Study species*

101 The European whitefish (*Coregonus lavaretus*), widely distributed in the freshwaters of Northern Europe and  
102 introduced into the major lakes of Northern Italy, are also present in the lakes of North and Central Italy, mainly as a  
103 result of stock enhancement practices (Orban *et al.* 2006). *C. lavaretus*, a stenothermic fish feeding mainly on  
104 zooplankton and benthos. This species lives in clear, cold and well-oxygenated waters, and is characterized by a rapid  
105 growth reaching sexual maturity in the second year of life, with a size of 30 cm circa. As a food source, whitefish is an  
106 appreciated and valuable freshwater fish species.

107 The European perch (*Perca fluviatilis* L.), a predatory freshwater fish species feeding on invertebrates and fish, was  
108 originally confined to the temperate waters of the northern hemisphere, mainly Europe and North America. In Italy, in  
109 the North-East sector of the Po basin, the presence of perch has been documented for centuries, while in Central Italy,  
110 this species is mainly present in the lakes of the Umbria and Latium regions (Orban *et al.* 2007). Within 2–3 years,

111 when the body length is 15–25 cm, the perch reaches sexual maturity. The average perch length is 20–25 cm, but it may  
112 grow up to 45 cm. On the other hand may be subjected to dwarfism when living in high density populations.

### 113 2.2 *Fish collection*

114 Fish were captured by gillnetting in Autumn 2012. The study included 40 specimens of European perch (*Perca*  
115 *fluviatilis*) and 50 specimens of European whitefish (*Coregonus lavaretus*).

116 Sampling locations (Figure 1) were selected according to the ecological needs and the seasonal distribution of the two  
117 species. In fact, the European perch specimens were caught in the mouth of the River Tresa and close to the ground at a  
118 water depth of approximately 25 m (46.00006, 8.72228), while the individuals of European whitefish were captured in  
119 the town of Luino (Province of Varese) in open water, at a water depth of approximately 30 m (46.01419, 8.74468).

120 Fish were captured in agreement with the animal welfare legislation procedure. Perch specimens (10 males and 30  
121 females) were collected at sizes of  $16.5 - 29.5 \pm 0.5$  cm, with weights ranging from  $53 - 371 \pm 1$  g. European whitefish  
122 specimens (20 males and 30 females) were collected at sizes of  $26.0 - 32.0 \pm 0.5$  cm, with weights ranging from 157  
123 to  $307 \pm 1$  g. Individual identification numbers were assigned to each sample.

124 Captured fish were preserved on ice and transported to the laboratory. Samples were sexed and dissected to obtain  
125 muscle tissues, which were immediately frozen and stored at  $-20$  °C.

### 126 2.3 *Reagents and Analytical methods*

127 Certified standards of PFOS (96%) and PFOA (98%) were purchased from Sigma Aldrich (Milan, Italy). The internal  
128 standard (PFOA 13C8, 99%) was purchased from Cambridge Isotope Laboratories (Highwood Drive Tewksbury, MA,  
129 USA).

130 Ammonium acetate, 37% hydrochloric acid, and sodium acetate were purchased from Sigma Aldrich. Sodium  
131 hydroxide was supplied by Carlo Erba (Milan, Italy), while 33% ammonium hydroxide was from RDH (Seelze,  
132 Germany). Methanol, with a degree of purity suitable for HPLC-MS/MS analysis, was purchased from Sigma Aldrich,  
133 and ultrapure water was produced by Maxima UltraPure Water (Thermo Scientific, MA, USA).

134 Purification of samples was performed by Oasis Wax SPE columns 6 cc cartridge 150 mg, 30  $\mu$ g of Waters (Milford,  
135 MA, USA).

136 Stock solutions of each analyte were prepared by dissolving the powder in methanol.

137 Diluted solutions for determining the matrix calibration line were prepared in methanol. The internal standard was also  
138 diluted in methanol.

139 Fish muscle (2.5 g) was homogenized with 2.5 ml of sodium hydroxide using Ultraturrax homogenizer (IKA, Staufen,  
140 Germany), after addition of IS (13C8PFOA). Methanol (10 mL) was added to each homogenate. The shacked extract  
141 was centrifuged and purified using Oasis Wax SPE columns, followed by conditioning with 4 mL of methanol and 4  
142 mL of water. The cartridge-purified extract, was then washed with 4 ml of 25 mM acetate buffer (pH 4-5), followed by

143 8 mL of methanol. Analytes were eluted with 1 mL of 2% ammonium hydroxide in methanol. The solvent was dried by  
144 evaporation by means of a nitrogen stream. The residue was reconstituted in the mobile phase and subjected to LC-  
145 MS/MS analysis.

146 LC-MS/MS analysis was performed by an Agilent HPLC 1100 procedure (Agilent Technologies, Palo Alto, CA, USA).  
147 Chromatographic separation was obtained by a reversed-phase Phenomenex Synergi Fusion column (150 x 2 mm, 4  
148  $\mu\text{m}$ ) coupled to a guard column (Phenomenex Security Guard 4.0 x 2.0 mm). Eluents used were 20 mM aqueous  
149 ammonium acetate (eluent A) and 2 mM ammonium acetate in methanol (eluent B), at a flow-rate of 350  $\mu\text{L}/\text{min}$ . The  
150 elution gradient consisted of the following steps: from 0 to 5 min, A at 50%; from 5 to 7 min, A at 100%; at 8 min, A%  
151 at 50; from 8 to 11 min, A% at 50%. The total run time for each injection was 11 min, with a 20  $\mu\text{L}$  injection volume.  
152 Mass spectral analyses were performed using an Applied Biosystems API 4000 triple quadrupole mass spectrometer  
153 (Applied Biosystems Sciex, Ontario, Canada) operating in electrospray ionization (ESI)negative ion mode. Detection  
154 and quantification of the two molecules were performed by selected reaction monitoring (SRM), as shown in Table 1.  
155 The method limit of quantification (LOQ) was 0.50  $\text{ng g}^{-1}$  for PFOA and 0.70  $\text{ng g}^{-1}$  for PFOS, with a recovery rate of  
156 99-102% and 96-108%, respectively. Commission Recommendation 2010/161 recommends to the Member States to  
157 apply methods of analysis with recovery rates ideally in the range of 70-120 %.

158 All analyses were carried out in duplicate. To check the purity of the reagents and any contamination, “blanks” were  
159 analyzed for each calibration run, using the same procedure.

### 160 *2.3 Statistical analysis*

161 Statistical analyses were performed in SPSS v. 20 (SPSS Inc., Chicago, IL, U.S.A.) for Macintosh. The analytical data  
162 (PFOS levels) were normally distributed in both species, either considering sexes together or separately (Kolmogorov–  
163 Smirnov test,  $p > 0.05$ ).Initially, a linear regression analysis was performed to study the associations between body  
164 weight and PFOS concentrations in both species. Subsequently, differences in mean concentrations of PFOS between  
165 species and sexes were tested using a Student’s t-test (all p-values were two-tailed)..

## 166 **3. Results and discussion**

### 167 *3.1. PFC concentrations in fish and PFOS levels in relation to the data from the literature*

168 Several studies have indicated that PFC concentrations are higher in fish caught from fresh water compared to marine  
169 water and in particular the PFC contamination levels of fish caught in waters affected by anthropogenic pollution is  
170 generally higher than concentrations in fish from open oceans (Berger et al., 2009; Schuetze et al., 2010). In the last  
171 EFSA Opinion concerning PFCs (2012), a constantly higher mean concentration in fish from fresh water was  
172 demonstrated, while diadromous fish had slightly higher mean concentrations, tending to be more similar to the marine  
173 fish. The highest contributors to dietary PFOS exposure across all age classes examined were ‘Fish and other

174 seafood'(50 to 80 %). Among PFCs, PFOS had the highest mean concentrations in all three fish categories and in the  
175 food group 'Fish and other seafood', the concentrations found in fish meat ranged from 0.04 to 211 ng g<sup>-1</sup>. In the  
176 present study, PFOS mean values were 22.20 ng g<sup>-1</sup> fw in *Perca fluviatilis* and 19.98 ng g<sup>-1</sup> fw in *Coregonus lavaretus*,  
177 while PFOA values were less than the limit of quantitation in all samples. All 90 samples had detectable levels of PFOS  
178 (> LOQ) and concentrations ranged from 5.0 to 45.8 ng g<sup>-1</sup> fw (Table 2).

179 The presence of PFCs in Lake Maggiore, and in drinking water produced from the lake, was previously observed by  
180 Loos (2007), who showed that these compounds were relatively equally distributed in the lake water. In fact, PFOS was  
181 detected at a concentration of 7.8 ng g<sup>-1</sup> fw (range 7.2-8.6 ng g<sup>-1</sup> fw) while PFOA was reported at a lower concentration  
182 2.4 ng g<sup>-1</sup> fw (range (1.8-2.9 ng g<sup>-1</sup> fw), while no biota data were reported. Even if the fate properties of PFOA are  
183 similar to those of PFOS, and once in the environment they both are extremely persistent and not known to undergo  
184 significant further abiotic or biotic degradation, the bio accumulation pattern in fish seems to differ. In fact, PFOS  
185 exhibits a higher tendency to bind to organic matter and bioaccumulate compared to PFOA, due to its longer  
186 perfluoroalkyl chain length (Conder et al. 2010). Moreover, PFOS has been shown to bioaccumulate and biomagnify in  
187 wildlife species such as fish and piscivorous birds and is the only PFC that has been shown to accumulate to levels of  
188 concern in fish tissue (EFSA 2008). Because PFOS is both hydrophobic and lipophobic it does not follow the typical  
189 pattern of partitioning into fatty tissues followed by accumulation, the typical pattern of many persistent organic  
190 pollutants. Instead, it binds to proteins in the plasma and, as a result, is present in highly perfused tissues such as the  
191 liver and kidneys rather than lipid tissue. Therefore, the mechanism of bioaccumulation likely differs from most other  
192 bioaccumulative chemicals (UNEP, 2006). However, monitoring data from top predators at various locations show  
193 highly elevated levels of PFOS and demonstrate the substantial bioaccumulation and biomagnification (BMF)  
194 properties of PFOS. Monitoring studies and field studies indicate that biomagnification in various terrestrial and marine  
195 mammals occur (Bossi et al., 2005a,b).

196 Most international studies on PFOS levels in fish have been focused on liver, blood, or whole body homogenates as  
197 analytical matrices, and only a few reports on levels in fish muscle are available for comparison with our data (Hoff *et*  
198 *al.*, 2003; Järnberg and Holmström, 2003; Kannan *et al.*, 2005; Gulkowska *et al.*, 2006; Tittlemier *et al.*, 2007; Ericson  
199 *et al.*, 2008).

200 A Swedish study of PFOS and PFOA in perch from waters without local pollution reported low PFOS levels in fish  
201 muscle tissue (1–2 ng g<sup>-1</sup> fw), (Järnberg and Holmström, 2003). Higher levels were found in perch caught in Lake  
202 Mälaren (20–44 ng g<sup>-1</sup> fw), an important lake for both commercial and recreational fishing, situated in one of the most  
203 densely populated areas in Sweden. PFOA levels were, however, below the method detection limit (MDL) of 0.5 ng g<sup>-1</sup>  
204 fw in all samples (Järnberg and Holmström, 2003). These data are comparable to ours that were obtained in Lake  
205 Maggiore, where PFOS levels in perch ranged from 11 to 46. No data are available in the literature about European



206 whitefish PFC levels. The concentrations of PFOS in the present study were higher than those reported in fish from  
207 other areas in different wild fish species from the Mediterranean Sea (Brambilla et al., 2009, Nania et al., 2009).  
208 Brambilla and coworkers analyzed wild and farmed fish from different areas in the Mediterranean and reported  
209 concentrations in wild fish ranging from 0.09 to 5.96 ng g<sup>-1</sup> w.w. Nania and coworkers analysed fish samples included  
210 pelagic fish and benthonic fish caught in the Mediterranean Sea, reporting PFOS mean values of 4 and 13 ng g<sup>-1</sup> w.w  
211 respectively. Benthonic fish showed PCFs levels, on average, higher than pelagic fish and this could be explained by the  
212 fact that benthonic fish can absorb contaminants both from water and from sandy and muddy sediments (Berger et al.,  
213 2004). From Nania results emerged the presence of few fish that showed an extremely high contamination by PFOS,  
214 maybe linked to “dot-like” pollutants release. In our findings, analytical data showed a normal distribution for PFOS in  
215 both species, either considering sexes together or separately.

### 216 3.2 *Intraspecies and between-species comparisons*

217 Gender related length, weight and PFOS levels in fish, are shown in Table 2. In European whitefish length and weight  
218 were not affected by gender, while highest PFOS levels were found in female specimens (42.3 ng g<sup>-1</sup> fw), although  
219 mean and median PFOS values were comparable in both sexes.  
220 In perch, highest PFOS levels (45.8 ng g<sup>-1</sup> fw) were still found in females with an average length and weight of 29.5 cm  
221 and 371 g, respectively. However, mean and median values were comparable in both sexes. In our sample, there was no  
222 correlation (R = 0.016) between fish weight and PFOS concentration in the muscle (Figure 2).  
223 The Student’s t-test showed that there was no significant difference in the mean concentration of POFS, neither between  
224 the two species (*Coregonus lavaretus* = 50, *Perca fluviatilis* = 40, t = 1.430, df = 88, p = 0.154), nor between sexes  
225 (*Coregonus lavaretus*: males = 30, females = 20, t = - 0.735, df = 48, p = 0.466; *Perca fluviatilis*: males = 30, females =  
226 10, t = 0.79, df = 38, p = 0.938). Our data confirmed previous findings (Giesy and Kannan, 2005), underlining that the  
227 variations in concentrations of perfluorochemicals found between sexes and among different age groups are not  
228 statistically significant.

### 229 3.3 *Fish as a source of human exposure to PFOS*

230 Diet, particularly consumption of fish and fishery products, is known to be a source of exposure to PFOS, PFOA, and  
231 other PFCs (Ericson et al., 2008; Guruge et al., 2008; Nania et al., 2009; Clarke et al., 2010). Public health concern  
232 regarding PFASs was raised after several studies indicated that PFOS and PFOA are present in the environment,  
233 including within the human body (EFSA 2012). Several adverse health effects, e.g. hepatotoxicity, developmental  
234 toxicity, neurobehavioral toxicity, immunotoxicity, reproductive toxicity, lung toxicity, hormonal effects, as well as a  
235 weak genotoxic and carcinogenic potential have been demonstrated in experimental studies in animals (Lau et al., 2006;  
236 Zhang et al., 2009; Pinkas et al., 2010). Recently, an epidemiological study performed on a cohort of children in the  
237 Faroe Islands indicated that high exposure to PFASs was associated with reduced humoral immune response to

238 immunizations in children (Grandjean et al., 2012). Moreover, the Italian project PREVIENI, funded by the Italian  
239 Environment Ministry, aims to link environment and human health through the investigation of exposure to selected  
240 endocrine disrupters (EDs) and associated biomarkers related to human infertility conditions. Preliminary results related  
241 to the metropolitan area indicate that subjects affected by infertility factors tend to have both higher PFOS levels and  
242 higher gene expression of specific nuclear receptors (La Rocca et al., 2012).

243 The Scientific Panel on Contaminants in the Food Chain (CONTAM) identified 30 ng g<sup>-1</sup> body weight (b.w.) per day as  
244 the lowest no-observed-adverse-effect level (NOAEL) (effect: changes in lipids and thyroid hormones) to derive a  
245 provisional TDI of 150 ng kg<sup>-1</sup> b.w. per day by applying an overall uncertainty factor of 200 to the NOAEL (EFSA,  
246 2008, Schuetze et al., 2010).

247 In an adult consumer with a body weight of 60 kg, this value is reached when consuming fish that contains 30 ng g<sup>-1</sup> of  
248 PFOS, considering a consumption of 300 g fish per day. We estimated human exposure from fish consumption by  
249 calculating the Estimated Human Daily Intake (EHDI), as follows:

$$250 \text{ EHDI} = (C \times DC) / \text{BW}$$

251 where C is the contaminant mean concentration, DC indicates the daily fish consumption for the Italian population, as  
252 reported by the National Research Institute for Food and Nutrition (Leclercq et al., 2009), and BW is the human body  
253 weight (60 kg).

254 The consumption figures used were the 50th and 95th percentile intakes for the total population in consumers of all  
255 ages; we obtained a value of 11.9 ng kg<sup>-1</sup> bw day and 54.39 ng kg<sup>-1</sup> bw day respectively. Comparing our EHDI with  
256 the Tolerable Daily Intake (TDI) established for PFOS (150 ng kg) we can conclude that the intake related to fish  
257 consumption from Lake Maggiore is below the tolerable daily intake (TDI).

258 However, other food and sources other than food may contribute to the total human exposure; than particularly for high  
259 fish consumers the intake from fish consumption may constitute a considerable contribute to the total daily intake of  
260 PFOS.

#### 261 **4. Conclusions**

262 Fish caught in polluted freshwater systems can be a significant source of dietary human PFOS exposure, and may  
263 continue to be for many years or decades to come; however the occurrence of suspected carcinogens, such as PFOS, in  
264 food for human consumption should not be tolerated, even at low levels. This study has contributed to the knowledge of  
265 PFOS levels in fish from an important Italian lake, and whilst our biomonitoring results did not show a particularly  
266 alarming level of pollution by PFCs, the PFOS contamination levels and the resulting consumer's intake were mostly  
267 higher than those found in similar European monitoring. Then, measures should therefore be taken to reduce the  
268 consumption of these damaging substances by humans.

269

270

271

272 **Acknowledgment**

273 This work was financially supported by the Italian Ministry of Health.

274 The authors are very grateful also to the referees and the editor for their constructive criticism and suggestions that

275 helped to greatly improve the paper.

276 **References**

- 277 Agency for Toxic Substance and Disease Registry (ATSDR). 2009. Draft Toxicological Profile for Perfluoroalkyls.  
278 [www.atsdr.cdc.gov/toxprofiles/tp200.pdf](http://www.atsdr.cdc.gov/toxprofiles/tp200.pdf)  
279
- 280 Berger, U., Järnberg, U., & Kallenborn, R. 2004. Perfluorinated alkylated substances (PFAS) in the European nordic  
281 environment. *Organohalogen Compounds* 66:4046–4052.  
282
- 283 Berger U, Glynn A, Holmström KE, Berglund M, Ankarberg EH, Törnkvist A. (2009) Fish consumption as a source of  
284 human exposure to perfluorinated alkyl substances in Sweden Analysis of edible fish from Lake Vättern and the Baltic  
285 Sea. *Chemosphere* 76:799–804.  
286
- 287 Bossi, R., Riget, F.F., Dietz, R., 2005a. Temporal and spatial trends of perfluorinated compounds in ringed seal (*Phoca*  
288 *hispidus*) from Greenland. *Environ. Sci. Technol.* 39 (19), 7416–7422.  
289
- 290 Bossi, R., Riget, F.F., Dietz, R., Sonne, C., Fauser, P., Dam, M., Vorkamp, K., 2005b.  
291 Preliminary screening of perfluorooctane sulfonate (PFOS) and other fluorochemicals in fish, birds and marine mammals  
292 from Greenland and the Faroe Islands. *Environ. Pollut.* 136, 323–329.  
293
- 294 Brambilla, G., Abate, V., Davoli, E., De Filippis, S.P., De Luca, S., Dellatte, E., Fanelli, R., Fattore, E., Fochi, I.,  
295 Fulgenzi, A.R., Iacovella, N., Iamiceli A.L., Lucchetti, D., Melotti, P., Miniero, R., Moret, I., Pignata, S., Roncarati, A.,  
296 Triboni P., Ubaldi, A., Zambon, S., di Domenico, A., August 20–25, 2009. Occurrence of persistent organic pollutants  
297 (POPs) in Italian wild and farmed fish in Mediterranean Sea 29. In: International Symposium on Halogenated Persistent  
298 Organic Pollutants  
299 (POPs) – DIOXIN 2009 Proceedings; Pechino (RPC).  
300
- 301 Conder, J.M., Wenning, R.J., Travers, M., and Blom, M. 2010. Overview of the Environmental Fate of Perfluorinated  
302 Compounds. Network for Industrially Contaminated Land in Europe (NICOLE) Technical Meeting. 4 November 2010.  
303
- 304 Chu S, Letcher RL. (2009) Linear and branched perfluorooctane sulfonate isomers in technical product and  
305 environmental samples by in-port derivatization-gas chromatography-mass spectrometry. *Analytical Chemistry* 81:  
306 4256-4262.  
307
- 308 Cornelis C, D'Hollander W, Roosens L, Covaci A, Smolders R, Van den Heuvel R, Govarts E, Van Campenhout K,  
309 Reynders H, Bervoets L. (2012) First assessment of population exposure to perfluorinated compounds in Flanders,  
310 Belgium. *Chemosphere* 86: 308-314.  
311
- 312 EFSA, 2008. European Food Safety Authority; opinion of the Scientific Panel on contaminants in the food chain on  
313 perfluorooctanesulfonate (PFOS), perfluorooctanoic acid (PFOA) and their salts. *EFSA J.* 653, 1–131, Available online:  
314 <http://www.efsa.europa.eu/en/efsajournal/doc/653.pdf>.  
315
- 316 EFSA, 2011. European Food Safety Authority; results of the monitoring of perfluoroalkylated  
317 substances in food in the period 2000–2009. *EFSA J.* 9 (2), 2016, Available online:  
318 <http://www.efsa.europa.eu/en/efsajournal/doc/2016.pdf>; [www.efsa.europa.eu/efsajournal.htm](http://www.efsa.europa.eu/efsajournal.htm).  
319
- 320 Ericson, I., Marti-Cid, R., Nadal, M., Van Bavel, B., Lindstrom, G., Domingo, J.L., 2008. Human exposure to  
321 perfluorinated chemicals through the diet: intake of per-fluorinated compounds in foods from the Catalan (Spain)  
322 market. *J. Agric. Food Chem.* 56, 1787–1794.  
323
- 324 DIRECTIVE 2006/122/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL  
325 of 12 December 2006 amending for the 30th time Council Directive 76/769/EEC on the approximation of the laws,  
326 regulations and administrative provisions of the Member States relating to restrictions on the marketing and use of  
327 certain dangerous substances and preparations (perfluorooctane sulfonates). *Official Journal of the European Union*, L  
328 372 pp 32-34.  
329
- 330 Fromme, H., Tittlemier, S.A., Völkel, W., Wilhelm, M., Twardella, D. 2009. Perfluorinated compounds—Exposure  
331 assessment for the general population in western countries. *Int. J. Hyg. Environ. Health*  
332 212:239-270.  
333
- 334 Governini, L., Guerranti, C., Focardi, S., Cuppone, A.M., Stendardi, A., De Leo, V., Piomboni,  
335 P., 2009. Impact of environmental exposure to perfluorinated compounds on sperm DNA quality. *Syst. Biol. Reprod.*  
336 *Med.* 2, 41–55.  
337

338 Grandjean, P., Wreford Andersen, E., Budtz-Jørgensen, E., Nielsen, F., Mølbak, K., Weihe, P., Heilmann, C., 2012.  
339 Serum Vaccine Antibody Concentrations in Children Exposed to Perfluorinated Compounds. *J Am. Med. Ass.* 307 n.4,  
340 *JAMA.* 2012;307(4):391-397. doi:10.1001/jama.2011.2034.

341

342 Gruber, L., Schlummer, M., Ungewiss, J., Wolz, G., Moeller, A., Weise, N., Sengl, M., Frey, S., Gerst, M., Schwaiger,  
343 J., 2007. Tissue distribution of perfluorooctanesulfonate (PFOS) and perfluorooctanoic acid (PFOA) in fish.  
344 *Organohalogen Comp.* 69, 2889–2891.

345

346 Gulkowska, A., Jiang, Q., So, M.K., Taniyasu, S., Lam, P.K., Yamashita, N., 2006. Persistent perfluorinated acids in  
347 seafood collected from two cities of China. *Environ. Sci. Technol.* 40, 3736–3741.

348

349 Guruge, K.S., Yeung, L.W., Yamanaka, N., Miyazaki, S., Lam, P.K., Giesy, J.P., Jones, P.D., Yamashita, N. 2006.  
350 Gene expression profiles in rat liver treated with perfluorooctanoic acid (PFOA). *Toxicol.Sci.* 89, 93-107.

351

352 Haug LS, Salihovic S, Jogsten IE, Thomsen C, Bavel B van, Lindström G, Becher G. (2010a) Levels in food and  
353 beverages and daily intake of perfluorinated compounds in Norway. *Chemosphere* 80:1137–1143.

354

355 Haug LS, Thomsen C, Brantsæter AL, Kvaalem HE, Haugen M, Becher G, Alexander J, Meltzer HM, Knutsen HK.  
356 (2010b). Diet and particularly seafood are major sources of perfluorinated  
357 compounds in humans. *Environment International* 36:772-778.

358

359 Hoff, P.T., Van de Vijver, K., Van Dongen, W., Esmans, E.L., Blust, R., De Coen, W.M., 2003. Perfluorooctane  
360 sulfonic acid in bib (*Trisopterus luscus*) and plaice (*Pleuronectes platessa*) from the Western Scheldt and the Belgian  
361 North Sea: distribution and biochemical effects. *Environ. Toxicol. Chem.* 22, 608–614.

362

363 Houde, M., Martin, J.W., Letcher, R.J., Solomon, K.R., Muir, D.C.G. 2006. Biological monitoring of polyfluoroalkyl  
364 substances: a review. *Environ Sci Technol* 40:3463–73.

365

366 Järnberg, U., Holmström, K., 2003. Perfluorooctane sulfonate concentrations in Swedish urban and background fish  
367 samples. SETAC Europe 13th Annual Meeting, Hamburg, Germany.  
368 <http://abstracts.co.allenpress.com/pweb/setaceu2003/document/?ID=2252.2>

369

370 La Rocca, C., Alessi, E., Bergamasco, B., Caserta, D., Ciardo, F., Fanello, E., Focardi, S., Guerranti, C., Stecca, L.,  
371 Moscarini, M., Perrad, G., Tait, S., Zaghi, C., Mantovani, A., 2012. Exposure and effective dose biomarkers for  
372 perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) in infertile subjects: Preliminary results of the  
373 PREVIENI project. *Int. J. Hyg. Environ. Health* 215 :206–211.

374

375 Lau, C., Thibodeaux, J.R., Hanson, R.G., Narotsky, M.G., Rogers, J.M., Lindstrom, A.B., Strynar, M.J., 2006.. Effects  
376 of perfluorooctanoic acid exposure during pregnancy in the mouse. *Toxicol Sci.* 90:510–518.

377

378 Leclercq, C., Arcella, D., Piccinelli, R., Sette, S., Le Donne, C., Turrini, A., 2009. The Italian National Food  
379 Consumption Survey INRAN-SCAI 2005–06: main results in terms of food consumption. *Public Health Nutr.* 12,  
380 2504–2532.

381

382 Loos, R., Wollgast, J., Huber, T., Hanke, G., 2007. Polar herbicides, pharmaceutical products, perfluorooctanesulfonate  
383 (PFOS), perfluorooctanoate (PFOA), and nonylphenol and its carboxylates and ethoxylates in surface and tap waters  
384 around Lake Maggiore in Northern Italy. *Anal Bioanal Chem.*, 387(4):1469-78.

385

386 Kannan, K., Tao, L., Sinclair, E., Pastva, S.D., Jude, D.J., Giesy, J.P., 2005. Perfluorinated compounds in aquatic  
387 organisms at various trophic levels in a Great Lakes food chain. *Archives of Environmental Contamination and*  
388 *Toxicology* 48, 559, 566.

389

390 Orban, E., Nevigato, T., Masci, M., Di Lena, G., Casini, I., Caproni, R., Gambelli, L., De Angelis, P., Rampacci, M.,  
391 2007. Nutritional quality and safety of European perch (*Perca fluviatilis*) from three lakes of Central Italy. *Food*  
392 *Chemistry* 100: 482–490.

393

394 Orban, E., Masci, M., Nevigato, T., Di Lena, G., Casini, I., Caproni, R., Gambelli, L., De Angelis, P., Rampacci, M.,  
395 2006. Nutritional quality and safety of whitefish (*Coregonus lavaretus*) from Italian lakes. *Journal of Food*  
396 *Composition and Analysis* 19: 737–746

397

398 Nania, V., Pellegrini, G.E., Fabirizi, L., Sesta, G., de Sanctis, P., Luchetti, D., Pasquale, M.D., Coni, E., 2009.  
399 Monitoring of perfluorinated compounds in edible fish from the Mediterranean Sea. *Food Chem.* 115, 951–957.

400

401 Paul, A.G., Jones, K.C., Sweetman, A.J., 2009. A first global production, emission, and environmental inventory for  
402 perfluorooctane sulfonate. *Environ. Sci. Technol.* 43, 386–392.  
403

404 Pinkas, A., Slotkin, T.A., Brick-Turin, Y., Zee, E.A. van der, Yanai, Y., 2010. Neurobehavioral  
405 teratogenicity of perfluorinated alkyls in an avian model. *Neurotoxicology and Teratology* 32:182–186.  
406

407 Post, G.B., Cohn, P.D., Cooper, K.R., 2012. Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant:  
408 a critical review of recent literature. *Environ Res.* 116,93-117.  
409

410 Schuetze, A., Heberer, T., Effkemann, S., Juergensen, S., 2010. Occurrence and assessment of perfluorinated chemicals  
411 in wild fish from Northern Germany. *Chemosphere* 78: 647–652.  
412

413 Tittlemier, S.A., Pepper, K., Seymour, C., Moisey, J., Bronson, R., Cao, X.L., Dabeka, R.W., 2007. Dietary exposure of  
414 Canadians to perfluorinated carboxylates and perfluorooctane sulfonate via consumption of meat, fish, fast foods, and  
415 food items prepared in their packaging. *J. Agric. Food Chem.* 55, 3203–3210.  
416

417 UNEP. 2006. Risk profile on perfluorooctane sulfonate. Stockholm Convention on Persistent Organic Pollutants  
418 Review Committee. Geneva, 21 November 2006.

419 USEPA (United States Environmental Protection Agency), 2003. Preliminary risk assessment of the developmental  
420 toxicity associated with exposure to perfluorooctanoic acid and its salts. US Environmental Protection Agency, Office  
421 of Pollution Prevention and Toxics, Risk Assessment Division, April 10, 2003. Washington, DC, USA.  
422

423 Zhang, X., Chen, L., Fei, X., Y Ma, Y., Gao, H., 2009. Binding of PFOS to serum albumin and DNA: insight into the  
424 molecular toxicity of perfluorochemicals. *BMC Molecular Biology* 10:16 doi:10.1186/1471-2199-10-16  
425

**Table 1.**  
**MS experimental condition of PFOA and PFOS and internal standard**

| <b>Analyte</b>    | <b>Parent ion</b> | <b>Product Ion</b> | <b>Decluring Potential (V)</b> | <b>Collision Energy (V)</b> |
|-------------------|-------------------|--------------------|--------------------------------|-----------------------------|
| PFOA              | 412.7             | 368.9              | -61                            | -14                         |
|                   |                   | 169.0              |                                | -26                         |
|                   |                   | 219.0              |                                | -22                         |
|                   |                   | 80.1               |                                | -94                         |
| PFOS              | 498.5             | 99.0               | -50                            | -72                         |
|                   |                   | 169.0              |                                | -51                         |
|                   |                   | 376.0              |                                | -14                         |
| Internal standard | 420.7             | 376.0              | -32                            | -14                         |

**Table 2**  
**Concentrations of PFOS in fish muscle from Lake Maggiore**

*Coregonis lavaretus*

|               | PFOS (ng g <sup>-1</sup> ) |        | length (cm) |        | weight (g) |        |
|---------------|----------------------------|--------|-------------|--------|------------|--------|
|               | male                       | female | male        | female | male       | female |
| <b>min</b>    | 9.4                        | 5.0    | 26.0        | 27.5   | 157.0      | 195.0  |
| <b>max</b>    | 29.5                       | 42.3   | 31.5        | 32.0   | 307.0      | 306.0  |
| <b>mean</b>   | 19.1                       | 20.6   | 29.1        | 30.0   | 2258       | 243.5  |
| <b>median</b> | 19.0                       | 19.0   | 29.3        | 30.0   | 229.0      | 238.5  |

*Perca fluviatilis*

|               | PFOS (ng g <sup>-1</sup> ) |        | length (cm) |        | weight (g) |        |
|---------------|----------------------------|--------|-------------|--------|------------|--------|
|               | male                       | female | male        | female | male       | female |
| <b>min</b>    | 11.0                       | 11.3   | 16.5        | 18.0   | 53.0       | 65.0   |
| <b>max</b>    | 31.8                       | 45.8   | 22.0        | 29.5   | 126.0      | 371.0  |
| <b>mean</b>   | 22.4                       | 22.1   | 20.6        | 21.8   | 101.4      | 120.3  |
| <b>median</b> | 20.9                       | 20.5   | 21.0        | 21.5   | 108.0      | 109.5  |



**Table 3**  
**Length and weight versus gender**  
**in *Coregonis lavaretus***


|               | <b>Length (cm)</b> |        | <b>Weight (g)</b> |        |
|---------------|--------------------|--------|-------------------|--------|
|               | male               | female | male              | female |
| <b>Min.</b>   | 26.0               | 27.5   | 157.0             | 195.0  |
| <b>Max.</b>   | 31.5               | 32.0   | 307.0             | 306.0  |
| <b>Mean</b>   | 29.1               | 30.0   | 225.8             | 243.5  |
| <b>Median</b> | 29.2               | 30.0   | 229.0             | 238.5  |

**Table 4**  
**Length and weight versus gender**  
**in *Perca fluviatilis***

|               | <b>Length (cm)</b> |        | <b>Weight (g)</b> |        |
|---------------|--------------------|--------|-------------------|--------|
|               | male               | female | male              | female |
| <b>Min.</b>   | 16.5               | 18.0   | 53.0              | 65.0   |
| <b>Max.</b>   | 22.0               | 29.5   | 126.0             | 371.0  |
| <b>Mean</b>   | 20.5               | 21.8   | 101.4             | 120.3  |
| <b>Median</b> | 21.0               | 21.5   | 108.0             | 109.5  |



**Swiss  
Confederation**



**Italy**


- 372 m

**Lake Maggiore**

**Luino**  
⊙ Tresa

1

2



### *Perca fluviatilis* PFOS levels (ng/g)

