AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Methodological problems related to spotlight count as census technique for *Lepus europaeus* in an alpine environment.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/149138> since 2016-10-06T16:34:17Z

Published version:

DOI:10.1007/s13364-013-0157-6

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Acta Theriologica, Volume 59, Issue 2, pp 271-276, 2014, 10.1007/s13364-013-0157-6

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://link.springer.com/journal/13364>

Methodological problems related to spotlight count as a census technique for *Lepus europaeus* in an alpine environment.

P. Tizzani . A. Menzano . A. Dematteis . P. G. Meneguz

P. Tizzani, A. Menzano, P. G. Meneguz: Dipartimento di Scienze Veterinarie, Università di Torino, V. Leonardo da Vinci 44-10095, Grugliasco, Torino, Italy.

A. Dematteis: Centro Ricerca sulla Gestione della Fauna Selvatica, fr. Rore 17-12020, Sampeyre, Cuneo, Italy

Abstract Population estimations are necessary for effective conservation management. In Italy, brown hare populations are commonly censused by spotlighting, but this method does not seem to suit an alpine environment due to its vegetational and orographical complexity. The aim of this study is to evaluate the critical aspects related to spotlight census method in an alpine environment. Spotlight was carried out along two transects of a typical alpine environment. Observed animals were used to define density (number of animal seen/100 ha) and method precision (coefficient of variation (CV) applied to monthly repeated observations). Animal detectability was evaluated using half-normal function with cosine expansion (Distance 5.0@). Animal observability was evaluated by analyzing density estimates related to habitat conformation (unseen areas or full visibility). The exact surface surveyed by spotlighting was evaluated, defining the observation spotlight beam range (OTA) and the land useful sighting (LUS). In the end, LUS was classified in three patches according to hare presence: no hares, occasional hare presence, and constant hare presence. To evaluate habitat influence onto CV, we used a bootstrap simulation. The results show that spotlighting alone is not the most suitable method to apply in the alpine environment because habitat structure highly influences census results. Recommendations to improve spotlight surveys for monitoring European brown hare populations are given.

Keywords Alpine environment. European brown hare . *Lepus europaeus* . Spotlighting

Introduction

Assessing and monitoring animal populations precisely and accurately are essential for research, management, and conservation (Caughley and Sinclair 1994; Wilson and Delahay 2001). European brown hare populations seem to decline in the Italian alps (Gobbi et al. 2001; De Battisti et al. 2002) mainly due to the disappearing of agricultural activities and to landscape transformation with increase of uncultivated areas and woodlands. Such situation has led to the fragmentation of hare habitat in suitable insular patches not always connected to each other. Heterogeneity generated by fragmentation can create barriers to movement (Arnold et al. 1993) and, ultimately, lead to the local extinction of the species (Burkley 1989; Soulé et al. 1992). In this context, it is of paramount importance to use accurate and precise census techniques and strict monitoring plans.

During the past 50 years, a great deal of literature on methods for estimating population density and demographic parameters has been published. Methods of censusing have become more refined and sophisticated, and they should be individually designed or adapted to species behavior and to study aims (Sutherland 1996; Krebs 1999; Southwood and Henderson 2000). An extensive review of methods used to census lagomorph populations is presented in Langbein et al. (1999). It includes indirect methods which rely on censusing signs of animal presence, e.g., faecal pellet count and track count, and direct methods which imply the possibility of seeing animals, e.g., clearance drives or total capture, belt assessment, drag line counts, spotlighting, vantage point counts, and markresight. Due to hares' nocturnal activity, one of the most used direct method is the spotlight count along line transect (Eltringham and Flux 1971; Frylestam 1981; Barnes and Tapper 1985). Spotlighting is also the main method used in Italy to census European brown hare *Lepus europaeus*, both in the plain and in the mountain and hill habitats (Trocchi and Riga 2005). It assumes that it is possible to observe animals in all directions and that the view should not be obstructed by obstacles. Very often, these assumptions are not met, and the validity of the data collected in the field has been questioned (Mahon et al. 1998; Sharp et al. 2001). Major concerns are (1) low precision (Edwards et al. 2000), (2) the species detection bias (Jones and Coman 1982; Mahon et al. 1998), and (3) habitat influence (Langbein et al. 1999). The observation of hares is not a problem in the plains, characterized by open habitats (Spagnesi and Trocchi 1993), but it is more difficult in the mountains and hills due to (1) a complex topography and (2) progressive loss of open areas caused by abandonment of traditional agriculture. This study evaluates the critical aspects related to spotlight census method in the alpine environment. In particular, we want to assess the validity of spotlighting techniques for detecting the abundance of brown hares by repeating spotlighting transects and calculating density estimates and associated levels of precision. We examine the consequences of two variability factors: detectability (probability of detecting a hare in the spotlight range) and observability (effects of nonvisible areas on density estimates). Finally, we evaluate the precision of the method considering the coefficient of variation (CV). We discuss reasons for our findings and make recommendations for future spotlight surveys.

Materials and methods

Study areas

Spotlight counts were carried out in two study areas located in the Western Alps in Cuneo Province (Varaita Valley, Piedmont region; 44°58'N, 7°22'E; 900-1,100 m a.s.l.). The valley has a glaciofluvial origin with a U-shaped conformation, and maximum precipitations occur during spring and autumn. The first area (SA1) presents a southern exposition and is part of a protected area devoted for game reproduction. It is characterized by pastures and mowed meadows, woods, and scrublands; open areas are set amidst wooded formations (*Quercus petraea*, *Fagus sylvatica*, *Tilia sylvestris*, *Fraxinus excelsior*, *Castanea sativa*, and *Rosa canina*). The second area (SA2) has the same topography and vegetation, but hunting is allowed. Both areas are characterized by a mosaic of farmland and woodland areas, representative of the alpine environment; they present a cutting meadow regime with two harvesting periods (June-July and August-September), followed by a period of cattle pasture.

Sampling design

A transect of 4.2 km and a transect of 14.6 km were selected, respectively, in SA1 and SA2 from the existing road network to sample the areas in a representative manner with respect to the available land use type. In SA1, the study was undertaken from September to April between 2003 and 2006; in SA2, during April between 2005 and 2009. We carried out three observation sessions in each month, also in case of snow cover, to test the precision of the method. Snow cover does not affect the census method. Since this method is highly affected by changes in visibility, foggy and misty conditions were avoided. The theory and assumptions of spotlight count for hares and practical aspects of survey design are described by Eltringham and Flux (1971), Frylestam (1981), and Barnes and Tapper (1985). In this study, a 4x4 car was driven along a transect, at low speed (15-20 km/h). Hares were searched for by two observers using a one million candlepower handheld spotlight. One observer swept the spotlight on both sides of the road, in front and behind the vehicle, and the other one scanned each lighted field for hares. The number of animals seen, the distance from the observer measured perpendicular to the route using a laser range finder (Leica LRF 900 scan), the distance travelled (kilometers from the beginning of the transect), and the GPS localization were recorded and reported in a CTR (Region Technical Map, Italy) in scale of 1:10,000 with ArcView v3.3 GIS software (ESRI, Redlands, USA). The transects were surveyed from 2 h after sunset.

To evaluate the detectability of animals at different distances from the observer, we used half-normal function with cosine expansion as the selected model (Distance 5.0®). The software uses the "distance sampling approach" based on the assumption that animal detectability is related to their distance from the observer (detection function). The detection function allows for correction for the missing individuals (Buckland et al. 1997).

To evaluate the observability of hare spotlight count in mountain habitat, we analyzed density estimates related to habitat structure, considering unseen areas or assuming full visibility. Density estimates were calculated as the number of animals seen/100 ha. We used geographic information systems (ArcView 3.2®), land use maps 1:10,000, and digital elevation model with 50-m resolution (Piedmont Region).

According to Barnes and Tapper (1985) and Pegel (1986), we calculated:

1. observation spotlight beam range: effective strip width in which hares could be detected; observation theoretical area applied to linear transect (OTAL): application of the hare observation range, as buffer, to hypothetical linear transects;
2. observation theoretical area (OTA): application of the hare observation range, as buffer, to the transects considering the bends of the roads;
3. Land useful sighting (LUS): utilization of land use maps, orthophotos (CGR, Parma, Italy) and digital elevation model to subtract from OTA the zones classified as wood and urban area and the zones over which the view is obstructed (presence of barriers as woods, scrubs, buildings, undulating terrain; Pegel 1986). Using digital elevation model and the view shed function in the Spatial Analyst package (Esri-ArcView 3.2®), we excluded areas not observable due to their orography.

LUS was classified in three patch typologies according to the presence of detected hares: patch 0 = LUS without detections (encounter rate = 0), patch 1 = LUS with an occasional detection ($0 < \text{encounter rate} < 1$), and patch 2 = LUS with animals constantly detected during the study period (i.e., animals aggregated; encounter rate = 1).

Precision was calculated using the coefficient of variation (Davis 1982) applied to monthly repeated observations ($CV = \text{standard deviation} \times 100 / \text{mean observed hares}$). The habitat influence on the CV and their relationship was modeled using bootstrap analyses (Sokal and Rohlf 1995) to identify a threshold area to improve precision. Random distribution of 10,000 observations was plotted in the study areas, and CV for each visible patch was calculated. Sampling intensity was determined arbitrarily at the point where CV reached an asymptote and did not increase markedly with an increase in sample size (Greig-Smith 1957).

The Mann-Whitney U test was used to assess differences among patch areas. Data were reported as means; statistical significance was set at $p < 0.05$. Statistical analyses were performed using R 2.6.0 software® (R Development Core Team 2007).

Results

Detectability

The observation spotlight beam range was evaluated during a preliminary sampling carried out in the same period and in the same areas of the present study during which 108 hares were observed by spotlighting. All these animals have

been observed within a distance of 120 m from the observer (unpublished data). This distance was used to define a buffer along line transects in both study areas, obtaining OTA and LUS values (Table 1 and Fig. 1). The 108 hares were also used to define the probability of observation at various distances (Fig. 2). Observation histograms fitted well the detection function (chi-square goodness of fit test = 14.3; $p=0.22$). Detection probability calculated on detection function was 33.4 %.

Observability and precision

SA1

During the study period, 194 hares were observed by spotlighting (Table 2). Rare density ranged from 1.8 to 12.3 animals/100 OTA ha (mean = 5.3) and from 7.5 to 51.9 animals/100 LUS ha (mean = 22.3). The highest density was recorded in November (mean = 36.8 animals/100 LUS ha) and the lowest in April (mean = 15.1 animals/100 LUS ha). CV ranged from 0 % in December 2006 to 109.0 % in February 2004 (mean = 43.9 %).

SA2

During the study period, 38 hares were observed by spotlighting (Table 2). The density per transect ranged from 0.1 to 1.5 animals/100 OTA ha (mean = 0.7) and from 0.4 to 6.2/100 LUS ha (mean 3.0). CV managed from 0 % in April 2006 to 173.2 % in April 2009 (mean = 63.7 %).

SA1 and SA2

The three patches identified in each study area presented different mean area dimensions: patch 0=0.59 ha, patch 1= 1.65 ha, and patch 2= 11.80 ha. Significant differences were recorded among areas of these patches (patch 0 vs. patch 1: $W = 228$ and $p<0.001$; patch 0 vs. patch 2: $W = 0$ and $p<0.001$; patch 1 vs. patch 2: $W = 0$ and $p<0.001$). Patch 2 represented the 13.6 % of OTA and the 55.8 % of LUS.

Rare observations were more frequent and continuous in some areas than in others, determining the variability of CV values (SA1 CV = 43.9 %; SA2 CV = 63.7 %), but excluding occasional detections from the survey, the CV decreased to 27.3 % (reduction of 37.8 %) in SA1 and to 20.2 % (reduction of 68.3 %) in SA2. The model equation is $y=1.62 \cdot x^{-0.71}$ (Fig. 3), and it suggests survey patch areas of about 9 ha to obtain acceptable CV

Discussion

Our results show that spotlighting, as described in the literature (Eltringham and Flux j 971; Frylestam 1981; Bames and Tapper 1985), is not the appropriate census method to apply in the alpine environment. The main problem is due to the alpine landscape in which hedge rows, undulating terrain, forest, buildings, and blind spot behind these often obstruct sight so that many hares can be missed, badly influencing the accuracy and precision of the method. Such habitat characteristics ("unseen") were very important in our study areas, involving 76.3 % of the environment in SA1 and 75.5 % in SA2.

Hare densities obtained considering OTA or LUS are very different, showing that habitat structure highly influences census results, and therefore, that population density comparisons between areas with unknown habitat composition are questionable. To obviate this problem, we propose to define a "habitat standardization coefficient" which considers the LUS/OTA density ratio (in this study = 24 %) to normalize OTA density estimations. Another way to normalize data is to use the detection probability value (detectability standardization coefficient; in this study = 33.4 %). Indeed, the study of the habitat structure (LUS/OTA) and the study of detection probability are two methods that, starting from different assumptions, have similar conclusions (similar standardization coefficients). From an "on field" point of view, both methods are not so easy to apply because they require access to sophisticated geographical information systems. We suggest the use of the habitat standardization coefficient because it defines the OTA and LUS parameters for each transect only once and can be applied year by year without further calculations.

To obtain a satisfactory precision is apparently a problem in the alpine terrain, and this was visible in high CV values caused by high habitat fragmentation. Applying a bootstrap simulation, we defined a threshold patch area over which the census method presents high precision. Such finding is confirmed by our data: considering patch 2, we obtained a CV of 27.3 % (with reduction of 37.8 %) in study area 1 and a CV of 20.2 % (with reduction of 68.3 %) in study area 2. These CV values (<22.5 %) are in accordance with the results of Marchandeu and Gaudin's study (1994) and are acceptable both for a hunting area characterized by low animal density and for a protected area characterized by medium-high density. Apparently, the CV reduction is due to an increase of the ratio perimeter/area of the patches: the smaller the surveyed patch area, the more casual is the possibility to observe an animal (in open areas of more than 9 ha, the probability to census a hare is statistically higher).

In conclusion, spotlight count is a simple method to obtain information of hare densities, but the interpretation of the data is highly complex and could lead to erroneous conclusions (Wincentz Jensen 2009). To achieve valid estimates in the alpine environment, a correction of the raw data is necessary. We, therefore, propose two types of counts: (1) point transect spotlight count in preselected open areas of at least 9 ha and (2) line transect spotlight count associated to distance sampling method which allows for correction for the missed individuals (Buckland et al. 2001).

Acknowledgments We would like to thank Comprensorio Alpino CN2, Valle Varaita for allowing us to gather field data and for the logistic support to Davide Bonetto, Marco Ghirardi, Ornar Giordano, Livio Martino, Massimiliano Pellerino, and Andrea Rimbotti for their help in hare surveys. Jesus Pérez Jiménez who furnished useful suggestions.

References

- Arnold GW, Steven DE, Weeldenburg JR, Smith EA (1993) influences of remnant size, spacing pattern and connectivity on population boundaries and demography in euros *Macropus robustus* Living in a fragmented landscape. *Biol Conserv* 64:219-230
- Barnes RFW, Tapper SC (1985) A method for counting hares by spotlight. *J Zool* 206:273-276
- Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L (2001) Introduction to distance sampling. Estimating abundance of biologic populations. Oxford University Press, Inc, New York
- Burldey TV (1989) Extinction in nature reserves: the effect of fragmentation and the importance of migration between fragments. *Oikos* 55:75--81
- Caughley G, Sinclair AR (1994) Wildlife ecology and management. Blackwell, Cambridge, UK
- Davis ED (1982) CRC handbook of census methods for terrestrial vertebrates. CRC, Boca Raton
- De Battisti R, Trocchi V, Zamboni U (2002) Indagini per la definizione di un modello gestionale della lepre (*Lepus europaeus*) in ambiente alpino: l'esperienza della Provincia di Trento. Associazione Cacciatori della Provincia di Trento, Trento
- Edwards GP, De Preu ND, Shakeshafi BJ, Crealy N (2000) An evaluation of two methods of assessing feral cat and dingo abundance in central Australia. *Wildl Res* 27:143-149
- Eltringham SK, Flux JEC (1971) Night counts of hares and other animals in East Africa. *Afr J Ecol* 9:67-72
- Frylestam B (1981) Estimating by spotlight the population density of the European hare. *Acta Theriol* 26:419-427
- Gobbi G, Grimod I, Mochet AM (2001) Ricerca sull'ecologia della lepre comune *Lepus europaeus* in Valle d'Aosta e sugli interventi di miglioramento ambientale finalizzati alla sua conservazione. Regione Autonoma Valle d'Aosta, Aosta
- Greig-Smith P (1957) Quantitative plant ecology. Butterworths, London
- Jones E, Coman BJ (1982) Ecology of the feral cat, *Felis catus* (L.) in south-eastern Australia. III. Home ranges and population ecology in semi-arid north-west Victoria. *Aust Wildl Res* 9:409-420
- Krebs CJ (1999) Ecological methodology. Longman, Menlow Park, California
- Langbein J, Hutchings MR, Harris S, Stoate C, Tapper SC, Wray S (1999) Techniques for assessing the abundance of brown hares *Lepus europaeus*. *Mammal Rev* 29(2):93-116
- Mahon PS, Banks PB, Dickman CR (1998) Population indices for wild camivores: a critical study in sand-dune habitat, south-western Queensland. *Wildl Res* 25:11-22
- Marchandau S, Gaudin JC (1994) Effets du sens du transect et de la période d'observation sur la valeur des indices kilométriques d'abondance de lapins de garenne (*Oryctolagus cuniculus*). *Gibier Faune Sauvage* 11 :85-91
- Pegel M (1986) Deer Feldhasen in Beziehungsfüge seiner Um und Mitweltfaktoren. Schriften des Arbeitskreises Wildbiologie und Jagdwissenschaft an der Justus. Liebig Universitiit Giessen. Hefi
- R Development Core Team (2007) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. Available from: [http:// www.R-projecLorg](http://www.R-projecLorg). Accessed 3 Oct 2007
- Sharp A, Northon M, Marks A, Holmes K (2001) An evaluation of two indices of red fox (*Vulpes vulpes*) abundance in an arid environmen!. *Wildl Res* 28:419-424
- Sokal RR, Rohlf FJ (1995) Biometry: the principles and practice of statistics in biological research. W.H. Freeman, New York
- Soulé ME, Alberts AC, Bolger DT (1992) The effects of habitat fragmentation on chaparral plants and vertebrates. *Oikos* 63:39-47
- Southwood R, Henderson PA (2000) Ecological methods. Blackwell, Oxford
- Spagnesi M, Trocchi V (1993) La lepre comune. Istituto Nazionale per la Fauna Selvatica. Documenti Tecnici 13, Bologna
- Sutherland WJ (1996) Who census? In: Sutherland WJ (ed) Ecological census techniques: a handbook. Cambridge University Press, Cambridge, pp 1-10
- Trocchi V, Riga F (2005) I lagomorfi in Italia. Linee guida per la conservazione e gestione. Istituto Nazionale per la Fauna Selvatica. Documenti Tecnici 25, Bologna
- Wilson GJ, Delahay RJ (2001) A review of methods to estimate the abundance of terrestrial caivores using field signs and observation. *Wildl Res* 28:151-164
- Wincentz Jensen TL (2009) Identifying causes for population decline of the brown hare (*Lepus europaeus*) in agricultural landscapes in Denmark. PhD Thesis. University of Copenhagen, Denmark

Fig. 1 Observation theoretical area (OTA) and land useful sighting (LUS) in study area 1.

Fig. 2 Number of hare sightings (about 8-m interval grouping; *blue columns*) at different distances from the observer using the spotlight census method and Distance 5.0@, where good model fit was obtained (detection function is uniform key with cosine expansion; *red line*). Sample size = 108 hares.

Fig. 3 Bootstrap analysis for the determination of threshold patch areas: relationship between dimension of open areas (ha) and CV (%).

Table 1 OTAL, OTA (observation theoretical area), land use characteristics, and LUS values for each study area.

Study area	OTAL (ha)	OTA (ha)	Land use characteristics		LUS (ha)	Total length of transect (km)
			Presence of wood and urban buildings (%)	Obstructed view (%)		
1	105.3	84.1	62.8	13.5	19.9	4.2
2	354.9	350.5	62.4	13.1	85.8	14.6

OTAL observation theoretical area applied to linear transect, OTA observation theoretical area, LUS land useful sighting

Table 2 Results of spotlight counts for each study area and for each study period.

Study area	Total number of hares seen	CV (%)	Theoretical density (OTAL)	Exact density (LUS)
1	194	43.9	5.27	22.27
2	38	63.7	0.7	3.0

CV coefficient of variation