

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Using digital camera images to analyse snowmelt and phenology of a subalpine grassland

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/150402> since 2015-12-03T10:29:40Z

Published version:

DOI:10.1016/j.agrformet.2014.08.007

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

**[*AGRICULTURAL AND FOREST METEOROLOGY* , anno 2014, Vol 198-199, pag.
116-125 DOI: 10.1016/j.agrformet.2014.08.007]**

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://dx.doi.org/10.1016/j.agrformet.2014.08.007>]

13 **Using digital camera images to analyse snowmelt and phenology of an**
14 **alpine grassland.**

15

16 Tommaso Julitta*, Edoardo Cremonese, Mirco Migliavaccac, Roberto Colombo, Marta Galvagno, Consolata
17 Siniscalco, Micol Rossini, Francesco Fava, Sergio Cogliati, Umberto Morra di Cella, Annette Menzel

18 Remote Sensing of Environmental Dynamics Lab., DISAT, Università di Milano-Bicocca, P.zza della Scienza
19 1, 20126 Milano, Italy

20 Environmental Protection Agency of Aosta Valley, ARPA VdA, Climate Change Unit, Aosta, Italy

21 Max Planck Institute for Biogeochemistry Department Biogeochemical Integration, Jena, Germany

22 Plant Biology Department, Università degli Studi di Torino, Torino, Italy

23 Institute for Advanced Study, Technische Universität München, Lichtenbergstrasse 2a, 85748 Garching,
24 Germany

25 **Abstract**

26 Plant phenology is a good indicator of the impact of climate change on vegetation. In mountain areas
27 phenology is in general governed by environmental constraints such as air temperature, photoperiod and
28 presence of snow. Several studies in literature showed the effectiveness of using automated repeat digital
29 photograph for monitoring plant phenology. Most of these works analyse a small region of interest of the
30 image do not considering the potential of using the entire image to exploit temporal and spatial evolution
31 of phenology of a heterogeneous vegetated target. In the present study, digital images collected with a
32 Campbell Scientific Camera (model CC640) over 3 years (2009, 2010 and 2011) in a subalpine grassland
33 were used to analyse the spatial patterns of phenological events and their relationship with the timing of
34 snowmelt.

35 Yearly time series of green chromatic coordinates (gcc) were computed from hourly images. In order to
36 analyse the spatial pattern of phenological metrics, gcc time series were computed for each 10x10 pixel
37 regions of the target ecosystem and the beginning of the season for the 10x10 regions was extracted.
38 Based on the same grid dimension a snowmelt date map corresponding to the day of the year in which the
39 snow disappears from the ground was obtained.

40 Despite the snowmelt occurs rapidly, as maximum in seven days, several distinct spatial patterns were
41 identified. A negative correlation was found between the two snowmelt dates and the beginning of the
42 season, meaning that the growing season begins later in areas characterized by an early snowmelt, and vice
43 versa. Moreover the early snowmelt occurred in 2011 (about 40 days earlier than the previous two years)
44 highlighted two main phenological patterns at the beginning of the season. A field vegetational analysis
45 revealed that these patterns were related to communities with different plant composition. In particular
46 differences in terms of species seem to be related to convex and concave areas, suggesting that different

47 patterns of snow and soil accumulation due to micromorphology affect the vegetation communities and so
48 indirectly phenology. These results support the possibility of using digital repeat photography to analyze
49 the spatial variability of phenological development of complex ecosystems such as alpine grasslands.

50 **1 Introduction**

51 Climate change, is expected to strongly influence high altitude ecosystems in Europe, by increasing
52 temperature, changes in timing of winter and summer precipitation and duration of snow-pack (Alcamo et
53 al., 2007; Auer et al., 2007; Beniston, 2005).

54 Plant phenology is highly sensitive to climate change (Menzel et al., 2006) and particularly in ecosystems
55 with seasonality a clear correspondence exists between climate and phenological patterns (Richardson et
56 al., 2013). Snow is among the most important environmental factors controlling high-altitude plant
57 phenology (Galvagno et al., 2013; Wipf et al., 2009) moreover snow provides frost protection for plants in
58 winter, and water supply and nutrient mobilisation at snowmelt (Keller and Körner, 2003). Any changes in
59 the date of snowmelt or snow season establishment can induce strong ecosystem responses: 1) differences
60 in seasonal patterns of carbon and water fluxes related to the time for plant photosynthesis and growth
61 (i.e. snow-free season, (Galvagno et al., 2013)), 2) changes in interaction between species (Wipf et al.,
62 2006), between plants and their pollinator, pests or pathogens (Roy et al., 2004), 3) the timing of soil
63 nutrient availability and ecosystem functioning promoting conservative species (Saccone et al., 2013;
64 Schimel et al., 2004). According to the literature the expected warming of the Alpine region lead to more
65 frequent earlier snowmelt (Foppa and Seiz, 2012) that might cause changes in phenological patterns in
66 alpine vegetation.

67 Efforts have been made to review in a comprehensive way the numerous studies conducted by means of
68 direct snow manipulation (Wipf and Rixen, 2010). In fact several studies have investigated the effect of
69 changing snow cover on alpine plant phenology and high variability in responses has been related both to
70 species, growth forms and habitat (e.g. snowbed, fellfield). Previous studies (Keller and Körner, 2003; Wipf
71 and Rixen, 2010; Wipf et al., 2006) proposed that species adapted to habitats characterised by different
72 time of snow lying on the ground (such as snowbed or fell-field) can show distinct responses to snowmelt
73 advancement or delay.

74 Because many vegetation-climate feedbacks (albedo, evapotranspiration and CO₂ fluxes) are mediated by
75 phenology (Richardson et al., 2013) the monitoring and modelling of vegetation phenological cycle has
76 become a major issue in global change research (Cleland et al., 2007). In this study we hypothesised that
77 even within the same ecosystem it is possible to distinguish separate phenological behaviours related to
78 site microheterogeneity and snow patches. Since mountain ecosystems are generally located in complex
79 topography, the presence of microhabitats and the related different snow patches may induce, also within
80 the same alpine plant community, several vegetation developmental strategies. The observed phenological

81 patterns may reflect distinct adaptations to resource use efficiency, reproductive competition, site
82 micromorphology, life cycle completion strategies (Keller and Körner, 2003). Monitoring of diverse
83 phenological patterns in a plant community is useful to interpret the whole ecosystem response to climate
84 variability. In heterogeneous plant communities, such as mountain grassland, disentangling the
85 contribution of single species to community phenology is onerous. For the time being, to our awareness no
86 methodology has been proposed for long-term monitoring community-level phenological responses to,
87 first, the snowmelt variability, and, second, to the snow patches due to complex topography.

88 Traditional methods used to monitor plant phenology mainly consist of field observations and remote
89 sensing data (Busetto et al., 2010; Linderholm, 2006). However, both strategies present some limits in term
90 of spatial and temporal resolution. Direct phenological surveys hardly provide continuous information on
91 plant phenology and do not allow covering vast areas (Schwartz et al., 2002). Moreover observations do not
92 refer to a land surface phenology but to single plant and are frequently affected by the observer
93 subjectivity. As regard the satellite products, while they can provide invaluable synoptic phenological
94 information, are often challenging in mountainous areas due to complex topography, and do not allow a
95 detailed evaluation of the variability in species responses (Ide and Oguma, 2013).

96 Several studies demonstrated the capability of RGB digital imagery to provide accurate phenological data
97 (Ahrends et al., 2009; Ahrends et al., 2008; Bater et al., 2011; Migliavacca et al., 2011; Mizunuma et al.,
98 2013; Richardson et al., 2007; Richardson et al., 2009). The digital cameras are usually fixed above a
99 vegetated target (e.g. forests, croplands, grasslands and peatlands) and the images are continuously
100 acquired during the season. The colour information contained in the images (chromatics coordinates) is
101 then used as index of the canopy development. The time series of the indices extracted can be used to
102 identify the timing of key phenological events. Most of the studies that use digital repeated photography
103 for phenological applications compute the analysis on a single region of the image, considering it as
104 representative of the entire monitored ecosystem. An interesting improvement in this technique is to
105 analyse the spatial information in the considered frame in order to identify spatial differences of phenology
106 within the image (Ide and Oguma, 2013).

107 The RGB images can also be used to retrieve important information of the snowmelt processes, as reported
108 in literature (Hinkler et al., 2002; Parajka et al., 2012). Therefore the combination of the spatial information
109 of phenology and the spatial characterization of the snowmelt process can improve the knowledge of the
110 dependencies of plant development on the snowmelt timing.

111 In this study we investigate temporal and spatial patterns of phenology and their relationship with
112 snowmelt using digital images. Therefore the objectives of the study are: (i) can we use digital camera
113 photograph to identify spatial patterns of phenology within the image? (ii) Does phenology change
114 according to the snowmelt? What is the effect on phenology during early snowmelt year?

115 To this aim snowmelt and spring phenology maps were computed using digital images collected at a
116 subalpine grassland during three years and spatial patterns of phenology and their relationship with spatial
117 patterns of snowmelt and species compositions was evaluated and discussed.

118 **2 Materials and Methods**

119 **2.1 Site description**

120 The study site is located in the northwestern Italian Alps (Aosta Valley, IT) at 2160 m a.s.l. (45° 50'40" N, 7°
121 34'41" E). The region is classified as intra-alpine with semi-continental climate. The mean annual
122 temperature is 3.1 °C while the mean annual precipitation is 880 mm. The site is generally covered by snow
123 from the end of October to late May. The site is a subalpine unmanaged pasture with two different plant
124 compositions related to micromorphological conditions: the first one with dominant *Nardus stricta* the
125 second with dominant forbs (*Crocus vernus*, *Geum urbanum verde*, *Ranunculus pyrenaicus*, *Poa alpina*).

126 **2.2 Data collection**

127 Digital images were collected using a Campbell digital camera (model CC640 Campbell Scientific, Logan, UT,
128 USA) installed in 2009 at the experimental site. Following Richardson et al. (2007) the camera was pointed
129 north, and set at an angle of about 20° below horizontal. Camera focal length was 3.5 mm and the field of
130 view was approximately 79.8°. The camera was fixed at 2.5 m above ground and the same view scene has
131 been captured (Fig 1). The JPEG images collected have a resolution of 0.3 megapixels, with three colours
132 channels, red, green and blue, of 8-bit. The images were collected hourly from 10 am to 5 pm and exposure
133 mode and white balance were set to automatic. The 5644 JPEG images analysed during the study were
134 collected between 21st May 2009 and the 22nd November 2011.

135 **2.3 Image analysis**

136 Usually digital images are used to track the phenological plant development by considering a single Region
137 of Interest (ROI) in the image as reference of the mean behaviour of the entire ecosystem analysed
138 (Migliavacca et al., 2011; Richardson et al., 2006; Sonnentag et al., 2012). Few published works use more
139 than one ROI to analyse phenological differences between plants contained in the same image (Ahrends et
140 al., 2008; Jacobs et al., 2009; Mizunuma et al., 2013) while Ide and Oguma (2013) are the first that perform
141 a pixel per pixel analysis. In our study we evaluated the spatial evolution of snowmelt in the time
142 considering the effect of the snowmelt patterns on the spring vegetation phenology. The same ROI of the
143 image was used to analyse snowmelt and vegetation phenology (Fig 1). ROI position was defined to select
144 only the foreground part of the grassland in the image, less affected by clouds sometimes present in the
145 background (Migliavacca et al., 2011). Pixel size in the ROI ranges from 1 to 4 cm and the extension of the
146 analysed area corresponds approximately to 150 m². Then to investigate the spatial variability of the

147 processes, the analysis was conducted resampling the original ROI on a 10x10 pixels grid (Fig 1) providing a
148 number of cells grid equal to 931 (19 rows and 49 columns).

149 2.3.1 Snowmelt date maps

150 Snowmelt maps were computed performing an unsupervised classification (K-means) on the grid for
151 identifying two classes, corresponding to “snow” / “no snow”. In a second step the date of snowmelt per
152 each cell of the grid was defined as the DOY when more than 50% of pixels of the cell are classified as “no
153 snow”. Finally spatial anomalies of snowmelt were computed for each year as the difference between cell
154 values and mean snowmelt date of the entire grid.

155 2.3.2 Spring phenology maps

156 The chromatic information contained in the collected images can be extracted using indices that synthesize
157 the primary colours content of the images: red (R), green (G), and blue (B). The time series of single
158 channels (R,G,B) digital number (DN) values are known to be sensitive to illumination condition and camera
159 acquisition setup. These influences can be minimized transforming the RGB DN to RGB chromatic
160 coordinates (Gillespie et al., 1987; Woebbecke et al., 1995) as follow:

$$161 \quad g_{CC} = G_{DN} / (R_{DN} + G_{DN} + B_{DN}) \quad (1)$$

$$162 \quad b_{CC} = B_{DN} / (R_{DN} + G_{DN} + B_{DN}) \quad (2)$$

163 where R_{DN} , G_{DN} , B_{DN} are the red, green and blue DN values of each color channel respectively. In literature
164 g_{cc} is widely used to track the greenness of the canopy and could be considered as a good descriptor of the
165 plant phenology (Migliavacca et al., 2011; Richardson et al., 2007; Richardson et al., 2009; Sonnentag et al.,
166 2012).

167 The green chromatic coordinates (g_{cc}) were computed for each pixel of the ROI and then averaged for each
168 cell for all the images recorded during the three years, resulting in a cell based g_{cc} time series. The
169 illumination changes in the field of view, especially due to cloudy sky conditions, affected the diurnal and
170 seasonal trend of the index producing spikes in the time series. The time series was consequently filtered to
171 reduce data noise. The filtering strategy could be divided into two steps.

172 I) The diffuse light conditions affects the blue channel more than the others, thus the presence of the
173 clouds could be detected considering the behaviour of the blue channel (Ide and Oguma, 2010).
174 When the weather conditions are not stable the blue channel value shows a higher daily variability.
175 Here a threshold on the b_{cc} trend has been chosen. The images were discarded if the b_{cc} value is
176 outside the range:

$$177 \quad b_{cc_{dm}} - 5^{th} \text{ percentile } (b_{cc_{sd}}) < b_{cc} < b_{cc_{dm}} + 5^{th} \text{ percentile } (b_{cc_{sd}}) \quad (3)$$

178 where b_{cc} is blue chromatic coordinate defined in formula 2, $b_{cc\ dm}$ is the daily mean value of b_{cc} and
179 $b_{cc\ sd}$ is the seasonal standard deviation of b_{cc} .

180 From a visual inspection the filter based on the use of the b_{cc} time series pointed out that this
181 strategy removed all the cloudy images but also images collecting during clear sky condition. This
182 conservative approach could be applied considering a time series with 7 daily images.

183 II) The approach suggested by (Sonnentag et al., 2012) that consists in a 3 days moving-window
184 assigning the 90th percentile to the center day (calculated using all the gcc values referred to the
185 images acquired in 3 days of measurements) was used to process the data. The effect of this filters
186 is to reduce the temporal resolution of gcc time series from daily to three daily.

187 A curve fitting (cubic smoothing spline following Migliavacca et al. (2011) was computed to the gcc time
188 series. The beginning of the season (BOS) was defined as the time where the greenness curve reaches the
189 half maximum of the spring growth (Bradley et al., 2007; Fisher et al., 2006; White et al., 1997). According
190 to the literature we used a fixed threshold on the curve (50% of the growth) to obtain the BOS. The defined
191 methodology was automatically applied first to the averaged ROI value and subsequently to all the cells of
192 the grid leading to a BOS map. A no value flag was assigned to grid cells occupied by the grey panel visible
193 in the images and to grid cells having BOS values outside the range snowmelt date - end of July. Yearly BOS
194 anomaly maps were obtained subtracting to each cell the mean BOS value of the entire grid.

195 **2.4 Comparison of the beginning of the season and the snowmelt dates**

196 In order to evaluate the influence of snowmelt temporal and spatial patterns on spring phenology, BOS and
197 snowmelt maps of the three years were compared performing the non-parametric Kendall correlation test
198 using all the values of the maps. Correlation analysis was also performed on data aggregated with the
199 following approach: grid cells were grouped in 15 classes according to snowmelt quantiles, for each cell
200 falling in a given class snowmelt and BOS median was computed. Lastly generalized linear models (glm,
201 (McCullagh and Nelder, 1989) were computed to analyse the sensitivity of the BOS to the snowmelt.

202 **3 Results**

203 **3.1 Spring phenology and snowmelt maps**

204 Air temperature trend was comparable during the 3 years. While a warm spring spell in April 2011 caused
205 an extremely early snowmelt (~40 days) if compared to both 2009 and 2010 or to long term (1928-2010)
206 data (Galvagno et al., 2013). The spatial distribution of snowmelt dates is highly heterogeneous (Fig 2).
207 Mean snow melt date occurred at DOY at 145, 143 and 103 in 2009, 2010, 2011 respectively even if the
208 time needed for a complete snowmelt on all the area analysed ranged between 3 days in 2009 and 2010

209 and 6 days in 2011 (Fig 3). Considering the BOS map a similar heterogeneous behaviour has been found.
210 Mean BOS occurred at DOY 172, 173, 162 in 2009, 2010, 2011 respectively while considering the variability
211 of the BOS dates in the analysed area of each year ranged between 66 days in 2009 and 111 days in 2011
212 (Fig 3). A consistent spatial dynamic can be seen in the 3 years both in the snowmelt and in the BOS maps.
213 The correlation analysis results performed between the snowmelt maps and the spring phenology maps are
214 reported in table 1 (expressed in terms of tau, Kendall rank correlation coefficient). The results showed a
215 significant inverse correlation between the two maps, meaning that later snowmelt areas correspond to
216 areas with earlier BOS.

217 **3.2 Phenology at community level**

218 The early spring spell occurred in 2011 gave us a unique opportunity to analyse the BOS distribution within
219 the image. The two modes highlighted in the 2011 BOS frequency distribution (Fig 3) have been used to
220 create two masks on the BOS map distinguishing the areas where the BOS occurred early or later.
221 Therefore the two masks have been considered to define plots at ground level for a phytosociological
222 surveys with the aim of characterizing plant species composition. The survey was conducted in May 2012. A
223 total of 8 early and 8 late phenology plots were selected for the observations. Species composition,
224 grouped into early or late BOS, are reported in table 2. Generally the areas characterized by an earlier BOS
225 correspond to areas with abundance of forbs while in areas with late BOS the main coverage is attributable
226 to grasses species. In figure 4 the yearly averaged BOS is shown grouped into forbs and grasses species
227 (according to the bimodal distribution of 2011 BOS) and considering the averaged BOS referring to the
228 overall ROI value as well. The variation in the grasses BOS within the 3 years varies of 7 days as maximum in
229 2011. On the contrary the averaged forbs BOS is similar in 2009 and 2010 while occurs 26 days before in
230 2011. Considering the ROI averaged value of BOS during the three year (2009: DOY mean 171 ± 13 , 2010:
231 DOY mean 173 ± 14 , 2011: DOY mean 165 ± 23) no significant advance can be noticed.

232 3.3 Phenology and snowmelt relationship

233 Due to the relationship detected between snowmelt and phenology and considering the differences
234 between the two main functional groups (grasses and forbs) we carried out the sensitivity analysis between
235 snowmelt and phenology using glm both on the entire community and on the two functional groups (Fig 5).
236 A strong negative correlation was found between BOS and snowmelt as reported in table 3. The table
237 reports separately the results obtained by (i) applying the model on all the grid cells or (ii) distinguishing
238 between the grid cells referred to each of the two plant functional group (forbs and grasses). The results
239 suggested that a constant inverse relation between snowmelt date and BOS existed, meaning that areas
240 characterized by early snowmelt corresponded to area where the BOS took place later (and vice versa). In
241 particular the results referred to the all community show a strong relationship in the 3 years considered
242 (high value of r^2) with a similar slope in the three years considered. On the contrary analysing separately

243 the two functional groups we obtained different results. The forbs BOS in 2011 took place before and the
244 slope of the relationship, constant in 2009 and 2010, decreased, meaning a less sensitivity to snowmelt on
245 the BOS when the snowmelt occurs before even if the r^2 values of these relationships are still high. For
246 grasses in 2011 the BOS took place few days later (as shown in Fig 4). While the BOS increased during the 3
247 years, the slopes of the relationship between snowmelt and BOS decreased suggesting a reduction of the
248 influence of snowmelt on grasses BOS depending on when the snowmelt takes place.

249 **4 Discussions**

250 **4.1 Snowmelt and spatial phenology**

251 We showed the suitability of using digital repeated images for analysing the phenological heterogeneity of
252 an ecosystem such as grasslands. The results show the potential of using a standard digital camera
253 collecting automatically repeated images to monitor patterns of phenology in relation to snowmelt
254 variability in subalpine grassland. The combination of temporal and spatial information gave important
255 insights on the grassland vegetation development dynamics, improving the characterization of the study
256 site. In turn, the spatial analysis of the snowmelt can provide useful information for interpreting the
257 complex relationships existing between this process and the phenological behaviour of a heterogeneous
258 ecosystem. The obtained outcomes of the analysis point out the importance of using spatial disaggregated
259 data. In figure 6 the gcc curves are separately plotted distinguishing between the ROI value, the forbs and
260 the grasses. Without disentangling the contribution of forbs and grasses in the phenological development
261 of the ROI we obtain a similar time series during the three considered years, while analysing separately the
262 information related to forbs and grasses a different behaviour can be noticed. In particular in 2011 the
263 differences between the two functional groups are more evident as expected.

264 The snowmelt occurred in latest May in 2009 and 2010 while in the middle of April in 2011, due to a strong
265 spring warm. The differences in terms of date also caused a different overall duration of the snowmelt
266 process that varies between 3 and 6 days, depending on the year (Fig 3). Even if the snowmelt took place
267 differently according to the year considered similar snowmelt spatial patterns have been found in all the
268 years considered, meaning that areas where the snow disappears earlier or later are constant among years.
269 These spatial patterns are related to the surface microtopography, in fact in concave areas the snow lasts
270 longer, while in the convex ones the snow disappears before.

271 Similar patterns exist in the spring phenology maps as well. We showed an interesting negative correlation
272 between snowmelt and BOS patterns (tab 1) indicating that the way in which snow melt takes place also
273 affects phenology as previously indicated by (Evans et al., 1989) and (Walker et al., 1995). The phenological
274 differences are strictly related to the presence of two functional groups. The presence of different species
275 is shown in Fig 3 where the bimodal distribution of BOS observed in 2011 underlines two different

276 phenological behaviours. In particular these phenological behaviours are related to the presence of grasses
277 and forbs (tab 2). This observation confirms the hypothesis that the spatial pattern of snowmelt, through
278 the influence exerted on species distribution, is indirectly responsible for the observed heterogeneity in
279 phenology (BOS).

280 **4.3 Ecological feedback**

281 The results point out that the forbs species, whose opportunistic behaviour is well known in literature
282 (Keller and Körner, 2003), are capable to immediately tune its growth even in very early years (Fig 4),
283 regardless of air temperature and photoperiod. Indeed in 2011, the year characterized by an extreme early
284 snowmelt, a strong anticipation of BOS is observed in areas of the grassland dominated by forbs.
285 Conversely the grasses (mainly *Nardus stricta*) showed a “safer” behaviour and started to develop in similar
286 periods among years and did not show any advancement in BOS neither in the extreme year 2011. The BOS
287 grasses seems to be more stable trough years (fig 4) and thus more affected by photoperiod (and
288 temperature) rather than the date of snowmelt. These results agree with (Wipf and Rixen, 2010), where
289 grasses did not show any consistent response to snow manipulations. Forbs are sensitive precursors of
290 snowmelt (Keller and Körner, 2003). Hence the fast BOS of forbs species in 2011 could be the response of a
291 species adapted to grow in late lying snow areas. While in habitats with early snowmelt (e.g. convex areas),
292 any advancement of snowmelt may lead to frost-damage, hence a photoperiod-dependent BOS may be a
293 safer strategy for this species. Differences in phenological development may also have an important role on
294 the whole ecosystem processes, especially during years characterised by extreme snowmelt. For example a
295 reduction in net CO₂ ecosystem uptake was observed in spring/summer 2011 (Galvagno et al., 2013): this
296 may in part be due to the later development of grasses compared to forbs (less abundant at ecosystem
297 level).

298 **5 Conclusions**

299 During this study we demonstrate that digital repeat photographs can provide reliable information on the
300 analysis of the effect of snowmelt on plant phenology in a heterogeneous ecosystem such as grassland
301 without any manipulation experiment. The repeated images can be used to describe the spatial evolution
302 of snow patches during the snowmelt phase. At the same time we show that by combining temporal and
303 spatial information of the images we're able to appreciate a phenological spatial heterogeneity within the
304 same ecosystem. We underline the importance of considering the spatial information of the images usually
305 not taken into account in most of the published studies that analyse digital repeated images for
306 phenological purposes. This analysis approach can provide a more detailed description of heterogeneous
307 ecosystems, helpful for a better knowledge of the phenological process.

308 The analysis show that plant phenology reflects the evolution of snowmelt (mainly driven by
309 microtopography), through the influence exerted on species distribution, confirming a dependency of the
310 plant development on the snowmelt.

311 We can conclude that digital repeat photographs can be a tool for long-term monitoring of the community-
312 level phenological responses to the snowmelt variability.

313 **Acknowledgments**

314 This work was supported by the PhenoALP project, an Inter-reg project co-funded by the European
315 Regional Development Fund, under the operational program for territorial cooperation Italy–France
316 (ALCOTRA) 2007–2013. The analysis was supported by the Aosta Valley Region Operative Program
317 “Occupation” co-funded by European Social Fund. We thank all the people that collaborate in maintaining
318 the study site working.

319

320 **TABLES**

	2009	2010	2011
Raw	-0.34	-0.25	-0.3
Aggregated	-0.89	-0.63	-0.73

321 Table 1 Kendall correlation coefficient (tau) computed between the snowmelt date and the BOS. The two lines refer to the
 322 correlation analysis conducted considering all the grid cell values (Raw line) and using only the aggregated data (Aggregated).

323

	Early beginning of season	Late beginning of season
Grasses	52.8±14	93.4±5
Forbs	47.2±14	6.6±5
<i>Nardus stricta</i>	51 ±17	93.4±5
<i>Poa alpine</i>	1.8±2	-
<i>Crocus vernus</i>	12.2±10	2.4±3
<i>Potentilla alpine</i>	3±4	2.2±2
<i>Trifolium alpinum</i>	3±4	-
<i>Arnica Montana</i>	5±8	-
<i>Polygonum bistorta</i>	2±4	-
<i>Hieracium pilosella</i>	1±2	-
<i>Geum urbanum verde</i>	16±9	2±2
<i>Ranunculus pyrenaicus</i>	3±4	-
<i>Leontodon</i>	1±2	-
<i>Ranunculus acris</i>	1±0.4	-

324 Table 2 Field phenological analysis results, May 2012. The table represent the averaged species composition distribution
 325 expressed in terms of percentage of cover. The plots considered in the field were defined according to areas identified in the
 326 2011 BOS anomaly map. 16 plots were analysed in the field distinguishing 8 areas characterized by an early beginning of the
 327 season and 8 areas were the season took place later.

328

	All				Forbs				Grasses			
	r2	m	St er	P	r2	M	St er	p	r2	M	St er	p
2009	0.9	-17.21	1.93	< 0.01	0.70	-14.48	3.56	< 0.01	0.74	-11.48	2,46	< 0.01
2010	0.75	-13.50	3.107	< 0.01	0.75	-14.4	3.22	< 0.01	0.56	-4.11	1.29	0.02
2011	0.8	-20.84	3.29	< 0.01	0.6	-3.93	0.82	< 0.01	0.63	-2.79	0.68	< 0.01

329 Table 3 Results of the analysis conducted using Generalized Linear Model. The relationship between the BOS and the snowmelt
 330 dates is here reported in terms of r², slope, standard error and p value. The table distinguish between analysis conducted on the
 331 all scene, or divided into the two main plant communities (forbs or grasses).

332

333

334 **FIGURES**

335




336

337 **Figure 1.** The study site JPEG image. The Region of interest considered during the analysis is bordered in red. It has been chosen
338 in the foreground avoiding problems related to distance, weather and excessive pixel size differences. The region selected has
339 been then divided using a grid of 10x10 pixel. As depicted in the figure 19 row and 49 columns result from the division. In the
340 right side the grey panel, considered as no value during the analysis

341


342



343


344 **Figure 2.** Left panel: snowmelt anomaly maps computed against the mean value of the entire grid; right panel: maps of the
345 anomaly of BOS computed against the mean BOS of the entire grid. Each row identifies a different year. The color scales
346 represent the range of variation of the snowmelt/BOS dates in the images. Dark green highlights areas where dates are
347 anticipated, while the light green refers to areas where the snowmelt or the BOS are postponed.

348



349

350 **Figure 3. Frequency distribution of snowmelt dates (light blue) and BOS (white) in the 3 years considered. The unimodal**
 351 **distribution in the first two years becomes bimodal in the third one. In the year 2011 the snowmelt date is strongly anticipated**
 352 **compared to 2009 and 2010 (~40 days before).**




353


354 **Figure 4. The graph shows the 3 years averaged BOS and related standard deviation obtained by distinguishing between the two**
 355 **main functional groups (grasses and forbs) and the overall ROI value. The BOS of the forbs species is strongly anticipated in 2011**
 356 **(26 days before the two previously years) while the variation of grasses BOS is stable in the 3 years considered. Analysing the**
 357 **averaged BOS value of the ROI a small advance with the higher standard deviation is recorded in 2011.**

358


359



360




361



362

363 **Figure 5. Relationship between snowmelt date and BOS analysed using the glm. The models have been applied considering the**
 364 **overall value of all the grid cells (A), and distinguishing the plant communities existing at the grassland site (B, C). In particular**
 365 **the second graph represents the analysis carried out considering only the grasses species, while the third one refer to the forbs**
 366 **community.**



367

368 **Figure 6. The 3 years GI time series are separately plotted considering averaged value of the overall ROI (red line), and the two**
 369 **functional groups (Forbs, dark green line and grasses, light green line). The 2011 is characterized by a strong advance in the**
 370 **growth of the forbs species while in 2009 and 2010 the difference of the two functional groups is not so emphasized.**

371

372

373 **REFERENCES**

- 374 Ahrends, H. et al., 2009. Tree phenology and carbon dioxide fluxes: use of digital photography for process-
375 based interpretation at the ecosystem scale. *Climate Research*, 39(3): 261-274.
- 376 Ahrends, H.E. et al., 2008. Quantitative phenological observations of a mixed beech forest in northern
377 Switzerland with digital photography. *Journal of Geophysical Research: Biogeosciences*, 113(G4):
378 G04004.
- 379 Alcamo, J., Flörke, M. and Märker, M., 2007. Future long-term changes in global water resources driven by
380 socio-economic and climatic changes. *Hydrological Sciences Journal*, 52(2): 247-275.
- 381 Auer, I. et al., 2007. HISTALP—historical instrumental climatological surface time series of the Greater
382 Alpine Region. *International Journal of Climatology*, 27(1): 17-46.
- 383 Bater, C. et al., 2011. Using digital time-lapse cameras to monitor species-specific understorey and
384 overstorey phenology in support of wildlife habitat assessment. *Environmental Monitoring and
385 Assessment*, 180(1-4): 1-13.
- 386 Beniston, M., 2005. Mountain Climates and Climatic Change: An Overview of Processes Focusing on the
387 European Alps. *pure and applied geophysics*, 162(8-9): 1587-1606.
- 388 Bradley, B.A., Jacob, R.W., Hermance, J.F. and Mustard, J.F., 2007. A curve fitting procedure to derive inter-
389 annual phenologies from time series of noisy satellite NDVI data. *Remote Sensing of Environment*,
390 106(2): 137-145.
- 391 Busetto, L. et al., 2010. Remote sensing of larch phenological cycle and analysis of relationships with climate
392 in the Alpine region. *Global Change Biology*, 16(9): 2504-2517.
- 393 Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A. and Schwartz, M.D., 2007. Shifting plant phenology in
394 response to global change. *Trends in Ecology & Evolution*, 22(7): 357-365.
- 395 Corripio, J.G., 2004. Snow surface albedo estimation using terrestrial photography. *International Journal of
396 Remote Sensing*, 25(24): 5705-5729.
- 397 Evans, E., Smith, C. and Gendron, R., 1989. Timing of reproduction in a prairie legume: seasonal impacts of
398 insects consuming flowers and seeds. *Oecologia*, 78(2): 220-230.
- 399 Fisher, J.I., Mustard, J.F. and Vadeboncoeur, M.A., 2006. Green leaf phenology at Landsat resolution: Scaling
400 from the field to the satellite. *Remote Sensing of Environment*, 100(2): 265-279.
- 401 Foppa, N. and Seiz, G., 2012. Inter-annual variations of snow days over Switzerland from 2000–2010
402 derived from MODIS satellite data. *The Cryosphere*, 6(2): 331-342.
- 403 Galvagno, M. et al., 2013. Phenology and carbon dioxide source/sink strength of a subalpine grassland in
404 response to an exceptionally short snow season. *Environmental Research Letters*, 8(2): 025008.
- 405 Gillespie, A.R., Kahle, A.B. and Walker, R.E., 1987. Color enhancement of highly correlated images. II.
406 Channel ratio and "chromaticity" transformation techniques. *Remote Sensing of Environment*,
407 22(3): 343-365.
- 408 Hinkler, J., Pedersen, S.B., Rasch, M. and Hansen, B.U., 2002. Automatic snow cover monitoring at high
409 temporal and spatial resolution, using images taken by a standard digital camera. *International
410 Journal of Remote Sensing*, 23(21): 4669-4682.
- 411 Ide, R. and Oguma, H., 2010. Use of digital cameras for phenological observations. *Ecological Informatics*,
412 5(5): 339-347.
- 413 Ide, R. and Oguma, H., 2013. A cost-effective monitoring method using digital time-lapse cameras for
414 detecting temporal and spatial variations of snowmelt and vegetation phenology in alpine
415 ecosystems. *Ecological Informatics*, 16(0): 25-34.
- 416 Jacobs, N. et al., 2009. The global network of outdoor webcams: properties and applications, *Proceedings of
417 the 17th ACM SIGSPATIAL International Conference on Advances in Geographic Information
418 Systems*. ACM, Seattle, Washington, pp. 111-120.
- 419 Keller, F. and Körner, C., 2003. The Role of Photoperiodism in Alpine Plant Development. *Arctic, Antarctic,
420 and Alpine Research*, 35(3): 361-368.
- 421 Lieth, H., 1974. Purposes of a Phenology Book. In: H. Lieth (Editor), *Phenology and Seasonality Modeling*.
422 *Ecological Studies*. Springer Berlin Heidelberg, pp. 3-19.

423 Linderholm, H.W., 2006. Growing season changes in the last century. *Agricultural and Forest Meteorology*,
424 137(1–2): 1-14.

425 McCullagh, P. and Nelder, J.A., 1989. *Generalized linear models (Monographs on statistics and applied*
426 *probability 37)*. Chapman Hall, London.

427 Menzel, A. et al., 2006. European phenological response to climate change matches the warming pattern.
428 *Global Change Biology*, 12(10): 1969-1976.

429 Migliavacca, M. et al., 2011. Using digital repeat photography and eddy covariance data to model grassland
430 phenology and photosynthetic CO₂ uptake. *Agricultural and Forest Meteorology*, 151(10): 1325-
431 1337.

432 Mizunuma, T. et al., 2013. The relationship between carbon dioxide uptake and canopy colour from two
433 camera systems in a deciduous forest in southern England. *Functional Ecology*, 27(1): 196-207.

434 Parajka, J., Haas, P., Kirnbauer, R., Jansa, J. and Blöschl, G., 2012. Potential of time-lapse photography of
435 snow for hydrological purposes at the small catchment scale. *Hydrological Processes*, 26(22): 3327-
436 3337.

437 Richardson, A. et al., 2007. Use of digital webcam images to track spring green-up in a deciduous broadleaf
438 forest. *Oecologia*, 152(2): 323-334.

439 Richardson, A.D., Bailey, A.S., Denny, E.G., Martin, C.W. and O'Keefe, J., 2006. Phenology of a northern
440 hardwood forest canopy. *Global Change Biology*, 12(7): 1174-1188.

441 Richardson, A.D., Braswell, B.H., Hollinger, D.Y., Jenkins, J.P. and Ollinger, S.V., 2009. Near-surface remote
442 sensing of spatial and temporal variation in canopy phenology. *Ecological Applications*, 19(6): 1417-
443 1428.

444 Richardson, A.D. et al., 2013. Climate change, phenology, and phenological control of vegetation feedbacks
445 to the climate system. *Agricultural and Forest Meteorology*, 169(0): 156-173.

446 Roy, B.A., Güsewell, S. and Harte, J., 2004. RESPONSE OF PLANT PATHOGENS AND HERBIVORES TO A
447 WARMING EXPERIMENT. *Ecology*, 85(9): 2570-2581.

448 Saccone, P. et al., 2013. The effects of snowpack properties and plant strategies on litter decomposition
449 during winter in subalpine meadows. *Plant and Soil*, 363(1-2): 215-229.

450 Schimel, J.P., Bilbrough, C. and Welker, J.M., 2004. Increased snow depth affects microbial activity and
451 nitrogen mineralization in two Arctic tundra communities. *Soil Biology and Biochemistry*, 36(2): 217-
452 227.

453 Schwartz, M.D., Reed, B.C. and White, M.A., 2002. Assessing satellite-derived start-of-season measures in
454 the conterminous USA. *International Journal of Climatology*, 22(14): 1793-1805.

455 Sonnentag, O. et al., 2012. Digital repeat photography for phenological research in forest ecosystems.
456 *Agricultural and Forest Meteorology*, 152(0): 159-177.

457 Walker, M.D., Ingersoll, R.C. and Webber, P.J., 1995. Effects of Interannual Climate Variation on Phenology
458 and Growth of Two Alpine Forbs. *Ecology*, 76(4): 1067-1083.

459 White, M.A., Thornton, P.E. and Running, S.W., 1997. A continental phenology model for monitoring
460 vegetation responses to interannual climatic variability. *Global Biogeochemical Cycles*, 11(2): 217-
461 234.

462 Wipf, S. and Rixen, C., 2010. A review of snow manipulation experiments in Arctic and alpine tundra
463 ecosystems. *Polar Research*, 29(1): 95-109.

464 Wipf, S., Rixen, C. and Mulder, C.P.H., 2006. Advanced snowmelt causes shift towards positive neighbour
465 interactions in a subarctic tundra community. *Global Change Biology*, 12(8): 1496-1506.

466 Wipf, S., Stoeckli, V. and Bebi, P., 2009. Winter climate change in alpine tundra: plant responses to changes
467 in snow depth and snowmelt timing. *Climatic Change*, 94(1-2): 105-121.

468 Woebbecke, D.M., Meyer, G.E., Von Bargen, K. and Mortensen, D.A., 1995. Color indices for weed
469 identification under various soil, residue, and lighting conditions. *Transactions of the American*
470 *Society of Agricultural Engineers*, 38(1): 259-269.

471

472