

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Cultural Adaptation of the Difficulties in Emotion Regulation Scale: Reliability and Validity of an Italian Version**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1508137> since 2016-11-29T10:25:41Z

*Published version:*

DOI:10.1002/jclp.21876

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Giromini L; Velotti P; de Campora G; Bonalume L; Zavattini GC. Cultural Adaptation of the Difficulties in Emotion Regulation Scale: Reliability and Validity of an Italian Version. *JOURNAL OF CLINICAL PSYCHOLOGY*. 68 (9) pp: 989-1007.  
DOI: 10.1002/jclp.21876

The publisher's version is available at:

<http://doi.wiley.com/10.1002/jclp.21876>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1508137>



**Cultural Adaptation of the Difficulties in Emotion Regulation Scale: Reliability and Validity of an Italian Version**

Journal:	<i>Journal of Clinical Psychology</i>
Manuscript ID:	JCLP-11-0038.R2
Wiley - Manuscript type:	Research Article
Keywords:	emotional dysregulation, emotion regulation, adaptation, eating disorders

SCHOLARONE™  
Manuscripts

Peer Review

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

**Abstract**

**Objective:** The aim of this study was to evaluate the reliability and validity of an Italian version of the Difficulties in Emotion Regulation Scale (DERS; Gratz & Roemer, 2004).

**Method:** Three studies were completed. First, factorial structure, internal consistency, and concurrent validity of our Italian version of the DERS were examined with a sample of 323 students (77% female; mean age 25.6). Second, test-retest analyses were completed using a different sample of 61 students (80% female; mean age 24.7). Third, the scores produced by a small clinical sample ( $N = 38$ ; mean age = 24.2) affected by anorexia, binge eating disorder, or bulimia were compared to those of an age-matched, nonclinical female sample ( $N = 38$ ; mean age = 24.7). **Results:** The factorial structure replicated quite well the six-factor structure proposed by Gratz and Roemer. The internal consistency and test-retest reliability were adequate and comparable to previous findings. The validity was good, as indicated by both the concurrent validity analysis and the clinical-nonclinical sample comparison. **Conclusions:** These studies provide further support for the multidimensional model of emotion regulation postulated by Gratz and Roemer and strengthen the rationale for cross-cultural utilization of the DERS.

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 In the past two decades, emotion regulation (ER) has been seen as what underlies  
4  
5 diverse symptom presentation. Elaborated emotion-dysregulation theories have been applied  
6  
7 to depression (Gross & Muñoz, 1995), generalized anxiety disorder (McLaughlin, Mennin, &  
8  
9 Farach, 2007; Mennin, Heimberg, Turk, & Fresco, 2005), alcoholism and substance abuse  
10  
11 (Fox, Axelrod, Paliwal, Sleeper, & Sinha, 2007; Fox, Hong, & Sinha, 2008; Gratz,  
12  
13 Bornovalova, Delany-Brumsey, Nick, & Lejuez, 2007), self-injury (Klonsky, 2009), suicide  
14  
15 (Zlotnick, Donaldson, Spirito, & Pearlstein, 1997), eating disorders (Sim & Zeman, 2005,  
16  
17 2006; Whiteside, Chen, Neighbors, Hunter, Lo &, Larimer, 2006), borderline personality  
18  
19 disorder (Glenn & Klonsky, 2009; Linehan, 1993), and post-traumatic stress symptoms (Tull,  
20  
21 Barrett, McMillan, & Roemer, 2007). Despite this increased attention, there remains some  
22  
23 confusion about what ER is (Gross, 2008). This is partly due to the several processes that  
24  
25 underlie diverse emotion regulation acts as well as the lack of consistent, agreed-upon  
26  
27 conceptualizations of ER.  
28  
29  
30

31  
32 Broadly defined, the construct of ER refers to the conscious or unconscious attempt of  
33  
34 an individual to influence the expression and type of emotion he or she feels (Mauss, Bunge  
35  
36 & Gross, 2007). ER also encompasses the extrinsic or intrinsic processes that are responsible  
37  
38 for the monitoring, evaluation, and modification of emotional responses, including the ability  
39  
40 to modulate and regulate behaviors when experiencing emotional distress (Gratz & Roemer,  
41  
42 2004; Thompson, 1994).  
43  
44

45  
46 According to Gross and Thompson (2007), there are five points in time in which  
47  
48 emotions could be regulated: situation selection, situation modification, attentional  
49  
50 deployment, cognitive change, and response modulation. Each of the five ER strategies has  
51  
52 received a great deal of interest by several researchers (Gross, 2008). Moreover, two ER  
53  
54 strategies have received major attention: *cognitive reappraisal* and *expressive suppression*.  
55  
56 The first consists of attempts to think about the situation as to alter its meaning and emotional  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 impact; the second consists of attempts to inhibit or reduce ongoing emotion-expressive  
4  
5 behavior (Gross, 1998).  
6

7  
8 An attempt to integrate all the features of emotion regulation and dysregulation into a  
9  
10 unitary, multidimensional conceptualization was made by Gratz and Roemer (2004). The  
11  
12 authors, after reviewing the relevant literature, described ER as “involving the (a) awareness  
13  
14 and understanding of emotions, (b) acceptance of emotions, (c) ability to control impulsive  
15  
16 behaviors and behave in accordance with desired goals when experiencing negative emotions,  
17  
18 and (d) ability to use situationally-appropriate emotion regulation strategies flexibly to  
19  
20 modulate emotional responses as desired in order to meet individual goals and situational  
21  
22 demands” (p. 42). On the basis of this conceptualization, Gratz and Roemer also proposed a  
23  
24 multidimensional self-report measure – the Difficulties in Emotion Regulation Scale (DERS;  
25  
26 Gratz & Roemer, 2004). This scale includes six clinically-relevant domains of emotion  
27  
28 dysregulation: nonacceptance of emotion responses (NONACCEPTANCE), lack of emotional  
29  
30 awareness (AWARENESS), limited access to emotion regulation strategies (STRATEGIES),  
31  
32 difficulties engaging in goal-directed behavior when emotionally aroused (GOALS), impulse  
33  
34 control difficulties (IMPULSE), and lack of emotional clarity (CLARITY).  
35  
36  
37

38  
39 The Gratz and Roemer (2004)’s model of ER has recently received increased attention,  
40  
41 and empirical findings have supported the validity and reliability of the DERS in different  
42  
43 samples (Fox et al., 2007; Gratz, 2007; Gratz & Gunderson, 2006; Gratz, Lacroce, &  
44  
45 Gunderson, 2006; Gratz & Roemer, 2004; Gratz, Rosenthal, Tull, Lejuez, & Gunderson,  
46  
47 2006; Gratz, Tull, Baruch, Bornovalova, & Lejuez, 2008; Salters-Pedneault, Roemer, Tull,  
48  
49 Rucker, & Mennin, 2006; Tull & Roemer, 2007). Nonetheless, more research on this topic is  
50  
51 needed, and the extent to which Gratz and Roemer’s results can be generalized to samples  
52  
53 from diverse cultural backgrounds remains an empirical question.  
54  
55

**Adaptability of the DERS to an Italian Sample.**  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 Investigating the diverse components of the DERS in different cultures provides a  
4 unique opportunity to identify the quintessential features of ER. The more reliable and valid a  
5 scale is across different cultures, the higher is its relevance and importance in defining and  
6 measuring the construct. Yet, the DERS has only been adapted for and administered to a few  
7 diverse cultural backgrounds thus far, and few studies provide information on the cross-  
8 cultural adaptability of the DERS. Specifically, Dutch (Neumann, van Lier, Gratz, & Koot,  
9 2010), Turkish (Rugancı & Gençöz, 2010), and Portuguese (Coutinho, Ribeiro, Ferreirinha, &  
10 Dias, 2010) versions have overall confirmed the factorial structure identified by Gratz and  
11 Roemer (2004), and shown good internal consistency and concurrent validity. An Italian  
12 version has obtained similar results, but the factorial structure of the original version has not  
13 been replicated by a confirmatory factor analysis (Sighinolfi, Norcini Pala, Chiri, Marchetti,  
14 & Sica, 2010). In regard to this aspect, however, it should be pointed out that the sample  
15 collected by Sighinolfi et al. (2010) was smaller ( $N = 190$  vs.  $N = 357$ ), older (mean age =  
16 30.8,  $SD = 9.7$  vs. mean age = 23.1,  $SD = 5.7$ ) and more heterogeneous (e.g., participants were  
17 recruited in job-related contexts, at the university or in open psychology meetings) than the  
18 student sample collected by Gratz and Roemer (2004). Also, as stated by the Italian authors  
19 themselves (Sighinolfi et al. 2010, p. 159), the results obtained by a subsequent exploratory  
20 factor analysis were not much different from the original structure proposed by Gratz and  
21 Roemer. Indeed, although two items were removed from the analysis (i.e., item 17 and item  
22 34) and the item composition of one factor (i.e., STRATEGIES) was slightly different from  
23 that obtained by Gratz and Roemer, the resulting six-factor model remained largely  
24 unchanged. The debate on the factorial structure of the DERS in Italy is not yet settled and  
25 more research on this topic is needed.

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54 According to Terracciano, McCrae, and Costa (2003), many stereotypes suggest that  
55 Italians are characterized by an “affective volatility.” More specifically, stereotypes indicate  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 that Italians tend to be romantic, jealous, and short-tempered, speak loudly in public,  
4  
5 communicate with hand gestures and facial expressions, have an exaggerated love of their  
6  
7 mothers, and so forth. Although some indirect empirical support exists for these suppositions  
8  
9 (McCrae, 2001; Terracciano, McCrae & Costa, 2003), little is known about ER in Italy and  
10  
11 possible differences in ER between Italian and American cultures. In fact, the studies that  
12  
13 examined ER cross-cultural differences primarily focused on the comparison between  
14  
15 collectivistic and individualistic cultures (Hofstede, 2001), and from this perspective Italy  
16  
17 seems to be more similar than different to North America (Nelson & Fuvish, 2004; Triandis,  
18  
19 1993). Hofstede and colleagues (Hofstede, 1980; Hofstede, Hofstede & Minkov, 2010), for  
20  
21 instance, reported that Italy is one of the seven countries that have individualism as their  
22  
23 highest dimension, and that individualism is a value shared by American, British, Dutch,  
24  
25 French, and also Italian populations. Though this may or not be true, Italian populations may  
26  
27 still have the “affective volatility” discussed above. Thus, more research on this topic is  
28  
29 needed as well.  
30  
31  
32  
33

34 Investigating the adaptability of the DERS in an Italian population would broaden our  
35  
36 knowledge of ER and its assessment in two ways. First, by administering the DERS to a large  
37  
38 sample of Italians, Italian research and knowledge on ER would be expanded. Second, and  
39  
40 most importantly, establishing the reliability and validity of the DERS also in Italy would  
41  
42 provide support to Gratz and Roemer (2004)’s conceptualization of ER, strengthen cross-  
43  
44 cultural utilization of the DERS, and make further cross-cultural comparisons possible.  
45  
46

47 For these aforementioned reasons, we conducted three studies. In study 1, we  
48  
49 translated the DERS into Italian and investigated its factorial structure, internal consistency,  
50  
51 and concurrent validity. In study 2, the test-retest reliability of our adapted version of the  
52  
53 DERS was examined. In study 3, the construct validity of our version of the DERS was  
54  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

further investigated by comparing the DERS scores produced by a nonclinical sample to those of a clinical sample characterized by deficits in ER.

**Study 1**

It has been reported that cultures differ in terms of appraisals that lead to emotion (Matsumoto, Kudoh, Scherer & Wallbot, 1988; Mauro, Sato & Tucker, 1992; Roseman, Dhawan, Rettek & Naidu, 1995; Scherer, 1997a, 1997b), emotional expression (Matsumoto & Kupperbusch, 2001), and coping (Bjorck, Cuthbertson, Thuman & Lee, 2001; Cole, Bruschi & Tamang, 2002; Hwang, Scherer, Wu, Hwang & Li, 2002; Morling, Kitayama & Miyamoto, 2003; Taylor, Sherman, Kim, Jarcho & Takagi, 2004; Tweed, White & Lehman et al., 2004; VanderVoort, 2001; Yeh & Inose, 2002). According to the most recent empirical findings, however, such cultural differences in ER are mostly accounted for by individual differences in personality traits. What in the past appeared to be “cultural” may in fact have been group differences on personality traits among different cultures (Matsumoto, 2006). Neuroticism and Extraversion, in this sense, are the two most relevant personality traits. Individuals with higher scores on Extraversion are more likely to use cognitive reappraisal and less likely to use expressive suppression, while individuals with higher scores on Neuroticism are more prone to lack ER abilities (Matsumoto, 2006).

On the basis of these considerations, though little is known about ER in Italians, some hypotheses regarding the DERS scores within an Italian sample could be formulated by considering existing cross-cultural studies on personality traits. For example, given that Italians score higher than Americans on the Revised NEO Personality Inventory's (NEO-PI-R; Costa & McCrae, 1992; see also Caprara, Barbaranelli, Hahn & Comrey, 2001) Neuroticism factor (McCrae, 2001), and that Neuroticism is associated with a lack of ER abilities, one may speculate that the mean values of the DERS in an Italian sample may be higher than those originally observed by Gratz and Roemer (2004). This would also be in

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 agreement with the notion reported by Terracciano et al. (2003), according to which Italians  
4  
5 may be characterized by “affective volatility”.  
6

7           Factorial structure, internal consistency, and concurrent validity should not be affected  
8  
9 by these potential mean differences. These psychometric properties are calculated through  
10  
11 statistical analyses that are based on correlations, and differences in the mean values do not  
12  
13 imply any changes in the correlation values themselves. In fact, other instruments that  
14  
15 measure constructs related to ER have been adapted for Italian samples beforehand, and high  
16  
17 congruence between the American and Italian versions was observed in terms of factorial  
18  
19 structure, internal consistency, and concurrent validity. For instance, Bressi et al. (1996)  
20  
21 worked on the Italian cross-validation of a measure of alexithymia, the 20-item Toronto  
22  
23 Alexithymia Scale (TAS-20; Bagby, Parker, & Taylor, 1994a, 1994b). A confirmatory factor  
24  
25 analysis confirmed the factor structure of the original instrument, and both the internal  
26  
27 consistency and the test-retest reliability resulted adequate. Similarly, Balzarotti, John, and  
28  
29 Gross (2010) adapted the Emotion Regulation Questionnaire (ERQ; Gross & John, 2003), a  
30  
31 measure of reappraisal and suppression. The two-scale ERQ structure was confirmed via  
32  
33 confirmatory factor analysis, and good indices of internal consistency and concurrent validity  
34  
35 were obtained. Also, Terracciano et al. (2003) evaluated an Italian version of the Positive and  
36  
37 Negative Affect Schedule (PANAS; Watson, Clark, & Tellegen, 1988), an instrument to  
38  
39 assess Positive and Negative Activation. Again, high congruence between the American and  
40  
41 Italian versions was obtained for factorial structure and construct validity.  
42  
43  
44  
45  
46

47           Although mean value differences may become apparent, the validity and reliability of  
48  
49 the DERS should carry over to Italy. Hence, aiming to investigate factorial structure, internal  
50  
51 consistency, and concurrent validity of an Italian version of the DERS, we hypothesized that:  
52  
53 (a) the factorial structure observed by Gratz and Roemer (2004) would be confirmed in our  
54  
55 Italian sample; (b) the internal consistency would be adequate and comparable to the previous  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 findings on the same subject; (c) the concurrent validity would support the construct validity  
4  
5 of the instrument.

**Method**

6  
7  
8  
9 The original version of the DERS was translated into Italian and administered, along  
10  
11 with some other questionnaires, to a sample of Italian students.

12  
13  
14 In order to produce equivalent versions of a measure across different languages and  
15  
16 cultures, several translation methods are used. One of the most common approaches is the  
17  
18 “back translation” method (Brislin, 1980; Geisinger, 2003; Van de Vijver & Hambleton,  
19  
20 1996) in which the measure is initially translated from a source language (e.g., English) into a  
21  
22 target language (e.g., Italian) by a bilingual individual and then back-translated into the source  
23  
24 language (e.g., English) by a second bilingual individual in order to address possible  
25  
26 inconsistencies. Another approach is referred to as the “double translation/double back-  
27  
28 translation” method, which is quite similar to the “back-translation” method though it requires  
29  
30 a greater number of independent translators (Kristjansson, Desrochers & Zumbo, 2003).  
31  
32 Conversely, “forward-translation” involves a group of bilingual individuals directly focusing  
33  
34 on both the source and the target language versions of a measure (Hambleton & Li, 2005).  
35  
36 Finally, a variation of the “back translation” and “forward translation” approaches is the  
37  
38 “review team” in which a team of individuals evaluates the translation at various steps of the  
39  
40 translation process. Regardless of the specific approach, the primary goal of each method is to  
41  
42 maximize the meaningful equivalence of the measure rather than to obtain a literal translation  
43  
44 (Brislin, 1970; Hambleton & Li, 2005; Kristjansson et al., 2003).  
45  
46  
47

48  
49 Given that the “back translation” procedure is a widely-used, practical translation  
50  
51 method, it was used for our study on the DERS. First, a bilingual researcher in Dynamic  
52  
53 Psychology at the University of Milano-Bicocca translated the DERS from English into  
54  
55 Italian. Then another bilingual individual (a professional translator) blind to the original  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

questionnaire re-translated this version back to the original language. Finally, the two English versions of the DERS were compared to each other to review and resolve any arising inconsistencies.

The congruence between the two English versions was evaluated by a bilingual researcher in Dynamic Psychology at the University of Milano-Bicocca and by two Italian collaborators that are fluent in English and have lived in the United States for more than one year for research. Except for a few slight discrepancies (e.g., the back translation for “I believe that I will remain that way for a long time” resulted in “I think I will stay that way for a long time”), most of the items did not have any significant inconsistencies. After the first evaluation, another bilingual individual – blind both to the original version and to the first back translation – was asked to provide a new back translation for any inconsistent items (e.g., the Italian word for “work” sometimes is translated into English as “job”, and the back translation for “getting work done” resulted in “to finish a job”). This final step confirmed that our Italian version of the DERS was consistent with the original English version.

**Participants**

The initial sample comprised of 351 Italian psychology students at the University of Milano-Bicocca. In order to be consistent with the first developmental study of the DERS (Gratz & Roemer, 2004), subjects with missing data on one or more item of the DERS were removed. The final sample consisted of 323 participants, ranging from 18 to 64 years of age with a mean age of 25.6 ( $SD = 8.6$ ). Seventy-seven percent ( $n = 249$ ) of the sample was female.

**Procedure**

During the 2007-2008 academic year, a large amount of students attending several different psychology classes at the University of Milano-Bicocca were informed about the opportunity to volunteer for this study. Eligibility criteria required that participants were

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 Italian<sup>1</sup>, that Italian was their first language, and that they were not currently receiving  
4  
5 psychiatric therapy or taking any psychiatric medications. Each individual filled out a  
6  
7 questionnaire with information on inclusion criteria prior to participant selection for the study.  
8  
9 Extra credit points were given to students to compensate for their participation in the study.  
10

11  
12 Students who decided to volunteer were required to give written consent for  
13  
14 participation and anonymously complete several questionnaires in a classroom setting. Seven  
15  
16 appointments were planned over the academic year with an average of 50 participants per  
17  
18 room. The average completion time was 60-90 minutes for all questionnaires, though the time  
19  
20 limit was 120 minutes.  
21

**Measures**

22  
23  
24  
25 When we began this study, no instruments measuring difficulties in ER were available  
26  
27 within the Italian context. Therefore, we decided (a) to administer the Toronto Alexithymia  
28  
29 Scale (TAS-20; Bagby, Parker & Taylor, 1994a, 1994b) which has been adapted for an Italian  
30  
31 population by Bressi et al., (1996), and (b) to adapt other questionnaires that had not yet been  
32  
33 validated for the Italian population. The original versions of all questionnaires utilized in this  
34  
35 study were the objects of previously published studies, those submitted for review, or those in  
36  
37 preparation for review. Each participant completed the following self-report questionnaires:  
38  
39

40  
41 **Difficulties in Emotion Regulation Scale (DERS; Gratz & Roemer, 2004).** The  
42  
43 DERS is a 36 item self-report measure developed to assess clinically-relevant difficulties in  
44  
45 emotion regulation. Items are scored on six scales: Nonacceptance of Emotional Responses  
46  
47 (NONACCEPTANCE, 6 items); Difficulties Engaging in Goal-Directed Behavior (GOALS, 5  
48  
49 items); Impulse Control Difficulties (IMPULSE, 6 items); Lack of Emotional Awareness  
50  
51 (AWARENESS, 6 items); Limited Access to Emotion Regulation Strategies (STRATEGIES,  
52  
53 8 items); and Lack of Emotional Clarity (CLARITY, 5 items). Participants are asked to  
54  
55

---

56  
57 <sup>1</sup> According to the Italian Census Bureau (ISTAT, [www.istat.it](http://www.istat.it)), the current ethnic make-up of the Italian  
58  
59 population consists primarily of Italians (more than 90%), with the remaining portion including Europeans  
60  
(around 4%, mostly Romanians and Albanians), North Africans (around 1%), and others.

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 indicate how often each of the 36 items applied to them on a 5-point Likert scale ranging from  
4  
5 1 (*almost never*) to 5 (*almost always*). Subscale scores are obtained by summing  
6  
7 corresponding items. Gratz and Roemer (2004) reported an internal consistency of .93, a test-  
8  
9 retest reliability of .88 during a 4- to 8-week interval, and a clear factor structure. The DERS  
10  
11 predicts significant correlations with several criterion variables, including experiential  
12  
13 avoidance and self-harm. Other empirical findings confirmed good construct validity and a  
14  
15 high internal consistency with both clinical and non-clinical populations (Fox et al., 2007;  
16  
17 Gratz et al., 2008; Johnson et al., 2008). The DERS has demonstrated sensitivity to changes  
18  
19 due to successful therapeutic outcome (Gratz & Gunderson, 2006; Gratz, Lacroce, &  
20  
21 Gunderson, 2006) and convergence with a behavioral measure of emotion dysregulation  
22  
23 (Gratz, et al., 2006). Higher scores on the DERS indicate greater difficulties in emotion  
24  
25 regulation.  
26  
27  
28

**Toronto Alexithymia Scale (TAS-20; Bagby, Parker & Taylor 1994a, 1994b).**

29  
30 Alexithymia is a multifaceted personality construct that represents a deficit in the cognitive  
31  
32 processing of emotion. Currently, the TAS-20 is the most common measure of alexithymia.  
33  
34 Participants are asked to rate 20 items on 5-point Likert scale ranging from 1 (*strongly*  
35  
36 *disagree*) to 5 (*strongly agree*). The TAS-20 yields a total score and three subscale scores  
37  
38 (i.e., Difficulty Identifying Feelings, Difficulty Describing Feelings, and Externally Oriented  
39  
40 Thinking). In this study, we used the Italian version of the TAS-20 (Bressi et al., 1996) which  
41  
42 showed good internal consistency (Cronbach's  $\alpha$  of .75 and .82 in normal and clinical groups,  
43  
44 respectively) and high test-retest reliability over 2 weeks ( $r = .86$ ).  
45  
46  
47  
48

49  
50 Given that higher scores on the TAS-20 indicate greater difficulties in the cognitive  
51  
52 processing of emotion, positive correlations between the TAS-20 and the DERS were  
53  
54 predicted.  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3           **Trait Meta-Mood Scale (TMMS; Salovey, Mayer, Goldman, Turvey, & Palfai,**  
4  
5           **1995).** The TMMS is a measure of emotional awareness and understanding that includes  
6  
7 ability to regulate, clarify, and attend to feelings. In the original validation study, participants  
8  
9 were asked to rate 48 items on a 5-point Likert scale ranging from 1 (*strongly disagree*) to 5  
10  
11 (*strongly agree*). According to the results of a factor analysis, Salovey et al., (1995) removed  
12  
13 18 items and offered directions for calculating a total score and three subscale scores (i.e.,  
14  
15 Attention to Feelings, Clarity of Experience of Feelings, and Repair of Emotions). This final  
16  
17 30-item version of the TMMS has shown adequate to good internal consistency, and high  
18  
19 scores are associated with less depression and more life satisfaction (Martinez- Pons, 1997).  
20  
21 Because no validated Italian versions of the TMMS were available, we adapted it using a  
22  
23 back-translation procedure. No noteworthy inconsistencies between the original and the final  
24  
25 back-translated version emerged. To examine the factorial structure, reliability, and validity of  
26  
27 our Italian version of the TMMS, all 48 items were administered. The total score and three  
28  
29 subscale scores were calculated using only the 30 items indicated by Salovey et al. (1995).  
30  
31  
32  
33

34           Given that the TMMS measures emotional awareness and understanding that includes  
35  
36 ability to regulate, clarify, and attend to feelings, negative correlations between TMMS and  
37  
38 DERS scores were predicted.  
39

40           **Scale of Dissociative Activities (SODAS; Mayer & Farmer, 2003).** The SODAS is a  
41  
42 relatively new measure which includes acting without awareness, a lack of perception of the  
43  
44 inner experience, memory disruptions, and perceptions of unreality. Participants are asked to  
45  
46 rate 35 items on a 5-point Likert scale ranging from 1 (*never*) to 5 (*very frequently*), and a  
47  
48 total score is calculated by summing items. Mayer and Farmer (2003) reported that the  
49  
50 SODAS has good internal consistency ( $\alpha = .95$ ) and test-retest stability during a 38-day  
51  
52 interval ( $r = .77$ ). Furthermore, the SODAS showed significant positive correlations with  
53  
54 other measures of dissociation, including experience sampling measures in naturalistic  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

environments. Because no validated Italian versions of the SODAS were available, we adapted it using a back-translation procedure. No noteworthy inconsistencies between the original and the final back-translated version emerged.

Given that dissociation and a lack of perception of inner experience are related to emotion dysregulation, we predicted positive correlations between the SODAS and the DERS.

### Results

Factorial structure, internal consistency, and concurrent validity of our Italian version of the DERS were analyzed.

#### DERS Scores

Descriptive statistics for males and females on the total DERS score as well as each subscale are shown in Table 1. No significant gender differences were found, and the DERS scores produced by our Italian sample were largely similar to those reported by Gratz and Roemer (2004). Two significant differences between our Italian sample and Gratz and Roemer (2004)'s original American sample were observed: Italian women scored significantly higher ( $M = 12.2$ ;  $SD = 4.5$ ) than American women ( $M = 10.8$ ;  $SD = 4.4$ ) on IMPULSE,  $t(507) = 3.49$ ,  $p < .005$ ,  $d = .31$ , and Italian men scored significantly lower ( $M = 14.4$ ;  $SD = 3.6$ ) than American men ( $M = 16.3$ ;  $SD = 4.6$ ) on AWARENESS,  $t(169) = 2.96$ ,  $p < .005$ ,  $d = .44$ .

Most of the scores presented low skew and kurtosis values. However, within the entire sample IMPULSE exhibited a slightly positive skew (skew = 1.2).

Correlations among the DERS subscales are shown in Table 2. As expected, the six subscales were correlated with one another. However, similar to Gratz and Roemer's findings (Gratz & Roemer, 2004), GOALS did not correlate with AWARENESS.

#### Factorial Structure



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

To test whether the factor structure of the DERS found for U.S. students was replicated in our Italian sample, we conducted a confirmatory factor analysis (CFA).

According to Curran, West, and Finch (1996) as well as several other researchers (e.g., Muthen & Kaplan, 1985, 1992), nonnormality concerns arise in CFA with a skewness of 2.0 and kurtosis of 7.0. Given that no skew or kurtosis values exceeded these thresholds, robust CFA methods were not implemented. As for the structural model, a correlation matrix was used. Six latent variables were specified, corresponding to the six subscales of the DERS, which were allowed to correlate. Lisrel 8.50 (Jöreskog & Sörbom, 2001) was used for the analysis.

We focused on four fit indices: the comparative fit index (CFI), the nonnormed fit index (NNFI), the root mean square error of approximation (RMSEA), and the ratio of the value of chi-square to its degrees of freedom ( $\chi^2 / df$ ). CFI and NNFI values greater than .90 are thought to indicate *good* fit between a model and the data (Bentler & Bonett, 1980); RMSEA values of .05 are thought to indicate *close* fit, .08 *fair* fit, and .10 *marginal* fit (Browne & Cudeck, 1993);  $\chi^2 / df$  values close to 2.00 or less are considered *good* while values under 5.00 are *quite promising* (Watkins, 1989).

According to these thresholds, the CFA performed on the Italian version of the DERS replicated quite well the factorial structure obtained by Gratz and Roemer (2004). The CFI and NNFI were not far from the threshold value of .90 (CFI = .86, NNFI = .84), the RMSEA was *fair* (RMSEA = .072; 90% confidence interval = .068 – .076), and the  $\chi^2 / df$  was *quite promising to good* ( $\chi^2 = 1547.0$ ;  $df = 579$ ;  $\chi^2 / df = 2.67$ ). Furthermore, all factor loadings were greater than .40, except for item 23 which yielded a factor loading of .37 (Table 3). Item 23 was kept in as its factor loading was only .03 points lower than a commonly-used threshold (.40) and its item-total and item-subscale correlations were relatively high (respectively,  $r = .46$ ,  $p < .001$  and  $r = .54$ ,  $p < .001$ ).

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

**Internal Consistency**

To determine the internal consistency of the DERS items, Cronbach's  $\alpha$  was calculated for the total DERS score and for each of the six subscales. Similar to Gratz and Roemer's findings (2004), the results indicated that the DERS has high internal consistency with a Cronbach's  $\alpha$  of .92 for the total DERS score and  $\alpha > .80$  for five of the six subscales. The sixth subscale yielded a .77 for AWARENESS (see Table 4 for additional information on the internal consistency of each subscale). Also similar to Gratz and Roemer's findings (2004), item-total correlations ranged from  $r = .15$  to  $r = .70$ , and thirty-four of the items had item-total correlations above  $r = .30$ .

**Concurrent Validity**

To provide data on the concurrent validity of the DERS, correlations between the DERS and other questionnaires were examined. All Cronbach's  $\alpha$  values for the compared questionnaires were higher than .70 except for the Externally Oriented Thinking scale of the TAS, for which it was .68.

Correlations between the DERS and other constructs are shown in Table 5. As predicted, the total DERS score yielded high positive correlations with the total TAS-20 score ( $r = .56, p < .01$ ) and with the SODAS ( $r = .62, p < .01$ ), and yielded high negative correlations with the total TMMS score ( $r = -.54, p < .01$ ).

**Study 2**

In order to assess the test-retest reliability of the DERS, a second sample was examined. As in study 1, we determined that while cultural differences in ER between Italy and North America may affect the mean values of the DERS, they should not affect its validity and reliability. Thus, we expected our Italian version of the DERS to yield adequate test-retest reliability.

**Method**

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

**Participants**

Participants were 61 Italian psychology students at the University of Rome – Sapienza who met the same eligibility criteria described in Study 1. Ages ranged from 20 to 60, with a mean age of 24.7 ( $SD = 5.8$ ). Eighty percent ( $n = 53$ ) were female.

**Procedure**

Before a psychology class, students were invited to complete a questionnaire. Upon giving their written consent, all class members decided to participate and complete our version of the DERS. After 4 weeks, the DERS was administered again in the same classroom setting before a subsequent lesson of the same psychology class. Test-retest reliability was investigated on the 61 students that were present at both of the administrations.

**Results**

The intraclass correlation coefficient (ICC) for the total DERS score was .78. The ICCs for the DERS' subscales ranged from .49 to .73. More specifically, the ICCs for the DERS' subscales were: .73 for NONACCEPTANCE; .72 for GOALS; .65 for IMPULSE; .68 for AWARENESS; .73 for STRATEGIES; and .49 for CLARITY. According to the suggested benchmarks (Cicchetti, 1994; Cicchetti & Sparrow, 1981; Fleiss, 1981), the test-retest reliability was *excellent* for the total DERS score, *fair* for CLARITY, and *good* for NONACCEPTANCE, GOALS, IMPULSE, AWARENESS, and STRATEGIES.

**Study 3**

To further investigate the construct validity of our version of the DERS, we compared the DERS scores of a nonclinical sample with a clinical sample characterized by deficits in ER. Thus, we administered the DERS to both a nonclinical sample and to a small clinical sample affected by anorexia, binge eating disorder, and bulimia.

Emotional problems and deficits in ER are thought to be a core feature of eating disorders (EDs): as compared to controls, individuals with EDs have more problems in ER

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 and tend to have lower emotional awareness (Gilboa-Schechtman, Avnon, Zubery, &  
4 Jeczmiem, 2006), demonstrate higher levels of alexithymia (see for example, Corcos et al.,  
5 2000; de Groot, Rodin, & Olmsted, 1995; Speranza, Corcos, Loas, Stèphan, Guilbaud, Perez-  
6 Diaz, et al., 2005; Taylor, Parker, Bagby, & Bourke, 1996), have more difficulty  
7 communicating feelings (Rizzuto, 1988), and experience more difficulties in recognizing  
8 facially-expressed emotions (Kucharska-Pietura, Nikolaou, Masiak, & Treasure, 2004;  
9 Zonnevijlle-Bender, van Goozen, Cohen-Kettenis, van Elburg, & van Engeland, 2002). A  
10 couple of recent studies have linked EDs and ER also using the DERS: Harrison, Sullivan,  
11 Tchanturia, and Treasure (2009) found that women with anorexia have more difficulties with  
12 emotional recognition and regulation than their nonclinical counterparts; Whiteside, Chen,  
13 Neighbors, Hunter, Lo, and Larimer (2007) found that binge eaters exhibit fewer ER skills  
14 than do controls. Moreover, according to several authors, symptoms of EDs may regulate  
15 negative affect (Abraham & Beaumont, 1982; Hayaki, Friedman & Brownell, 2002;  
16 Heatherton, & Baumeister, 1991; Herman & Polivy, 1988; Match & Simons, 2000;  
17 McCarthy, 1990), and ER problems may mediate the relationship between body  
18 dissatisfaction and bulimic symptoms such as binge eating (e.g., Leon, Fulkerson, Perry &  
19 Early-Zald, 1995; Sim, & Zeman, 2005; Stice, Nemeroff, & Shaw, 1996).

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Given the association between EDs and emotion dysregulation, we expected that the DERS scores of our clinical sample would exceed those of a nonclinical sample matched by age and gender.

### Method

#### Participants

The clinical sample was composed of 38 Italian women affected by either bulimia nervosa ( $N = 22$ ), anorexia ( $N = 10$ ) or binge eating disorder ( $N = 6$ ). All diagnoses were made

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 by expert clinicians that have been in practice for many years. Their ages ranged from 16 to  
4  
5 54 years ( $M = 24.2$ ,  $SD = 8.7$ ) and they had an average to high level of education.

6  
7 The nonclinical sample was composed of 38 Italian women screened to ensure the  
8  
9 absence of potential eating or other disorders that feature emotion dysregulation. Their ages  
10  
11 ranged from 16 to 48 years ( $M = 24.7$ ,  $SD = 8.1$ ) and they had an average to high level of  
12  
13 education.  
14

**Procedure**

15  
16  
17 The participants in the clinical sample were recruited in either a public hospital ( $N =$   
18  
19 20) or private clinic ( $N = 18$ ). Before beginning the standard screening procedure, all patients  
20  
21 were informed that they could also complete an additional questionnaire for research purposes  
22  
23 in addition to the routine test battery. They were handed the DERS along with other  
24  
25 questionnaires and documents (including a demographic form), with the instruction that if  
26  
27 they wanted to participate in the research project they had to complete both the DERS and the  
28  
29 informed consent. No one, among the patients contacted, refused to participate.  
30  
31  
32

33  
34 The participants in the nonclinical sample were recruited using a snowball strategy:  
35  
36 Initial participants asked other people (relatives, friends, and acquaintances) to take part in a  
37  
38 psychological study by completing questionnaires at home and possibly recruiting further  
39  
40 participants. All volunteer participants were directly contacted by the authors, who  
41  
42 administered them the assigned questionnaires and a demographic form. Any prospective  
43  
44 participant that reported receiving psychiatric medications or was at risk for an eating  
45  
46 disorders or other psychopathology (see below), his or her data was excluded from the  
47  
48 analysis and other participants were recruited to replace them. This led to the exclusion of five  
49  
50 persons before achieving the desired sample size.  
51  
52

53  
54 To ensure that participants in the two groups were matched on age between-samples,  
55  
56 four age ranges were included: 15-24, 25-34, 35-44, and 45-54 years of age. For every woman  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 in the clinical group in one of these age ranges, there was one woman in the same age range in  
4  
5 the nonclinical group.  
6

**Measures**

7  
8  
9 In an effort to ensure the absence of an eating or other disorder that featured emotion  
10 dysregulation within the nonclinical sample, the nonclinical volunteers were administered two  
11  
12 other questionnaires in addition to the DERS: the Eating Disorder Inventory (EDI-3; Garner,  
13  
14 2004) and the Symptom Checklist 90 (SCL-90; Derogatis, 1977).  
15  
16  
17

18  
19 **Eating Disorder Inventory (EDI-3; Garner, 2004).** The EDI-3 is a 91 item self-  
20 report measure developed to assess three eating disorder-specific scales and nine general  
21  
22 psychological scales that are highly relevant (but not specific) to eating disorders. The EDI-3  
23  
24 also includes six composite scores: the Eating Disorder Risk scale, which is eating-disorder  
25  
26 specific; the Ineffectiveness scale; the Interpersonal Problems scale; the Affective Problems  
27  
28 scale; the Overcontrol scale; and the General Psychological Maladjustment scale. Garner  
29  
30 (2004) reported that the Eating Disorder Risk scale's reliability ranges from .90-.97 (median =  
31  
32 .94) across four diagnostic and normative groups. For the three Eating Disorder Risk scales,  
33  
34 all reliability values are generally in the high .80s to low .90s across normative groups, and  
35  
36 the test-retest stability coefficients are excellent. In addition, the EDI-3 produced adequate  
37  
38 convergent and discriminant validity indices (Cumella, 2006) and excellent sensitivity and  
39  
40 specificity indices (Clausen, Rosenvinge, Friborg, & Rokkedal, 2011).  
41  
42  
43  
44

45 For this study, the Italian version of the EDI-3 (Giannini, Pannocchia, Dalle Grave &  
46  
47 Muratori, 2008) was used with the intent to identify for exclusion participants with potential  
48  
49 eating disorders in the nonclinical group. Before achieving the desired sample size, two  
50  
51 potential participants were excluded from the study because their scores on the Eating  
52  
53 Disorder Risk scale were above the 85<sup>th</sup> percentile of Italian normative data.  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3       **Symptom Checklist 90 (SCL-90; Derogatis, 1977).** The SCL-90 is a 90 item self-  
4 report measure that is widely-used in both its original and Italian forms (Magni, Messina, De  
5 Leo, Mosconi & Carli, 1983) to assess a broad range of psychopathology symptoms  
6 (Derogatis, 1994). It includes nine dimensions: somatization; obsessive-compulsive;  
7 interpersonal sensitivity; depression; anxiety; hostility; phobic anxiety; paranoid ideation; and  
8 psychoticism. According to European validations of the instrument (Holi, 2003; Schmitz,  
9 Hartkamp & Franke , 2000), a global total score covering all 90 items – the Global Severity  
10 Index (GSI) – is a valid, reliable, and commonly-used criterion for screening  
11 psychopathology.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23       In this study, the GSI of the Italian version of the SCL-90 (Magni et al., 1983) was  
24 used to exclude potential psychopathology from the nonclinical group. Prior to achieving the  
25 desired sample size, four prospective participants (one of whom was also above the cut-off on  
26 the EDI-3) were excluded from the study because their GSI scores were equal to or greater  
27 than 1.  
28  
29  
30  
31  
32

### Results

33  
34  
35  
36       As expected, the total DERS score for the clinical sample ( $M = 111.6$ ;  $SD = 24.8$ ) was  
37 significantly higher than the total DERS score for the nonclinical sample ( $M = 73.6$ ;  $SD =$   
38  $16.1$ ),  $t(74) = 7.93$ ,  $p < .005$ ,  $d = 1.82$ . Significant differences were also found for each DERS  
39 subscale (shown in Table 6). All comparisons remain significant even after Bonferroni's  
40 correction, and the 95% confidence intervals clearly demonstrate that the two groups produce  
41 distinct mean values on each subscale. Neither age nor education significantly differed  
42 between the clinical and nonclinical groups.  
43  
44  
45  
46  
47  
48  
49  
50

### Discussion

51  
52       The importance of emotion regulation (ER) in human wellbeing is acknowledged by  
53 several authors and Gratz and Roemer (2004) have recently proposed the DERS, a  
54  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 multidimensional self-report measure of difficulties in ER. To supplement the literature on the  
4  
5 cross-cultural validity of this scale, we adapted the DERS for an Italian sample and evaluated  
6  
7 its reliability and validity. Three studies were conducted to complete this task.  
8

9  
10 In study 1, we translated the DERS into Italian and investigated its factorial structure,  
11  
12 internal consistency, and concurrent validity. According to our hypotheses, we expected the  
13  
14 factorial structure obtained by Gratz and Roemer (2004) to be confirmed and show adequate  
15  
16 internal consistency and concurrent validity.  
17

18  
19 The Italian data replicated quite well the factorial structure obtained by Gratz and  
20  
21 Roemer (2004), thus confirming that ER is a multidimensional construct. As in Gratz and  
22  
23 Roemer's original findings (Gratz & Roemer, 2004), GOALS did not correlate with  
24  
25 AWARENESS. This may indicate that there is "a difference between being aware of  
26  
27 emotional responses and having a clear understanding of the nature of these responses" (p.  
28  
29 47). All internal consistency alpha indices were adequate and comparable with those reported  
30  
31 by Gratz and Roemer (2004) and by other validation studies (Coutinho, et al., 2010; Ruganci  
32  
33 & Gençöz, 2010). The total DERS score was positively correlated with the total TAS-20  
34  
35 score, the total SODAS score, and negatively correlated with the total TMMS score, thus  
36  
37 supporting the construct validity of the DERS. In addition, each of the subscales of the DERS  
38  
39 correlated, both significantly and in the expected direction, with at least one of the other  
40  
41 subscales that measured similar constructs. For example, the subscales obtaining the highest  
42  
43 correlation values with the DERS subscale CLARITY were the TMMS subscale Clarity of  
44  
45 Experience of Feelings and the TAS-20 subscale Difficulty Identifying Feelings. Examined  
46  
47 together, these results confirm our hypotheses and support the reliability and validity of the  
48  
49 DERS.  
50  
51  
52

53  
54 A couple of discrepancies between the results of this study and Gratz and Roemer's  
55  
56 findings (Gratz & Roemer, 2004) warrant mentioning. First, while Gratz and Roemer (2004)  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 report that in their sample men scored higher than women on AWARENESS, no gender  
4  
5 differences were observed in our Italian sample. Second, women in our sample scored  
6  
7 significantly higher than women in Gratz and Roemer's sample (2004) on IMPULSE, while  
8  
9 men in our sample scored significantly lower than men in Gratz and Roemer's sample (2004)  
10  
11 on AWARENESS. Given that the subscale AWARENESS is the source of score differences  
12  
13 between Italian and American men and between American men and women, it is a possibility  
14  
15 that AWARENESS is more affected by culture and personality traits than are the other five  
16  
17 subscales. As this is pure conjecture at this point (these effects may also be due to translation  
18  
19 inequities, different response styles, or sample characteristics), more research on this topic is  
20  
21 needed.  
22  
23

24  
25 Another interesting finding of study 1 is that the total DERS score of Italians was not  
26  
27 significantly different from the total DERS score of Americans. From this point of view, the  
28  
29 hypothesis that Italians may be characterized by "affective volatility" remains unsupported.  
30  
31

32 In study 2, we investigated the test-retest reliability of the DERS. According to our  
33  
34 hypotheses, we expected the ICCs between test and retest to be adequate and comparable to  
35  
36 previous findings on this subject (Gratz & Roemer, 2004; Ruganci & Gençöz, 2010). The  
37  
38 results confirmed our hypotheses, showing that the test-retest reliability of the Italian version  
39  
40 of the DERS is excellent for the total score and fair to good for the subscale scores.  
41  
42

43 In study 3, a construct validity analysis comparing clinical and nonclinical samples  
44  
45 was conducted. We predicted the DERS scores (both total and subscale) of a clinical sample  
46  
47 characterized by emotion dysregulation (i.e., a sample affected by EDs) would be significantly  
48  
49 higher than those produced by a nonclinical, age-matched sample. As shown on table 6, the  
50  
51 results confirm our hypotheses: individuals affected by EDs scored significantly higher on our  
52  
53 Italian version of the DERS than did healthy controls. All effect size indices qualified as *large*  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 effect sizes, which further supported the construct validity of the DERS and the association  
4  
5 between EDs and deficits in ER.  
6

7  
8 Taken together, the results of these three studies indicate that the DERS is valid and  
9  
10 reliable in an Italian context, thus supporting the cross-cultural adaptability of the DERS. The  
11  
12 factorial structure replicated the six-factor structure proposed by Gratz and Roemer (2004),  
13  
14 the internal consistency and test-retest reliability were adequate and comparable to previous  
15  
16 findings, and the validity was good as indicated by both the concurrent validity analysis and  
17  
18 the comparison between a clinical and a nonclinical sample.  
19

20  
21 Several limitations warrant mentioning, however. First, the clinical sample we used in  
22  
23 study 3 was small ( $N = 38$ ) and only focused on one specific clinical population, chosen in  
24  
25 part because of the association between EDs and emotion dysregulation and in part because of  
26  
27 our personal interest in EDs. Results, therefore, may not be applicable to other clinical  
28  
29 diagnostic categories associated with emotion dysregulation. Future research with larger  
30  
31 sample sizes and including other clinical populations is necessary. Second, the participants in  
32  
33 the clinical and control groups of study 3 were taking the DERS under unique circumstances  
34  
35 (i.e., part of an admission procedure for a treatment program vs. volunteer participation upon  
36  
37 invitation of family or friends, respectively). Coupled with a possible selection bias due to the  
38  
39 sampling procedure utilized for the nonclinical sample, this may have had a noteworthy effect  
40  
41 on the results. Third, two of the three scales administered in study 1 to measure DERS-related  
42  
43 constructs have not yet been cross-validated for Italian use. It should be noted, however, that:  
44  
45 (a) these two instruments were translated with a back-translation procedure and obtained high  
46  
47 internal consistency indices and (b) there is a lack of cross-validated instruments for Italian  
48  
49 use and the fact that appropriate alternatives were not available is one of the reasons for this  
50  
51 study. Fourth, divergent validity has not been addressed by this study. Fifth, given that in  
52  
53 study 1 the participants answered a great deal of items concerning relatively similar concepts,  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 potential respondent fatigue may have occurred. Despite the generous time limit given to  
4  
5 participants completing questionnaires, several missing data were observed. Although the  
6  
7 presentation order of the questionnaires was randomized and the participants should have  
8  
9 been sufficiently motivated by the class credit they received, fatigue may have affected the  
10  
11 results of study 1 on some level. Sixth, the factorial structure and test-retest analyses of study  
12  
13 1 and 2 only refer to student samples. Further examinations should consider analysis of  
14  
15 clinical data as well as replication of our results.  
16  
17

18  
19 Despite all these limitations, however, our investigation provides unique contributions  
20  
21 to ER research for several reasons. First, all six components of ER proposed by Gratz and  
22  
23 Roemer (2004) were valid and reliable. This provides further support for Gratz and Roemer  
24  
25 (2004)'s model of ER, and suggests that each component of the model is important in defining  
26  
27 and measuring ER. Second, the congruence between our results and other international  
28  
29 findings provides support for cross-cultural utilization of the DERS and facilitates Italian  
30  
31 research on ER. Third, the DERS scores of a clinical sample characterized by difficulties in  
32  
33 ER were higher than those of a nonclinical sample, which provides further support for the  
34  
35 utilization of the DERS in a clinical context, especially for ER assessment and treatment  
36  
37 planning. Due to its multidimensional approach, the DERS allows the clinician to capture  
38  
39 patient functioning in a number of ER content areas, assess his or her strengths and  
40  
41 weaknesses, and identify relevant targets of therapy. For example, in cases of limited  
42  
43 emotional awareness (i.e., high scores on AWARENESS) one might adopt exercises such as  
44  
45 the analysis of emotions and cognitions in everyday emotion-inducing situations (Legenbauer,  
46  
47 Vocks, & Ruddle, 2008). In cases of nonacceptance of emotional responses (i.e., high scores  
48  
49 on NONACCEPTANCE), emphasizing a nonjudgmental acceptance of thoughts and feelings  
50  
51 during the therapeutic process may be recommended (Hayes, et al., 1999; Hayes, Wilson,  
52  
53 Gifford, Follette, & Strosahl, 1996). In all cases of deficits in ER (i.e., high scores on the total  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

1  
2  
3 score), improving clients' capacity to tolerate and regulate emotions should be a critical  
4  
5 therapeutic goal (Gutwill & Gitter, 1994; Kearney-Cooke & Striegel Moore, 1994). All these  
6  
7 suggestions are applicable to individuals with EDs, in which the desire to escape from self-  
8  
9 awareness is particularly relevant (Heatherton & Baumeister, 1991; de Groot and Rodin,  
10  
11 1998). Evidence supports the clinical value of incorporating emotion regulation training in  
12  
13 the treatment of EDs (Telch, Agras, & Linehan, 2001).  
14  
15

16  
17 On the basis of all these considerations, our findings provide further evidence  
18  
19 supporting the utility, validity, and reliability of the DERS in Italy in both research and  
20  
21 clinical settings.  
22

**Acknowledgments**

23  
24 We thank Gherardo Amadei, Margherita Lang, Laura Parolin, Angela Tagini, Laura  
25  
26 Rivolta, Caterina Giovannini, and Paola De Cesare for their help with working on the  
27  
28 translation of the questionnaires and collecting the data. We also heartily thank Harpreet  
29  
30 Malla of the CLASS center at Alliant International University – San Diego for proofreading  
31  
32 the manuscript.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

## References

- 1  
2  
3  
4  
5 Abraham, S. F., & Beaumont, P. J. V. (1982). How patients describe bulimia or binge eating.  
6  
7 *Psychological Medicine, 12*, 625–635.  
8
- 9  
10 Bagby, R. M., Parker, J. D. A. & Taylor, G. J. (1994a). The twenty-item Toronto Alexithymia  
11  
12 Scale: I. Item selection and cross-validation of the factor structure. *Journal of*  
13  
14 *Psychosomatic Research, 38*, 23-32.  
15
- 16  
17 Bagby, R. M., Parker, J. D. A. & Taylor, G. J. (1994b). The twenty-item Toronto Alexithymia  
18  
19 Scale: II. Convergent, discriminant, and concurrent validity. *Journal of Psychosomatic*  
20  
21 *Research, 38*, 33-40.  
22
- 23  
24 Balzarotti, S., John, O.P. & Gross, J. (2010). An Italian Adaptation of the Emotion Regulation  
25  
26 Questionnaire. *European Journal of Psychological Assessment, 26*, 61-67.  
27
- 28  
29 Bentler, P.M., & Bonett, D.G. (1980). Significance tests and goodness-of-fit in the analysis of  
30  
31 covariance structures. *Psychological Bulletin, 88*, 588-600.  
32
- 33  
34 Bjorck, J. P., Cuthbertson, W., Thurman, J. W., & Lee, Y. S. (2001). Ethnicity, coping, and  
35  
36 distress among Korean Americans, Filipino Americans, and Caucasian Americans.  
37  
38 *Journal of Social Psychology, 141*, 421-442.  
39
- 40  
41 Bressi, C., Taylor, G. J., Parker, J. D. A., Bressi, S., Brambilla, V., Aguglia, E., Allegranti, I.,  
42  
43 Bongiorno, A., Giberti, F., Bucca, M., Todarello, O., Callegari, C., Vender, S., Gala,  
44  
45 C. & Invernizzi, G. (1996), Cross validation of the factor structure of the 20-Item  
46  
47 Toronto Alexithymia Scale: an Italian multicenter study. *Journal of Psychosomatic*  
48  
49 *Research, 41*, 551-559.  
50
- 51  
52 Brislin, R.W. (1980). Translation and content analysis of oral and written material. In H.C.  
53  
54 Triandis & J.W. Berry (Eds.), *Handbook of cross-cultural psychology (Vol. 1)* (pp.389-  
55  
56 444). Boston: Allyn & Bacon.  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Browne, M.W., & Cudeck, R. (1993). Alternative ways of assessing model fit. In K. A Bollen  
4 & J. S. Long (Eds.), *Testing Structural Equation Models* (pp. 136–162). Beverly Hills,  
5 CA: Sage.  
6  
7  
8  
9  
10 Caprara, G.V., Barbaranelli, C., Hahn, R. & Comrey, A. (2001). Factor analyses of the NEO-  
11 PI-R Inventory and the Comrey Personality Scales in Italy and the United States.  
12 *Personality and individual differences*, 30 (2), pp. 217-228.  
13  
14  
15  
16 Cicchetti, D. V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and  
17 standardized assessment instruments in psychology. *Psychological Assessment*, 6,  
18 284– 290.  
19  
20  
21  
22  
23 Cicchetti, D. V., & Sparrow, S. S. (1981). Developing criteria for establishing the interrater  
24 reliability of specific items in a given inventory. *American Journal of Mental*  
25 *Deficiency*, 86, 127–137.  
26  
27  
28  
29  
30 Clausen, L., Rosenvinge, J. H., & Friborg, O. (2011). Validating the Eating Disorder  
31 Inventory-3 (EDI-3): A Comparison Between 561 Female Eating Disorders Patients  
32 and 878 Females from the General Population. *Journal of Psychopathology and*  
33 *Behavioral Assessment* 33:101–110.  
34  
35  
36  
37  
38  
39 Cole, P. M., Bruschi, C. J., & Tamang, B. L. (2002). Cultural differences in children's  
40 emotional reactions to difficult situations. *Child Development*, 73, 983-996.  
41  
42  
43 Corcos, M., Guilbaud, O., Speranza, M., Patemiti, S., Loas, G., Stephan, P., et al. (2000).  
44 Alexithymia and depression in eating disorders. *Psychiatry Research*, 93, 263–266.  
45  
46  
47  
48 Costa, P.T., Jr., & McCrae, R.R. (1992). *Revised NEO Personality Inventory (NEOPI- R) and*  
49 *NEO Five-Factor Inventory (NEOFFI) – Professional Manual*. Odessa, FL:  
50 Psychological Assessment Resources, Inc  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Coutinho, J., Ribeiro, E., Ferreirinha, R., & Dias, P. (2010). The Portuguese version of the  
4  
5 Difficulties in Emotion Regulation Scale and its relationship with psychopathological  
6  
7 symptoms. *Revista de Psiquiatria clinica*, 37, 4: 145-151.  
8  
9  
10 Cumella, E.J. (2006). Review of the Eating Disorder Inventory-3. *Journal of Personality*  
11  
12 *Assessment*, 87 (1), 116-117.  
13  
14 Curran, J. P., West, S. G. & Finch, J. F. (1996). The Robustness of Test Statistics to  
15  
16 Nonnormality and Specification Error in Confirmatory Factor Analysis. *Psychological*  
17  
18 *Methods*, 1, 16-29.  
19  
20  
21 de Groot, J., & Rodin, G. (1998). Coming alive: The psychotherapeutic treatment of patients  
22  
23 with eating disorders. *Canadian Journal of Psychiatry*, 43, 359–366.  
24  
25 de Groot, J., Rodin, G., & Olmsted, M. (1995). Alexithymia, depression, and treatment  
26  
27 outcome in bulimia nervosa. *Comprehensive Psychiatry*, 36(1), 53–60.  
28  
29  
30 Derogatis, L. R. (1977). The SCL-90 Manual I: Scoring, administration and procedures for the  
31  
32 SCL-90. Baltimore: Johns Hopkins University School of Medicine, Clinical  
33  
34 Psychometrics Unit.  
35  
36  
37 Fleiss, J. L. (1981). *Statistical methods for rates and proportions (2nd ed.)*. New York: Wiley.  
38  
39  
40 Fox, H.C., Axelrod, S.R., Paliwal, P., Sleeper, J. & Sinha, R. (2007). Difficulties in emotion  
41  
42 regulation and impulse control during cocaine abstinence. *Drug and alcohol*  
43  
44 *dependence*, 89, 298-301.  
45  
46  
47 Fox, H.C., Hong, K.A., & Sinha, R. (2008). Difficulties in emotion regulation and impulse in  
48  
49 recently abstinent alcoholics compared with social drinkers. *Addictive Behaviors*, 33,  
50  
51 388-394.  
52  
53  
54 Garner, D.M. (2004). *Eating Disorder Inventory-3 professional manual*. Odessa, FL:  
55  
56  
57  
58  
59  
60 Psychological Assessment Resources.

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Geisinger, K.F. (2003). Testing and assessment in cross-cultural psychology. In J.R. Graham,  
4  
5 J.A. Naglieri & I.B. Weiner (Eds.), *Handbook of psychology (Vol.10). Assessment*  
6  
7 *Psychology* (pp. 95-118). New Jersey: John Wiley & Sons.  
8  
9  
10 Giannini, M., Pannocchia, L., Dalle Grave, R. & Muratori, F. (2008). Versione Italiana *Eating*  
11  
12 *Disorder Inventory-3*. Edizioni O.S.- Giunti, Firenze  
13  
14 Gilboa-Schechtman, E., Avnon, L., Zubery, E., & Jeczmiern, P. (2006). Emotional processing  
15  
16 in eating disorders: Specific impairment or general distress related deficiency?  
17  
18 *Depression and Anxiety*, 23(6), 331–339.  
19  
20  
21 Glenn, C., & Klonsky, E.D. (2009). Emotion dysregulation as a core feature of borderline  
22  
23 personality disorder. *Journal of Personality Disorders*, 23, 20-28.  
24  
25 Gratz, K. L. (2007). Targeting emotion dysregulation in the treatment of deliberate self-injury.  
26  
27 *Journal of Clinical Psychology*, 63, 1091-1103.  
28  
29  
30 Gratz, K.L., & Gunderson, J.G. (2006). Preliminary data on an acceptance-based emotion  
31  
32 regulation group intervention for deliberate self-harm among women with borderline  
33  
34 personality disorder. *Behavior Therapy*, 37, 25-35.  
35  
36  
37 Gratz, K.L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and  
38  
39 dysregulation: Development, factor structure, and initial validation of the difficulties  
40  
41 in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment*,  
42  
43 26, 41-54.  
44  
45  
46 Gratz, K.L., Bornoalova, M.A., Delany-Brumsey, A., Nick, B., & Lejuez, C.W. (2007). A  
47  
48 laboratory-based study of the relationship between childhood abuse and experiential  
49  
50 avoidance among inner-city substance users: The role of emotional nonacceptance.  
51  
52 *Behavior Therapy*, 38, 3: 256-268.  
53  
54  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Gratz, K.L., Lacroce, D., & Gunderson, J.G. (2006). Measuring changes in symptoms relevant  
4  
5 to borderline personality disorder following short term treatment across partial hospital  
6  
7 and intensive outpatient levels of care. *Journal of Psychiatric Practice, 12*, 153-159.  
8  
9  
10 Gratz, K.L., Rosenthal, M.Z., Tull, M.T., Lejuez, C.W., & Gunderson, J.G. (2006). An  
11  
12 experimental investigation of emotion dysregulation in borderline personality disorder.  
13  
14 *Journal of Abnormal Psychology, 115*, 850–855.  
15  
16 Gratz, K.L., Tull, M.T., Baruch, D.E., Bornovalova, M.A., & Lejuez, C.W. (2008). Factors  
17  
18 associated with co-occurring borderline personality disorder among inner-city  
19  
20 substance users: The roles of childhood maltreatment, negative affect  
21  
22 intensity/reactivity, and emotion dysregulation. *Comprehensive Psychiatry, 49*, 603-  
23  
24 615.  
25  
26  
27 Gross, J. J., & Thompson, R. A. (2007). Emotion regulation: Conceptual foundations. In J. J.  
28  
29 Gross (Ed.), *Handbook of emotion regulation* (pp. 3-24). New York: Guilford Press.  
30  
31  
32 Gross, J.J. (1998). The emerging field of emotion regulation: An integrative review. *Review of*  
33  
34 *General Psychology, 2*, 271-299.  
35  
36  
37 Gross, J.J. (2008). Emotion Regulation. In Lewis, M., Haviland-Jones, J.M. & Feldman  
38  
39 Barrett, L. (Eds.), *Handbook of Emotions*, 3rd ed. (pp. 497-512). New York: Guilford  
40  
41 Press.  
42  
43 Gross, J.J., & John, O.P. (2003). Individual differences in two emotion regulation processes:  
44  
45 Implications for affect, relationships, and well-being. *Journal of Personality and*  
46  
47 *Social Psychology, 85*, 348–362.  
48  
49  
50 Gross, J.J., & Muñoz, R.F. (1995). Emotion regulation and mental health. *Clinical*  
51  
52 *Psychology: Science and Practice, 2*, 151-164.  
53  
54  
55 Gutwill, S., & Gitter, A. (1994). Eating problems and sexual abuse: Treatment considerations.  
56  
57 In C. Bloom, A. Gitter, S. S. Gutwill, L. Kogel, & L. Zaphiropoulous (Eds.), *Eating*  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 *problems: A feminist psychoanalytic perspective* (pp. 205–26). New York: Basic  
4  
5 Books.  
6  
7 Hambleton, R. K. & Li, S. (2005). Translation and Adaptation Issues and Methods for  
8  
9 Educational and Psychological Tests. In Frisby, C. L., Reynolds, C. R.  
10  
11 (Eds.), *Comprehensive handbook of multicultural school psychology* (pp. 881-903).  
12  
13 Hoboken, NJ, US: John Wiley & Sons Inc.  
14  
15  
16 Harrison, A., Sullivan, S., Tchanturia, K. & Treasure, J. (2009). Emotion recognition and  
17  
18 regulation in Anorexia Nervosa. *Clinical Psychology and Psychotherapy*, 16, 348-356.  
19  
20  
21 Hayaki, J., Friedman, M., & Brownell, K. (2002). Shame and severity of bulimic symptoms.  
22  
23 *Eating Behaviours*, 3(1), 73–83.  
24  
25  
26 Hayes, S. C., Bissett, R. T., Zettle, R. D., Rosenfarb, I. S., Cooper, L. D., & Grundt, A. M.  
27  
28 (1999). The impact of acceptance versus control rationales on pain tolerance. *The*  
29  
30 *Psychological Record*, 49, 33-47.  
31  
32  
33 Hayes, S. C., Wilson, K. W., Gifford, E. V., Follette, V. M., & Strosahl, K. (1996).  
34  
35 Experiential avoidance and behavioral disorders: A functional dimensional approach  
36  
37 to diagnosis and treatment. *Journal of Consulting and Clinical Psychology*, 64(6),  
38  
39 1152-1168  
40  
41  
42 Heatherton, T. F., & Baumeister, R. F. (1991). Binge eating as an escape from selfawareness.  
43  
44 *Psychological Bulletin*, 110, 86–108.  
45  
46  
47 Herman, C. P. & Polivy, J.(1988). Psychological factors in the control of appetite. In Winick,  
48  
49 M. (Ed.). *Control of Appetite* (pp.41-51). Oxford, England: John Wiley & Sons.  
50  
51  
52 Hofstede, G. (2001). *Culture's consequences: Comparing values, behaviors, institutions and*  
53  
54 *organizations across nations* (2nd ed.). Beverly Hills, CA: Sage.  
55  
56  
57 Hofstede, G. (1980). *Culture's consequences: International differences in Work-Related*  
58  
59 *Values*.Beverly Hills CA: Sage Publications.  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Hofstede, G., Hofstede, G. J., Minkov, M. (2010). *Cultures and Organizations: Software of the Mind*. (3rd ed.). McGraw-Hill USA.

Holi, M. (2003). *Assessment of psychiatric symptoms using the SCL-90*. (Doctoral dissertation, Department of Psychiatry, Helsinki University, Finland). Retrieved from <http://ethesis.helsinki.fi/julkaisut/>.

Hwang, C.-E., Scherer, R. F., Wu, Y., Hwang, C.-H., & Li, J. (2002). A comparison of coping factors in western and non-western cultures. *Psychological Reports, 90*, 466-476.

John, O.P., & Gross, J.J. (2004). Healthy and unhealthy emotion regulation: Personality processes, individual differences, and lifespan development. *Journal of Personality, 72*, 1301-1334.

Johnson, K.A, Zvolensky, M.J., Marshall, E.C., Gonzalez, Abrams, A., & Vujanovic, A.A. (2008). Linkages between cigarette smoking outcome expectancies and negative emotional vulnerability. *Addictive Behaviours, 33*, 1416-1424.

Jöreskog, K., & Sörbom, D. (2001). *LISREL 8.51 for Windows. Computer Software*. Lincolnwood, IL: Scientific Software International, Inc.

Kearney-Cooke, A., & Striegel Moore, R. H. (1994). Treatment of childhood sexual abuse in anorexia nervosa and bulimia nervosa: A feminist psychodynamic approach. *International Journal of Eating Disorders, 15*, 305–319.

Klonsky, E.D. (2009) The functions of self-injury in young adults who cut themselves: clarifying the evidence for affect-regulation. *Psychiatry Research, 166*, 260-268.

Kristjansson, E.A., Desrochers, A. & Zumbo, B. (2003). Translating and Adapting Measurement Instruments for Cross-Linguistic and Cross-Cultural Research: A Guide for Practitioners. *Canadian Journal of Nursing Research, Vol 35(2)*, pp. 127-142

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Kucharska-Pietura, K., Nikolaou, V., Masiak, M., & Treasure, J. (2004). The recognition of  
4  
5 emotion in the faces and voice of anorexia nervosa. *International Journal of Eating*  
6  
7 *Disorders*, 35, 42–47.  
8
- 9 Leon, G. L., Fulkerson, J. A., Perry, C. L., & Early-Zald, M. B. (1995). Prospective analysis  
10  
11 of personality and behavioral vulnerabilities and gender influences in the later  
12  
13 development of disordered eating. *Journal of Abnormal Psychology*, 104, 140-149.  
14  
15
- 16 Linehan, M.M. (1993). *Cognitive-behavioral treatment of borderline personality disorder*.  
17  
18 New York: Guilford Press.  
19
- 20 Macht, M., & Simons, G. (2000). Emotions and eating in everyday life. *Appetite*, 35, 65–71.  
21  
22
- 23 Magni, G., Messina, C., De Leo, D., Mosconi, A., & Carli, M. (1983). Psychological distress  
24  
25 in parents of children with acute lymphatic leukemia. *Acta Psychiatrica Scandinavica*,  
26  
27 68, 297–300.  
28
- 29 Martinez- Pons, (1997). The relation of emotional intelligence with selected areas of personal  
30  
31 functioning. *Imagination, Cognition, and Personality*, 17, 3-13.  
32  
33
- 34 Matsumoto, D. (2006). Are cultural differences in emotion regulation mediated by personality  
35  
36 traits?. *Journal of Cross-Cultural Psychology*, 37 (4), 421-437.  
37  
38
- 39 Matsumoto, D., & Kupperbusch, C. (2001). Idiocentric and allocentric differences in  
40  
41 emotional expression and experience. *Asian Journal of Social Psychology*, 4, 113-131.  
42
- 43 Matsumoto, D., Kudoh, T., Scherer, K., & Wallbott, H. (1988). Antecedents of and reactions  
44  
45 to emotions in the United States and Japan. *Journal of Cross-Cultural Psychology*, 19,  
46  
47 267-286.  
48
- 49 Mauro, R., Sato, K., & Tucker, J. (1992). The role of appraisal in human emotions: A cross-  
50  
51 cultural study. *Journal of Personality and Social Psychology*, 62, 301-317.  
52  
53
- 54 Mauss, I. B., Bunge, S. A., & Gross, J. J. (2007). Automatic emotion regulation. *Social and*  
55  
56 *Personality Psychology Compass*, 1, 146-167.  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Mayer, J. L., & Farmer, R. F. (2003). The development and psychometric evaluation of a new  
4  
5 measure of dissociative activities. *Journal of Personality Assessment*, *80*, 185-196.  
6  
7  
8 McCarthy, M. (1990). The thin ideal, depression and eating disorders in women. *Behavior*  
9  
10 *Research and Therapy*, *28*, (3), 205-215.  
11  
12 McCrae, R.R. (2001). Trait psychology and culture: Exploring intercultural comparisons.  
13  
14 *Journal of Personality*, *69*, 819-846.  
15  
16 McLaughlin, K.A., Mennin, D.S., & Farach, F.J. (2007). The contributory role of worry in  
17  
18 emotion generation and dysregulation in generalized anxiety disorder. *Behaviour*  
19  
20 *Research and Therapy*, *45*, 1735-1752.  
21  
22  
23 Mennin, D.S., Heimberg, R.G., Turk, C.L., & Fresco, D.M. (2005). Emotion regulation  
24  
25 deficits as a key feature of generalized anxiety disorder: Testing a theoretical model.  
26  
27 *Behaviour Research and Therapy*, *43*, 1281-1310.  
28  
29  
30 Morling, B., Kitayama, S., & Miyamoto, Y. (2003). American and Japanese women use  
31  
32 different coping strategies during normal pregnancy. *Personality and Social*  
33  
34 *Psychology Bulletin*, *29*, 1533-1546.  
35  
36  
37 Muthen, B., & Kaplan, D. (1985). A Comparison of Some Methodologies for the Factor  
38  
39 Analysis of Non-normal Likert Variables. *British Journal of Mathematical and*  
40  
41 *Statistical Psychology*, *38*, 171-189.  
42  
43  
44 Muthen, B., & Kaplan, D. (1992). A Comparison of Some Methodologies for the Factor  
45  
46 Analysis of Non-normal Likert Variables: A Note on the Size of the Model. *British*  
47  
48 *Journal of Mathematical and Statistical Psychology*, *45*, 19-30.  
49  
50  
51 Nelson, K. & Fivush, R. (2004). The emergence of autobiographical memory: A social  
52  
53 cultural developmental theory. *Psychological review*, *111*, 486-511.  
54  
55  
56  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Neumann, A., van Lier, P. A. C., Gratz, K. L., & Koot, H. M. (2010). Multidimensional  
4  
5 assessment of emotion regulation difficulties in adolescents using the Difficulties in  
6  
7 Emotion Regulation Scale. *Assessment, 17, 138-149.*  
8  
9  
10 Rizzuto, A. M. (1988). Transference, language, and affect in the treatment of bulimarexia.  
11  
12 *International Journal of Psychoanalysis, 69, 369-387.*  
13  
14 Roseman, I. J., Dhawan, N., Rettek, S. I., & Naidu, R. K. (1995). Cultural differences and  
15  
16 cross-cultural similarities in appraisals and emotional responses. *Journal of Cross-*  
17  
18 *Cultural Psychology, 26, 23-48.*  
19  
20  
21 Rugancı, R.N., & Gençöz, T. (2010). Psychometric properties of a Turkish version of the  
22  
23 difficulties in emotion regulation scale. *Journal of Clinical Psychology, 66, 4: 442-*  
24  
25 *455.*  
26  
27  
28 Salovey, P., Mayer, J.D., Goldman, S.L., Turvey, C., & Palfai, T.P. (1995). Emotional  
29  
30 attention, clarity, and repair: Exploring emotional intelligence using the Trait Meta-  
31  
32 Mood Scale. In J.W. Pennebaker (Ed.), *Emotion, disclosure, and health* (pp. 125-154).  
33  
34 Washington, DC: American Psychological Association.  
35  
36  
37 Salters-Pedneault, K., Roemer, L., Tull, M., Rucker, L., & Mennin, D.S. (2006). Evidence of  
38  
39 broad deficits in emotion regulation associated with chronic worry and generalized  
40  
41 anxiety disorder. *Cognitive Therapy and Research, 30, 469-480.*  
42  
43  
44 Scherer, K. R. (1997a). Profiles of emotion-antecedent appraisal: Testing theoretical  
45  
46 predictions across cultures. *Cognition & Emotion, 11, 113-150.*  
47  
48  
49 Scherer, K. R. (1997b). The role of culture in emotion-antecedent appraisal. *Journal of*  
50  
51 *Personality & Social Psychology, 73, 902-922.*  
52  
53  
54 Schmitz, N., Hartkamp, N., Franke, G. H. (2000). Assessing clinically significant change:  
55  
56 application to the SCL-90-R. *Psychological Reports, 86 :263-274.*  
57  
58  
59  
60

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Sighinolfi C., Norcini Pala A., Chiri L. R., Marchetti I. e Sica C. (2010). Difficulties in  
4  
5 Emotion Regulation (DERS): traduzione e adattamento italiano. *Psicoterapia*  
6  
7 *Cognitiva Comportamentale*, 16, 141-170.  
8  
9  
10 Sim, L., & Zeman, J. (2005). Emotion regulation factors as mediators between body  
11  
12 dissatisfaction and bulimic symptoms in early adolescent girls. *Journal of Early*  
13  
14 *Adolescence*, 25, 478–496.  
15  
16 Sim, L., & Zeman, J. (2006). The contribution of emotion regulation to body dissatisfaction  
17  
18 and disordered eating in early adolescent girls. *Journal of Youth and Adolescence*, 35,  
19  
20 207–216.  
21  
22  
23 Speranza M, Corcos M, Loas G, Stephan P, Guilbaud O, Perez- Diaz F, et al.(2005).  
24  
25 Depressive personality dimensions and alexithymia in eating disorders. *Psychiatry*  
26  
27 *Research* 135, 153–163.  
28  
29  
30 Stice, E., Nemeroff, C., & Shaw, H. (1996). A test of the dual pathway model of bulimia  
31  
32 nervosa: Evidence for restrained eating and affect regulation mechanisms. *Journal of*  
33  
34 *Social and Clinical Psychology*, 15, 340-363.  
35  
36  
37 Taylor, G., Parker, J., Bagby, M., & Bourke, M. (1996). Relationships between alexithymia  
38  
39 and psychological characteristics associated with eating disorders. *Journal of*  
40  
41 *Psychosomatic Research*, 41(6), 561–568.  
42  
43  
44 Taylor, S. E., Sherman, D. K., Kim, H. S., Jarcho, J., & Takagi, K. (2004). Culture and social  
45  
46 support: Who seeks it and why? *Journal of Personality and Social Psychology*, 87,  
47  
48 354-362.  
49  
50  
51 Telch, C. F., Agras, W. S., & Linehan, M. M. (2001). Dialectical behavior therapy for binge  
52  
53 eating disorder. *Journal of Consulting and Clinical Psychology*, 69, 1061-1065.  
54  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

- 1  
2  
3 Terracciano A, McCrae RR, & Costa, PT Jr. (2003). Factorial and construct validity of the  
4  
5 Italian Positive and Negative Affect Schedule (PANAS). *European Journal of*  
6  
7 *Psychological Assessment, 19*, 131-141.
- 9  
10 Thompson, R. (1994). Emotion regulation: A theme in search of definition. *Monographs of*  
11  
12 *the Society for Research in Child Development, 59*, 25-52.
- 14  
15 Triandis, H. (1993). Collectivism and individualism as cultural syndromes. *Cross- Cultural*  
16  
17 *Research: The Journal of comparative Social Science, 27*, 155-180.
- 18  
19 Tull, M.T., & Roemer, L. (2007). Emotion regulation difficulties associated with the  
20  
21 experience of uncued panic attacks: Evidence of experiential avoidance, emotional  
22  
23 nonacceptance, and decreased emotional clarity. *Behavior Therapy, 38*, 378–391.
- 25  
26 Tull, M.T., Barrett, H.M., McMillan, E.S., & Roemer, L. (2007). A preliminary investigation  
27  
28 of the relationship between emotion-regulation difficulties and posttraumatic stress  
29  
30 symptoms. *Behavior Therapy, 38*, 303–313.
- 32  
33 Tweed, R. G., White, K., & Lehman, D. R. (2004). Culture, stress, and coping: Internally- and  
34  
35 externally-targeted control strategies of European Canadians, east Asian Canadians,  
36  
37 and Japanese. *Journal of Cross-Cultural Psychology, 35*, 652-668.
- 38  
39 Van de Vijver, F., & Hambleton, R.K. (1996). Translating tests: Some practical guidelines.  
40  
41 *European Psychologist, 1*, 89-99.
- 42  
43 VanderVoort, D. J. (2001). Cross-cultural differences in coping with sadness. *Current*  
44  
45 *Psychology: Developmental, Learning, Personality, and Social, 20*, 147-153.
- 47  
48 Watson, D., Clark, L.A., & Tellegen, A. (1988). Development and validation of brief measures  
49  
50 of positive and negative affect: The PANAS scales. *Journal of Personality and Social*  
51  
52 *Psychology, 54*, 1063–1070.
- 53  
54  
55  
56  
57  
58  
59  
60



## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Whiteside, U., Chen, E., Neighbors, C., Hunter, D., Lo, T., & Larimer, M. (2007). Difficulties regulating emotions: Do binge eaters have fewer strategies to modulate and tolerate negative affect? *Eating Behaviors*, 8, 162–169.

Yeh, C., & Inose, M. (2002). Difficulties and coping strategies of Chinese, Japanese and Korean immigrant students. *Adolescence*, 37, 69-82.

Zlotnick, C., Donaldson, D., Spirito, A., Pearlstein, T. (1997). Affect regulation and suicide attempts in adolescent inpatients. *Journal of the American Academy of the Child & Adolescent Psychiatry*, 36, 793-798.

Zonnevylle-Bender, M.J., van Goozen, S.H., Cohen-Kettenis, P.T., van Elburg, A., & van Engeland, H. (2002). Do adolescent anorexia nervosa patients have deficits in emotional functioning? *European Child & Adolescent Psychiatry*, 11, 38–42.

RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 1

Descriptive Statistics for DERS Scales Among Women (N = 249), Men (N = 74), and Entire Sample (N = 323).

Scale	Women (N = 249)				Men (N = 74)				Entire Sample (N = 323)			
	Mean	SD	Skew	Kurtosis	Mean	SD	Skew	Kurtosis	Mean	SD	Skew	Kurtosis
NONACCEPTANCE	12.4	4.4	.9	.6	12.4	4.6	.6	-.5	12.4	4.5	.8	.3
GOALS	14.7	4.4	.3	-.5	14.6	4.4	.2	-.3	14.6	4.4	.2	-.4
IMPULSE	12.2	4.5	1.2	1.4	11.6	3.8	1.0	1.2	12.1	4.3	1.2	1.5
AWARENESS	14.5	4.1	.3	-.2	14.4	3.6	.1	.1	14.5	4.0	.3	-.1
STRATEGIES	17.1	6.3	.8	0	17.2	5.9	.9	1.2	17.1	6.2	.8	.2
CLARITY	10.3	3.3	1.0	1.5	10.6	3.4	.5	.7	10.4	3.3	.9	1.3
TOTAL	81.1	18.8	.6	.7	80.8	17.8	.3	.3	81.1	18.5	.6	.6

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 2

*Correlations Among Subscales of the DERS (N = 323).*

	NONACCEPTANCE	GOALS	IMPULSE	AWARENESS	STRATEGIES	CLARITY
NONACCEPTANCE	–					
GOALS	.35**	–				
IMPULSE	.52**	.56**	–			
AWARENESS	.12*	-.03	.15**	–		
STRATEGIES	.59**	.56**	.57**	.12*	–	
CLARITY	.34**	.21**	.35**	.52**	.33**	–

*Note.* \*  $p < .05$ ; \*\*  $p < .01$ .

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 3

*Factor Loadings for DERS Items Obtained From CFA among Italian Students (N = 323).*

NONACCEPTANCE	
Item 29	.76
Item 21	.84
Item 12	.74
Item 11	.69
Item 25	.77
Item 23	.37
GOALS	
Item 26	.89
Item 18	.88
Item 13	.85
Item 33	.73
Item 20	.47
IMPULSE	
Item 32	.81
Item 27	.79
Item 14	.87
Item 19	.87
Item 3	.55
Item 24	.46
AWARENESS	
Item 2	.86
Item 6	.81
Item 10	.47
Item 17	.40
Item 8	.65
Item 34	.45
STRATEGIES	
Item 16	.76
Item 15	.75
Item 31	.62
Item 35	.75
Item 28	.76
Item 22	.52
Item 36	.76
Item 30	.72
CLARITY	
Item 5	.70
Item 4	.51
Item 9	.62
Item 7	.80
Item 1	.85

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 4

*Internal Consistency Reliability Analyses for DERS Subscales (N = 323).*

Subscale	No. of items	Cronbach's alpha	Range of item-total correlations	Range of interitem correlations	Mean interitem correlation
NONACCEPTANCE	6	.83	.54 – .82	.22 – .73	.51
GOALS	5	.87	.64 – .89	.36 – .81	.58
IMPULSE	6	.86	.64 – .86	.29 – .80	.52
AWARENESS	6	.77	.59 – .80	.20 – .71	.37
STRATEGIES	8	.89	.63 – .81	.35 – .69	.50
CLARITY	5	.83	.64 – .84	.34 – .70	.49

For Peer Review

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 5

*Concurrent Validity Analyses for the DERS.*

	DERS TOTAL	DERS NONACCEPTANCE	DERS GOALS	DERS IMPULSE	DERS AWARENESS	DERS STRATEGIES	DERS CLARITY
TAS-20 ( <i>N</i> = 310)							
DIF	.64**	.50**	.28**	.50**	.41**	.43**	.56**
DDF	.37**	.22**	.13*	.15**	.42**	.22**	.50**
EOT	.27**	.16**	.01	.13*	.51**	.09	.31**
TOTAL	.56**	.38**	.19**	.35**	.57**	.33**	.59**
TMMS( <i>N</i> = 305)							
ATTENTION	-.16**	-.07	.04	.05	-.49**	-.04	-.26**
CLARITY	-.57**	-.37**	-.23**	-.35**	-.52**	-.31**	-.73**
REPAIR	-.47**	-.31**	-.25**	-.30**	-.17**	-.56**	-.25**
TOTAL	-.54**	-.33**	-.19**	-.26**	-.58**	-.37**	-.60**
SODAS ( <i>N</i> = 279)							
TOTAL	.62**	.45**	.45**	.56**	.23**	.49**	.42**

*Note.* TAS-20 = 20-item Toronto Alexithymia Scale; DIF = Difficulty Identifying Feelings; DDF = Difficulty Describing Feelings; EOT = Externally Oriented Thinking; TMMS = Trait Meta-Mood Scale; ATTENTION = Attention to Feelings; CLARITY = Clarity of Experience of Feelings; REPAIR = Repair of Emotions; SODAS = Scale of Dissociative Activities; \*  $p < .05$ ; \*\*  $p < .01$ .

## RELIABILITY AND VALIDITY OF THE DERS IN ITALY

Table 6

*Comparison Between Clinical (N = 38) and Nonclinical Age-Matched (N = 38) Female Samples*

DERS Scale	Sample	M	SD	Confidence Interval (95%)		t	df	p	d
				Min	Max				
NONACCEPTANCE	Nonclinical	11.2	3.7	10.0	12.4	7.93	74	<.005	1.17
	Clinical	17.7	7.0	15.4	20.1				
GOALS	Nonclinical	13.6	4.4	12.1	15.0	5.09	55.5*	<.005	.95
	Clinical	18.1	5.2	16.4	19.8				
IMPULSE	Nonclinical	11.6	3.4	10.4	12.7	4.13	74	<.005	1.39
	Clinical	18.3	5.9	16.3	20.2				
AWARENESS	Nonclinical	14.0	3.7	12.8	15.3	6.08	59.3*	<.005	.96
	Clinical	18.6	5.6	16.8	20.5				
STRATEGIES	Nonclinical	14.1	5.0	12.5	15.8	4.18	64.3*	<.005	1.69
	Clinical	24.6	7.1	22.2	26.9				
CLARITY	Nonclinical	9.2	2.6	8.3	10.0	7.35	74	<.005	1.49
	Clinical	14.9	4.8	13.4	16.5				
TOTAL	Nonclinical	73.6	16.1	68.3	78.9	6.50	57.7*	<.005	1.82
	Clinical	111.6	24.8	103.5	119.8				

*Note.* \* Since homoscedasticity could not be assumed, Welch-Satterthwaite method was used to adjust degrees of freedom.