

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Re: El-Garem et al.: Seminal Helicobacter pylori Treatment Improves Sperm Motility in Infertile Asthenozoospermic Men (Urology 2014;84:1347-1350).

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1509267> since 2016-05-10T11:06:00Z

Published version:

DOI:10.1016/j.urology.2015.01.040

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Urology. 2015 May;85(5):1217. doi: 10.1016/j.urology.2015.01.040.

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://www.goldjournal.net/issue/S0090-4295%2815%29X0004-5?page=1>

On *Helicobacter pylori* treatment: involvement in male infertility

Gian Paolo Caviglia¹, Sharmila Fagoonee², Rinaldo Pellicano³

¹Department of Medical Sciences, University of Turin, Turin, Italy

²Institute for Biostructures and Bioimages-CNR c/o Molecular Biotechnology Center, University of Turin, Turin, Italy

³Department of Gastroenterology, Città della Salute e della Scienza Hospital, Turin, Italy

Address correspondence to: Gian Paolo Caviglia, Department of Medical Sciences, University of Turin, Turin, 10100, Italy. E-mail: caviglia.giampi@libero.it

We have read the paper by El-Garem et al reporting the effect of *Helicobacter pylori* (*H. pylori*) treatment on sperm motility in infertile asthenozoospermic men.¹ Evaluating seminal antibodies (IgA) to *H. pylori* by enzyme-linked immunosorbent assay on a total of 223 men with progressive and nonprogressive sperm motility, authors found elevated levels of seminal *H. pylori* IgA antibody in 22 (9.8%) subjects. These cases were treated with anti-*H. pylori* triple therapy based on omeprazole, tinidazole and clarithromycin. After treatment, a significant decrease of antibody levels was observed in association with a significant increase in sperm motility and normalization of morphology. Considering a potential antigenic mimicry mechanism, inducing an autoimmune cross-reaction between antibody directed to bacterial antigens and β -tubulin protein of human spermatozoa, authors concluded that *H. pylori* treatment improved sperm motility suggesting the research of seminal antibodies to *H. pylori* in asthenozoospermic men.¹

H. pylori is a Gram-negative bacterium, its niche is the stomach where can cause gastritis and peptic disease. Although since two decades extra-gastric manifestations have been attributed to this bacterium but only for a small part of these a role has been proved.²

Although the authors correctly reported the lack of association between serum and seminal IgA, a clear clinical message should be highlighted. In this study the presence of *H pylori* has not been shown because the assessment of antibodies to the bacterium in semen is not an appropriate test. In fact, according to International guidelines the non-invasive diagnosis of *H pylori* infection should be based on either ¹³C-Urea breath test (sensitivity 88-95% and specificity 95-100%) or monoclonal stool antigen test (sensitivity 94% and specificity 92%), called direct tests; as last resort, only validated IgG serology should be used (indirect test).³ Indeed, the presence of antibodies to *H pylori* could represent just a marker of previous exposure without necessarily indicating a current infection. In the context of interventional studies the use of a direct diagnostic test both prior to and after treatment is mandatory.

Since the message highlighted in this paper¹ could carry relevant implications in clinical practice, we suggest that future studies on this issue should use appropriate and validated methods for *H pylori* diagnosis in order to avoid unnecessary antibiotic treatments in potential uninfected individuals.

References

1. El-Garem Y, El-Sawy M, Mostafa T. Seminal Helicobacter pylori Treatment Improves Sperm Motility in Infertile Asthenozoospermic Men. *Urology*. 2014;84:1347–1350.
2. Marietti M, Gasbarrini A, Saracco G, Pellicano R. Helicobacter pylori infection and diabetes mellitus: the 2013 state of art. *Panminerva Med*. 2013;55:277-281.
3. Malfertheiner P, Megraud F, O’Morain CA, et al. Management of Helicobacter pylori infection--the Maastricht IV/ Florence Consensus Report. *Gut*. 2012;61:646–664.