AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Lp microlocal properties for vector weighted pseudodifferential operators with smooth symbols

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/151610> since 2016-02-03T16:52:25Z

Publisher:

PKJS

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

■ **Gianluca Garelo** Dipartimento di Matematica, Università di Torino, Italy,
email: gianluca.garelo@unito.it,

Alessandro Morando Dipartimento di Matematica, Università di Brescia, Italy,
email: alessandro.morando@ing.unibs.it

L^p microlocal properties for vector weighted pseudodifferential operators with smooth symbols

Abstract

The authors introduce a class of pseudodifferential operators, whose symbols satisfy completely inhomogeneous estimates at infinity for the derivatives, namely:

$$|\partial_\xi^\alpha \partial_x^\beta a(x, \xi)| \leq c_{\alpha, \beta} m(\xi) \Lambda(\xi)^{-\alpha},$$

where $m(\xi)$ is a suitable positive continuous weight function, which indicates the “order” of the symbol, and $\Lambda(\xi) = (\lambda_1(\xi), \dots, \lambda_n(\xi))$ is a weight vector.

Continuity properties in suitable weighted Sobolev spaces of L^p type are given and L^p microlocal properties studied.

BIBLIOGRAPHY

- [1] Beals, R., *A general calculus of pseudodifferential operators*, Duke Math. J. **42**, 1-42, (1978).
- [2] Garelo, G., Morando, A., *L^p microlocal properties for multi-quasi-elliptic pseudodifferential operators*, Pliska Stud.Math. Bulgar. To appear (2013)
- [3] M.E. Taylor M.E., *Pseudodifferential Operators*, Princeton Univ. Press, 1981.