



since

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Urban air and tobacco smoke as conditions that increase the risk of oxidative stress and respiratory response in youth

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/151735

Published version:

DOI:10.1016/j.envres.2014.12.008

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in ENVIRONMENTAL RESEARCH, 137, 2015, 10.1016/j.envres.2014.12.008.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), 10.1016/j.envres.2014.12.008

The definitive version is available at: http://linkinghub.elsevier.com/retrieve/pii/S0013935114004575

Urban air and tobacco smoke as conditions that increase the risk of oxidative stress and respiratory response in youth

Roberto Bono^{1*}, Roberta Tassinari¹, Valeria Bellisario¹, Giorgio Gilli¹, Marco Pazzi², Valentina Pirro², Giulio Mengozzi³, Massimiliano Bugiani⁴, Pavilio Piccioni⁴

¹ Department of Public Health and Pediatrics. University of Torino, Italy

² Department of Chemistry, University of Torino, Italy

³ Clinical Chemistry Laboratory, San Giovanni Battista Hospital, Torino, Italy

⁴ Unit of Respiratory Medicine, National Health Service (ASL TO2), Torino, Italy

* Correspondig author: prof. Roberto Bono. University of Turin, ITALY. Department of Public Health and Pediatrics. Via Santena 5 bis. 10126 Turin, Italy. Phone: (+39) 011-670-5818 Fax: (+39) 011-236-5818. e-mail: roberto.bono@unito.it

ABSTRACT

BACKGROUND. Air pollution and tobacco smoke can induce negative effects on the human health and often leads to the formation of oxidative stress.

OBJECTIVE. The purpose of this study was to clarify the role of the urbanization degree and of passive exposure to tobacco smoke in the formation of oxidative stress. Thus, a group of non-smoking adolescents was recruited among those who live and attend school in areas with three different population densities. To each subject a spot of urine was collected to quantify 15-F_{2t} isoprostane as a marker of oxidative stress and cotinine as a marker of passive exposure to tobacco smoke. Furthermore, respiratory functionality was also measured.

RESULTS. Multiple Linear Regression analysis results showed a direct correlation (p<0.0001) of 15-F_{2t} isoprostane with both the urbanization and passive smoke. Lung function parameters proved significantly lower for the subjects living in the most populous city of Torino.

CONCLUSION. This remarks the negative effect that urbanization has on the respiratory conditions. Lastly, lung functionality presented a low inverse correlation with 15- F_{2t} isoprostane, suggesting an independent mechanism than that of the urban factor.

KEY WORDS: tobacco smoke, urban pollution, adolescents, oxidative stress, respiratory fluxes.

ABBREVIATIONS

PM = particulate matter 15-F_{2t} IsoP = 15-F_{2t}-isoprostane SIDRIA = Italian Studies on Respiratory Disorders of Childhood and the Environment CREA = creatinine E.L.I.S.A. = Enzyme-Linked Immuno Sorbent Assay FVC = Forced vital capacity, FEV1 = forced expiratory volume in one second PEF = maximal expiratory flows at peak of FVC, FEF50 = maximal expiratory flows at 50% of FVC, FEF25 = maximal expiratory flows at 25% of FVC, FEF25-75 = maximal expiratory flows among 25-75% of FVC MLR = Multiple linear regression C.I. = confidence interval

Ethical considerations

Since the subjects were underage, during a public meeting, parents and teachers were informed on the objective of this study. A written informed consent was signed and delivered by each the participants' parents. Thus, the participation of all the human subjects did not occur until after informed consent was obtained. However, the local ethics committee (ASL TO2, Turin Italy) has expressed a favorable opinion with practice number 826/13/08.

1. INTRODUCTION

The airborne particulate matter (PM) has several origins, is formed in different places where its precursors may be different; thus it possesses various physico-chemical and toxicological properties (Götschi T. 2005; Hazenkamp-Von Arx, 2004; Traversi et al., 2008). Depending on type and quantity, the presence of airborne PM can determine deleterious effects on the global environment, cultural heritage, human activities and health (Fang et al., 2010; Henschel et al., 2012; Katsouyanni et al., 2009; Levy et al., 2012; Poschl, 2005; Raaschou-Nielsen et al., 2013; Strak et al., 2012). To contain the problem, the European Union established air quality guidelines for PM as well as for other risky air pollutants (UNION, 2008). At the same time, the research activities of the scientific community were focused on the urban air pollution and its potential risk for health (Bono et al., 2010; Bono R., 2001; Bono R., 2014; Cohen et al., 2005; Fraser et al., 2003; Tzivian, 2011), in search of the best preventive techniques against the onset of diseases related to air pollution.

Exposure to urban air pollutants, whose concentration is partly dependent on proximity and intensity of traffic, is connected with the onset of asthma, development of respiratory allergies (Badyda et al., 2013; Ghio et al., 2012; Laumbach and Kipen, 2012), lung dysfunction (Kelly and Fussell, 2011; Wright and Brunst, 2013), inflammation, and exacerbation of other respiratory and cardiovascular problems (Mills et al., 2009). Numerous among these pathological conditions can be preceded or highlighted by the presence of internal dose markers, by biosynthesis of biological effects markers or, in some cases, by the formation of oxidative stress (Castro-Giner et al., 2009; Patel et al., 2013). An imbalance of the oxidative status is often a condition that precedes the onset of these respiratory diseases, and it is due to the exposure to airborne oxidants (Kelly et al., 2011; Sava and Carlsten, 2012) and a decreasing biosynthesis of endogenous antioxidant molecules (Yang W., 2009). To date, the mechanisms by which oxidants interact with molecules, cells, and tissue remain largely unclear. Remarkably, oxidative stress is also related to the

inflammatory response due to tobacco smoke (Doruk et al., 2011; Howard et al., 1998), which contains a complex mixture of mutagenic chemicals (Granella et al., 1996) able to promote lipid peroxidation (Kalra et al., 1991), protein and DNA oxidation (Vadhanam et al., 2012; van Rijt et al., 2012).

F2-isoprostanes, specific and stable products of lipid peroxidation (Basu, 2009), are noninvasive biomarkers for *in vivo* investigations of oxidative stress status (Roberts and Morrow, 2000; Romanazzi et al., 2013), airways inflammation (Basu, 2008) and asthma (Wedes et al., 2009). They can also be implicated in a larger number of human diseases, even if a clear correlation between many of these pathological conditions and oxidative stress is far from being proven (Giustarini et al., 2009). The determination of F2-isoprostanes levels in selected populations may help understanding the role that some environmental factors play in the expression of oxidative stress. In particular, the 15-F_{2t}-isoprostane (15-F_{2t} IsoP) can be monitored, since it has been proven capable to highlight different biological responses to environmental stimuli, particularly those concerning airborne chemicals (Bono 2014).

Quantification of oxidative stress by means of F2-IsoPs has several advantages if compared to other biomarkers, including the one that its levels are unaffected by diet (Gopaul 2000, Jacob 2013). At this concern, Roberts and Morrow reported that urinary F2-IsoPs, in subjects consuming a normal diet, does not decrease after a four days diet consisting only of glucose (Roberts and Morrow, 2000), and Richelle refers that the lipid content of the diet does not affect the level of urinary F2-IsoPs (Richelle 1999). This aspect of F2-IsoPs is particularly useful when, as in this case, the role of the diet is not object of interest, although it is very important in the manifestation of oxidative stress.

Finally, the relationship between biosynthesis of 15- F_{2t} IsoP and levels of respiratory functionality, in relation to the environmental conditions of life, are still largely to deepen. That is, the purpose of this study was to clarify the role that some independent

environmental, individual, and physiological variables have on the oxidative stress status of

a large population of healthy non-smoking adolescents, living in three different areas of the Piedmont region (northwestern Italy).

2. MATERIALS AND METHODS

15-F_{2t} IsoP levels were studied in relation to the urbanization degree of the selected areas where the adolescents live and attend school, in order to understand the role that urbanization might play on oxidative stress formation. Any additional information, essential for the study, was collected through a questionnaire filled out by all the adolescents, after their parents or legal tutors had signed an informed consent. In detail:

2.1. Sampling Sites. As shown in **Figure 1**, three geographic areas with different levels of urbanization and anthropization were chosen in the Piedmont region (northwestern Italy, 25.401,56 km²): Torino, capital of Piedmont, a urbanized city with almost 900.000 inhabitants (6.700 inhabitants/km², 130,2 km², 240 m. above sea level); Chivasso, a smaller and less urbanized city with about 26.000 inhabitants (507 inhabitants/km², 51,3 km² 183 m above sea level); and Casalborgone, a rural site with 1880 inhabitants (93,3 inhabitants/km², 20,2 km², 205 m above sea level). Due to the relative proximity with one another, the three locations do not have significant differences in climate, geography, altitude or social habits.

2.2. Epidemiological sample. The epidemiological sample was prepared with the aim to represent the young population of the three locations of the Piedmont region. All subjects were volunteers recruited in lower secondary schools. In more detail, three schools were located in residential and commercial areas of the city of Torino and 214 subjects were recruited from there; one school was in Chivasso, where 119 subjects were recruited; one school was in the rural area of Casalborgone and 57 subjects were enrolled from there. Since all the students were minors, parents or legal tutors were asked to sign an informed consent. Sampling was carried out over the period from March to May 2012. Each adolescent was asked to fill out a questionnaire, perform a spirometry test to evaluate their

respiratory functionality, and provide a urine sample for the determination of 15- F_{2t} lsoP and cotinine.

Questionnaire. For each student, a short version of the questionnaire "SIDRIA" was prepared to acquire information on age, sex, place of residence, hobbies, therapies, and parent's smoking habits (SIDRIA 1997). An interviewer administered the questionnaire during school hours, the same day the urine sampling and the spirometry took place.

2.3. Biological samples and statistical analysis.

2.3.1. Urinary cotinine. Cotinine measurement was carried out to quantify the passive exposure to tobacco smoke, which represents a possible factor of oxidative stress formation. A specimen of the first morning urine was collected from each volunteer and stored at -80 °C until analysis. Cotinine was measured by gas chromatography-mass spectrometry. The analytical procedure has been described in detail elsewhere (Bono R., 2014). Cotinine concentrations were normalized to the urinary creatinine (CREA) levels, as usual for every urinary measurement.

2.3.2. Urinary Isoprostane. 15- F_{2t} IsoP was measured in urine by ELISA, as previously described (Romanazzi et al., 2013). A microplate kit specific for urinary 15- F_{2t} IsoP (Oxford, MI, USA) was used following manufacturers' instructions. The declared limit of detection is 0.2 ng/ml and the possible cross-reactivity of this method is fixed below 3%. To achieve better accuracy by the ELISA method, a dilution rate of 1:4 (v/v) was adopted (Romanazzi et al., 2013). 15- F_{2t} IsoP concentrations were normalized to the CREA levels.

2.3.3. Spirometry. According to the current standards (ATS/ERS 2005), maximal expiratory flow-volume curves were obtained while the subjects were in a standing position, wearing a nose clip and breathing into a pneumotachograph (Medicalgraphics). The instrument was calibrated with a 3-liter syringe. The measurements were repeated until the volume variability did not exceed 150 ml for at least 2 times. Forced vital capacity (FVC),

forced expiratory volume in one second (FEV1) and maximal expiratory flows at peak 50%, 25% and among 25-75% of FVC (PEF, FEF50, FEF25, FEF 25-75) were recorded (Bono et al., 1998; Miller M.R. and McKay R., 2005).

2.3.4. Statistical analysis. Statistical analysis was carried out with the statistical package "Stata" (version 12 SE for MS Windows® 64 bit). In **table 1** descriptive statistics was reported *per* each location of sampling. A Box-Cox regression (Box GEP, 1964) was performed to find the power transformation that stabilize the variance and normalize the distribution.

A Multiple linear regression (MLR) was carried out to assess the effect on covariates on 15-F2t IsoP and lung function parametrers respectively, using 15-F_{2t} IsoP or lung function parametrers as dependent variable, and age, height, weight, gender (female as reference value), urinary cotinine, and sampling location as independent variables. A significant level (a two tailed P-value) of 0.05 (CI = 95%) was chosen for the statistical tests. For the final regression model, only variables that proved to be significant were selected.

3. RESULTS

The characteristics of the population enrolled in the study are described in **table 1**. Cohort numerousness, mean, and standard deviation (sd) and percentage (%) for gender, height and weight, age, and passive smoking exposure are reported per each investigated location where the adolescents live and attend school. All these parameters proved not to be statistically different among locations. Therefore, we could consider these individual characteristics as homogeneous in the three different sampling sites.

In **table 2**, means and sd, minimum and maximum values of 15- F_{2t} IsoP concentrations, normalized to the urinary creatinine values (ng/mg), are listed *per* sampling location. Torino is the area with highest mean value of 15- F_{2t} IsoP in comparison to Chivasso e Casalborgone (p<0.001). Since Torino is the most densely populated site, this result

suggests the presence in the city of a possible "urban factor": the greater the urbanization level, the higher the 15-F_{2t} IsoP concentrations. According to the Box-Cox regression results, the values of 15-F_{2t} IsoP and urinary cotinine were subjected to a logarithmic transformation prior to execute the multiple linear regression (MLR) analysis.

Running the MLR test allowed us to observe that sex, height and weight are not statistically significant in the model (p>0.050) and, therefore, were excluded from the computation of the final regression model. On the contrary, urinary cotinine, sampling location and age had a significant relationship with log 15-F_{2t} IsoP (r^2 =0.37; *p*<0,001) and, thus used to compute the model (**table 3**). In particular, log 15-F_{2t} IsoP adjusted for age and sampling site, proved to be positively correlated to cotinine, as shown in **figure 2**, with an estimated increase of 17% for every increasing unit of cotinine concentration in a log scale.

The mean value of log transformed 15- F_{2t} IsoP concentrations, referring to the entire population, and adjusted for log cotinine and sampling site, significantly decreases of 6% for every year of age (**figure 3**).

As stated above, the mean value of 15- F_{2t} IsoP concentrations was significantly higher in the adolescents of Torino when compared to those who live in Chivasso (+12%) and Casalborgone (+51%). The adolescents living in Chivasso also presented higher concentrations in respect to those who live in Casalborgone (+34%). All the effects are orthogonal like (**figure 4**).

Table 4 shows the marginal geometric means, adjusted for the covariates (age, gender, BMI), and the lower and upper limits at a 95% confidence interval (C.I.) of the lung function parameters *per* sampling location. All lung function parameters (volumes and flows), were significantly lower for the adolescents of Torino when compared to the other locations. Adjusting the concentrations for age, gender, BMI, and sampling location, the middle volume flow rates (FEF50 and FEF 25-75) and FEV1/FVC% proved to be negatively correlated with the population density of the three sampling sites (**figure 5**); while the volumes FVC (mean

+0.002, CI 95% -0022 +0026) and FEV1 (CI 95% -0019 -0048 +0009) did not. This evidence can be accepted considering that, compared to the volumes, the flows are more sensitive, especially in the pediatric age.

4. DISCUSSION.

The main goal of this work was to highlight the role that the urbanization level of the location where people inhabit may have in the oxidative stress formation. Healthy non-smoking adolescents were chosen as target population. Three areas of the Piedmont region with different demographic and road-traffic intensity, though not very far from each other, were investigated: Torino (a big city), Chivasso (a small town), and Casalborgone (a small rural village). The oxidative stress level was monitored through the quantification of $15-F_{2t}$ IsoP urinary concentration. Levels of this biomarker are unaffected by diet, an antioxidant factor, potentially confounding the relationship we have investigated. (Gopaul et al., 2000; Roberts and Morrow, 2000, Jacob et al. 2013, Richelle 1999). In particular, the diet is very similar among all the students because: a) they benefit from the same school lunch prepared by the same company according to the requirements imposed by nutritionists working at the local health authority to minimize oxidant food, b) all the students are white and of Caucasian ethnicity. This may mean that the diet consumed the previous evening at home is likely to be similar, c) although the three groups of students are different for population density, the distance between them does not exceed 50 kilometers and the altitude above sea level is the same.

Since the passive exposure to tobacco smoke can also influence the oxidative stress level, urinary cotinine was measured to know the role of the tobacco in the onset of 15-F_{2t} IsoP values and used it to adjust the relationship between 15-F2t ISoP and the urban factor. Finally, the respiratory function has also been taken into account as physiological factor

potentially able to be altered by the two environmental aspects regarded in the present study. The results showed the presence of a direct correlation between 15-F_{2t} IsoP and the degree of urbanization of the areas where the adolescents live and attend school. This suggests that an "urban factor" plays a direct role in the synthesis of 15-F_{2t} IsoP inducing its increase up to about 50%. Thus, the level of urbanization highlights a role of risk factor able to increase oxidative stress in adolescents, which proved to be a population particularly sensitive to even small environmental differences.

Passive exposure to tobacco smoke and age of the subject proved to be other factors that can significantly influence the 15-F_{2t} IsoP concentrations but while the exposure to passive smoke increases 15-F_{2t} IsoP levels, the age leads to the opposed result. The latter effect has been recently observed in an independent population of a similar age (Bono R., 2014) but opposes to the general trend observed in adults. Indeed, recent scientific studies showed an increase in the intensity of oxidative stress with aging, and with the onset of many age-related diseases, including Alzheimer (Bouzid MA, 2014; Jacob et al., 2013; Montine et al., 2011).

Another finding of this study is the significant lower level of respiratory fluxes in the adolescents living in Torino, in comparison to those living in less urbanized locations. This finding shows the responsibility of the higher level of urbanization of Turin in the reduction of respiratory flows. This aspect highlights, at the same time, an increase of respiratory risk. Furthermore, the three measures of fluxes (FEF 50, FEF 25-75, and FEV1/FVC), adjusted for age, gender, and BMI were negatively correlated to 15-F₂ lsoP (p<0.047, p<0.013, p<0.005 respectively) when compared per sampling location (**figure 5**). This allows us to consider the low values of flux intensity as a respiratory condition inversely correlated with the onset of oxidative stress. This is true even after removing the effect of 15-F₂ lsoP, which highlights the relationship of the respiratory effects from urbanization independent from inflammation and oxidative stress. Finally, we can conclude that the adolescents studied

show an increase in oxidative stress and a decrease in respiratory flow dependent from the urbanization and the tobacco smoke passively breathed. Thus, the evidence of this risky condition for public health may represent a platform for designing new preventive strategies against tobacco smoke exposure and urban pollution.

5. ACKNOWLEDGEMENTS

The authors kindly thank to all the students who have generously collaborated on this study.

This study was made possible by a grant of University of Turin, Italy to Roberto Bono (ex

60% 2013) and a grant of Regione Piemonte to Pavilio Piccioni (Ricerca finalizzata 2011).

6. REFERENCES

ATS/ERS statement: raised volume forced expirations in infants: guidelines for current practice. American Thoracic Society; European Respiratory Society. Am J Respir Crit Care Med. 2005 Dec 1; 172(11):1463-71.

Badyda A.J., Dabrowiecki P., Lubinski W., Czechowski P.O., Majewski G., Chcialowski A. and Kraszewski A., 2013. Influence of traffic-related air pollutants on lung function. Adv Exp Med Biol, 788, 229-35.

Basu S., 2008. F-2-isoprostanes in human health and diseases: from molecular mechanisms to clinical implications. Antioxid Redox Signal, 10, 1405-1432.

Basu S., Helmersson J., Jarosinska D., Sallsten G., Mazzolai B., Barregard L., 2009. Regulatory Factors of basal F2-isoprostane formation: population, age, gender and smoking habits in humans. free Radic. Res., 43, 85-91.

Bono R., Bellisario V., Romanazzi V., Pirro V., Piccioni P., Pazzi M., Bugiani M. and Vincenti M., 2010. Oxidative stress in adolescent passive smokers living in urban and rural environments. Int J Hyg Environ Health, 217, 287-93.

Bono R., Nebiolo F., Bugiani M., Meineri V., Scursatone E., Piccioni P., Caria E., Gilli G. and Arossa W., 1998. Effects of tobacco smoke exposure on lung growth in adolescents. J Expo Anal Environ Epidemiol, 8, 335-45.

Bono R., Bugliosi E.H., Schilirò T., Gilli G., 2001. The Lagrange Street story: The prevention of aromaticsair pollution during the last nine years in a European city. Atmospheric Environment, 35, S107-S113.

Bono R., Bellisario V., Romanazzi V., Pirro V., Piccioni P., Pazzi M., Bugiani M., Vincenti M., 2014. Oxidative stress in adolescent passive smokers living in urban and rural environments. Int J Hyg Envir Heal 217, 287-293.

Bouzid MA, Hammouda O., Matran R, Robin S, Fabre C., 2014. Changes in oxidative stress markers and biological markers of muscle injuery with aging at rest and in response to an exhaustive exercise. PLoS One, 11, 9(3):e90420.

Box GEP, Cox D.R., 1964. An analysis of transformations. Journal of the Royal Statistical Society Series B 211-252.

Castro-Giner F., Kunzli N., Jacquemin B., Forsberg B., de Cid R., Sunyer J., Jarvis D., Briggs D., Vienneau D., Norback D., Gonzalez J.R., Guerra S., Janson C., Anto J.M., Wjst M., Heinrich J., Estivill X. and Kogevinas M., 2009. Traffic-related air pollution, oxidative stress genes, and asthma (ECHRS). Environ Health Perspect, 117, 1919-24.

Cohen A.J., Ross Anderson H., Ostro B., Pandey K.D., Krzyzanowski M., Kunzli N., Gutschmidt K., Pope A., Romieu I., Samet J.M. and Smith K., 2005. The global burden of disease due to outdoor air pollution. J Toxicol Environ Health A, 68, 1301-7.

Doruk S., Ozyurt H., Inonu H., Erkorkmaz U., Saylan O. and Seyfikli Z., 2011. Oxidative status in the lungs associated with tobacco smoke exposure. Clin Chem Lab Med, 49, 2007-12.

Fang S.C., Cassidy A. and Christiani D.C., 2010. A systematic review of occupational exposure to particulate matter and cardiovascular disease. Int J Environ Res Public Health, 7, 1773-806.

Fraser M.P., Buzcu B., Yue Z.W., McGaughey G.R., Desai N.R., Allen D.T., Seila R.L., Lonneman W.A. and Harley R.A., 2003. Separation of fine particulate matter emitted from gasoline and diesel vehicles using chemical mass balancing techniques. Environ Sci Technol, 37, 3904-9.

Ghio A.J., Carraway M.S. and Madden M.C., 2012. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. J Toxicol Environ Health B Crit Rev, 15, 1-21.

Giustarini D., Dalle-Donne I., Tsikas D. and Rossi R., 2009. Oxidative stress and human diseases: Origin, link, measurement, mechanisms, and biomarkers. Crit Rev Clin Lab Sci, 46, 241-81.

Gopaul, N.K., Halliwell B., Anggard, E.E. 2000. Measurement of plasma F2-Isoprostanes as an index of lipid peroxidation does not appear to be counfounded by diet. Free Rad. Res. 33:115-127.

Götschi T., Hazenkamp-Von Arx M.E., Heinrich J., Bono R., Burney P., Forsberg B., Jarvis D., Maldonado J., Norbäck D., Stern W.B., Sunyer J., Torén K., Verlato G., Villani S., Künzli N., 2005. Elemental composition and reflectance of ambient fine particles at 21 European locations. Atmospheric Environment, 39, 5947-5958.

Granella M., Priante E., Nardini B., Bono R. and Clonfero E., 1996. Excretion of mutagens, nicotine and its metabolites in urine of cigarette smokers. Mutagenesis, 11, 207-11.

Hazenkamp-Von Arx M.E., Gotschi T., Ackermann-Liebrich U., Bono R., Burney P., Cyrys J., Jarvis D., Lillienberg L., Luczynska C., Maldonado JA, Jaen A., De Marco R., Mi Y., Modig L., Bayer- Oglesby L., Payo F., Soon A., Sunyer J., Villani S., Weyler J., Kunzli N., 2004. PM2.5 and NO2 assessment in 21 european study centres of ERCHSII: annual means and seasonal differences. Atmospheric Environment, 38, 1943-1953.

Henschel S., Atkinson R., Zeka A., Le Tertre A., Analitis A., Katsouyanni K., Chanel O., Pascal M., Forsberg B., Medina S. and Goodman P.G., 2012. Air pollution interventions and their impact on public health. Int J Public Health, 57, 757-68.

Howard D.J., Ota R.B., Briggs L.A., Hampton M. and Pritsos C.A., 1998. Oxidative stress induced by environmental tobacco smoke in the workplace is mitigated by antioxidant supplementation. Cancer Epidemiol Biomarkers Prev, 7, 981-8.

Jacob K.D., Noren Hooten N., Trzeciak A.R. and Evans M.K., 2013. Markers of oxidant stress that are clinically relevant in aging and age-related disease. Mech Ageing Dev, 134, 139-57.

Kalra J., Chaudhary A.K. and Prasad K., 1991. Increased production of oxygen free radicals in cigarette smokers. Int J Exp Pathol, 72, 1-7.

Katsouyanni K., Samet J.M., Anderson H.R., Atkinson R., Le Tertre A., Medina S., Samoli E., Touloumi G., Burnett R.T., Krewski D., Ramsay T., Dominici F., Peng R.D., Schwartz J. and Zanobetti A., 2009. Air pollution and health: a European and North American approach (APHENA). Res Rep Health Eff Inst, 5-90.

Kelly F.J. and Fussell J.C., 2011. Air pollution and airway disease. Clin Exp Allergy, 41, 1059-71.

Laumbach R.J. and Kipen H.M., 2012. Respiratory health effects of air pollution: update on biomass smoke and traffic pollution. J Allergy Clin Immunol, 129, 3-11; quiz 12-3.

Levy J.I., Diez D., Dou Y., Barr C.D. and Dominici F., 2012. A meta-analysis and multisite time-series analysis of the differential toxicity of major fine particulate matter constituents. Am J Epidemiol, 175, 1091-9.

Miller M.R., Hankinson J., Brusasco V., Burgos F., Casaburi R., Coates A., Crapo R., Enright P., Van der Grinten C.P.M., Gustafsson P., Jensen R., Johnson D.C., MacIntyre N., McKay R., Navajas D., Pedersen O.F., Pellegrino R., Viegi G. and Wanger J., 2005. Series "ATS/ERS Task Force: Standardisation Of Lung Function Testing". Standardization of spirometry. Eur Respir J, 26, 319-338.

Mills N.L., Donaldson K., Hadoke P.W., Boon N.A., MacNee W., Cassee F.R., Sandstrom T., Blomberg A. and Newby D.E., 2009. Adverse cardiovascular effects of air pollution. Nat Clin Pract Cardiovasc Med, 6, 36-44.

Montine T.J., Peskind E.R., Quinn J.F., Wilson A.M., Montine K.S. and Galasko D., 2011. Increased cerebrospinal fluid F2-isoprostanes are associated with aging and latent Alzheimer's disease as identified by biomarkers. Neuromolecular Med, 13, 37-43.

Patel M.M., Chillrud S.N., Deepti K.C., Ross J.M. and Kinney P.L., 2013. Traffic-related air pollutants and exhaled markers of airway inflammation and oxidative stress in New York City adolescents. Environ Res, 121, 71-8.

Poschl U., 2005. Atmospheric aerosols: composition, transformation, climate and health effects. Angew Chem Int Ed Engl, 44, 7520-40.

Raaschou-Nielsen O., Vineis P., Brunekreef B., Nieuwenhuijsen M., Hoffmann B., Forastiere F., Oudin A., Hoek G., Katsouyanni K., Schwarze P. and Beelen R., 2013. Air pollution and lung cancer in Europe - authors' reply. Lancet Oncol, 14, e440.

Richelle, M., Turini, M. E., Guidoux, R., Tavazzi, I., Metairon, S., Fay, L. B. 1999. Urinary isoprostane excretion is not confounded by the lipid content of the diet. FEBS Lett. 459:259-262.

Roberts L.J. and Morrow J.D., 2000. Measurement of F(2)-isoprostanes as an index of oxidative stress in vivo. Free Radic Biol Med, 28,4, 505-513.

Romanazzi V., Pirro V., Bellisario V., Mengozzi G., Peluso M., Pazzi M., Bugiani M., Verlato G. and Bono R., 2013. 15-F(2)t isoprostane as biomarker of oxidative stress induced by tobacco smoke and occupational exposure to formaldehyde in workers of plastic laminates. Sci Total Environ, 442, 20-5.

Sava F. and Carlsten C., 2012. Respiratory health effects of ambient air pollution: an update. Clin Chest Med, 33, 759-69.

SIDRIA. 1997. Asthma and respiratory symptoms in 6-7 yr old Italian children: gender, latitude, urbanization and socioeconomic factors. SIDRIA (Italian Studies on Respiratory Disorders in Childhood and the Environment). Eur Respir J 1997, 10:1780-6

Strak M., Janssen N.A., Godri K.J., Gosens I., Mudway I.S., Cassee F.R., Lebret E., Kelly F.J., Harrison R.M., Brunekreef B., Steenhof M. and Hoek G., 2012. Respiratory health effects of airborne particulate matter: the role of particle size, composition, and oxidative potential-the RAPTES project. Environ Health Perspect, 120, 1183-9.

Traversi D., Degan R., De Marco R., Gilli G., Pignata C., Ponzio M., Rava M., Sessarego F., Villani S. and Bono R., 2008. Mutagenic properties of PM2.5 air pollution in the Padana Plain (Italy) before and in the course of XX Winter Olympic Games of "Torino 2006". Environ Int, 34, 966-70.

Tzivian L., 2011. Outdoor air pollution and asthma in children. J Asthma, 48, 470-81.

UNION E., 2008. Directive 2008/50/EC of the European Parliament and of the Council Of 21 May 2008 on ambient air quality and cleaner air for Europe.

Vadhanam M.V., Thaiparambil J., Gairola C.G. and Gupta R.C., 2012. Oxidative DNA adducts detected in vitro from redox activity of cigarette smoke constituents. Chem Res Toxicol, 25, 2499-504.

van Rijt S.H., Keller I.E., John G., Kohse K., Yildirim A.O., Eickelberg O. and Meiners S., 2012. Acute cigarette smoke exposure impairs proteasome function in the lung. Am J Physiol Lung Cell Mol Physiol, 303, L814-23.

Wedes S.H., Khatri S.B., Zhang R., Wu W., Comhair S.A., Wenzel S., Teague W.G., Israel E., Erzurum S.C. and Hazen S.L., 2009. Non invasive markers of airway inflammation in asthma. Clin Transl Sci, 2, 112-7.

Wright R.J. and Brunst K.J., 2013. Programming of respiratory health in childhood: influence of outdoor air pollution. Curr Opin Pediatr, 25, 232-9.

Yang W., Omaye S.T., 2009. Air pollutants, oxidative stress and human health. mutat. Resp., 674, 45-54.

Table 1. Gender, age, height, weight, and number of passive smokers in the whole population and in three groups divided according to the three sites where the adolescents live and attend school.

	TORI (urban	NO site)	CHIVASSO (intermediate site)		CASALBORGONE (rural site)		TOTAL	
N	214		119		57		390	
Gender N (%)	male n. (%)	122 (57%)	male n. (%)	64 (53.8%)	male n. (%)	25 (43.9%)	male n. (%)	211 (54.1%)
	female n. (%)	92 (43%)	female n. (%)	55 (46.2%)	female n. (%)	32 (55.1%)	female n. (%)	179 (45.9%)
Height cm ± sd	149.6 ± 9,3		154.0 ± 8.9		153.1 ± 9.8		151.7 ± 9.5	
Weight kg ± sd	43.53 ± 11.0		47.3 ± 12.3		47.7 ± 12.2		45.4 ± 11.7	
Age years ± sd	11.5	± 0.8	12.7 ± 0.8		12.5 ± 0.6		12.0 ± 1.0	
Smoking habits N (%)	passive	70 (32.7)	Passive	52 (43.7%)	passive	24 (42.1%)	passive	146 (37.4%)
	not exposed	144 (67.3%)	not exposed	67 (56.3%)	not exposed	33 (57.9%)	not exposed	244 (62.6%)

Table 2. 15- F_{2t} IsoP values in the three sampling locations. The three concentrations prove a direct relationship to population density: Turin, the most populated city, displays the highest mean value, Casalborgone, the rural site, the lowest.

	15-F _{2t} IsoP [ng/mg]			
	Mean ± sd	min	max	
TORINO	7.2 ± 4.0	< LOD *	37.0	
CHIVASSO	6.4 ± 5.4	< LOD *	39.8	
CASALBORGONE	4.8 ± 3.0	1.5	14.7	
TOTAL	6.5 ± 4.4	< LOD *	39.8	

* LOD = limit of detection. 15-F2t IsoP fixed to limit of detection (0.2 ng/ml) if \leq of LOD.

min = minimum value, max = maximum value, sd = standard deviation. Units of 15- F_{2t} IsoP is nanograms of 15- F_{2t} IsoP every 1 mg of urinary creatinine.

Table 3. Multiple linear regression parameters, with 95% confidence interval (C.I.), of log 15-F2t IsoP as dependent variable and log (cotinine); age and sampling site as predictors. Note: gender, height, weight, and diet indicators, not signicant at 5% level, were excluded from the model. 15-F2t IsoP fixed to limit of detection (0.2 ng/ml) if \leq of LOD (limit of detection).

	Regression coefficient (95% C.I.)		
Urinary cotinine		+0.158 (0.119-0.197)	0.0001
	Chivasso vs Casalborgone	+0.301 (0.152 –0.561)	0.0001
Sampling sites	Torino <i>v</i> s Casalborgone	+0.414 (0.268 - 0.561)	0.0001
	Torino vs Chivasso+Casalborgone	+0.224 (0.108 - 0.338)	0.0001
Age		-0.058 (-0.1150.001)	0.050

Table 4. Marginal geometric means with 95% confidence intervals (C.I.) of the lung function parameters by sampling site, as estimated by multiple regression analysis adjusted by sex, age, height, BMI index and log (cotinine).

		Moone	C.I.	n -		
		Wiedlis	Lower limit	Upper limit	μ<	
	Casalborgone	2.91	2.82	2.99		
FVC (liters)	Chivasso	2.88	2.82	2.95	0.22	
	Torino	2.77	2.73	2.82	0.01	
	Casalborgone	2.62	2.53	2.71		
FEV1 (liters)	Chivasso	2.58	2.51	2.65	0.52	
	Torino	2.39	2.34	2.44	0.01	
	Casalborgone	89.08	87.57	90.59		
FEV1/FVC (%)	Chivasso	89.54	88.31	90.77	0.37	
	Torino	86.45	85.45	87.44	0.04	
	Casalborgone	3.53	3.30	3.75		
MEF50 (liters/sec.)	Chivasso	3.66	3.51	3.81	0.16	
	Torino	3.19	3.07	3.32	0.02	
	Casalborgone	3.15	2.96	3.34		
FEF25-15 (liters/sec)	Chivasso	3.25	3.11	3.38	0.24	
	Torino	2.80	2.68	2.91	0.02	

Figure 1. Map of the sampling sites. Torino, the capital of Piedmont region in Italy is a urbanized city with almost 900.000 inhabitants. Chivasso is a smaller and less urbanized city with about 26.000 inhabitants; Casalborgone is a rural site with 1880 inhabitants.





Figure 2. Plot of the relation between log 15-F2t IsoP and log cotinine, given age and sampling site.





Figure 4. Marginal means and confidence intervals of 15- F_{2t} IsoP levels measured in the three sampling sites, adjusted for log (cotinine) and age by means of multiple regression model (with log link). Casalborgone is the rural site, Chivasso the medium size city, Torino the big city.



Figure 5. Marginal means and confidence intervals of lung function parameters by the three sampling sites, adjusted for age, height, gender, BMI and log (cotinine) by means of multiple regression model. A) FEF 50 - Forced Expiratory Flow at 50% of Forced Vital Capacity (FVC), B) FEF 25-75 mean forced expiratory flows at 25/75% of FVC, C) FEV1/FVC% = Forced expiratory volume in 1 second as % of FVC.



