

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Serotonin transporter gene: a new polymorphism may affect response to antidepressant treatments in major depression

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/151749> since 2016-12-13T11:43:15Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Yoshihiko Matsumoto; Chiara Fabbri; Silvia Pellegrini; Stefano Porcelli; Pierluigi Politi; Silvio Bellino; Veronica Mariotti; Cristina Demi; Marco Menchetti; Valentina Martinelli; Marco Cappucciati; Paola Bozzatello; Elena Brignolo; Paolo Brambilla; Matteo Balestrieri; Diana De Ronchi Alessandro Serretti. Serotonin transporter gene: a new polymorphism may affect response to antidepressant treatments in major depression. *MOLECULAR DIAGNOSIS & THERAPY*. 18 (5) pp: 567-577.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/151749>

Serotonin transporter gene: a new polymorphism may affect response to antidepressant treatments in major depressive disorder

Yoshihiko Matsumoto· Chiara Fabbri· Silvia Pellegrini· Stefano Porcelli· Pierluigi Politi· Silvio Bellino· Caterina Iofrina· Veronica Mariotti· Erika Melissari· Marco Menchetti· Valentina Martinelli· Marco Cappucciati· Paola Bozzatello· Elena Brignolo· Paolo Brambilla, Matteo Balestrieri· Alessandro Serretti

Running title: serotonergic genes in response to antidepressants and psychotherapy

Y. Matsumoto· C. Fabbri· S. Porcelli· A. Serretti (✉)

Department of Biomedical and Neuromotor Sciences, University of Bologna, Viale Carlo Pepoli 5, 40123 Bologna, Italy.

Tel: +39 051 6584233

Fax: +39 051 521030

e-mail: alessandro.serretti@unibo.it

S. Pellegrini· C. Iofrina· V. Mariotti· E. Melissari

Surgical, Medical and Molecular Pathology, Medical Biotechnology, Epidemiology and Infectious Diseases and of Critical Area, University of Pisa, Pisa, Italy

P. Politi· V. Martinelli· M. Cappucciati

Department of Brain and Behavioral Sciences, University of Pavia, Italy

S. Bellino· P. Bozzatello· E. Brignolo

Neuroscience Department, Turin University, Italy

Y. Matsumoto

Department of Psychiatry, Yamagata University School of Medicine, Japan

M. Menchetti

Department of Medical and Surgical Sciences, University of Bologna, Italy

P. Brambilla· M. Balestrieri

Department of Experimental and Clinical Medical Sciences (DISM), ICBN, Unit of Psychiatry, University of Udine, Udine, Italy

P. Brambilla

Department of Psychiatry and Behavioral Sciences, UTHouston Medical School, Houston, TX, USA

Abstract

Objective: Several gene variants have been related to major depressive disorder (MDD) treatment outcomes, however few studies investigated a possible different effect on pharmacotherapy and brief psychotherapy response.

Methods: 137 MDD patients were randomized to either inter-personal counseling (IPC n=40) or antidepressant pharmacological treatment (n=97). Outcomes were remission, response, and symptom improvement at week 8. Five genetic variants were investigated (*5HTR2A* rs6314, *BDNF* rs6265, *SLC6A4* rs8076005, *CREB1* rs2253206, and *TPH2* rs11179023) as possible modulators of outcomes.

Results: *SLC6A4* rs8076005 AA genotype and A allele were associated with response rate in the antidepressant group (p=0.015 and p=0.005, respectively) and in the whole sample (p=0.03 and p=0.02, respectively). In the IPC group a non significant trend in the same direction was observed. *TPH2* rs11179023 A allele showed a marginal association with symptom improvement only in the IPC group. Other gene variants did not impact on outcomes in any treatment group.

Conclusion: Our study suggests that rs8076005 in *SLC6A4* gene may be a modulator of antidepressant response especially when pharmacological treatment is used.

Keywords: antidepressant, psychotherapy, *HTR2A*, *BDNF*, serotonin transporter, *SLC6A4*, *CREB1*, *TPH2*, gene, polymorphism, pharmacogenetics, major depressive disorder.

Key Points

- The role in treatment response of five genes involved in the modulation of the serotonergic system (*HTR2A*, *BDNF*, *SLC6A4*, *CREB1*, and *TPH2*) was examined in 137 major depressive disorder patients treated with inter-personal counseling (IPC) or antidepressant drugs.
- Our findings supported the association of *SLC6A4* rs8076005 with response in both treatment groups.
- A marginal association between *TPH2* rs11179023 and symptom improvement was observed only in the IPC group.

1. Introduction

Major depressive disorder (MDD) is one of the most common psychiatric disorder (life time prevalence: 12.8%) [1], and ~~major depression will be the leading cause of disability by 2030~~ it is among the major causes of worldwide disability [2]. The heavy personal and socio-economic burden of MDD is partly due to the still unsatisfactory response and remission rates (47 and 33%, respectively) [3]. Antidepressant drugs and psychotherapy are thought to be equally efficacious in moderate MDD [4, 5]. However, predictors of response to antidepressant treatments to guide treatment choice in the individual patient are not available. Among predictors of clinical outcomes, genetic variants are estimated to contribute substantially to the interindividual variability [6-9]. Serotonergic genes represent ideal candidates since current antidepressant drugs mainly act by the modulation of the serotonergic system.

The serotonin receptor gene 2A (*HTR2A*) is expressed in all neocortex, hippocampus, and prefrontal cortex [10]. 5HT-induced Ca^{2+} response in platelets was found to be enhanced in MDD [11] after remission during imipramine treatment [12]. The missense T allele of *HTR2A* rs6314 showed lower Ca^{2+} response to 5-HT than those without the mutation [13]. Following studies found that this polymorphism may predict SSRIs efficacy [14, 15].

Serotonin transporter (*SLC6A4* gene) plays a pivotal role in the regulation of serotonin concentration in the synaptic cleft and extrasynaptic sites. Promoter polymorphisms (5-HTTLPR and rs25531) have been extensively investigated, leading to replicated even if not univocal results [16]. The intronic rs8076005 has been previously associated with IL-6 levels in MDD and depressive symptom severity [17], but has not been studied as possible modulator of treatment response.

Tryptophan hydroxylase (*TPH*) catalyzes the rate-limiting step in synthesis of 5-HT. *TPH2* is the main isoform expressed in the CNS, but studies focused on its role in antidepressant response reported inconsistent results [18-33]. *TPH2* rs1179023 variation has not been investigated in the field before, thus represents a new candidate within a gene with strong biological rationale of involvement in antidepressant action.

During antidepressant treatment, the enhancement of 5-HT availability increases brain-derived neurotrophic factor (*BDNF*) synthesis and signaling, a major event in the stimulation of adult neurogenesis [34]. *BDNF* serum levels are associated with MDD and antidepressant response [35]. The rs6265 (Val66Met) was associated with *BDNF* secretion [36], *BDNF* serum levels [37], and antidepressants response [38-40].

Cyclic AMP response element-binding protein1 (*CREB1*) is a transcription factor activated by a number of signals, among which neuronal growth factors and stress-induced signals [41, 42]. *CREB1* gene has been associated with recurrent early-onset MDD [43, 44]. The G allele of the promoter polymorphism rs2253206 has been associated with lower promoter activity [45] and the same SNP may affect the risk of MDD, depression severity [46], and risk of treatment resistant depression [47].

Given the data supporting the involvement of these genes in antidepressant action, the present paper aims to study the role of *HTR2A* rs6314, *BDNF* rs6265, *SLC6A4* rs8076005, *CREB1* rs2253206, and *TPH2* rs11179023 in a sample of 137 MDD Italian patients treated with IPC or antidepressants in a multicenter randomized trial.

2. Methods

2.1. Sample

Subjects aged 18 years or older with diagnosis of MDD (DSM-IV TR) were recruited. Inclusion and exclusion criteria were detailed elsewhere [48]. Eligible patients were divided into inter-personal counseling (IPC, a brief psychotherapy derived from Interpersonal Psychotherapy [49-51]) or into pharmacological antidepressant treatment [52]. Subjects were collected in three centers (Neuroscience Department, Turin University; Department of Health Sciences, University of Pavia; and Institute of Psychiatry, Bologna University), as previously reported [48]. All patients recruited at Bologna University were treated with antidepressant drugs. Despite the design of the study has been planned as randomized, the failure in developing a psychotherapy arm at Bologna University (the centre that recruited the higher number of patients) resulted in different size between the two treatment arms. In all cases of pharmacological treatment, antidepressant was chosen and titrated according to current guidelines up to the effective dose range within the first week of treatment. Ethical approval was obtained from the ethics committees of all participating centers.

2.2. Evaluations

All subjects were evaluated with both structured and unstructured interviews to obtain diagnoses assigned by two independent experienced psychiatrists. Depressive states were assessed at baseline and after 8 weeks through Hamilton Depression Rating Scale (HDRS, 21 items version). Assessments were performed by trained psychiatrists and inter-rater reliability was $k=0.8$. A monitoring of treatment adherence through drug plasma level determination was not performed, but treatment adherence was assessed weekly by a trained psychiatrist and side effects were monitored weekly using the Dosage Record & Treatment Emergent Symptom scale (DOTES) [53]. Thus, we can suppose that non-responders at week 8 were not false non-responders due to overlooked side effects and poor treatment adherence.

2.3. Outcomes

The primary outcome was remission at week 8, defined according to standard criteria (HDRS score ≤ 7), and the secondary outcomes were response (at least 50% reduction in HDRS score) and HDRS percent improvement at week 8.

2.4. Polymorphism selection

The genotyped SNPs were selected according to the following criteria: 1) tagging approach ($R^2=0.08$); 2) reported prevalence of at least 5% for the minor allele among Caucasians (data from <http://hapmap.ncbi.nlm.nih.gov/>); 3) availability of a validated assay in our laboratory; and 4) previous literature findings. In detail, the Tagger software was used to test SNPs downloaded from HapMap (Rel 28 Phase II+III, August 2010, NCBI B36 assembly, dbSNP b126) and that satisfied some quality criteria (maximum number of Mendel errors=1; p value for Hardy-Weinberg equilibrium >0.001, MAF>5%). The Tagger program was used to identify SNPs that tagged at least other 2 SNPs with r^2 and LOD thresholds of 0.8 and 3, respectively. The selected polymorphisms ~~were~~ are shown in Supplementary Table 1 together with polymorphisms in complete linkage disequilibrium ($R^2>0.8$) with ~~them, the following SNPs: *TPH2* rs11179023 with rs12229394 and rs1487278; *SLC6A4* rs8076005 with rs12150214, rs11080122, rs8071667, rs2020936, rs25528, and rs6354; *BDNF* rs6265 with rs2030323.~~

2.5. Genotyping

DNA was purified from whole blood using the QIAamp DNA Blood Kit (Qiagen, CA, USA), according to the manufacturer's protocol. rs6314 (*HTR2A*, Chr 13q14-q21), rs6265 (*BDNF*, Chr 11p13), rs8076005 (*SLC6A4*, Chr 17q11.2), rs2253206 (*CREB1*, Chr 2q34), rs11179023 (*TPH2*, Chr 12q21.1) were genotyped by High Resolution Melting (HRM)-PCR. All the experiments were performed by the Rotor Gene Q instrument (Qiagen, CA, USA), using the Type-it HRM PCR kit (Qiagen, CA, USA). All primers were designed by Beacon Designer v. 7.9 (PREMIER Biosoft, CA, USA). After an initial step of enzyme activation at 95°C for 5 min, PCRs were carried out by 40 cycles as follows: denaturation at 95°C for 10 sec and annealing at 60°C for 30 sec. The HRM analysis was performed with a temperature resolution of 0.1°C and a temperature range between 70°C and 85°C. Data collection and genotype calls were obtained by the Rotor-Gene 6000 series software v. 1.7 (QIAGEN, CA, USA) using as reference genotypes DNA samples sequenced by ABI Prism[®] 310 Genetic Analyzer (Applied Biosystems, CA, USA).

2.6. Statistical analysis

The effect of genotypes and alleles on response and remission ~~rates~~ was tested by Pearson's chi-squared tests. A Multivariate Analysis of Variance (MANOVA) One-way ANOVA was used ~~within a linear regression model~~ to study the effect of genotypes and alleles on improvement at week 8 adjusting for center, gender, age, and baseline severity. Analyses were carried out only in completers, in the two individual treatment groups and in the whole sample.

Despite The level of significance was conservatively set to 0.05 since the selected candidate genes show a strong biological rationale for involvement in antidepressant mechanisms of action, ~~thus and the~~ pre-test probability of association is favorable. Bonferroni correction was applied to reduce the risk of false

positives. Thus, alpha value was set to 0.005 considering 5 independent polymorphisms that were tested in two independent groups of patients (IPC- and antidepressant-treated). The whole sample was not considered as an independent group, as well as the three analyzed phenotypes (response, remission and symptom improvement) were considered as highly related one to each other. Further, we stated only remission as primary outcome (see paragraph 2.3.).

Our whole sample setting alpha value to 0.005 two tailed provides a power of 0.80 to detect a risk allele (for non-remission) with OR \geq 4.0~~observe a difference between two variants with effect size $d=0.26$ (Gpower 3.1 software), which corresponds to a difference of at least 1.5 points in the HDRS at week 8 and to an explained variance of 2.2%.~~

3. Results

SCL6A4 rs8076005 slightly deviated from Hardy-Weinberg equilibrium (HWE) ($p=0.03$), while the other polymorphisms satisfied HWE (*HT2A* rs6314: $p=0.61$; *BDNF* rs6265: $P=0.34$; *CREB1* rs2253206: $P=0.52$; *TPH2* rs11179023: $P=0.45$). For 1 subject in the IPC group and 2 in the antidepressant group genotyping was not successful thus they were excluded from the analysis.

Clinical-demographic characteristics of the sample were reported in Table 1. The major part of patients was treated with SSRIs (80.4%). Further clinical data were reported in a previous paper [48]. Non completers (12.74%) did not show any difference in age, gender, and baseline severity when compared to the 8 week-completers [48]. As previously reported [48], no difference was observed in clinical outcomes between the two treatment groups (Table 1). In the IPC group the number of remitters remission rate was slightly higher than that of responders the response rate because the IPC group had lower baseline severity than the antidepressant group. Nevertheless, the difference was not so pronounced to be clinically relevant and analyses were adjusted for baseline severity.

Effect of polymorphisms on outcomes

SLC6A4 rs8076005 AA genotype and A allele were reached the significance threshold for association with associated with better response in the antidepressant group ($p=0.015$ and $p=0.005$, respectively) and showed a consistent trend in the whole sample ($p=0.03$ and $p=0.02$, respectively) (Tables 2 and 3). Further, rs8076005 A allele carriers showed a trend of higher symptom improvement at week 8 in the whole sample ($F=3.29$, $df=2$, $p=0.04$). The association between rs8076005 AA genotype-A allele (genotypes and alleles) and response was still observed more evident after correction for age, gender, baseline severity and center in the antidepressant group ($Z=2.81$, $p=0.005$; $Z=3.10$, $p=0.002$, respectively), as well as while trends were found in the whole group ($Z=2.51$, $p=0.012$; $Z=2.54$, $p=0.011$, respectively).

TPH2 rs11179023 A allele showed a mild trend of selective association with symptom improvement in the IPC group ($F=4.29$, $df=1$, $p=0.042$), but no effect was found on response and remission ~~rates in any group analyzed~~. The other investigated polymorphisms did not show evidence or trends of impact on ~~affect~~ clinical outcomes nor in individual treatment groups neither in the whole sample (Tables 2 and 3; data pertaining % improvement not shown).

4. Discussion

This study investigated the effect of five serotonergic polymorphisms on clinical outcomes of a brief psychotherapy, IPC, versus antidepressant pharmacotherapy. As previously reported in this sample [48], IPC is effective in the treatment of mild-moderate MDD as well as antidepressants.

SLC6A4 rs8076005 AA genotype and A allele were associated with response ~~rates~~ in the antidepressant group as well as in the whole sample (Table 2 and 3), with consistent trends in the whole sample (Table 2 and 3). Rs8076005 has been previously suggested as a risk factor for MDD through the neural-immune interaction, ~~particularly IL-6 levels~~ [17], even if this does not necessarily imply an impact of the SNP on antidepressant response. In detail, rs8076005 G allele carriers showed higher depressive symptom severity and higher IL-6 levels, as well as demonstrated for the minor allele of SNPs that are in linkage disequilibrium (LD) with rs8076005 (rs12150214, rs11080122, rs8071667, rs2020936, rs25528, and rs6354). Interestingly, IL-6 levels were previously demonstrated to negatively affect antidepressant response [54-56]. Two SNPs in LD with rs8076005 (rs6354 and rs2020936) were associated with anxiety, MDD and neuroticism [57].

On the other hand, rs8076005 A allele did not affect clinical outcomes in the IPC group, that instead may be influenced by *TPH2* rs11179023. Inconsistent results were found about the role of *TPH2* variants in antidepressant response [14, 20, 29, 31, 32, 58], but rs11179023 has not been investigated previously. In this study, rs11179023 A allele ~~was associated with~~ showed a trend of higher symptom improvement at week 8 only in the IPC group, while the genotypic analysis could not be performed since only 1 subject carried the AA genotype. Clearly, the present result did not allow to trace any statement but suggested a new polymorphism within a known candidate gene for further investigation ~~confirmation of the result in a larger sample is needed~~.

In the present study, *HTR2A* rs6314 was not associated with antidepressants or IPC outcomes. Despite some inconsistent results, previous studies suggested that this polymorphism may be associated with SSRIs response [14, 15, 59]. Thus, the use of different treatments in our study or the small sample size may partly explain the reported negative findings.

In this study, also *BDNF* rs6265 did not affect treatment outcomes. Recent meta-analyses reported that the heterozygous genotype may be a predictor of better antidepressant response [38-40], but negative findings

have been reported as well [60-71]. The inconsistency in previous results may be at least partly due to different rs6265 allele and genotype distribution across populations. Indeed, the frequency of the Met allele is higher in Asians than in Caucasians, and the Met allele may be protective against MDD and associated with better antidepressant response only in Asian subjects [38, 40, 72].

Phosphorylation of CREB is related to cellular resilience and neuroplasticity [73]. Phosphorylated CREB (pCREB) is positively correlated with antidepressant and interpersonal psychotherapy (IPT) response in MDD [73, 74]. Previous association studies suggested that the *CREB1* gene may show a selective effect on the risk of treatment resistance in MDD [47] and paroxetine response [71]. Our negative results are in line with other studies focused on response to mixed antidepressant treatments [61, 75].

Among the limitations of the present study, the first is the limited statistical power compared to current pharmacogenetic studies, particularly in the IPC group. Thus, the present findings should be cautiously considered. Anyway, the low sample size mainly affects the risk of false negative findings given the lack of power to find small effect sizes. Further, the choice of candidate genes with strong biological plausibility for involvement in antidepressant mechanisms of action contributes to reduce the risk of false positives. Other limitations are represented by the use of the present sample in a previous pharmacogenetic study [48] and the recruitment of patients in different centers (even if center of recruitment was used as covariate and inter-rater reliability was good). Another issue is the deviation of *SLC6A4* rs8076005 from HWE, despite the MAF was in line with that reported in the HapMap database for the CEU population (0.22 in our study and 0.18 according to HapMap Phase III release on NCBI B36 assembly, dbSNP b126). The availability of more evaluation time points could have provided information about response timing in the two treatment groups. Finally, the use of a naturalistic setting in the pharmacological arm of the study could be seen as a limitation, but also provided a perspective that is more near to the real clinical practice and the major part of patients were treated with SSRIs.

Despite these limitations, the present study is among the few that have investigated the impact of genetic variants on brief psychotherapy response. Previous studies were mainly focused on the effect of genetic variants on personality characteristics that may in turn affect psychotherapy outcome. Further genetic studies directly focused on the efficacy of psychological interventions in MDD are clearly needed. Personality features should be considered as modulators. However, in the present sample the comorbidity with personality disorders did not affect IPC outcomes [48], as found by an independent study [76].

5. Conclusion

Our study suggests that *SLC6A4* rs8076005 may impact on antidepressant response ~~rate~~ at week 8, especially in patients treated with antidepressant drugs. *TPH2* may be a selective modulator of IPC response, even if this result is even more preliminary given the limited size of the IPC-treated group and non-significance of the finding after Bonferroni correction. IPC, an effective treatment in mild-moderate MDD, is worthy of further investigation in genetic studies.

Acknowledgments

The study was funded by MIUR (2008XWR4SB).

Disclosure statement

Dr Serretti is or has been a consultant/ speaker for: Abbott, Astra Zeneca, Clinical Data, Boheringer, Bristol Myers Squibb, Eli Lilly, GlaxoSmithkline, Janssenn, Lundbeck, Pfizer, Sanofi, and Servier.

Other authors have no conflicts of interest.

References

1. Alonso J, Angermeyer MC, Bernert S, Bruffaerts R, Brugha TS, Bryson H, et al. Disability and quality of life impact of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr Scand Suppl* 2004; 420: 38-46
2. World Health Organization. The global burden of disease: 2004 update. Geneva: World Health Organization; 2008.
3. Trivedi MH, Rush AJ, Wisniewski SR, Nierenberg AA, Wardren D, Ritz L, et al. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry* 2006; 163: 28-40
4. Cuijpers P, Sibrandij M, Koole SL, Andersson G, Beekman AT and Reynolds CF 3rd. The efficacy of psychotherapy and pharmacotherapy in treating depressive and anxiety disorders: a meta-analysis of direct comparisons. *World Psychiatry* 2013; 12: 137-148
5. Unutzer J and Park M. Strategies to improve the management of depression in primary care. *Prim Care* 2012; 39: 415-431
6. Crisafulli C, Fabbri C, Porcelli S, Drago A, Spina E, De Ronchi D and Serretti A. Pharmacogenetics of antidepressants. *Front Pharmacol* 2011; 2: 6. doi: 10.3389/fphar.2011.00006
7. Porcelli S, Drago A, Fabbri C, Gibiino S, Catali R and Serretti A. Pharmacogenetics of antidepressant response. *J Psychiatry Neurosci* 2011;36: 87-113
8. Porcelli S, Fabbri C, Spina E, Serretti A and De Ronchi D. Genetic polymorphisms of cytochrome P450 enzymes and antidepressant metabolism. *Expert Opin Drug Metab Toxicol* 2011;7: 1101-1115
9. Narasimhan S and Lohoff FW. Pharmacogenetics of antidepressant drugs: current clinical practice and future directions. *Pharmacogenomics* 2012; 13: 441-464.
10. Burnet PW, Eastwood SL, Lacey K and Harrison PJ. The distribution of 5-HT1A and 5-HT2A receptor mRNA in human brain. *Brain Res* 1995;676: 157-168
11. Kusumi I, Koyama T and Yamashita I. Serotonin-induced platelet intracellular calcium mobilization in depressed patients. *Psychopharmacology* 1994; 113: 322-327
12. Tomiyoshi, R, Kamei K, Muraoka S, Muneoka K and Takigawa M. Serotonin-induced platelet intracellular Ca²⁺ responses in untreated depressed patients and imipramine responders in remission. *Biol Psychiatry* 1999; 45: 1042-1048
13. Ozaki N, Manji H, Lubierman V, Lu SJ, Lappalainen J, Rosenthal NE and Goldman D. A naturally occurring amino acid substitution of the human serotonin 5-HT2A receptor influences amplitude and timing of intracellular calcium mobilization. *J Neurochem* 1997; 68:2186-2193
14. Peters EJ, Slager SL, McGrath PJ, Knowles JA and Hamilton SP. Investigation of serotonin-related genes in antidepressant response. *Mol Psychiatry* 2004; 9:879-889

15. Wilkie MJ, Smith G, Day RK, Matthews K, Smith D, Blackwood D, Reid IC and Wolf CR . Polymorphisms in the SLC6A4 and HTR2A genes influence treatment outcome following antidepressant therapy. *Pharmacogenomics J* 2009; 9: 61-70
16. Porcelli S, Fabbri C and Serretti A. Meta-analysis of serotonin transporter gene promoter polymorphism (5-HTTLPR) association with antidepressant efficacy. *Eur Neuropsychopharmacol* 2012; 22: 239-258
17. Su S, Zhao J, Bremner JD, Miller AH, Tang W, Bouzyk M, Snieder H, Novik O, Afzal N, Goldberg J and Vaccarino V. Serotonin transporter gene, depressive symptoms and interleukin-6. *Circ Cardiovasc Genet.* 2009; 2: 614-620
18. Zill P, Banghai TC, Zwanzger P, Schule C, Eser D, Rupprecht R, Moller HJ, Bondy B and Ackenheil M. SNP and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene provide evidence for association with major depression. *Mol Psychiatry* 9: 2004; 1030-1036
19. Garriock HA, Allen JJ, Delgado P, Nahaz Z, Kling MA, Carpenter L, Burke M, Burke W, Schwartz T, Marangell LB, Husain M, Erickson RP and Moreno FA. Lack of association of TPH2 exon XI polymorphisms with major depression and treatment resistance. *Mol Psychiatry* 2005; 10: 976-977
20. Zhang X, Gainetdinov RR, Beaulieu JM, Sotnikova TD, Burch LH, Williams RB, Schwartz DA, Krishnan KR and Caron MG. Loss-of-function mutation in tryptophan hydroxylase-2 identified in unipolar major depression. *Neuron* 2005; 45: 11-16
21. Zhou Z, Roy A, Lipsky R, Kuchipudi K, Zhu G, Taubman J, Enoch MA, Virkkunen M and Goldman D. Haplotype-based linkage of tryptophan hydroxylase 2 to suicide attempt, major depression, and cerebrospinal fluid 5-hydroxyindoleacetic acid in 4 populations. *Arch Gen Psychiatry* 2005; 62: 1109-1118
22. Bicalho MA, Pimenta GJ, Neves FS, Correa H, de Moraes EN, Da Marco L and Romano-Silva MA. Genotyping of the G1463A (Arg441His) TPH2 polymorphism in a geriatric population of patients with major depression. *Mol Psychiatry* 2006; 11: 799-800
23. Delorme R, Durand CM, Betancur C, Wagner M, Ruhrmann S, Grabe HJ, Nygren G, Gillberg C, Leboyer M, Bourgeron T, Courtet P, Jollant F, Buresi C, Aubry JM, Baud P, Bondolfi G, Bertschy G, Perroud N and Malafosse A. No human tryptophan hydroxylase-2 gene R441H mutation in a large cohort of psychiatric patients and control subjects. *Biol Psychiatry* 2006; 60: 202-203
24. Van Den Bogaert A, Slegers K, De Zutter S, Heyrman L, Norrback KF, Adolfsson R, Van Broeckhoven C and Del-Favero J. Association of brain-specific tryptophan hydroxylase, TPH2, with unipolar and bipolar disorder in a Northern Swedish, isolated population. *Arch Gen Psychiatry* 2006; 63: 1103-1110
25. Gizatullin R, Zaboli G, Jonsson EG, Asperg M and Leopardi R. The tryptophan hydroxylase (TPH) 2 gene unlike TPH-1 exhibits no association with stress-induced depression. *J Affect Disord* 2008; 107: 175-179

26. Haghghi F, Bach-Mizrachi H, Huang YY, Arango V, Shi S, Dwork AJ, Rosoklija G, Sheng HT, Morozova I, Ju J, Russo JJ and Mann JJ. Genetic architecture of the human tryptophan hydroxylase 2 gene: existence of neural isoforms and relevance for major depression. *Mol Psychiatry* 2008; 13: 813-820
27. Mann JJ, Currier D, Murphy L, Huang YY, Galfalvy H, Brent D, Greenhill L, Oquendo M. No association between a TPH2 promoter polymorphism and mood disorders or monoamine turnover. *J Affect Disord* 2008; 106: 117-121
28. Illi A, Setala-Soikkeli E, Vikki M, Poutanen O, Huhtala H, Mononen N, Lehtimäki T, Leinonen E and Kampan O. 5-HTR1A, 5-HTR2A, 5-HTR6, TPH1 and TPH2 polymorphisms and major depression. *Neuroreport* 2009; 20: 1125-1128
29. Tsai SJ, Hong CJ, Liou YJ, Yu YW, Chen TJ, Hou SJ and Yen FC. Tryptophan hydroxylase 2 gene is associated with major depression and antidepressant treatment response. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33: 637-641
30. Claes S, Myint AM, Domschke K, Del-Favero J, Entrich K, Engelborghs S, De Deyn P, Mueller N, Baune B and Rothermundt M. The kynurenine pathway in major depression: haplotype analysis of three related functional candidate genes. *Psychiatry Res* 2011; 188: 355-360
31. Serretti A, Chiesa A, Porcelli S, Han C, Patkar AA, Lee SJ, Park MH and Pae CU. Influence of TPH2 variants on diagnosis and response to treatment in patients with major depression, bipolar disorder and schizophrenia. *Psychiatry Res* 2011; 189: 26-32
32. Shen X, Wu Y, Qian M, Wang X, Hou Z, Liu Y, Sun J, Zhong H, Yang J, Lin M, Li L, Liang L, Guan T, Shen Z and Yuan Y. Tryptophan hydroxylase 2 gene is associated with major depressive disorder in a female Chinese population. *J Affect Disord* 2011; 133: 619-624
33. Gao J, Pan Z, Jiao Z, Li F, Zhao G, Wei Q, Pan F and Evangelou E. TPH2 gene polymorphisms and major depression- a meta-analysis. *Plos one* 2012; 7: e36721. doi: 10.1371/journal.pone.0036721
34. Quesseveur G, David DJ, Gaillard MC, Pla P, Wu MV, Nguyen HT, Nicolas V, Auregan G, David I, Dranovsky A, Hantraye P, Hen R, Gardier AM, Deglon N and Guiard BP. BDNF overexpression in mouse hippocampal astrocytes promotes local neurogenesis and elicits anxiolytic-like activities. *Transl Psychiatry* 2013; 3: e253. doi: 10.1038/tp.2013.30
35. Aydemir O, Deveci A and Taneli F. The effect of chronic antidepressant treatment on serum brain-derived neurotrophic factor levels in depressed patients: a preliminary study. *Prog Neuropsychopharmacol Biol Psychiatry* 2005; 29: 261-265
36. Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS and Bertolino A et al. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 2003; 112: 257-269
37. Ozan E, Okur H, Eker C, Eker OD, Gonul AS and Akarsu N. The effect of depression, BDNF gene val66met polymorphism and gender on serum BDNF levels. *Brain Res Bull* 2010; 81: 61-65

38. Zou YF, Ye DQ, Feng XL, Su H, Pan FM and Liao FF. Meta-analysis of BDNF val66met polymorphism association with treatment response in patients with major depressive disorder. *Eur Neuropsychopharmacol* 2010; 20: 535-544
39. Kato M and Serretti A. Review and meta-analysis of antidepressant pharmacogenetic findings in major depressive disorder. *Mol Psychiatry* 2010; 15: 473-500
40. Niitsu T, Fabbri C, Bentini F, Serretti A. Pharmacogenetics in major depression: a comprehensive meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry* 2013; 45: 183-194
41. Mayr B and Montminy M. Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol* 2001; 2: 599-609
42. Lonze BE and Ginty DD. Function and regulation of CREB family transcription factors in the nervous system. *Neuron* 2002; 35: 605-623
43. Zubenko GS, Hughes HB, Stiffler JS, Zubenko WN and Kaplan BB. Genome survey for susceptibility loci for recurrent, early-onset, major depression: results at 10cM resolution. *Am J Med Genet* 2002; 114:413-422
44. Zubenko GS, Hughes III HB, Stiffler JS, Zubenko WN, Kaplan BB. D2S2944 identifies a likely susceptibility locus for recurrent, early-onset, major depression in women. *Mol Psychiatry* 2002; 7: 460-467
45. Rankinen T, Argyropoulos G, Rice T, Rao DC and Bouchard C. CREB1 is a strong genetic predictor of the variation in exercise heart rate response to regular exercise: the HERITAGE Family Study. *Circ Cardiovasc Genet* 2010; 3: 294-299
46. Juhasz G, Dunham JS, Mckie S, Thomas E, Downey D, Chase D, Lloyd-Williams K, Toth ZG, Platt H, Mekli K, Payton A, Elliott R, Williams SR, Anderson IM and Deakin JF. The CREB1-BDNF-NTRK2 Pathway in depression: multiple gene-cognition-environment interactions. *Biol psychiatry* 2011; 69: 762-771
47. Serretti A, Chiesa A, Calati R, Massat I, Linotte S, Kasper S, Lecrubier Y, Antonijevic I, Forray C, Snyder L, Bollen J, Zohar J, De Ronchi D, Souery D and Mendlewicz J. A preliminary investigation of the influence of CREB1 gene on treatment resistance in major depression. *J Affect Disord* 2011; 128: 56-63.
48. Serretti A, Fabbri C, Pellegrini S, Porcelli S, Politi P, Bellino S, Menchetti M, Mariotti V, Demi C, Martinelli V, Cappucciati M, Bozzatello P, Brignolo E, Brambilla P, Pae CU, Balestrieri M and De Ronchi D. No effect of Serotonergic Gene Variants on Response to Interpersonal Counseling and Antidepressants in Major Depression. *Psychiatry Investig* 2013; 10: 180-189
49. Judd FK, Piterman L, Cockram AM, McCall L and Weissman MM. A comparative study of venlafaxine with a focused education and psychotherapy program versus venlafaxine alone in the treatment of depression in general practice. *Hum Psychopharmacol* 2001; 16: 423-428
50. Oranta O, Luutonen S, Salokangas RK, Vahlberg T, Leino-Kilpi H. The outcomes of interpersonal

counselling on depressive symptoms and distress after myocardial infarction. *Nord J Psychiatry* 2010; 64:78-86

51. Weissman M. *Interpersonal Counselling for Stress and Distress in Primary Care: A Treatment Manual*. New York: Columbia University; 1988.
52. Menchetti M, Bortolotti B, Rucci P, Scocco P, Bombi A, Berardi D, et al. Depression in primary care: interpersonal counseling vs selective serotonin reuptake inhibitors. The DEPICS study. A multicenter randomized controlled trial. Rationale and design. *BMC Psychiatry* 2010; 10: 97. doi: 10.1186/1471-244X-10-97
53. Guy W. Dosage record and treatment emergent symptom scale [DOTES]. In: US Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse and Mental Health Administration, NIMH Psychopharmacology Research Branch, Division of Extramural Research Programs, editors. *ECDEU Assessment Manual for psychopharmacology Revised*. Rockville: National Institute of Mental Health; 1976. pp. 223-244.
54. Frommberger UH, Bauer J, Haselbeuer P, Fraulin A, Riemann D and Berger M. Interleukin-6-(IL-6) plasma levels in depression and schizophrenia: comparison between the acute state and after remission. *Eur Arch Psychiatry Clin Neurosci* 1997; 247: 228-233
55. Lanquillon S, Krieg JC, Bening-Abu-Shach U and Vedder H. Cytokine production and treatment response in major depressive disorder. *Neuropsychopharmacology* 2000; 22: 370-379
56. Yoshimura R, Hori H, Ikenouchi-Sugita A, Uneme-Nakano W, Ueda N and Nakamura J. Higher plasma interleukin-6 (IL-6) level is associated with SSRI- or SNRI-refractory depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33: 722-726
57. Wray NR, James MR, Gordon SD, Dumenil T, Ryan L, Coventry WL, Statham DJ, Pergadia ML, Madden PA, Heath AC, Montgomery GW and Martin NG. Accurate, Large-Scale Genotyping of 5HTTLPR and Flanking Single Nucleotide Polymorphisms in an Association Study of Depression, Anxiety and Personality Measures. *Biol Psychiatry* 2009; 66: 468-476
58. Tzvetkov MV, Brockmoller J, Roots I, Kirchheiner J. Common genetic variations in human brain-specific tryptophan hydroxylase-2 and response to antidepressant treatment. *Pharmacogenet Genomics* 2008; 18: 495-506
59. Minov C, Baghai TC, Schule C, Zwanzger P, Schwarz MJ, Zill P, Rupprecht R and Bondy B. Serotonin-2A-receptor and -transporter polymorphisms: lack of association in patients with major depression. *Neurosci Lett* 2001; 303:119-122
60. Tsai SJ, Cheng CY, Yu YW, Chen TJ, Hong CJ. Association study of a brain-derived neurotrophic-factor genetic polymorphism and major depressive disorders, symptomatology, and antidepressant response. *Am J Med Genet B Neuropsychiatr Genet* 2003; 123B: 19-22
61. Wilkie MJ, Smith D, Reid IC, Day RK, Matthews K, Wolf CR, Blackwood D and Smith G. A splice site polymorphism in the G-protein beta subunit influences antidepressant efficacy in

- depression. *Pharmacogenet Genomics* 2007; 17: 207-215
62. Kang RH, Chang HS, Wong ML, Choi MJ, Park JY, Lee HY, Jung IK, Joe SH, Kim L, Kim SH, Kim YK, Han CS, Ham BJ, Lee HJ, Ko YU, Lee MS and Lee MS. Brain-derived neurotrophic factor gene polymorphisms and mirtazapine responses in Koreans with major depression. *J Psychopharmacol* 2010; 24: 1755-63
 63. Licinio J, Dong C, Wong ML. Novel sequence variations in the brain-derived neurotrophic factor gene and association with major depression and antidepressant treatment response. *Arch Gen Psychiatry* 2009; 66: 488-497
 64. Domschke K, Lawford B, Laje G, Berger K, Young R, Morris P, Deckert J, Arolt V, McMahon FJ and Baune BT. Brain-derived neurotrophic factor (BDNF) gene: no major impact on antidepressant response. *Int J Neuropsychopharmacol* 2010; 13: 93-101
 65. Kocabas NA, Antonijevic I, Faghel C, Forray C, Kasper S, Lecrubier Y, Linotte S, Massat I, Mendlewicz J, Noro M, Montgomery S, Oswald P, Snyder L, Zohar J and Souery D. Brain-derived neurotrophic factor gene polymorphisms: influence on treatment response phenotypes of major depressive disorder. *Int clin Psychopharmacol* 2011; 26: 1-10
 66. Yoshimura R, Kishi T, Suzuki A, Uneme-Nakano W, Ikenouchi-Sugita A, Hori H, Otani K, Iwata N and Nakamura J. The brain-derived neurotrophic factor (BDNF) polymorphism val66met is associated with neither serum BDNF level nor response to selective serotonin reuptake inhibitors in depressed Japanese patients. *Prog Neuropsychopharmacol Bio Psychiatry* 2011; 35: 1022-1025
 67. Musil R, Zill P, Seemuller F, Bondy B, Obermeier M, Spellmann I, Bender W, Adli M, Heuser I, Zeiler J, Gaebel W, Maier W, Rietschel M, Rujescu D, Schennach R, Moller HJ and Riedel M. No influence of brain-derived neurotrophic factor (BDNF) polymorphisms on treatment response in a naturalistic sample of patients with major depression. *Eur Arch Psychiatry Clin Neurosci* 2013 ;263:405-12 .
 68. Tsuchimine S, Yasui-Furukori N, Nakagami T, Sato Y and Kaneko S. DRD3 but not BDNF, genotype affects treatment response to paroxetine in major depressive disorder: a preliminary study. *J Clin Psychopharmacol* 2012; 32: 724-726
 69. Xu G, Lin K, Rao D, Dang Y, Ouyang H, Mai G and Zhang M. Brain-derived neurotrophic factor gene polymorphism (Val66Met) and the early response to antidepressant in Chinese Han population. *Psychiatr Genet* 2012; 22: 214-215
 70. Brunoni AR, Kemp AH, Shiozawa P, Cordeiro Q, Valiengo LC, Goulart AC, Coprerski B, Lotufo PA, Brunoni D, Perez AB, Fregni F and Bensenor IM (2013) Impact of 5-HTTLPR and BDNF polymorphisms on response to sertraline versus transcranial direct current stimulation: Implications for the serotonergic system. *Eur Neuropsychopharmacol* 2013; 23:1530-40.
 71. Murphy GM Jr, Sarginson JE, Ryan HS, O'Hara R, Schatzberg AF and Lazzaroni LC. BDNF and CREB1 genetic variants interact to affect antidepressant treatment outcomes in geriatric depression.

Pharmacogenet Genomics 2013; 23:301-313

72. Verhagen M, van der Meij A, van Deurzen PAM, Janzing JGE, Arias-Vasquez A, Buitelaar JK and Franke B. Meta-analysis of the BDNF Val66Met polymorphism in major depressive disorder: effects of gender and ethnicity. *Mol Psychiatry* 2010; 15: 260-271
73. Koch JM, Hinze-Selch D, Stingele K, Huchzermeier C, Goder R, Seeck-Hirschner M and Aldenhoff JB. Changes in CREB Phosphorylation and BDNF plasma levels during psychotherapy of depression. *Psychother Psychosom* 2009; 78:187-192
74. Koch JM, Kell S, Hinze-Selch D and Aldenhoff JB. Changes in CREB- phosphorylation during recovery from major depression. *J Psychiatr Res* 2002; 36:369-375
75. Crisafulli C, Shim DS, Andrisano C, Pae CU, Chiesa A, Han C, Patkar AA, Lee SJ, Serretti A and De Ronchi D. Case-control association study of 14 variants of CREB1, CREBBP and CREM on diagnosis and treatment outcome in major depressive disorder and bipolar disorder. *Psychiatry Res* 2012; 198: 39-46
76. Jorgensen AL, Williamson PR. Methodological quality of pharmacogenetic studies: issues of concern. *Stat Med* 2008; 27: 6547-69.
77. Levenson JC, Wallace ML, Fournier JC, Rucci P and Frank E. The role of personality pathology in depression treatment outcome with psychotherapy and pharmacotherapy. *J Consult Clin Psychol* 2012; 80: 719-729