

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Identification of fungal ene-reductase activity by means of a functional screening

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/152111> since 2016-06-29T15:42:23Z

*Published version:*

DOI:10.1016/j.funbio.2015.01.006

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *FUNGAL BIOLOGY*, 119, 2015, 10.1016/j.funbio.2015.01.006.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.funbio.2015.01.006

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S1878614615000173>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/152111>

1 **Identification of fungal ene-reductase activity by means of a functional screening.**

2 Alice Romagnolo<sup>a</sup>, Federica Spina<sup>a</sup>, Elisabetta Brenna<sup>b</sup>, Michele Crotti,<sup>b</sup> Fabio

3 Parmeggiani<sup>b</sup> and Giovanna Cristina Varese<sup>a\*</sup>

4 <sup>a</sup> Department of Life Science and Systems Biology, University of Turin, viale P. A.

5 Mattioli 25, 10125 Turin, Italy

6 <sup>b</sup> Department of Chemistry, Materials and Chemical Engineering “G. Natta”,

7 Politecnico di Milano, via L. Mancinelli 7, 20131 Milan, Italy

\*Corresponding author (G.C. Varese)

Tel.: +39 0116705984;

Fax: +39 0116705962

E-mail address: [cristina.varese@unito.it](mailto:cristina.varese@unito.it)

Postal address: viale P.A. Mattioli 25, 10125 Turin, Italy

8 E-mail: [alice.romagnolo@unito.it](mailto:alice.romagnolo@unito.it); [federica.spina@unito.it](mailto:federica.spina@unito.it); [elisabetta.brenna@polimi.it](mailto:elisabetta.brenna@polimi.it);

9 [michele.crotti@chem.polimi.it](mailto:michele.crotti@chem.polimi.it); [cristina.varese@unito.it](mailto:cristina.varese@unito.it)

10

11 **Abstract**

12 Bioeconomy stresses the need of green processes promoting the development of  
13 new methods for biocatalyzed alkene reductions.

14 A functional screening of 28 fungi belonging to Ascomycota, Basidiomycota  
15 and Zygomycota isolated from different habitats was performed. Their capability to  
16 reduce C=C double bonds was evaluated towards three substrates (cyclohexenone,  $\alpha$ -  
17 methylnitrostyrene and  $\alpha$ -methylcinnamaldehyde) with different electron-withdrawing  
18 groups, i.e., ketone, nitro and aldehyde, respectively.

19 Almost all the fungi showed this reducing activity. Noteworthy *Gliomastix*  
20 *masseei*, *Mucor circinelloides* and *Mucor plumbeus* resulted very versatile and  
21 effective, being able to reduce all the model substrates quickly and with high yields.

22 **Keywords**

23 Biocatalysis, filamentous fungi, ene-reductases,  $\alpha,\beta$ -unsaturated compounds,  
24 bioreduction.

25 **1. Introduction**

26 Nowadays, the synthesis of molecules with biotechnological exploitations is  
27 mainly done by traditional chemical processes, which generally have high costs and  
28 environmental impact. The growing awareness about safety problems brought to  
29 restrain the use of chemical catalysts as heavy metals or unsafe gasses that require harsh  
30 working conditions in terms of temperature and pressure (Faber, 2011).

31 On the other hand, biocatalysts seem to be a viable alternative to traditional  
32 methods for the minimal environmental impact due to the low energy demand, waste  
33 and by-products formations and for the reduced process costs. Moreover, biocatalysis is  
34 a powerful tool to obtain chiral molecules in enantiomerically pure form, which are  
35 highly valued for instance in the pharmaceutical field (Soartet and Vandamme, 2010).

36 The reduction of C=C double bonds conjugated with different electron-  
37 withdrawing groups (EWG) such as carbonyl, nitro and ester can be catalyzed by Ene-  
38 Reductases (E.C. 1.6.99.1, ERs). Most of the known ERs are flavin-dependent  
39 oxidoreductases belonging to the Old Yellow Enzyme family, which require NAD(P)H  
40 as cofactor (Stuermer et al., 2007). They were first described in *Saccharomyces*  
41 *pastorianus* (Stott et al., 1993) and *S. cerevisiae* (Karplus et al., 1995); in the following  
42 years, many ERs were described in other yeasts, bacteria, plants and animals, but still  
43 little is known about their occurrence in filamentous fungi (Stuermer et al., 2007). To  
44 date, these enzymes were poorly investigated at molecular and structural level in these  
45 microorganisms. These studies are complex because only a few genomes are completely  
46 sequenced and hence it is very difficult to investigate the presence of genes coding for  
47 ERs in filamentous fungi. Moreover, no information is available relating to the structure

48 of these proteins in these microorganisms and to date no ERs have been purified and  
49 characterized from this source. Otherwise, for some yeasts, bacteria and plants the  
50 presence of ERs at molecular level, their structure and their characterization have been  
51 extensively investigated. Their biological role is still unknown although some authors  
52 suggested their involvement in the stress response (Brigè et al., 2006).

53 Several authors described the capability of filamentous fungi to reduce the C=C  
54 double bonds of a single substrate or of a set of compounds belonging to the same  
55 structural class (Arnone et al., 1990; Fuganti et al., 1998 a; Hall et al., 2006;  
56 Skrobiszewski et al., 2013). To date, the main functional screening of filamentous fungi  
57 was performed by Carballeira et al. (2004). Among the 241 fungi, only 3 were capable  
58 to reduce the C=C double bond of carvone.

59 Considering the natural biodiversity and the broad heterogeneous enzymatic  
60 pattern, filamentous fungi are indeed excellent biocatalysis agents. Actually, some  
61 strains or their enzymes are formerly used in the production of building blocks of  
62 pharmaceuticals, agrochemicals or fragrances (Colwell, 2002; Gavrilescu and Chisti,  
63 2005). On the whole, there is a strong need to identify potential biocatalysts to enlarge  
64 the portfolio of microorganisms and enzymes to be used for synthetic applications.

65 Regarding ERs, a validated enzymatic assay has not yet been developed, to date.  
66 So far, the only available method is related to the oxidation of the NAD(P)H by purified  
67 enzymes (Gao et al., 2012). Its applicability is limited because many unsaturated  
68 substrates commonly used in biocatalysis absorb at the same wavelength of the cofactor.  
69 Since it is not possible to verify directly the presence of these enzymes in fungi, either  
70 with molecular methods or enzymatic assays, a screening to identify the products of the  
71 reaction that probably involved ERs is a convenient approach. In addition, this method  
72 also allows to analyze a wide biodiversity of microorganisms.

73 Data comparison to literature is difficult due to the few model substrates used to  
74 investigate ERs activity, since this enzymatic activity has been poorly deepened. Few  
75 studies take into consideration more than one compound. For example, Goretti et al.  
76 (2011) described a whole-cell system of non-conventional yeasts in the bioconversion  
77 of  $\alpha,\beta$ -unsaturated ketones and aldehydes.

78 The present study aims to identify filamentous fungi showing ER activity. A  
79 functional screening was set up using 28 fungi belonging to Ascomycota,  
80 Basidiomycota and Zygomycota, isolated from different habitats. Three representative  
81 model substrates characterized by different EWGs (ketone, nitro and aldehyde)  
82 conjugated with the C=C double bond were selected. The reduction of the C=C double  
83 bonds was followed by GC/MS analysis.

## 84 **2. Materials and Methods**

### 85 2.1 Fungi

86 The fungi belong to different physiological and taxonomical groups and were  
87 isolated from different habitats (Table 1). They are preserved at the *Mycotheca*  
88 *Universitatis Taurinensis* (MUT, Department of Life Sciences and Systems Biology,  
89 University of Turin).

### 90 2.2 Chemicals

91 Cyclohexenone (CE) and  $\alpha$ -methylcinnamaldehyde (MCA) were purchased from  
92 Sigma-Aldrich (Italy). (*E*)- $\alpha$ -methylnitrostyrene (MNS) was synthesized according to  
93 the literature (Kawai et al., 2001).

94 Stock solutions (500 mM) of each substrate were prepared by dissolving them in  
95 dimethyl sulfoxide (DMSO).

### 96 2.3 Biotransformation experiments

97 Fungal strains were pre-grown in Petri dishes containing malt extract solid  
98 medium (MEA: 20 g/l glucose, 20 g/l malt extract, 20 g/l agar, 2 g/l peptone) from

99 which the fungal inoculum for liquid cultures was set up. When possible, a conidia  
100 suspension was prepared ( $1 \cdot 10^6$  conidia final concentration in flask). Otherwise, the  
101 inoculum was made by homogenizing agar squares derived from the margins of an  
102 overgrown colony together with sterile water ( $1 \text{ cm}^2/\text{ml}$ ). Fungi were inoculated in 50  
103 ml flasks containing 30 ml of malt extract liquid medium. Flasks were incubated at 25  
104 °C and were maintained in agitation (110 rpm) in the dark.

105 After two days of pre-growth, the substrates were separately added (5 mM final  
106 concentration). For each substrate, three biological replicates were run.

107 The experiment was run for 7 days: 1 ml of cultural broth was collected after 2,  
108 4 and 7 days and extracted by two-phase separation using 0,5 ml of methyl *t*-butyl ether  
109 (MTBE) as solvent. The organic phases were dried over anhydrous  $\text{Na}_2\text{SO}_4$  and  
110 analyzed by means of GC/MS.

111 After two days, one flask for each fungus was sacrificed to measure the initial  
112 biomass and pH. Those parameters were also evaluated for all the flasks at the end of  
113 the experiment. The liquid was separated from the biomass by filtration and was used  
114 for pH measurement. The mycelia were dried at 60 °C for 24 h to measure the  
115 biomasses dry weight.

#### 116 2.4 GC/MS analysis

117 GC/MS analyses were performed on an Agilent HP 6890 gas chromatograph  
118 equipped with a 5973 mass detector and an HP-5-MS column ( $30 \text{ m} \times 0.25 \text{ mm} \times 0.25$   
119  $\mu\text{m}$ , Agilent), employing the following temperature program: 60 °C (1 min) / 6 °C  $\text{min}^{-1}$   
120 / 150 °C (1 min) / 12 °C  $\text{min}^{-1}$  / 280 °C (5 min). GC retention times: cyclohexenone  
121 (CE) 5.40 min, cyclohexanone (CO) 4.65 min, cyclohexanol (COH) 4.45 min, (*E*)- $\alpha$ -  
122 methylnitrostyrene (MNS) 17.7 min, (*Z*)- $\alpha$ -methylnitrostyrene (MNS) 15.6 min, 2-  
123 nitropropylbenzene (NPB) 14.8 min,  $\alpha$ -methylcinnamaldehyde (MCA) 14.7 min,  $\alpha$ -  
124 methylcinnamyl alcohol (MCOH) 15.5 min,  $\alpha$ -methyl dihydrocinnamyl alcohol

125 (MDHCOH) 13.6 min. The enantiomeric excess (ee) values of MDHCOH was  
126 determined by GC analysis, performed using a Chirasil Dex CB column (0.25  $\mu\text{m} \times$   
127 0.25 mm  $\times$  25 m, Varian), according to the following conditions: 60  $^{\circ}\text{C} / 5^{\circ}\text{C min}^{-1} / 95$   
128  $^{\circ}\text{C}$  (25 min) / 50  $^{\circ}\text{C min}^{-1} / 220^{\circ}\text{C}$  (10 min). GC retention times: (*R*)-enantiomer 26.6  
129 min, (*S*)-enantiomer 27.9 min.

### 130 **3. Results and Discussion**

131 The results of the biotransformation of the three substrates by the 28 fungi are  
132 shown in Table 2, the maximal percentage of C=C double bonds reduction and the  
133 timing of the reaction are reported. Four groups were established on account of the rate  
134 of substrates transformation by means of a putative ER activity: group A 100-75 %;  
135 group B 74-50 %; group C 49-25 % and group D 24-0 %.

136 Regarding pH measurements, the cultural broth of each fungus remained  
137 unchanged during the experiments. The pH values ranged between 3 and 6 and seemed  
138 to depend on the metabolism of each fungus; variations due to the addition of substrates  
139 were not detected.

#### 140 3.1 CE biotransformation

141 Almost all the fungi (96,4 %) were able to reduce the C=C double bond of CE  
142 among which 75 % could be listed in group A. The other fungi were 3,6 % in group B,  
143 11 % in group C and 11 % in group D. As it can be seen in Table 2, 11 out of 28 fungi  
144 were able to completely transform this substrate within two days (*C. herbarum*, *G.*  
145 *masseei*, *P. citrinum*, *S. fimicola*, *T. viride*, *A. cylindracea*, *A. splendida*, *Coprinellus*  
146 *sp.*, *T. pubescens*, *M. circinelloides*, *M. plumbeus*). On the whole, the majority of the  
147 fungi were not only able to reduce the C=C double bond of this substrate but also  
148 reached the complete biotransformation of this molecule.

149 CE is a well-accepted substrate; in this study only one fungus out of 28 was  
150 ineffective towards this compound. Ketonic substrates have been frequently considered



151 in the literature; for example, Gatti et al. (2014) discussed several ketone substrates such  
152 as carvone or ketoisophorone used in bioconversions that involved ERs.

153 The results obtained in this study may be compared with literature. In particular,  
154 two fungi belonging to the genus *Mucor* were very effective toward CE confirming the  
155 results obtained by Fuganti and Zucchi (1998 b).

156 Other authors analyzed the biotransformation of molecules with ketonic EWG  
157 with a basic scaffold similar to CE. Skrobiszewski et al. (2013) described a strain of *P.*  
158 *ostreatus* effective towards C=C double bonds reduction; this data was confirmed also  
159 by the strain used in this study. *Absidia glauca* and *Beauveria bassiana* were poorly  
160 effective towards C=C double bonds reduction of CE while the strains used by  
161 Carballeira et al. (2004) and Fuganti and Zucchi (1998 b) reduce ketonic substrate with  
162 high yield in benzalacetones derivatives.

163 CE biotransformation led to the identification of two products (Figure 1):  
164 cyclohexanone (CO) in which ER activity is involved in the reduction of C=C double  
165 bond, and cyclohexanol (COH) in which an alcohol dehydrogenase (ADH) reduces the  
166 C=O bond. Most of the fungi (67 %) convert CE into COH showing the action of ERs  
167 and ADHs, while 26 % preferentially reduced the C=C double bond, producing only  
168 CO. When the reduction was slow, it was possible to define a putative reaction profile  
169 in which the two enzymes act in cascade (Figure 2). In most cases, the reaction was  
170 very fast and only the formation of COH was detected.

171 The same reaction profile was hypothesized by other authors in the reduction of  
172 analogous substrates of CE (Fuganti et al., 1998 a; Fuganti and Zucchi, 1998 b;  
173 Carballeira et al., 2004; Hall et al., 2006; Stuermer et al., 2007; Skrobiszewski et al.,  
174 2013).

175 3.2 MNS biotransformation

176 The C=C double bond of this substrate was reduced by 82 % of the fungi (Table  
177 2), among which 14 % could be listed in group A. The other fungi were 7,1 % in group  
178 B, 25 % in group C, and the majority (53,6 %) in group D. As it can be seen in Table 2,  
179 two fungi out of 28 (*A. niger* and *M. circinelloides*) were very active and reached an  
180 almost complete conversion of the substrate within 2 days.

181 MNS biotransformation led to the identification of 2-nitropropylbenzene (NPB)  
182 as the sole product through the reduction of a C=C double bond by ER activity (Figure  
183 3). Since the stereogenic center of the reduced product is too labile under the reaction  
184 conditions no reliable ee values could be obtained.

185 Although nitrostyrene derivatives are good substrates for ERs (Toogood et al.,  
186 2008; Gatti et al., 2014), to our knowledge, this is the first report of the reduction of the  
187 C=C double bonds of nitroalkenes by filamentous fungi. A whole-cell system using  
188 *Saccharomyces cerevisiae* in the reduction of the C=C double bond of MNS was  
189 reported by Kawai et al. (2001). These authors obtained yields comparable to the fungi  
190 gathered in group A.

### 191 3.3 MCA biotransformation

192 The C=C double bond of this substrate was reduced by 35,7 % of the fungi  
193 (Table 2) among which only 7,1 % in group A. The other fungi were 3,6 % in group B  
194 and 89,3 % in group D. Two fungi, *M. circinelloides* and *M. plumbeus*, completely  
195 converted MCA within 2 days into the (*S*)-enantiomer of the corresponding saturated  
196 alcohol MDHCOH, showing an ee value of 80 %. This result is promising compared to  
197 literature: Fronza et al. (2009) reported a conversion rate of 12 % and ee value of 70 %  
198 ((*S*)- enantiomer) in the reduction of MCA with *S. cerevisiae* whole-cell.

199 The difficulty to reduce MCA has been also found by other authors. For example  
200 Goretti et al. (2011) screened non-conventional yeasts but only *Kazachstania*  
201 *spencerorum* out of 23 microorganisms was able to convert this substrate (60 %).

202           The MCA biotransformation led to the identification of two products, probably  
203 involving two enzymes (Figure 4). The reduction of C=C and C=O double bonds led to  
204 the formation of  $\alpha$ -methyl dihydrocinnamyl alcohol (MDHCOH) by means of ERs and  
205 ADHs. The reduction of the aldehyde EWG by ADHs before the C=C reduction could  
206 take place, led to the formation of  $\alpha$ -methylcinnamyl alcohol (MCOH). Since the  
207 MCOH lacks the necessary EWG, it is not a substrate for ERs and consequently  
208 accumulates in the medium. In our experiments, two fungi (*A. glauca* and *E. nigrum*)  
209 formed MCOH showing only ADH activity. By contrast, 8 fungi formed a mixture of  
210 MCOH and MDHCOH, preferentially reducing the aldehydic group. This reaction  
211 profile was noticed also by other authors for *S. cerevisiae* (Gatti et al. 2014).

#### 212 3.4 General considerations

213           The screening clearly showed that the EWG on the C=C double bonds strongly  
214 influenced the reaction rate of the various strains tested. Probably the catalytic activity  
215 of the enzymes was also affected by the steric hindrance of the other substituents and by  
216 electronic effects, as widely discussed by Stuermer et al. (2007) and Gatti et al. (2014).  
217 In detail, the substrates were transformed following this outline: CE>MNS>MCA. CE  
218 was the most easily converted substrate, due to the presence of a strong EWG (ketone)  
219 and only two substituents on the double bond, both with a modest steric hindrance.  
220 MNS and MCA share the same basic scaffold, with higher steric hindrance due to the  
221 aromatic ring, but they differ in the EWGs (nitro and aldehyde, respectively): the higher  
222 conversion of MNS can be justified by the much higher electron-withdrawing power of  
223 the nitro group.

224           It has to be noticed that this enzymatic activity appears to be genus specific. For  
225 instance, the strains of *Mucor* reached the same conversion yields in the  
226 biotransformation of all the substrates. Whereas, the strains of *Penicillium* behave  
227 differently towards the substrates analyzed (Table 2). This consideration makes the

228 screening for the selection of strains for definite reactions a required step for the  
229 analysis of the intraspecific variability.

230 The fungi afforded very different yields in the conversion of these substrates. This  
231 may not only be due to the different affinity of the enzyme for the substrate but also to  
232 an activation of the secondary metabolism of the fungus that would lead to the  
233 production of putative ERs in the presence of different substrates.

### 234 3.5 Biomasses

235 On the whole, all fungi were able to grow with the substrates. In some cases  
236 weight differences have been reported in the three biological replicates despite the  
237 biomass seems to be very similar morphologically. For this reason, some standard  
238 deviations were high. In particular, as it can be seen in Figure 5 each substrate had a  
239 different effect on the biomass production. It was not possible to measure the biomass  
240 of the yeast *G. cucujoidarum*. The growth of this fungus was measured by cell counting  
241 using a Burker chamber and the growth was about  $3,6 \cdot 10^8$  cells/ml in each of the  
242 replicates.

243 Interestingly, CE seems to stimulate the production of biomass in almost all the  
244 fungi. For almost all the fungi, the presence of MNS seems to cause a scarce  
245 development of the biomass. In fact, at the end of the experiment they were lower than  
246 the initial growth. For example, *P. citrinum* and *C. funicola* displayed a biomass loss of  
247 80 % suggesting that this compound may inhibit the biomass growth. Despite this  
248 strong biomass decrease, many fungi reached very high conversion yields.

249 The addition of MCA partially inhibited the development of biomass in almost all  
250 the fungi. The fungus that displayed the highest biomass loss (– 91 %) was *C. funicola*.

251 In general, the biomass growth is an important parameter that most often is not  
252 taken into account by other authors, reason for which it is not easy to make comparisons  
253 with other studies. Regarding CE, a correlation between biomass and ER activity is not

254 possible because the majority of the fungi increased the biomass independently of the  
255 biotransformation yield. For example, *A. glauca* displayed a biomass loss of 40 % but  
256 reached the total biotransformation of CE. On the other hand, *B. bassiana* displayed a  
257 growth of over 80 %, although it was not able to convert this substrate.

258 Even in the case of MNS and MCA a proper correlation between the biomass  
259 growth and the biotransformation yields was not possible. In particular, for these  
260 compounds the majority of the fungi displayed a decrease in biomass. Nonetheless, the  
261 fungi that reached the highest conversion rate had an important biomass loss. For  
262 example *M. circinelloides* almost totally converted MNS but during the trial this fungus  
263 had a weight loss of 40 %. Similarly, *M. plumbeus* almost totally converted MCA and  
264 had a biomass decrease of 45 %. The few fungi that increased the biomass showed two  
265 behaviors: either partial conversion of the two substrates with low yields or  
266 ineffectiveness towards these compounds.

267 Since the biomass development implicates the primary metabolism, and a  
268 correlation between the biomass development and the biotransformation yields cannot  
269 be drawn, the involvement of the secondary metabolism in the conversion of CE, MNS  
270 and MCA may be hypothesized.

#### 271 **4. Conclusions**

272 The screening highlighted that ER activity is widespread in filamentous fungi. In  
273 fact, 27 out of 28 microorganisms reduced at least one substrate. Consequently,  
274 although the biological role of ER is still unknown, this activity may be involved in the  
275 secondary metabolism of the microorganisms analyzed.

276 *M. circinelloides*, *M. plumbeus* and *G. marseeii* resulted the most versatile strains  
277 converting all the substrates analyzed, with the highest yields. Moreover, this study also  
278 highlighted problems related to substrate selection: by now, several chemical classes

279 have been investigated, but these activities are difficult to compare due to the lack of  
280 validated model compounds.

## 281 **References**

282 **Arnone A, Cardillo R, Nasini G, Vajna de Pava O, 1990.** Enantioselective reduction  
283 of racemic abscisic acid by *Aspergillus niger* cultures. J. Chem. Soc., Perkin Trans. 1:  
284 3061-3063.

285 **Brigé A, Van den Hemel D, Carpentier W, De Smet L, Van Beeumen JJ, 2006.**  
286 Comparative characterization and expression analysis of the four Old Yellow Enzyme  
287 homologues from *Shewanella oneidensis* indicate differences in physiological function.  
288 Biochem. J. 394: 335-344.

289 **Carballeira JD, Valmaseda M, Alvarez E, Sinisterra-Gago JV, 2004.** *Gongronella*  
290 *butleri*, *Schizosaccharomyces octosporus* and *Diplogelasinospora grovesii*: novel  
291 microorganisms useful for the stereoselective reduction of ketones. Enz. Microb.  
292 Technol. 34: 611-623.

293 **Colwell RR, 2002.** Fulfilling the promise of biotechnology. Biotechnol. Adv. 20: 215–  
294 228.

295 **Faber K, 2011.** Biotransformations in organic chemistry. A textbook. Springer-Verlag,  
296 Berlin, Heidelberg 2011.

297 **Fronza G, Fuganti C, Serra S, 2009.** Stereochemical course of baker's yeast mediated  
298 reduction of the tri- and tetrasubstituted double bonds of substituted cinnamaldehydes.  
299 Eur. J. Org. Chem. 6160-6171

300 **Fuganti C, Minut J, Pedrocchi Fantoni G, Servi S, 1998 a.** On the microbial  
301 transformation of  $\alpha,\beta$ -unsaturated aryl ketones by the fungus *Beauveria bassiana*. J.  
302 Mol. Catal. B: Enzym. 4: 47-52.

303 **Fuganti C, Zucchi G, 1998 b.** Product distribution in the microbial biogeneration of  
304 raspberry ketone from 4-hydroxybenzalacetone. J. Mol. Catal. B: Enzym. 4: 289-293.

305 **Gao X, Ren J, Qiaqing W, Dunming Z, 2012.** Biochemical characterization and  
306 substrate profiling of a new NADH-dependent enoate reductase from *Lactobacillus*  
307 *casei*. *Enz. Microb. Technol.* 51: 26-34.

308 **Gavrilescu M, Chisti Y, 2005.** Biotechnology a sustainable alternative for chemical  
309 industry. *Biotech. Adv.* 23: 471-499.

310 **Gatti FG, Parmeggiani F, Sacchetti A, 2014.** Synthetic strategies based on C=C  
311 bioreductions for the preparation of biologically active molecules, in: “Synthetic  
312 methods for biologically active molecules – Exploiting the potential of bioreductions”,  
313 Brenna E. (Ed.), Wiley-VCH, Weinheim.

314 **Goretti M, Ponzoni C, Caselli E, Marchegiani E, Cramarossa MR, Turchetti B,**  
315 **Forti L, Buzzini P, 2011.** Bioreduction of  $\alpha$ ,  $\beta$ -unsaturated ketones and aldehydes by  
316 non-conventional yeast (NCY) whole-cell. *Biores. Technol.* 102: 3993-3998.

317 **Hall M, Hauer B, Stuermer R, Kroutil W, Faber K, 2006.** Asymmetric whole-cell  
318 bioreduction of an  $\alpha$ ,  $\beta$ -unsaturated aldehyde (citral): competing prim-alcohol  
319 dehydrogenase and C–C lyase activities. *Tetrahedron: Asymmetry* 17: 3058-3062.

320 **Karplus PA, Fox KM, Massey V, 1995.** Structure-function relations for old yellow  
321 enzyme. *FASEB J.* 9: 1518-1526.

322 **Kawai Y, Inaba Y, Tokitoh N, 2001.** Asymmetric reduction of nitroalkenes with  
323 baker’s yeast, *Tetrahedron: Asymmetry* 12: 309-318.

324 **Skrobiszewski A, Ogorek R, Plaskowska E, Gladkowski W, 2013.** *Pleurotus*  
325 *ostreatus* as a source of enoate reductase. *Biocat. Agric. Biotech.* 2: 26-31.

326 **Soartet W, Vandamme EJ, 2010.** Industrial biotechnology in the chemical and  
327 pharmaceutical industries. In: *Industrial biotechnology. Sustainable growth and*  
328 *economic success.* Wiley-VCH, Weinheim.

329 **Stott K, Saito K, Thiele DJ, Massey V, 1993.** Old yellow enzyme. The discovery of  
330 multiple isozyme and a family of related proteins. *J. Biol. Chem.* 268: 6097-6106.

331 **Stuermer R, Hauer B, Hall M, Faber K, 2007.** Asymmetric bioreduction of activated  
332 C=C bonds using enoate reductases from the old yellow enzyme family. *Curr. Opin.*  
333 *Chem. Biol.* 11: 203-213.

334 **Toogood HS, Fryzkowska A, Hare V, Fisher K, Roujeinikova A, Leys D,**  
335 **Gardiner GM, Stephens JM, Scrutton NS, 2008.** Structure-based insight into the  
336 asymmetric bioreduction of the C=C double bond of  $\alpha,\beta$ -unsaturated nitroalkenes by  
337 pentaerythritol tetranitrate reductase. *Adv. Synth. Catal.* 350: 2789-2803.

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355



356 **Table 1:** list of the strains analyzed during the screening and their isolation site (MUT:  
 357 accession number).

| <b>Fungi</b>    | <b>MUT</b>                | <b>Species</b>                   | <b>Isolation site</b>                                 |
|-----------------|---------------------------|----------------------------------|---|
| <b>Asco-</b>    | 3874                      | <i>Aspergillus niger</i>         | air   |
|                 | 1720                      | <i>Beauveria bassiana</i>        | air   |
|                 | 1087                      | <i>Botrytis cinerea</i>          | fresco of Botticelli                                  |
|                 | 3726                      | <i>Chaetomium funicola</i>       | dried <i>Boletus</i> fungi from Europe                |
|                 | 3856                      | <i>Cladosporium herbarum</i>     | air   |
|                 | 3848                      | <i>Epicoccum nigrum</i>          | air   |
|                 | 4824                      | <i>Geotrichum cucujoidarum</i>   | wastewater of a tanning industry                      |
|                 | 4855                      | <i>Gliomastix masseei</i>        | <i>Flabelia petiolata</i> (marine algae)              |
|                 | 281                       | <i>Mesobotryis simplex</i>       | cultivated soil                                       |
|                 | 1749                      | <i>Myxotrichum deflexum</i>      | air   |
|                 | 1381                      | <i>Oidiodendron maius</i>        | roots of <i>Vaccinium myrtillus</i> (black raspberry) |
|                 | 4862                      | <i>Penicillium citrinum</i>      | <i>Flabelia petiolata</i> (marine algae)              |
|                 | 4831                      | <i>Penicillium purpurogenum</i>  | wastewater of a tanning industry                      |
|                 | 4892                      | <i>Penicillium vinaceum</i>      | <i>Padina pavonica</i> (marine algae)                 |
|                 | 4833                      | <i>Scopulariopsis</i> sp.        | wastewater of a tanning industry                      |
|                 | 1148                      | <i>Sordaria fimicola</i>         | <i>Picea abies</i> (norway spruce)                    |
|                 | 1166                      | <i>Trichoderma viride</i>        | tallus of <i>Parmelia taractica</i> (lichen)          |
| 3788            | <i>Trichurus spiralis</i> | book pages                       |   |
| <b>Basidio-</b> | 2753                      | <i>Agrocybe cylindracea</i>      | carpophore  |
|                 | 2755                      | <i>Agrocybe farinacea</i>        | carpophore  |
|                 | 3696                      | <i>Agrocybe splendida</i>        | carpophore  |
|                 | 4897                      | <i>Coprinellus</i> sp.           | <i>Padina pavonica</i> (marine algae)                 |
|                 | 2976                      | <i>Pleurotus ostreatus</i>       | carpophore on <i>Populus</i> sp. (poplar)             |
|                 | 2400                      | <i>Trametes pubescens</i>        | carpophore on <i>Populus</i> sp. (poplar)             |
| <b>Zygo-</b>    | 1157                      | <i>Absidia glauca</i>            | tallus of <i>Peltigera praetextata</i> (lichen)       |
|                 | 2769                      | <i>Mucor plumbeus</i>            | air   |
|                 | 44                        | <i>Mucor circinelloides</i>      | -   |
|                 | 2770                      | <i>Syncephalastrum racemosum</i> | air   |

358

359

360

361

362


363 **Table 2:** maximal percentage of C=C double bond reduction. According to the  
 364 conversion of the substrates, four groups were defined: group A: 100-75 %; group B:  
 365 74-50 %; group C: 49-25 %; group D: 24-0 %. The table shows also the timing of the  
 366 reactions.

| Fungi                     | Conversion % |     |     | Group |     |     | Days |     |     |
|---------------------------|--------------|-----|-----|-------|-----|-----|------|-----|-----|
|                           | CE           | MNS | MCA | CE    | MNS | MCA | CE   | MNS | MCA |
| <i>A. niger</i>           | 57           | 97  | 0   | B     | A   | D   | 2    | 2   | 7   |
| <i>B. bassiana</i>        | 0            | 0   | 0   | D     | D   | D   | 7    | 7   | 7   |
| <i>B. cinerea</i>         | 10           | 0   | 0   | D     | D   | D   | 2    | 2   | 2   |
| <i>C. funicola</i>        | 89           | 30  | 0   | A     | C   | D   | 2    | 7   | 7   |
| <i>C. herbarum</i>        | 100          | 24  | 3   | A     | D   | D   | 2    | 7   | 2   |
| <i>E. nigrum</i>          | 30           | 32  | 0   | C     | C   | D   | 7    | 7   | 7   |
| <i>G. cucujoidarum</i>    | 32           | 26  | 0   | C     | C   | D   | 7    | 7   | 7   |
| <i>G. massei</i>          | 100          | 72  | 50  | A     | B   | B   | 2    | 2   | 4   |
| <i>M. simplex</i>         | 100          | 18  | 0   | A     | D   | D   | 7    | 7   | 7   |
| <i>M. deflexum</i>        | 100          | 44  | 0   | A     | C   | D   | 7    | 7   | 7   |
| <i>O. maius</i>           | 100          | 20  | 12  | A     | D   | D   | 4    | 2   | 2   |
| <i>P. citrinum</i>        | 100          | 98  | 3   | A     | A   | D   | 2    | 7   | 2   |
| <i>P. purpurogenum</i>    | 85           | 0   | 0   | A     | D   | D   | 7    | 7   | 7   |
| <i>P. vinaceum</i>        | 23           | 11  | 0   | D     | D   | D   | 7    | 7   | 7   |
| <i>Scopulariopsis</i> sp. | 100          | 17  | 0   | A     | D   | D   | 7    | 7   | 7   |
| <i>S. fimicola</i>        | 100          | 32  | 18  | A     | C   | D   | 2    | 2   | 7   |
| <i>T. viride</i>          | 100          | 30  | 10  | A     | C   | D   | 2    | 7   | 7   |
| <i>T. spiralis</i>        | 100          | 0   | 0   | A     | D   | D   | 7    | 7   | 7   |
| <i>A. cylindracea</i>     | 100          | 0   | 0   | A     | D   | D   | 2    | 7   | 7   |
| <i>A. farinacea</i>       | 100          | 13  | 0   | A     | D   | D   | 4    | 7   | 7   |
| <i>A. splendida</i>       | 100          | 34  | 0   | A     | C   | D   | 2    | 7   | 7   |
| <i>Coprinellus</i> sp.    | 100          | 11  | 5   | A     | D   | D   | 2    | 7   | 7   |
| <i>P. ostreatus</i>       | 100          | 19  | 0   | A     | D   | D   | 7    | 7   | 7   |
| <i>T. pubescens</i>       | 100          | 52  | 14  | A     | B   | D   | 2    | 4   | 4   |
| <i>A. glauca</i>          | 35           | 22  | 0   | C     | D   | D   | 2    | 7   | 7   |
| <i>M. circinelloides</i>  | 100          | 82  | 100 | A     | A   | A   | 2    | 2   | 2   |
| <i>M. plumbeus</i>        | 100          | 79  | 98  | A     | A   | A   | 2    | 7   | 2   |
| <i>S. racemosum</i>       | 100          | 16  | 0   | A     | D   | D   | 7    | 7   | 7   |

367


368

369 **Figure 1:** putative CE reaction profile. ER: ene-reductase, ADH: alcohol  
 370 dehydrogenase, CE: cyclohexanone, CO: cyclohexanone, COH: cyclohexanol.




371

372 **Figure 2:** products formation profile of *P. citrinum* during the experiment (2, 4 and 7  
 373 days). CE: cyclohexanone, CO: cyclohexanone, COH: cyclohexanol.




374

375 **Figure 3:** putative MNS reaction profile. ER: ene-reductase, MNS: (*E*)- $\alpha$ -  
 376 methylnitrostyrene, NPB: 2-nitropropylbenzene.




377

378 **Figure 4:** putative MCA reaction profile. ER: ene-reductase, ADH: alcohol  
 379 dehydrogenase, MCA:  $\alpha$ -methylcinnamaldehyde, MCOH:  $\alpha$ -methylcinnamyl alcohol,  
 380 MSHCOH:  $\alpha$ -methyldihydrocinnamyl alcohol.



381

382 **Figure 5:** biomass dry weight measurement. Comparison between the pre-growth and  
 383 the end of the trial for each substrate. The y axis represent the biomass weight (mg) and  
 384 the x axis represent the fungi used in this study. Since it was not possible to measure the  
 385 biomass (mg) of the yeast *G. cucujoidarum*, this fungus was not included in the figure.



386