AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Electrochemistry of *Canis familiaris* cytochrome P450 2D15 with gold nanoparticles: An alternative to animal testing in drug discovery

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1528389> since 2019-03-12T14:06:24Z

Published version:

DOI:10.1016/j.bioelechem.2015.03.012

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12

This is the accepted version of the following article:

[Francesco Rua, Sheila J. Sadeghi, Silvia Castrignanò, Francesca Valetti, and Gianfranco Gilardi.
Electrochemistry of *Canis familiaris* cytochrome P450 2D15 with gold nanoparticles: an alternative to
animal testing in drug discovery. Bioelectrochemistry (2015) 105: 110-116.],

which has been published in final form at
[<http://www.sciencedirect.com/science/article/pii/S1567539415000766>]

Electrochemistry of *Canis familiaris* cytochrome P450 2D15 with gold nanoparticles: an alternative to animal testing in drug discovery

Francesco Rua^a, Sheila J. Sadeghi^{a,b}, Silvia Castrignanò^a, Francesca Valetti^a, and Gianfranco Gilardi^{a,b,*}

^aDepartment of Life Sciences and Systems Biology, University of Torino, Italy.

^bCentre for Nanostructured Interfaces and Surfaces, University of Torino, Italy.

CORRESPONDING AUTHOR

* Gianfranco Gilardi, Department of Life Sciences and Systems Biology, Via Accademia Albertina 13, 10123 Turin, Italy. Phone: +39-011-6704593, Fax: +39-011-6704643, E-mail: gianfranco.gilardi@unito.it

ABSTRACT

This work reports for the first time the direct electron transfer of the *Canis familiaris* cytochrome P450 2D15 on glassy carbon electrodes to provide an analytical tool as an alternative to P450 animal testing in the drug discovery process. Cytochrome P450 2D15, that corresponds to the human homologue P450 2D6, was recombinantly expressed in *Escherichia coli* and entrapped on glassy carbon electrodes (GC) either with the cationic polymer polydiallyldimethylammonium chloride (PDDA) or in presence of gold nanoparticles (AuNPs). Reversible electrochemical signals of P450 2D15 were observed with calculated midpoint potentials ($E_{1/2}$) of -191 ± 5 and -233 ± 4 mV vs Ag/AgCl for GC/PDDA/2D15 and GC/AuNPs/2D15, respectively.

These experiments were then followed by the electro-catalytic activity of the immobilized enzyme in presence of metoprolol. The latter drug is a beta-blocker used for the treatment of hypertension and is a specific marker of the human P450 2D6 activity. Electrocatalysis data showed that only in the presence of AuNPs the expected α -hydroxy-metoprolol product was present as shown by HPLC. The successful immobilization of the electroactive *Canis familiaris* cytochrome P450 2D15 on electrode surfaces addresses the ever increasing demand of developing alternative *in vitro* methods for a more detailed study of animal P450 enzymes' metabolism, reducing the number of animals sacrificed in preclinical tests.

KEYWORDS

CYP, animal testing, carbon electrode, gold nanoparticle, immobilization, metoprolol.

1. Introduction

In view of the huge numbers of animals used during preclinical tests in drug discovery [1], both the European Union (European Commission and European Parliament, Directive 2010/63/EU) [2-3] and the US Food and Drug Administration (FDA) [4] are strongly supporting the development of alternative *in vitro* methods to implement the three Rs, that is “reduce, refine and replace” animal models [5].

Hepatic cytochromes P450 are central to toxicological studies due to their primary role in phase I metabolism of more than 80% of marketed drugs [6]. However to date, very little data is published on comparative *in vivo* interspecies studies with even less data available on the different animal P450 expression and/or their metabolic profile [7-10]. To this end, a fast and reliable *in vitro* method would offer the possibility of a more detailed study of the interaction of new drugs with the animal P450 enzymes therefore increasing the predictive value of these preclinical trials.

Canis familiaris is one of the most widely studied animal models used in safety determination of new pharmaceuticals [11-12] and, although the major isoforms of the human cytochromes P450 2D6, 3A4, 2E1, 2C19 and 1A2 have been identified in *C. familiaris* [13-16], there is still a lack of knowledge on their pharmacogenomic/metabolic diversity [9].

Electrochemical techniques already developed in our lab for human hepatic monooxygenases including cytochromes P450 [17-22] represent the ideal approach for a sensitive, accurate and rapid evaluation of animal P450-drug interactions obviating both the requirement for a redox partner and the addition of NADPH cofactor as already reported for some animal P450 enzymes recombinantly expressed in a soluble form [23-25].

In this work, *Canis familiaris* P450 2D15 was chosen as a model for the investigation of canine cytochromes P450 by adopting electrochemical approaches to provide a method for the screening of the safety of new chemical entities/drugs. This enzyme shares 75% identity with the human cytochrome P450 2D6 which alone is responsible for the metabolism of 20–25% of commonly used therapeutic drugs including antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants [6, 26]. The high P450 2D6 polymorphism profile is strictly related to either adverse drug reactions or no drug response [26-27] and to date six different variants have been already identified in the corresponding homologous *C. familiaris* P450 2D15 [28-30].

Generally, the study of the catalytic properties of canine P450 2D enzymes is hampered by the difficulty in their preparation from liver microsomes [14] or by their recombinant expression and purification in a stable form when not associated to the membranes [14-15, 28, 30]. To this end, in this work we report for the first time the recombinant expression and purification of the *Canis familiaris* P450 2D15 in a N-terminally modified form maintaining its structural integrity and function. Furthermore, the electrocatalytic functionality of the purified enzyme is tested whilst immobilized on glassy carbon electrodes (GC).

Two different immobilization strategies were adopted: a) entrapment within the cationic polymer polydiallyldimethylammonium chloride (PDDA) (GC/PDDA/2D15); b) immobilization in presence of gold nanoparticles (AuNPs) stabilized with didodecyldimethylammonium bromide (DDAB) (GC/AuNPs/2D15). Gold nanoparticles are widely used in electrochemical applications [31] especially because they act as excellent electron transfer relays by enhancing the electron transfer from the electrode to the protein [32].

Electrochemical properties of *C. familiaris* P450 2D15 immobilized on glassy carbon electrodes were characterized by cyclic voltammetry and the activity of this enzyme in presence of metoprolol, a selective β_1 receptor blocker used in treatment of cardiovascular disease and a marker for human P450 2D6 activity, assayed by chronoamperometry. The separation and identification of the product

formed was carried out by HPLC leading to the calculation of the K_M value of metoprolol for the first time through the electrochemical system GC/AuNPs/2D15.

2 Material and methods

2.1 Reagents

Kits for plasmid and gene purification were purchased from Sigma Aldrich (Italy). Restriction enzymes, T4 DNA ligase, Vent Polymerase and dNTPS were from New England Biolabs (UK). Chromatographic resins were purchased from GE healthcare (Italy). Quinidine ((S)-[(2R,4S,5R)-5-ethenyl-1-azabicyclo[2.2.2]octan-2-yl](6-methoxyquinolin-4-yl)methanol) and (±)-Metoprolol ({2-hydroxy-3-[4-(2-methoxyethyl)phenoxy]propyl}(propan-2-yl)amine) (+)-tartrate salt, racemic mixture, were purchased from Sigma Aldrich (Italy). Analytical grade chemicals tetrachloroauric (III) acid (gold(III) chloride), DDAB (didodecyl-dimethylammonium bromide) and PDDA (poly(dimethyldiallylammonium chloride)) were all purchased from Sigma-Aldrich and their solutions prepared prior to their use in appropriate solvent for electrochemical experiments.

2.2 Cloning of recombinant P450 2D15

The gene coding for cytochrome P450 2D15 was amplified from the liver cDNA of *C. familiaris* (Biochain, UK) using the primers Fw: 5'-AGACAGCTATGGGGCTGCTG-3' and Rv: 5'-TGGTTTATTGTACCTCGGGCC-3'. The entire gene was blunt end ligated into pBS SK II(+) cloning vector using the *EcoRV* restriction enzyme.

Subsequently, the full-length cDNA coding for the canine P450 was cloned into a pCW expression vector [33] fused in frame with the sequence coding for the leader peptide of the bacterial OmpA protein (MKKTAIAIAVALAGFATVAQA). This system has been previously used to enhance the expression of native mammalian cytochromes P450 in *E. coli* [14, 34]. This leader peptide is cleaved in the bacterial membrane to yield the native P450 enzymes. A code of four histidine residues was added at the C-terminus before the STOP codon to facilitate the protein purification by affinity chromatography.

2.3 Expression and purification of *C. familiaris* P450 2D15

Large-scale expression (4 liters) of P450 2D15 in *E. coli* DH5 α cells transformed with pCW- 2D15 was carried out as described previously for other P450 cytochromes [35] decreasing the post-induction expression time and temperature to 24 hours and 24 °C, respectively.

Purification of P450 2D15 was carried out starting from the isolated membrane fraction using an anion exchange DEAE sepharose column (GE-healthcare, Italy) and followed by a nickel ion affinity chromatography step (GE-healthcare, Italy) where the protein remained bound through the engineered His-tag and subsequently eluted using a 0–40 mM linear gradient of histidine. In order to preserve the P450 2D15 stability the protein was purified in presence of the human P450 2D6 inhibitor quinidine [36]. The UV-visible spectra of the oxidized, reduced and reduced-carbon monoxide bound forms of the protein were recorded on a Hewlett-Packard 8453 diode array spectrophotometer. The P450 concentration was calculated by the method described by Omura and Sato [37].

2.4 Synthesis of AuNPs

DDAB stabilized AuNPs were synthesized following the procedure described by Castrignanò et al. in 2012 [32]. In particular, 0.5 mL of 10 mM tetrachloroauric (III) acid aqueous solution were

mixed under vigorous stirring to 1 mL of a chloroform 0.1 M DDAB solution for 20 minutes to allow the complete transfer of Au(III) ions from the aqueous to the chloroform. After the two phases were separated, 0.2 mL of freshly prepared aqueous solution of 0.4 M sodium borohydride were slowly added. After additional two hours of vigorous stirring, the red organic phase containing the DDAB stabilized AuNPs was collected and stored at 4 °C before use. TEM morphological characterization showed that the synthesized DDAB stabilized AuNPs mainly consist of spherical particles of about 6.5-8.5 nm diameter, ranging from 2.5 to 13 nm in size.

2.5 Immobilization of P450 2D15 on glassy carbon electrode and electrochemical measurements

All electrochemical experiments were carried out at room temperature (25 °C) and in 50 mM phosphate buffer pH 7.4, containing 100 mM KCl as supporting electrolyte, using an Autolab PGSTAT12 potentiostat (Ecochemie, The Netherlands) controlled by GPES3 software. A conventional three-electrode glass cell of 0.5 mL volume, equipped with a platinum wire counter electrode, an Ag/AgCl (3 M NaCl) reference electrode and 3 mm diameter glassy carbon working electrode (BASi, UK), was also used. Before enzyme immobilization, GC electrodes were mechanically polished with alumina and subsequently rinsed and sonicated in ultra pure deionized water.

The P450 2D15 enzyme was immobilized on glassy carbon (GC) with two different approaches: a) P450 2D15 entrapment in PDDA film by mixing equal volumes of 30 µM protein and surfactant solutions before drop-coating on the electrode surfaces (GC/PDDA/2D15); b) P450 2D15 immobilization on GC electrodes using didodecyldimethylammonium bromide (DDAB) stabilized gold nanoparticles (AuNPs) (GC/AuNPs/2D15). 5 µL of 5 mM colloidal gold in 0.1 M DDAB/chloroform was placed on the electrode surface. After evaporation of the chloroform (10 min), 10 µL of 30 µM P450 2D15 solution was added onto the electrode surface. The electrodes were kept overnight at 4 °C in a humid chamber to prevent their total drying.

Cyclic voltammetry experiments were carried out under anaerobic conditions in a glove box with < 5 ppm oxygen (Belle Technologies, UK). Cyclic voltammograms were collected between +100 and -500 mV (vs Ag/AgCl) at a scan rate range of 20–120 mVs⁻¹ in the supporting electrolyte solution. Electrocatalysis experiments were carried out using chronoamperometry applying a potential bias of -650 mV (vs Ag/AgCl) or in cyclic voltammetry at 25°C in presence of metoprolol. The product formed was separated and analyzed by HPLC as previously reported [38]. All electrocatalytic experiments were carried out in triplicates and performed at 200 rpm rotation speed using a RDE-2 rotator system (BASi, USA).

2.6 HPLC analysis of electrocatalysis products

After chronoamperometry in the presence of metoprolol substrate, electrocatalysis solution was collected and an aliquot of 100 µL was injected in HPLC for product quantification. Electrocatalysis products were identified by comparison of retention times to those obtained with authentic standards. For this purpose, standard solutions of metoprolol, α-OH-metoprolol and O-desmethyl-metoprolol were separated isocratically by HPLC (Agilent Technologies-1200 series, Italy) coupled with diode array detector using a 4.6 × 150 mm 5 µm Eclipse XDB-C18 column (Agilent Technologies, USA). Retention times were 3.8 min, 4.7 min and 10.5 min respectively for α-OH-metoprolol, O-desmethyl-metoprolol and metoprolol.

3. Results and discussion

3.1 Recombinant P450 2D15

203 The gene coding for the full length cytochrome P450 2D15 was amplified from *Canis familiaris*
204 liver cDNA and three mutations were identified by full DNA sequencing. A silent substitution of an
205 adenine in guanine was found 325 bases after the 2D15 start with two other mutations leading to the
206 amino acid substitutions of Ile109Val and Phe115Leu in the full-length protein (g345c and g738a).
207 Cytochromes P450 variants in *C. familiaris* 2D15 are not unusual and have been previously
208 reported [28-30] as they are a consequence of the genetic variability existing between dog colonies
209 [39].
210

211 “Here Figure 1”
212

213 The gene encoding for the P450 2D15 variant with the three above-mentioned mutations was cloned
214 in the expression vector pCW and the OmpA leader peptide sequence was introduced at its N-
215 terminus to enhance the protein expression levels as described in the materials and methods. The
216 protein was heterologously expressed in *E. coli* and purified from the membrane fraction with a
217 yield of 4.1 mg of P450 per liter of culture. Cytochrome P450 2D15 (57 kDa) (Fig.1A) was purified
218 in presence of quinidine, a known inhibitor of human P450 2D6. The purified protein was also
219 stored in the presence of this inhibitor which has been shown to preserve the stability of the protein
220 [36]. The inhibitor was removed prior to each experimental assay through buffer exchange.
221 Contrary to all previous literature data, in this work the purified *C. familiaris* P450 2D15 protein
222 and not the microsomal or membrane-associated protein was studied.

223 Binding of carbon monoxide to the dithionite reduced protein confirmed the presence of the
224 correctly incorporated haem cofactor inside the protein scaffold. The protein in its oxidized form
225 presents a Soret peak at 417 nm, the α and β bands at 570 nm and 535 nm, respectively (Fig.1B). As
226 expected, the Soret peak shows the characteristic shift to 450 nm upon reduction and bubbling of
227 carbon monoxide due to the reduced and carbon monoxide-bound adduct and the α and β bands are
228 replaced by a single broad peak centered at 550 nm as shown in Fig.1B. Similarly to the
229 microsomal P450 2D15 reported by Roussel and colleagues [29], a partial amount of inactive
230 protein was also detected at 420 nm (Fig.1B).
231

232 3.2 Direct electrochemistry of P450 2D15 on glassy carbon electrodes

233 Recombinant P450 2D15 was immobilized on glassy carbon electrodes in a non-oriented fashion
234 either by entrapment within the cationic polymer polydiallyldimethylammonium chloride (PDDA)
235 (GC/PDDA/2D15) or in the presence of gold nanoparticles (AuNPs) (GC/AuNPs/2D15) which are
236 one of the most stable metal nanoparticles that provide an effective electron transfer to-and-from the
237 glassy carbon electrode [32, 40].

238 Once the protein was successfully immobilized on glassy carbon electrodes, cyclic voltammetry
239 experiments were carried out at 25°C under anaerobic conditions (< 5 ppm O_2) to prevent the
240 formation of the Fe^{2+} -dioxygen complex related to the second electron transfer step within the
241 catalytic cycle of cytochromes P450. Cyclic voltammograms of P450 2D15 entrapped with PDDA
242 (GC/PDDA/2D15) or in presence of AuNPs (GC/AuNPs/2D15) are shown in Fig.2A and B,
243 respectively. In both cases the ratio between the anodic and cathodic peak currents were found to be
244 one but the magnitude of the current observed was nearly 4x higher in the presence of AuNPs. As
245 stated by Laviron's theory [41], the linear dependence of the anodic and cathodic peak currents on
246 the scan rate (within the range 20-120 mV) for both GC/PDDA/2D15 and GC/AuNPs/2D15,
247 indicated that the electron transfer from/to the modified GC electrode is quasi-reversible and it is a
248 surface-controlled process as expected for an immobilized cytochrome P450 on electrode surfaces.
249

“Here Figure 2”

The resulting anodic and cathodic peak currents for GC/PDDA/2D15 were detected at -256 ± 6 mV (Ea) and -127 ± 5 mV (Ec), respectively, with a calculated midpoint potential (Em) of -191 ± 5 mV (vs Ag/AgCl). In the GC/AuNPs/2D15 immobilization strategy, the anodic and cathodic peak currents were measured at -267 ± 5 mV (Ea) and -200 ± 4 mV (Ec) (Fig.2D) and the resulting midpoint potential was calculated to be -233 ± 4 mV. These values are in the range of other electrochemically determined midpoint potentials reported in the literature for different human P450 enzymes [21].

Peak-to-peak separation value was calculated to be -129 ± 6 for GC/PDDA/2D15 and -67 ± 5 mV for GC/AuNPs/2D15 indicating that the presence of the gold nanoparticles facilitates a much faster electron transfer of P450 2D15 when immobilized on glassy carbon electrodes.

Apparent surface coverage of the electroactive protein was calculated from the slope of I_p versus v plot in accordance with the Brown Anson model using the equation:

$$i_p = n^2 F^2 A \Gamma v / 4RT$$

where n represents the number of electrons involved in the reaction, A is the electrode surface area (0.071 cm^2), Γ is the surface coverage, v is the scan rate, F is the Faraday constant, T is the temperature and R is the gas constant. The surface coverage value on GC/PDDA was calculated to be 1.1×10^{12} molecules/ cm^2 in the same range previously reported for the *Macaca fascicularis* P450 2C20 immobilized in the same manner [25]. As expected the same calculation of the surface coverage in the presence of AuNps resulted in a value of 4.6×10^{12} molecules/ cm^2 , around 4 times higher than using PDDA alone.

The electron transfer efficiency of the GC/AuNPs/2D15 was further investigated recording a series of cyclic voltammograms varying the pH within the range $6.5 < \text{pH} < 8.5$ (Fig. 3). A negative shift in peak potential upon lowering pH was observed which is indicative of proton-coupled electron transfer [42]. The measured shift in the formal potential was around 60 mV per pH unit, close to the theoretical value of 59 mV/pH expected at 25°C for one proton, one electron transfer. The protonation site of the reduced P450 remains unknown but it is reasonable to assume that it is either an amino acid in close proximity of the haem iron or the water bound in the sixth coordination, as has been suggested in the case of other cytochromes P450 [42-43]. Nocera's group [44] have also suggested that proton-coupled electron transfer in P450 enzymes is from a structured water, which is the sixth coordinated H_2O mentioned above.

“Here Figure 3”

3.3 Electrocatalysis of 2D15 immobilized on glassy carbon electrodes

Electrocatalysis by the immobilized P450 2D15 (GC/AuNPs/2D15) was followed by both chronoamperometry and cyclic voltammetry in the presence of metoprolol. The latter drug (Scheme 1) is a cardioselective β_1 -adrenergic blocking agent metabolized principally by the human P450 2D6 and converted into the α -OH-metoprolol and O-desmethyl-metoprolol products [38]. The amount of O-desmethyl-metoprolol (scheme1-left) after electrocatalysis was found to be below the

detection limits. For this reason, only α -OH-metoprolol (scheme 1-right) was considered as a product formed.

“Here Scheme 1”

As can be seen in Fig. 4A, a catalytic current was observed in air saturating conditions in the presence of metoprolol in cyclic voltammetry experiments. Furthermore, using chronoamperometry at regular intervals additions of metoprolol were carried out in the range of 12.5-250 μ M under aerobic conditions (Fig. 4B) with an applied bias of -650 mV. The change in the chronoamperometric currents registered due to the different concentrations of the drug showed a Michaelis–Menten kinetic mechanism with a plateau or saturation at higher than 70 μ M concentrations of metoprolol. The K_{Mapp} calculated from these currents was 39.5 $\pm$ 3.5 μ M (Fig. 5A).

“Here Figure 4”

Further electrocatalysis experiments were carried out by chronoamperometry for both the GC/AuNPs/2D15 and the GC/PDDA/2D15 systems applying a potential bias of -650 mV (vs Ag/AgCl) for 30 min at 25 °C in presence of different concentrations of metoprolol (Fig. 5B). For each concentration of the drug tested, the contents of the electrochemical cell were analyzed after the 30 min reaction by HPLC in order to separate and quantify the product(s) formed by the immobilized enzyme. No products were detected from the HPLC analysis for the *C. familiaris* P450 2D15 enzyme entrapped with the polycationic agent PDDA. On the other hand, the enzyme immobilized in the presence of AuNps (GC/AuNPs/2D15) was capable of hydroxylating the metoprolol producing 30.7 $\pm$ 2.3 pmol of product at the higher metoprolol concentrations used. No O-desmethylated product was observed in line with previously published data on microsomal P450 2D15 [28].

“Here Figure 5”

Control experiments were also carried out in the absence of the protein and as expected no product was detected (data not shown). Quantification of the reaction products in the presence of increasing amounts of metoprolol (Fig. 5B) resulted in a calculated apparent K_M value of 36.2 $\pm$ 1.4 μ M for GC/AuNPs/2D15 (Table 1) very similar to the K_{Mapp} obtained by titrating different concentrations of metoprolol (see Fig. 5A). The electrochemically determined K_M values are also in the same range as those reported by Ellis and colleagues for human P450 2D6 [45] therefore it can be concluded that the *C. familiaris* P450 2D15 possesses similar enzymatic activity to human P450 2D6, at least for metoprolol metabolism.

“Here Table 1”

4. Conclusions

The development of an electrochemical platform for the screening of animal cytochromes P450 represents an interesting and feasible solution to the ever increasing demand to reduce animal testing, maintaining a high level of predictivity of the animal models used in preclinical tests and providing the opportunity to study the P450 metabolism of new drugs and chemical entities.

To this end, the cloning of a new *Canis familiaris* P450 2D15 variant and its expression in a pure form for its electrochemical application reported in this work, brings us a step closer to the

342 realization of an *in vitro* electrochemical toxicity screening, an alternative to *in vivo* animal testing
343 and sacrifice.

344 The P450 2D15 enzyme was successfully immobilized on glassy carbon electrodes in a stable form
345 achieving an efficient electron transfer between the electrode and the enzyme. The presence of gold
346 nanoparticles enhanced the sensitivity of the electrode-protein system and allowed for the
347 estimation of kinetic parameters for metoprolol, a typical drug marker of the human P450 2D6.

348 These results represent the first step towards the development of electrochemical platform for the
349 rapid exploration of the metabolic diversity existing between animals and humans cytochromes
350 P450 contributing to a more reliable interpretation of the data generated in canine species during the
351 pharmacological preclinical studies.

352

353

354 **ACKNOWLEDGEMENTS**

355 The authors wish to acknowledge financial support from the Regione Piemonte CIPE 2006
356 (CYPTECH-project, Italy) and Progetto Ateneo-San Paolo 2012 (awarded to S. Sadeghi).

357

358

REFERENCES

- [1] M. Hudson-Shore. Statistics of Scientific Procedures on Living Animals 2012: another increase in experimentation - genetically-altered animals dominate again. *Altern Lab Anim.* 41 (2013) 313-319.
- [2] O. Varga, A.K. Hansen, P. Sandøe, I.A. Olsson, Validating animal models for preclinical research: a scientific and ethical discussion. *EMBO Rep.* 11 (2010) 500–503.
- [3] J.E. May, J. Xu, H.R. Morse, N.D. Avent, C. Donaldson, Toxicity testing: the search for an in vitro alternative to animal testing. *Br. J. Biomed. Sci.* 66 (2009) 160–165.
- [4] L.M. Schechtman, W.S. Stokes. The FDA's regulatory role in the ICCVAM process. *Altern Lab Anim.* (2004) 32, 663-668.
- [5] C.A. Schuppli, D. Fraser, M. McDonald. Expanding the three Rs to meet new challenges in humane animal experimentation. *Altern Lab Anim.* (2004) 32: 525-532.
- [6] F.P. Guengerich. Human cytochrome P450 enzymes. Ortiz de Montellano, P. R., editor. (2005) *Cytochrome P450: Structure, Mechanism, and Biochemistry*, 3rd ed. pp. 377–463. Plenum, New York.
- [7] J.J. Bogaards, M. Bertrand, P. Jackson, M.J. Oudshoorn, R.J. Weaver, P.J. van Bladeren, B. Walther. Determining the best animal model for human cytochrome P450 activities: a comparison of mouse, rat, rabbit, dog, micropig, monkey and man. *Xenobiotica.* 30 (2000) 1131-52.
- [8] J. Tibbitts. Issues related to the use of canines in toxicologic pathology--issues with pharmacokinetics and metabolism. *Toxicol Pathol.* 31 (2003) 17-24.
- [9] M. Martignoni, G.M. Groothuis, R. de Kanter. Species differences between mouse, rat, dog, monkey and human CYP-mediated drug metabolism, inhibition and induction. *Expert Opin Drug Metab Toxicol.* 6 (2006) 875-894.
- [10] F. Rua, G. Di Nardo, S.J. Sadeghi, G. Gilardi. Towards reduction in animal sacrifice for drugs: Molecular modelling of *Macaca fascicularis* P450 2C20 for virtual screening of *Homo sapiens* P450 2C8 substrates. *Biotechnol. Appl. Biochem.* (2012) 59, 479-489.
- [11] M.N. Martinez, L. Antonovic, M. Court, M. Dacasto, J. Fink-Gremmels, B. Kukanich, C. Locuson, K. Mealey, M.J. Myers, L. Trepanier. Challenges in exploring the cytochrome P450 system as a source of variation in canine drug pharmacokinetics. *Drug Metab Rev.* 45 (2013) 218-230.
- [12] S. Fleischer, M. Sharkey, K. Mealey, E.A. Ostrander, M. Martinez. Pharmacogenetic and metabolic differences between dog breeds: their impact on canine medicine and the use of the dog as a preclinical animal model. *AAPS* 10 (2008) 110-119.
- [13] M. Mise, T. Hashizume, S. Komuro, Characterization of substrate specificity of dog CYP1A2 using CYP1A2-deficient and wild-type dog liver microsomes. *Drug Metab Dispos.* 36 (2008) 1903-1908.

- [14] C.W. Locuson, B.T. Ethell, M. Voice, D. Lee, K.L. Feenstra, Evaluation of Escherichia coli membrane preparations of canine CYP1A1, 2B11, 2C21, 2C41, 2D15, 3A12, and 3A26 with co-expressed canine cytochrome P450 reductase. *Drug Metab Dispos.* 37 (2009) 457-61.
- [15] M. Shou, R. Norcross, G. Sandig, P. Lu, Y. Li, Y. Lin, Q. Mei, A.D. Rodrigues, T.H. Rushmore, Substrate specificity and kinetic properties of seven heterologously expressed dog cytochromes p450. *Drug Metab Dispos.* 31 (2003) 1161-1169.
- [16] K.L. Mealey, M. Jabbes, E. Spencer, J.M. Akey. Differential expression of CYP3A12 and CYP3A26 mRNAs in canine liver and intestine. *Xenobiotica.* 38 (2008) 1305-1312.
- [17] V. Dodhia, C. Sassone, A. Fantuzzi, G. Di Nardo S.J. Sadeghi, G. Gilardi, Modulating the coupling efficiency of human cytochrome P450 CYP3A4 at electrode surfaces through protein engineering. *Electrochem. Commun.* 10 (2008) 1744–1747.
- [18] A. Fantuzzi, E. Capria, L.H. Mak, V.R. Dodhia, S.J. Sadeghi, S. Collins, G. Somers, E. Huq, G. Gilardi, An electrochemical microfluidic platform for human P450 drug metabolism profiling. *Anal Chem.* 82 (2010) 10222-10227.
- [19] S. Castrignanò, S.J. Sadeghi, G. Gilardi. Electro-catalysis by immobilised human flavin-containing monooxygenase isoform 3 (hFMO3), *Anal Bioanal Chem*, 398 (2010), 1403-1409.
- [20] S.J. Sadeghi, R. Meirinhos, G. Catucci, V.R. Dodhia, G. Di Nardo, G. Gilardi. Direct electrochemistry of drug metabolizing human flavin-containing monooxygenase: electrochemical turnover of benzydamine and tamoxifen, *J Am Chem Soc.* 132 (2010), 458-9.
- [21] S.J. Sadeghi, A. Fantuzzi A, G. Gilardi. Breakthrough in P450 bioelectrochemistry and future perspectives. *Biochim Biophys Acta.* 1814 (2011) 237-248.
- [22] S. Castrignanò, A. Ortolani, S.J. Sadeghi, G. Di Nardo, P. Allegra, G. Gilardi. Electrochemical detection of human cytochrome P450 2A6 inhibition: a step toward reducing dependence on smoking. *Anal.Chem.* 86 (2014) 2760-2766.
- [23] D.L. Johnson, A.J. Conley, L.L. Martin. Direct electrochemistry of human, bovine and porcine cytochrome P450c17. *J Mol Endocrinol.* 36 (2006) 349-359.
- [24] V.V. Shumyantseva, T.V. Bulko, T.T. Bachmann, U. Bilitewski, R.D. Schmid, A.I. Archakov. Electrochemical reduction of flavocytochromes 2B4 and 1A2 and their catalytic activity. *Arch Biochem Biophys.* 377 (2000) 43-48.
- [25] F. Rua, S.J. Sadeghi, S. Castrignanò, G. Di Nardo, G. Gilardi. Engineering *Macaca fascicularis* cytochrome P450 2C20 to reduce animal testing for new drugs. *J Inorg Biochem.* 117 (2012) 277-284.

456 [26] S.F. Zhou, Polymorphism of human cytochrome P450 2D6 and its clinical significance: Part I.
 457 Clin Pharmacokinet. 48 (2009) 689-723.

458

459 [27] Neafsey P, Ginsberg G, Hattis D, Sonawane B. Genetic polymorphism in cytochrome P450
 460 2D6 (CYP2D6): Population distribution of CYP2D6 activity. J Toxicol Environ Health B Crit Rev.
 461 12 (2009) 334-361.

462

463 [28] K. Sakamoto, S. Kirita, T. Baba, Y. Nakamura, Y. Yamazoe, R. Kato, A. Takanaka, T.
 464 Matsubara, A new cytochrome P450 form belonging to the CYP2D in dog liver microsomes:
 465 purification, cDNA cloning, and enzyme characterization. Arch Biochem Biophys. 1 (1995) 372-82.

466

467 [29] F. Roussel, D.B. Duignan, M.P. Lawton, R.S. Obach, C.A. Strick, D.J. Tweedie, Expression
 468 and characterization of canine cytochrome P450 2D15. Arch Biochem Biophys. 357 (1998) 27-36.

469

470 [30] T. Tasaki, H. Iwata, A. Kazusaka, S. Fujita, Regio- and stereoselectivity in propranolol
 471 metabolism by dog liver microsomes and the expressed dog CYP2D15. J Biochem. 123 (1998) 747-
 472 751.

473

474 [31] S. Guo, E. Wang. Synthesis and electrochemical applications of gold nanoparticles. Anal.
 475 Chim. Acta. 598 (2007) 181-192.

476

477 [32] S. Castrignanò, S.J. Sadeghi, G. Gilardi. Entrapment of human flavin-containing
 478 monooxygenase 3 in the presence of gold nanoparticles: TEM, FTIR and electrocatalysis. Biochim
 479 Biophys Acta. 1820 (2012) 2072-2078.

480

481 [33] EM. Gillam, Z. Guo, FP Guengerich. Expression of modified human cytochrome P450 2E1 in
 482 Escherichia coli, purification, and spectral and catalytic properties. Arch. Biochem. Biophys 312
 483 (1994), 59-66.

484

485 [34] M.P. Pritchard, M.J. Glancey, J.A. Blake, D.E. Gilham, B. Burchell, C.R. Wolf, T. Friedberg,
 486 Functional co-expression of CYP2D6 and human NADPH-cytochrome P450 reductase in
 487 Escherichia coli. Pharmacogenetics. 8 (1998) 33-42.

488

489 [35] P. Panicco, VR Dodhia, A. Fantuzzi, G. Gilardi. First enzyme-based amperometric platform to
 490 determine the polymorphic response in drug metabolism by cytochromes P450, Anal. Chem. 83
 491 (2011) 2179-2186.

492

493 [36] K. Berka, E. Anzenbacherová, T. Hendrychová, R. Lange, V. Mašek, P. Anzenbacher, M.
 494 Otyepka. Binding of quinidine radically increases the stability and decreases the flexibility of the
 495 cytochrome P450 2D6 active site. J Inorg Biochem. 110 (2012) 46-50.

496

497 [37] T. Omura, R. Sato, The Carbon Monoxide-Binding Pigment of Liver Microsomes. I. Evidence
 498 for Its Hemoprotein Nature. J Biol Chem 239 (1964) 2370-8.

499

500 [38] I.H. Hanna, J.A. Krauser, H. Cai, M.S. Kim, F.P. Guengerich. Diversity in mechanisms of
 501 substrate oxidation by cytochrome P450 2D6. Lack of an allosteric role of NADPH-cytochrome
 502 P450 reductase in catalytic regioselectivity. J Biol Chem. 276 (2001) 39553-395561.

503

504 [39] M.N. Martinez, L. Antonovic, M. Court, M. Dacasto, J. Fink-Gremmels, B. Kukanich, C.
505 Locuson, K. Mealey, M.J. Myers, L. Trepanier. Challenges in exploring the cytochrome P450
506 system as a source of variation in canine drug pharmacokinetics. *Drug Metab Rev.* 45 (2013) 218-
507 230.

508 [40] R. Sardar, A.M. Funston, P. Mulvaney, R.W. Murray. Gold nanoparticles: past, present, and
509 future. *Langmuir.* 15 (2009) 13840-13851.

511 [41] E. Laviron, General expression of the linear potential sweep voltammogram in the case of
512 diffusionless electrochemical systems. *J. Electroanal. Chem.* 101 (1979) 19-28.

514 [42] B.D. Fleming, S.G. Bell, L-L. Wong, A.M. Bard, The electrochemistry of a heme-containing
515 enzyme, CYP199A2, adsorbed directly onto a pyrolytic graphite electrode, *J. Electroanal. Chem.*
516 611 (2007) 149-154.

518 [43] A. Shukla, E.M. Gillam, D.J. Mitchell, P.V. Bernhardt, Direct electrochemistry of enzymes
519 from the cytochrome P450 2C family, *Electrochem. Commun.* 7 (2005) 437-442.

521 [44] S.Y. Reece, J.M. Hodgkiss, J. Stubbe, D.G. Nocera, Proton-coupled electron transfer: the
522 mechanistic underpinning for radical transport and catalysis in biology, *Phil. Trans. R. Soc. B* 361
523 (2006) 1351-1364.

525 [45] S.W. Ellis, K. Rowland, M.J. Ackland, E. Rekka, A.P. Simula, M.S. Lennard, C.R. Wolf, G.T.
526 Tucker. Influence of amino acid residue 374 of cytochrome P-450 2D6 (CYP2D6) on the regio- and
527 enantio-selective metabolism of metoprolol. *Biochem J.* 316 (1996) 647-654.

529
530
531

Figure legends:

Fig. 1: Purification and spectral characterization of P450 2D15. (A) SDS-PAGE gel, lane 1: molecular weight markers, lane 2: purified protein (57 kDa). The absorbance spectra of the purified P450 2D15 (B) in the oxidized (solid black line), dithionite reduced (dotted line), and the reduced carbon monoxide-bound form (dashed line).

Fig. 2: Electrochemical characterization of P450 2D15 immobilized on glassy carbon electrodes (A) entrapped with the cationic surfactant PDDA (GC/PDDA/2D15), and (B) in presence of gold nanoparticles (GC/AuNPs/2D15). Cyclic voltammogram of P450 2D15 measured at a scan rate of 120 mV/s in 50 mM potassium phosphate buffer pH 7.4 with 100 mM KCl at 25 °C. Shown are the original and baseline corrected (upper traces) cyclic voltammograms. Bottom: plot of cathodic (filled symbols) and anodic (empty symbols) peak currents versus scan rate. $R^2 > 0.99$.

Fig. 3: Cyclic voltammograms of GC/AuNPs/2D15 at a scan rate of 120 mVs⁻¹ under anaerobic conditions in different pH value buffer. The cyclic voltammogram at pH 7.4 is shown in black.

Fig. 4: Electrochemical responses of GC/AuNPs/2D15 to metoprolol. (A) Cyclic voltammograms of the GC/AuNPs/2D15 in the air-saturated buffer in presence (black) and absence (grey) of metoprolol shown together with control trace obtained in the absence of the protein (dashed); (B) Chronoamperometric response of GC/AuNPs/2D15 at -650 mV followed for 600 sec in 50 mM potassium phosphate buffer pH 7.4. The arrows indicate the addition of different concentrations of metoprolol (12.5-250 µM).

Fig. 5: Michaelis–Menten plot of the turnover of P450 2D15 immobilized on glassy carbon electrode in presence of AuNps (GC/AuNPs/2D15) and metoprolol obtained from: (A) the current registered in the titration experiments (Δ : current increase upon metoprolol addition) and, (B) product quantified by HPLC after 30 min electrocatalysis.

Figure 1

562
563

Figure 2

564
565

Figure 3

566
567

Schematic 1

568
569

Figure 4

570
571

Figure 5