

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Therapeutic use of human renal progenitor cells for kidney regeneration

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1530426> since 2015-11-30T17:23:59Z

*Published version:*

DOI:10.1038/nrneph.2015.126

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

Nowadays, kidney diseases are considered a research and public health priority, as their incidence is increasing worldwide with augmented morbidity and mortality<sup>1</sup>. Acute kidney injury (AKI) is characterized by an acute loss of kidney cells and functions, resulting from ischemic, toxic or inflammatory insults, and associates with in-hospital morbidity and mortality, despite the supportive therapies. In addition, acute insults or chronic injury may lead to the progressive loss of nephrons and renal functions, requiring renal replacement therapy by dialysis or transplantation.

The development of acute and chronic renal disease is strictly correlated with an impaired ability of renal repair to regain function after injury. Neo-nephrogenesis does not occur in the adult human kidney. However, the kidney retains some regenerative potential. Several studies have provided evidence to suggest that repair activities in adult kidneys are supported by resident nephron cells surviving injury, with a dedifferentiated phenotype and a clonal proliferative potential<sup>2, 3</sup>. However, the transcription patterns that characterize nephron generation, known to be present in foetal renal progenitors, are only partially activated during adult kidney repair, indicating differences and weaknesses of the repair process<sup>4</sup>. Consequently, the repair ability in adult tissue is limited and, after repeated injury, results in a maladaptive response, ensuing in fibrosis and loss of function. In fact, it is well established that patients with AKI or subclinical rejections may develop chronic kidney disease (CKD)<sup>5</sup>. Finally, chronic renal failure is characterized by a loss of regenerative processes.

In the present review, we will examine the data relative to the presence, the isolation and the *in vitro* generation of renal progenitor-like cells (RPCs), focusing on results obtained in humans, and discuss their potential therapeutic applications.

## **A. Presence of progenitors or progenitor-like cells (RPCs) in the renal tissue**

### *RPCs in the foetal kidney*

In all mammals, the development of mature nephrons involves the differentiation and organization of RPCs of mesenchymal origin. From the embryonic metanephros, reciprocal inductive interactions between the metanephric mesenchyme and the ureteric bud are required for kidney formation<sup>6</sup>. The cap mesenchyme represents the niche where renal progenitors

1 undergo mesenchymal-to-epithelial transition<sup>7, 8</sup>. These committed progenitors sequentially  
2 form pre-tubular aggregates, renal vesicles and C- and S-shaped bodies, leading to the  
3 glomerular and renal tubule compartment<sup>6</sup>.

4 Embryonic progenitors represent the prototypes of the renal cell population that are able  
5 to proliferate and differentiate towards different types of nephron epithelia. Therefore, defining  
6 the cell programs involved in renal progenitor function may provide important knowledge for the  
7 mechanisms of renal regeneration in the adult kidney. Several transcription factors and  
8 markers have been identified to characterize RPCs in the developing kidney, and most of our  
9 knowledge derives from mouse studies. In particular, RPCs in the mouse embryo were first  
10 characterized as a multipotent population expressing Six2 with self-renewing capability<sup>8</sup>. In  
11 addition, fate mapping in the mouse embryo led to the identification of several genes marking  
12 progenitors and involved in their maintenance and differentiation<sup>9</sup>, including the Sall1<sup>10</sup>,  
13 EYA1<sup>11</sup>, OSR1<sup>12</sup>, CITED1<sup>7</sup> genes and the transcriptional factors PAX2<sup>13</sup> and WT1<sup>14</sup>. Other  
14 markers were reported as being involved in segment specification, such as the Wnt target gene  
15 *Lgr5* responsible for generating the thick ascending limb of Henle's loop and the distal  
16 convoluted tubule<sup>15</sup>, or Notch2, which appears to regulate the shift from self-renewal to  
17 differentiation, specifying proximal tubules<sup>16</sup>.

18 In humans, foetal RPCs have been characterized to a lesser extent. Many results were  
19 obtained from an extensive gene-expression profile analysis in Wilm's tumour, as this  
20 paediatric tumour is characterized by loss of terminal progenitor differentiation<sup>17</sup>. The  
21 population expressing foetal renal progenitor markers Six2, Wt1, Cited1, and Sall1 was shown  
22 to co-express NCAM and FZD7<sup>18</sup>. By contrast, the CD133 marker appeared to be expressed  
23 by Epcam<sup>+</sup> cells with a more differentiated phenotype<sup>18</sup>. Indeed, these two surface markers,  
24 NCAM and CD133, were successfully exploited for the isolation of RPCs from the foetal renal  
25 kidney. NCAM1 has been reported to facilitate selection for a nephron lineage that includes  
26 SIX2-positive cap mesenchyme cells with *in vitro* clonogenic and stem/progenitor properties<sup>19</sup>.  
27 While NCAM1 is absent in the normal adult human kidney, CD133 is expressed from the  
28 vesicle state onwards<sup>20,21</sup>, and is maintained in scattered cells in the adult renal tissue<sup>22</sup>.

29 In the adult kidney, the nephrogenic mesenchymal progenitor population disappears,  
30 maybe due to the loss of its niche<sup>23</sup>. RPCs are only induced in the nephrogenic pathway until  
31 the 34th week of gestation<sup>19</sup>, and no equivalent cell types can be identified in the adult kidney.

1 Even during repair, only a partial reactivation of certain foetal transcriptional programs occurs,  
2 such as induction of Pax2, BMI-1 and Six2<sup>4, 24</sup>.

3

#### 4 *RPCs in the adult kidney*

5 Even though the ability of whole nephron regeneration is lost in the human adult kidney,  
6 organ homeostasis and the repair process require cells with progenitor-like characteristics. This  
7 population is required for the continuous replacement of cells lost during physiological  
8 processes. In fact, it is estimated that about 6000 cells, which derive from different nephron  
9 segments, are lost every hour in the urine (Prescott). In addition, repair after acute injury  
10 requires repopulation of tubular compartments. Initial investigations in the stem cell field  
11 supported the search for residual stem cells involved in these processes, whereas subsequent  
12 studies failed to identify cells with multipotent nephron-generating ability.

13

#### 14 Murine studies

15

16 The presence of RPCs and their possible role in renal regeneration has been investigated  
17 thanks to lineage tracing and fate-mapping techniques. The involvement of nephron resident  
18 cells in tubule regeneration was first demonstrated by Humphrey and colleagues<sup>25</sup>. Using  
19 genetic label of mesenchyme-derived renal epithelial cells, they could exclude the role of  
20 interstitial or bone marrow derived stem cells in repair after acute injury<sup>25</sup>. In a subsequent  
21 study, double tracking of tubular cells following acute ischaemic damage showed a random  
22 proliferation of tubular epithelial cells. These results have been attributed to a proliferation of all  
23 cells within a specific tubular segment. However, they do not exclude the possibility that a  
24 subpopulation of cells with progenitor properties may proliferate and repair the tubular  
25 segments, as suggested by other reports<sup>2</sup>.

26 Using rainbow mice, the group of Dekel clearly identified that cells with segment-specific clonal  
27 and extensive proliferation ability can support kidney epithelia self-renewal throughout an entire  
28 lifetime<sup>2</sup>. Indeed, single colour-expressing cell colonies were detectable along kidney segments  
29 at 6 months after cell tagging, thus indicating the presence of clone-forming cells<sup>2</sup>. In addition,  
30 segment-specific cells with clonogenic ability also contributed to the repair of specific nephron  
31 segments, suggesting that nephron regeneration is granted by segment-committed clonogenic  
32 RPCs.

1 From all these observations, it can be inferred that the process of regeneration occurring in  
2 renal tubules relies on resident intratubular cells with a limited segment specific regenerative  
3 potential and with clonogenic and proliferative ability, thus representing a “functional”  
4 population of renal progenitor cells. However, the identity of these cells remains elusive, as  
5 they may possibly derive either by a pre-determined population of RPCs, or by a plastic  
6 phenomenon in which cells transiently acquire a RPC phenotype.

7 Recently, a subpopulation of proximal tubular cells (“scattered tubular cells”, STC) was  
8 identified and genetically labelled. STCs appeared as an injury-inducible population, as their  
9 number did increase when labelled during ischaemic injury<sup>27</sup>. Although these data do not  
10 exclude a concomitant activation of renal progenitors in the regenerative process, they  
11 underline the capability of tubular differentiated cells to acquire a progenitor-like function and  
12 suggest that the plasticity of the tubular phenotype might be involved in this process (review  
13 Berger).

14 In contrast to the tubules, the glomerular regenerative potential is quite limited. RPCs identified  
15 in Bowman's capsule appeared to contribute to podocyte regeneration, migrating onto the  
16 glomerular tuft, only during infancy to adolescence, whereas this process was not observed in  
17 adult life<sup>28</sup>. In addition, using label retention, intratubular slow cycling populations have been  
18 confirmed in proximal and distal tubules and in the thick ascending limb of the Henle's loop<sup>29</sup>  
19 as well as in the papilla<sup>30</sup>. Furthermore, by expression of stem-cell markers, specific  
20 populations of progenitors have been detected as nFatC1+ cells in proximal tubules<sup>31</sup> and as  
21 Lgr5 or c-kit expressing cells in the thick ascending limb of Henle's loop and distal convoluted  
22 tubules<sup>15,32</sup>. It is possible that distinct populations of segment progenitors might be  
23 characterized by different marker expression (Table I).

24

#### 25 Human studies

26 In human tissue, using the AC133 antibody, which recognizes a glycosylation-  
27 dependent epitope of prominin (CD133), expressed by haemopoietic stem cells<sup>33,34</sup>, a discrete  
28 population of CD133+ cells was detected in different segments of the nephron, namely the  
29 Bowman's capsule, proximal and distal tubules, as well as in the inner medullary papilla region,  
30 including Henle's loop and the S3 limb segment<sup>35-42</sup> (Table I). CD133+ cells co-expressed the  
31 mesenchymal marker vimentin and cytokeratins 7 and 19, not expressed by the bulk of tubular  
32 epithelial cells<sup>37,39</sup> (Table II). In the Bowman's capsule, CD133+CD24+ cells localized at the

1 urinary pole, whereas cells expressing differentiation markers of podocytes increased toward  
2 the vascular pole<sup>40</sup>. In proximal tubules, CD133+ cells were mainly located in the tubular plicae  
3 where proximal convoluted tubules make hairpin turns<sup>3</sup>, and co-expressed megalin and  
4 aquaporin-1<sup>36,41</sup>. In the distal nephron, small clusters of CD133+CD24+ cells were detected in  
5 the portion of the distal convoluted tubule that makes contact with the vascular pole of the  
6 glomerulus, and these cells also co-stained for the thiazide-sensitive Na/Cl cotransporter<sup>41</sup>. At  
7 variance, CD133+ cells in the medulla lacked expression of differentiation markers<sup>36</sup>.  
8 Moreover, the CD24+, CD133+, or vimentin-expressing scattered cells observed within the  
9 proximal tubules were shown to co-express known markers of scattered cells, such as annexin  
10 2, claudin-3 and 7, S100A6, and CXCR4<sup>37</sup>, as well as collagen 7A1 and the tight junction  
11 protein claudin-1<sup>3</sup> (Table II). As most experiments have been performed on normal portions of  
12 kidneys derived from tumour-bearing patients, it has been suggested that CD133 expression  
13 can be induced. Recently, CD133-expressing cells (from 3 to 12%) have also been shown to  
14 be present in normal renal tissue of pre-transplant biopsies from native donors of all ages,  
15 indicating their presence at a physiological level<sup>3</sup>. However, this result conflicts with the  
16 observation that scattered tubular cells are absent in normal juvenile tissue in rodents<sup>37</sup>.

17 A detailed analysis of scattered intratubular cell morphology, by electron microscopy,  
18 indicated a striking difference with respect to adjacent 'normal' proximal tubular cells, as they  
19 appeared smaller, lacked brush border microvilli and showed fewer mitochondria<sup>37</sup>. The lack of  
20 mitochondria in scattered cells has been further confirmed by electron microscopy using gold-  
21 conjugated vimentin antibodies, as well as by double staining of renal tissue with CD133 and  
22 mitochondrial markers<sup>3</sup>. In the light of the role of CD133 in glucose uptake<sup>42,43</sup>, the CD133-  
23 expressing population may possess a particular anaerobic metabolism, which could explain  
24 their ability to survive following injury<sup>3</sup>. Indeed, Bourseau- Guilmain *et al.*<sup>42</sup> showed that CD133  
25 inhibits endocytosis of the transferrin receptor, thus resulting in inhibition of iron uptake and  
26 consequently of mitochondrial activity. In turn, CD133 expression has been related to glucose  
27 uptake in glioma cells<sup>43</sup>. Taken together, these data lead to the proposal that CD133 might be  
28 involved in the regulation of glycolytic metabolism. Indeed, in an explant model in which human  
29 renal tissue was subjected to ischaemia reperfusion damage, the CD133+ scattered cells did  
30 not undergo necrosis and remained anchored to the basal membranes, whereas the detached  
31 cells were negative for these markers<sup>3</sup>. This observation supports the notion of a  
32 physiologically distinct population with high resistance to damage, rather than an injury-induced

1 population. It does not address, however, whether the CD133+ cells are a fixed population or a  
2 transient functional state modulated by microenvironmental stimuli.

3

#### 4 *RPCs in renal pathology*

5        Whatever their origin, the CD133 population appears to play a role in acute tissue repair  
6 of the tubular compartment. Conversely, their activation in glomeruli appears to be related to  
7 glomerular diseases<sup>20</sup>. Loverre et al. first reported that CD133+ cells were significantly  
8 increased in patients with delayed graft function resulting from an acute injury, compared to the  
9 pre-transplant biopsy or to patients with early graft function<sup>44</sup>. Three other studies reported an  
10 increase of tubular CD133+ cells with respect to normal tissue, after acute tubular necrosis in  
11 transplanted patients or in proteinuric glomerular diseases<sup>3,37,41</sup>. The involvement of these cells  
12 in renal repair is strongly supported by the evidence that almost all proliferating cells in acute  
13 tubular injury biopsies were CD133/CD24 positive. In addition, these cells co-expressed the  
14 KIM injury marker<sup>37</sup>. Interestingly, the number of CD133+ cells in patients with acute tubular  
15 necrosis appeared to be a predictive marker of tissue recovery<sup>45</sup>.

16        The presence of cells expressing progenitor markers was also reported in chronic tissue  
17 injury. Stripes and clusters of proliferating tubular CD133+ cells were detected in biopsies of  
18 patients with chronic tubule-interstitial damage due to urinary obstruction or focal segmental  
19 glomerulosclerosis (FSGS), suggesting the activation of mechanisms for tissue repair<sup>3,41</sup>. It  
20 would be interesting to understand whether the loss of the regenerative ability in chronically  
21 damaged human tissue could possibly be ascribed to cell exhaustion, or to mechanisms of cell  
22 senescence. Interesting, blocking the G2/M phase of the cycle was related to a senescent  
23 secretory phenotype occurring in tubular cells after repeated injury<sup>46</sup>.

24        The involvement of progenitor-like cells with enhanced proliferative ability also occurs when  
25 abnormal proliferation underlines pathological settings, such as in the glomerular tuft during  
26 acute glomerulonephritides and in cells forming cysts in polycystic kidney disease. Indeed,  
27 proliferation of CD133+ cells was reported in glomerular hypercellular lesions, and  
28 (pseudo)crescents leading to Bowman's space obliteration occurring in collapsing  
29 glomerulopathy and crescentic glomerulonephritis, and within glomerular tip lesions at the  
30 urinary pole of the Bowman's capsule<sup>20,47</sup>. Similarly, in polycystic kidney disease, all cyst-lining  
31 cells strongly expressed CD133<sup>48</sup>.

32



1 The results of the above-mentioned studies in human renal tissue clearly indicated the  
2 existence, in different nephron segments, of cells that can be defined as adult RPCs by a  
3 functional definition of progenitor cells, as they are able to proliferate, participate in tissue  
4 homeostasis and repair, and survive to injury. In addition, from a phenotypic point of view, they  
5 lack a differentiated phenotype and express stem-related markers. It seems plausible that adult  
6 human CD133+ RPCs represent the same population described as segment-restricted  
7 progenitors in mice. The debate on the fixed nature of renal progenitors or on the induction of  
8 their phenotype is of difficult answer, as no lineage tracing can be applied to human studies.  
9 However, cellular plasticity is the most likely explanation. This has been suggested for organs  
10 other than kidney, such as the lung, bone marrow and pancreas. Indeed, increasing body of  
11 evidence supports the argument that the stem/progenitor-related phenotype is not static but  
12 rather a dynamic process characterized by continuous cycle and regulated by microvesicle-  
13 related microenvironmental stimuli (Quesemerry Leukemia. 2014 Apr;28(4):813-22.).

14  
15

## 16 **B. RPC cultures**

17 Most of the data supporting the phenotypic and functional characteristics of RPCs have  
18 derived from the analysis of their behaviour and phenotype once cultured and isolated. Many  
19 different RPCs with repair ability have been isolated from adult mouse tissue using stem-  
20 related characteristics such as marker expression, BrdU label retention, side population  
21 labelling or clonogenicity<sup>49-53</sup>. These cell lines showed variability in marker expression, origin,  
22 potency and differentiation ability. More homogeneous and unifying results were obtained in  
23 humans, using the strategies depicted in the following paragraphs.

24

### 25 *Isolation and characterization of RPCs from human renal tissue*

26 We first isolated and characterized cells expressing the AC133-detected CD133 stem  
27 cell marker from the human renal cortex<sup>54</sup>. CD133<sup>+</sup> cells were subsequently isolated from the  
28 Bowman's capsule of human glomeruli<sup>35</sup>, as well as from the inner medulla<sup>36,38</sup>. These CD133<sup>+</sup>  
29 populations, isolated from different nephron segments, shared similar phenotypic properties, as  
30 they all expressed foetal renal markers, such as Pax-2, Wt-1, and CD24, along with several  
31 mesenchymal stem cell markers, such as CD29, CD73 and CD90, with minor variations in  
32 expression levels, possibly related to cell culture conditions. This was further confirmed by

1 analysis of a gene expression profile, which clustered CD133+ cells from glomeruli and tubules  
2 against epithelial differentiated cells<sup>55</sup>. However, some genes were differentially expressed,  
3 including VCAM- 1, also known as CD106, which was expressed approximately 300-fold more  
4 in CD133+CD24+ cells obtained from glomerular outgrowths than from tubules<sup>41</sup>. Functionally,  
5 CD133+ RPCs were clonogenic, and, most importantly, differentiated into tubular cells of  
6 different nephron segments and glomerular podocytes after appropriate stimulation<sup>35,54</sup>. When  
7 injected within Matrigel *in vivo* in immunodeficient mice, RPCs also generated tubular  
8 structures expressing nephron segment markers<sup>36,54</sup>. In addition, CD73<sup>+</sup>/CD133<sup>+</sup> RPCs  
9 isolated from the inner medulla were shown to synthesize and release erythropoietin under  
10 hypoxia<sup>56</sup>.

11 As CD133 cultures were generated by cell sorting, they should represent the CD133  
12 expressing cells in the renal tissue. However, it has been noticed that total cell cultures of  
13 kidney cells may express CD133 at fairly high levels<sup>57</sup>. A possible explanation is the selective  
14 ability of CD133+ RPCs to survive in culture. Another possibility is that CD133 could also be  
15 acquired by tubular cells due to culture-induced dedifferentiation. This has been recently  
16 excluded for podocytes that did not participate in the generation of CD133+ cells cultured from  
17 urine<sup>58</sup> but it could be a possible mechanism in tubular-derived CD133+ cell culture. In any  
18 case, plasticity of cultured CD133+ RPCs may provide insights in the mechanisms of kidney  
19 repair and could be exploited for renal regeneration.

20 Other selection strategies based on functional assays were utilized to identify and select  
21 RPCs from human renal tissue. For example, cells were sorted based by means of a functional  
22 assay that which evaluates the stem-related activity of ALDH<sup>39</sup>. ALDH<sup>high</sup> cells expressed  
23 CD133 and vimentin appeared were similar in appearance to CD133- sorted cells<sup>39</sup>. Another  
24 functional assay commonly used for stem cell isolation is the ability to growth as spheres. The  
25 so-called nephrospheres contained cells at different levels of maturation<sup>59</sup>. Selection of the  
26 quiescent cells within the nephrospheres using retention of the PKH<sub>26</sub> dye identified a self-  
27 renewing population capable of multiple differentiations into cells with epithelial, endothelial  
28 and podocytic characteristics. Interestingly, these cells also expressed a CD133<sup>+</sup>/CD24<sup>-</sup>  
29 phenotype<sup>59</sup>.

30 A different approach was proposed by the group of Dekel, who exploited the neo-  
31 expression of the foetal marker NCAM, by a fraction of human renal cells in culture<sup>57</sup>. Indeed,  
32 this marker is absent in normal adult renal tissue, but it can be reactivated after ischaemia-

1 reperfusion injury<sup>60</sup>. NCAM+ RPCs were shown to be enriched in renal progenitor/stem  
2 transcriptional factors (SIX2, SALL1, PAX2, and WT1 and Oct4) and to present a  
3 mesenchymal phenotype. In addition, NCAM positive cells showed enhanced functional  
4 properties related to stemness (clonogenicity, sphere-forming ability and tubulogenic  
5 differentiation on single-cell grafting in chick chorioallantoic membranes and mice)<sup>57</sup>.  
6 All the described cell lines of human RPCs possess a limited life span and undergo  
7 senescence over time. It is reasonable to suggest that no robust, long-term culture system  
8 currently exists that is capable of maintaining potent, clonal expansion of adult progenitors over  
9 long periods of time under defined conditions.

10

11 *Isolation of RPCs from urine*

12 It has been shown that freshly voided urine<sup>66</sup> and urine from the upper urinary tract<sup>67</sup>  
13 contain multipotent stem cells. In addition, CD133/CD24 RPCs have been isolated and cultured  
14 from the urine of normal subjects as well as from patients with kidney disorders<sup>58</sup>. Urine-  
15 derived RPCs were similar to tissue-derived RPCs at a phenotypic and functional level, and  
16 differentiated into both tubular cells and podocytes. In addition, urine-derived CD133/CD24  
17 RPCs were used for the diagnosis of glomerular genetic disorders<sup>58</sup>.  
18 The release of foetal urine into the amniotic fluid represented the rationale in the search for  
19 RPCs in the amniotic fluid. Da Sacco *et al.*<sup>68</sup> have successfully isolated, from the amniotic fluid,  
20 a sub-population of metanephric mesenchyme-like cells committed to nephron lineages, and  
21 capable of differentiating into podocytes.

22 Urine-derived cells may represent an attractive source of autologous cells for cell therapy. In  
23 addition, they could be exploited as diagnostic tool for genetic glomerular disorders.

24

25 *Isolation of renal mesenchymal stromal cells from renal tissue*

26 Cells with a mesenchymal origin and multipotent properties are known to be present in a  
27 population adjacent to endothelial cells of the microvasculature, on the basis of marker  
28 expression (CD146+ NG2+ CD140b+), and may possibly contribute to tissue support<sup>61</sup>. Renal  
29 mesenchymal stromal cells (MSCs) isolated from the mouse kidney, differentiated *in vivo* into  
30 renin-producing cells<sup>62</sup>, indicating a commitment towards renal cell types. Interestingly, a  
31 recent paper from Little's group showed that MSC-like cells from mouse collecting ducts can be  
32 isolated from adult murine kidneys due to a phenomenon of epithelial-mesenchymal transition

1 after long-term culture *in vitro*<sup>63</sup>. The obtained cells give rise to a long-term self-renewing MSC  
2 population, which shows the characteristic pan-mesodermal differentiation. In addition, renal  
3 MSCs possess the ability to integrate into aquaporin 2–positive medullary collecting ducts  
4 when injected into the kidneys of neonatal mice, unlike bone marrow-derived MSCs<sup>64</sup>. The  
5 differentiation of renal MSCs into renal tissue is supported by data showing the expression of a  
6 selected pattern of genes in renal MSCs, compared with MSCs of marrow origin, possibly  
7 related to a memory of tissue origin<sup>64</sup>. This suggests that renal MSCs may display organ-  
8 specific regenerative capacities that are highly superior to those of MSCs from unrelated  
9 organs, which could thus be exploited for therapeutic applications<sup>64</sup>.

10 In human kidneys, resident MSCs were investigated in glomeruli, based on their vascular  
11 nature, and characterized as CD133-/CD146+ cells<sup>65</sup>. Glomeruli-derived MSCs displayed self-  
12 renewal capability, clonogenicity, and the ability to differentiate into the different glomerular cell  
13 types (endothelial, mesangial and epithelial cells). Evidence that glomerular MSCs do not  
14 derive from the bone marrow has been provided by detection of MSCs derived from donors in a  
15 sex-unmatched transplant<sup>65</sup>.

16  
17

### 18 **C. Modulation or induction of an RPC phenotype**

#### 19 *Micro-environmental stimuli for RPC modulation*

20 As described above, the nature of the cells involved in kidney homeostasis/regeneration  
21 is unclear, and further studies are required to define the mechanisms involved in their possible  
22 activation<sup>69</sup>. According to the idea of a plastic phenotype of RPCs, it can be speculated that the  
23 progenitor phenotype could be a consequence of a cell cycle state, or rather it may result from  
24 the modulation of growth and differentiation of resident cells by external stimuli. Interestingly,  
25 soluble signals released by a diseased kidney were shown to induce the generation of  
26 scattered tubular cells in the contralateral kidney in mice, supporting the hypothesis of cell  
27 reprogramming by the microenvironment<sup>27,70</sup>. Knowledge of the activating mechanisms  
28 involved in the modulation of the RPC phenotype could be relevant, both in terms of a  
29 pharmacological approach for local cell modulation, and in terms of possible *ex vivo* treatment  
30 of cells before their *in vivo* injection.

1 A hypoxic environment appears to activate RPC properties. Hypoxia was reported to  
2 increase proliferation, clonogenicity and CD133 expression in RPCs isolated from the inner  
3 medulla<sup>36</sup>. Indeed, consistent HIF-1 activation was shown in these RPCs, possibly related to  
4 the hypoxic condition of the tissue of origin. The mechanism of hypoxia-related acquisition of  
5 progenitor properties may rely on the hypoxia-induced expression of the stem cell transcription  
6 factor OCT4A and its regulation by micro-RNA 145<sup>36</sup>. In addition, in cortical RPCs, a  
7 comparative analysis between RPCs and differentiated tubular cells identified two microRNAs  
8 involved in the regulation of stemness and differentiation of RPCs. In particular, miR-1915  
9 targeted the progenitor factors CD133 and PAX2, and miR-1225-5p targeted toll-like receptor 2  
10 (TLR2) expression, which is a key gene of damage signalling, which is up-regulated in RPCs<sup>71</sup>.  
11 In addition, Wnt signalling has been linked to RPC maintenance in the mouse, as renal cells  
12 with cloning ability showed increased activation of this pathway<sup>2</sup>.

### 13 *RPC generation from iPS or direct generation of inducible-RPC*

14 Recent advances in the stem cell field could facilitate the generation of RPCs with  
15 nephrogenic potential (metanephric mesenchyme-like progenitors) from somatic cells, through  
16 the generation of inducible pluripotent stem cells (iPS) or directly from embryonic stem cells.  
17 The obvious advantage in the use of iPS is the possible generation of autologous cells and the  
18 lack of ethical issues. IPS were successfully shown to generate entire renal structures as well  
19 as renal organoids (ref). Moreover, two recent investigations refined the methods for human  
20 intermediate mesoderm progenitor generation from iPS using small chemical inhibitors,  
21 obtaining high efficiency and rapidity<sup>73,74</sup>. Araoka *et al*<sup>73</sup> successfully generated human  
22 intermediate mesenchymal progenitors (80% efficiency in 5 days) from iPS using retinoids and  
23 an activator of the Wnt pathway. The resulting cells showed the ability to differentiate into  
24 multiple cell types, and to form renal tubule-like structures. Similarly, Lam *et al*.<sup>74</sup>, by using  
25 retinoic acid and Wnt activator plus FGF2, produced intermediate mesenchymal cells within 3  
26 days. In addition, intermediate progenitors, if treated with FGF9 and activin, further  
27 differentiated into Six2+ metanephric mesenchyme progenitors. The generation of autologous  
28 cells with nephrogenic potential is highly promising in regenerative medicine of the kidney.  
29 However, tumorigenic potential of these cells and the scarce proliferation of iPS-differentiated  
30 cells still represent a significant limitation for their application.

1 An alternative approach for RPC generation is the direct reprogramming of renal cells  
2 into progenitors. The group of Little successfully identified, by means of sequential screening, a  
3 cocktail of six genes (SIX1, SIX2, OSR1, EYA1, HOXA11, and SNAI2) able to reprogram  
4 human tubular cells into metanephric mesenchymal progenitors<sup>75</sup>. These progenitors showed  
5 the capacity to contribute to the nephrogenic compartment of a developing embryonic kidney  
6 *ex vivo*. Although this study represents a huge step towards the generation of progenitors for *in*  
7 *vivo* use, improved optimization of transfection, including the use of non-integrating or inducible  
8 vectors, as well as establishing ideal growth conditions is still required for therapeutic  
9 applications.

#### 10 **D. Therapeutic use of RPCs in experimental models of kidney injury**

11 Cell therapy is clearly the most promising tool to provide regenerating cells/stimuli to the renal  
12 tissue (Table III, IV and Figure 1). In zebrafish, Diep and colleagues<sup>76</sup> clearly showed that  
13 transplanting RPCs with nephrogenic potential generated multiple nephrons. In the fish the  
14 mesenchymal nephron progenitor population persists supporting *de novo* nephrogenesis.  
15 In human, this potential is not present in the adult and therefore RPCs do not generate new  
16 nephrons, but rather may support the process of nephron repair. However, a mesenchymal  
17 nephron progenitor population could facilitate the creation and integration of new  
18 nephrons.

#### 19 *Cell therapy with embryonic/ foetal RPCs*

20 Human RPCs isolated from foetal kidneys were shown to exert a therapeutic effect in  
21 acute and chronic renal injury. CD24+/CD133+ RPCs isolated from human foetal tissue  
22 ameliorated renal function after glycerol-induced AKI<sup>21</sup>. Using this model, the effects were  
23 comparable to those of adult RPCs. Human foetal nephron progenitor NCAM+ cells were  
24 tested in the 5/6-nephrectomy kidney injury model characterized by loss of renal mass and  
25 progression. In this model, cells were administered directly into the renal parenchyma in three  
26 consecutive injections, and were shown to engraft and integrate in diseased murine kidneys,  
27 with resulting beneficial effects on renal function and arrest of disease progression<sup>19</sup>.  
28 Metanephric mesenchyme-like RPCs generated by pluripotent stem cells also showed a

1 considerable differentiation potential *in vitro* and *in vivo* in the embryonic kidney<sup>72</sup>. Recently,  
2 RPCs derived from human iPSCs were tested in a murine model of AKI. Interestingly, they not  
3 only restored renal function but also showed a robust engraftment into damaged tubuli  
4 (Imberti).

#### 5 *Cell therapy with adult RPCs*

6 Pioneer experiments in acute models of tubular or glomerular damage showed a  
7 potential therapeutic effect of the administration of adult rat/murine RPCs of different type and  
8 origin<sup>49,51-53,77</sup> (Table III). Similarly, administering adult CD24+/CD133+ RPCs from the  
9 Bowman's capsule, cortical tubular compartment, and inner medulla to immunodeficient mice  
10 with rhabdomyolysis-induced AKI was shown to have therapeutic benefits<sup>35,41,54,78</sup> (Table IV).  
11 In these experiments, a single dose of CD133+ cells (ranging from  $5 \times 10^5$  to  $1 \times 10^6$  cells) was  
12 administered, after induction of renal damage. This cell therapy improved renal function, as  
13 shown by the rapid decrease of creatinine and blood urea, together with amelioration of tissue  
14 histology<sup>35,41,54,78</sup>. These results correlated closely with those commonly obtained using MSCs,  
15 and comparison of MSCs with medulla-derived CD133+ cells demonstrated an equivalent  
16 therapeutic effect<sup>78</sup>. However, a bio-distribution study showed that CD133+ cells preferentially  
17 localized to the kidney region, with less cell dispersion in other parenchymal organs, with  
18 respect to MSCs, when analysed by optical imaging 48 hours after injection<sup>78</sup>. Similarly,  
19 administration of CD133+CD24+ cells isolated from the Bowman's capsule promoted renal  
20 amelioration in a model of FSGS, characterized by podocyte injury, reducing proteinuria and  
21 improving chronic glomerular damage<sup>40</sup>. This effect was restricted to undifferentiated  
22 CD133+CD24+PDX- cells, whereas no effect was observed using partly differentiated  
23 CD133+CD24+PDX+ renal cells, or CD133 negative cells<sup>40</sup>. Moreover, CD133+CD24+ cells  
24 isolated from urine similarly differentiated into podocytes and reduced proteinuria in the  
25 adriamycin model<sup>58</sup>.

26 Using the same glycerol-induced AKI model used for CD133+ RPCs, the therapeutic  
27 effect of NCAM+ RPC overexpressed markers of fetal RPCs was compared with that of the  
28 total human kidney epithelial cell culture, uniformly positive for CD24/CD133. Both populations  
29 displayed a therapeutic effect, with no difference between NCAM+ and NCAM- cells<sup>57</sup>.

1 This observation may suggest a possible effect for cell therapy using renal epithelial  
2 cells, as reported below. By analogy mature tubular epithelial renal-like cells, obtained from  
3 differentiation of spermatogonial derived embryonic-like stem cells, protected against acute and  
4 subsequent chronic damage *in vivo* in a model of kidney ischaemia<sup>79</sup>. However, the he  
5 administration of human CD133 negative cells did not provide benefit in tubular or glomerular  
6 acute damage<sup>35,40,41</sup>. As the CD133 negative population was not fully characterized, it can not  
7 be determined whether the lack of a therapeutic effect may be related to the activity of  
8 fibroblasts or rather of renal epithelial cells.

9 *Mechanisms of action of RPCs*

10 An intriguing point of discussion in renal regenerative medicine is the mechanism of  
11 effect of the administered cells. While the effect ascribed to MSCs is mainly related to  
12 paracrine delivery of soluble mediators and bioactive vesicles containing genetic material  
13 (mRNA and microRNA),<sup>80</sup> mechanisms of both paracrine stimulation and tissue integration  
14 have been described for human RPCs. Detection of the injected adult RPCs in the kidneys of  
15 AKI mice was reported in both proximal and distal tubules<sup>35,41,54</sup>. Analysis of cell integration  
16 after 11 days showed that around 7% of the total number of tubular cells co-expressed the  
17 labelling dye and tubule markers<sup>35</sup>. Similarly, in a model of podocyte damage,  
18 CD133+CD24+PDX- cells generated new podocytes, representing of all podocytes after 7 days  
19  $11.08 \pm 3.3\%$ , and maintaining their engraftment after 45 days<sup>40</sup>. Interestingly, the ability to  
20 differentiate into podocytes was restricted to Bowmann's capsule isolated RPCS, that also  
21 differentiated into tubular cells, whereas tubular derived RPCS cells did not<sup>41</sup>. However, RPCs  
22 isolated from inner medulla showed a limited persistence within the murine tissue at 1 month  
23 after the injection<sup>78</sup>. High tissue integration has been also reported for foetal RPCs directly  
24 injected into the renal parenchyma in a 5/6 nephrectomized kidney<sup>19</sup>. Tissue analysis showed  
25 three distinct patterns of engraftment: initially, RPCs were located within existing mouse  
26 tubules or within the interstitium, and after 3 months they tended to organise into large mature-  
27 appearing epithelial tubules with a patent lumen, resembling segment nephrogenesis<sup>19</sup>.

28 In addition, RPCs were described to release a large variety of growth factors and vesicles that  
29 may contribute to tissue repair. In particular IL-15, a kidney specific factor involved in renal



1 differentiation<sup>81</sup>, was selectively released by CD133+ cells<sup>78</sup>. RPC-derived extracellular  
2 vesicles may also be involved in the process of repair, as they promoted *in vitro* survival and  
3 proliferation in target epithelial cells<sup>82</sup>.

4         The data from the literature and also from our laboratory indicate a variable number of  
5 RPC engraftment within the tubules. Whether the cell engraftment in the kidney entirely  
6 accounts for repopulation of tubular cells is questionable as no more than 10-11% human cells  
7 are detected within regenerating tubules and the evidence for their differentiation in mature  
8 epithelial cells are weak. The relevance of RPC integration would require dedicated long term  
9 experiments. In fact, cell integration both in AKI and in CKD models was assessed only up to 2  
10 weeks. We suggest a combined mechanism of paracrine support and integration in the effects  
11 of RPCs observed in pre-clinical studies. It is conceivable that the engrafted RPCs may  
12 coordinate the endogenous repair.

13 It could be hypothesized that a paracrine mechanism, as described for MSCs, could be  
14 sufficient for a therapeutic effect in AKI. This was also confirmed for kidney-derived MSCs as  
15 the therapeutic effects were recapitulated by administration of kidney-MSC-derived  
16 extracellular vesicles<sup>83,84</sup>. Indeed, in acute damage, activation of resident responsive  
17 regenerative cells could be sufficient to support repair. At variance, the integration mechanisms  
18 of RPCs find a specific need in CKD, as regenerating responsive cells could be exosted and  
19 the integration of new functional cells to regain function would be required.

20         Therefore, in acute settings, the use of MSCs appears clinically more applicable than  
21 that of RPCs, which show limited cell sources and expansion, undefined culture conditions, and  
22 poorly accessible or inadequate autologous tissue. In chronic settings, RPCs that are  
23 committed to the generation of nephron cells would represent the ideal cell type to obtain  
24 clinically-relevant beneficial effects. In addition, the use of patient-derived RPCs, from tissue or  
25 from urine, could have significant relevance in CKD as a possible autologous therapy that does  
26 not require immunosuppression. It would be of interest to test the effect of adult RPCs in a  
27 model of CKD characterized by nephron loss such as the 5/6 nephrectomy used for foetal  
28 RPCs<sup>19</sup>. In addition, data on the application of RPCs of adult and foetal sources should be  
29 obtained, and comparatively analysed.

1 Differential results may also relay on different modalities of cell delivery. Is it well  
2 established that i.v. injected cells may be entrapped in the lungs and liver and thus release  
3 pacrine factors from this distal sites. In addition, i.v. or intra-arterially delivered cells need to  
4 overcome two basement membranes to reach the tubular compartment. This appears  
5 achievable in acute injury, when those barriers are partly damaged due to tissue injury. At  
6 variance, during chronic injury, an intra-parenchymal application may be necessary for  
7 localization of injected cells within the nephron. Indeed, intra-parenchymal delivery was chosen  
8 to administer RPCs or differentiated tubular cells in models of murine CKD (Table III and IV).  
9 This route of administration presents the advantage to require a low number of cells and to  
10 avoid cell dispersion in undesired organs. In contrast, it might be associated with adverse  
11 events such as local necrosis, thrombosis and limited distribution of the cells within the kidney.

#### 12 *Cell therapy with adult renal-derived cells*

13 Interestingly, a therapeutic effect of administering adult differentiated cell populations  
14 isolated from normal kidneys has been reported by a number of studies. Human renal cultures  
15 enriched in erythropoietin-positive cells were administered to the renal parenchyma in a model  
16 of chronic renal damage<sup>85</sup>. Results showed that these cells improved renal function and  
17 reduced renal fibrosis and inflammation, unlike erythropoietin-depleted cells. However, cells did  
18 not survive up to 12 weeks, suggesting a rapid paracrine effect. Other studies indicated the  
19 therapeutic effect of a tubular cell-enriched renal cell population<sup>86</sup>. Indeed, intra-parenchymal  
20 injection of isolated and cultured epithelial cells to mice with established chronic kidney injury  
21 improved renal function, extended survival and reduced the severity of kidney tissue pathology.  
22 This epithelial cell subpopulation was mainly composed of cells from the tubular and collecting  
23 duct and was depleted of glomerular, vascular and erythropoietin-producing cells. Similarly, a  
24 rat tubule-enriched epithelial cell population promoted regeneration in diabetic rats when  
25 injected intra-parenchymally, and stabilized disease progression up to 1-year post-treatment<sup>87</sup>.  
26 The mechanism ascribed to the therapeutic effect of the intra-parenchymal injected cells in  
27 CKD mice kidney was the activation of a regenerative response process. In particular, after 3 to  
28 6 weeks, a number of genes were upregulated including the pluripotency-associated SOX2  
29 gene and the RPC gene CD24<sup>88</sup>. These data could possibly support the regenerative capacity  
30 of differentiated cells, as proposed by Kusaba et al<sup>26</sup>.

1 The strategy of intraparenchymal injection of adult renal cells is under investigation in a phase I  
2 trial in human patients suffering from CKD, using expanded autologous renal cells (Neo-Kidney  
3 Augment™ Cellular Therapy; ClinicalTrials.gov Identifier: NCT01846715). Human cells were  
4 isolated by biopsies from patients and expanded following a described protocol that allowed  
5 generation of an injectable large number of cells<sup>89</sup>. The obtained bioactive renal cells  
6 embedded into gelatine hydrogels were intra-parenchymally administered to patients (5 to  
7 date). The results of this trial have not yet been published. It will certainly be of interest to  
8 acquire knowledge on the mechanisms involved in the therapeutic effects of the administered  
9 cells, taking into account a possible, although limited, cell dedifferentiation into RPCs.

10

## 11 **Conclusions and perspectives**

12 While foetal cells with nephron-forming activity have been characterized in terms of  
13 molecular profile, the nature of the cells involved in kidney homeostasis/regeneration is  
14 unclear<sup>67</sup>, and further studies are required to understand the mechanisms involved in their  
15 possible activation. Adult human RPCs isolated from different nephron compartments appear  
16 to display progenitor-like properties, and provided a therapeutic effect in models of acute and  
17 chronic renal damage<sup>35,40,41,54,76</sup>. In acute pre-clinical models, their therapeutic efficacy was  
18 comparable to MSCs<sup>76</sup>, which, in turn, may benefit from more readily-available sources and  
19 reduced immune compatibility issues (Figure 1).

20 Although adult RPCs successfully engrafted within the renal tissue in acute models, their  
21 effect in a model of CKD with nephron loss needs to be established. Indeed, as they are  
22 committed to the kidney cell progeny, they could obviate the potential development of  
23 inappropriate cell types, and they would represent an ideal cell type to obtain clinically-relevant  
24 beneficial effects in patients with renal failure. However, the source of these cells is limited and  
25 they possess a short life span, undergo senescence over time and have a restricted  
26 nephrogenic potential.

27 Foetal-like RPCs with nephrogenic potential seem to be the ideal cell type for cell-based  
28 therapy in CKD. The in vitro generation of renal progenitors by iPS differentiation is of particular  
29 interest to overcome ethical issue and to get autologous tissue. However, it not only requires

1 complex and long differentiation passages, but it also has the limitation that once differentiated,  
2 progenitors do not proliferate in culture. The direct reprogramming of renal cells into induced  
3 RPCs may overcome this issue and generate expandable progenitors with high transfection  
4 efficacy. However, at present, no long-term culture system has proven to be capable of  
5 maintaining potent expansion of human adult progenitors over an extended period. Finally,  
6 adult renal cells unexpectedly appeared to be of therapeutic potential, and could be considered  
7 as a possible simple alternative for kidney regeneration. Increasing information on the nature of  
8 this population, including the presence of RPCs, and on the mechanism of action, is required.

9  
10

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

References

1. Perico, N. & Remuzzi, G. Chronic kidney disease: a research and public health priority. *Nephrol. Dial. Transplant.* **27**, iii19-26 (2012)
- Jha V, Garcia-Garcia G, Iseki K, et al. Chronic kidney disease: global dimension and perspectives. *Lancet* 2013;382:260-72
2. Rinkevich Y, et al. In vivo clonal analysis reveals lineage-restricted progenitor characteristics in mammalian kidney development, maintenance, and regeneration. *Cell Rep.* **7**, 1270-83 (2014)
3. Hansson, J. et al. Evidence for a morphologically distinct and functionally robust cell type in the proximal tubules of human kidney. *Hum. Pathol.* **45**, 382-93 (2014).
4. Metsuyanin, S. et al. Accumulation of malignant renal stem cells is associated with epigenetic changes in normal renal progenitor genes. *Stem Cells* **26**, 1808-17 (2008).
5. Chawla, L.S., Eggers, P.W., Star, R.A. & Kimmel, P.L. Acute kidney injury and chronic kidney disease as interconnected syndromes. *N. Engl. J. Med.* **371**, 58-66 (2014).
6. Costantini, F. & Kopan R. Patterning a complex organ: branching morphogenesis and nephron segmentation in kidney development. *Dev. Cell.* **18**, 698-712 (2010).
7. Boyle, S. et al. Fate mapping using Cited1-CreERT2 mice demonstrates that the cap mesenchyme contains self-renewing progenitor cells and gives rise exclusively to nephronic epithelia. *Dev. Biol.* **313**, 234-45 (2008).
8. Kobayashi, A. et al. Six2 defines and regulates a multipotent selfrenewing nephron progenitor population throughout mammalian kidney development. *Cell Stem Cell* **3**, 169-181 (2008).
9. Kopan, R., Chen, S. & Little, M. Nephron progenitor cells: shifting the balance of self-renewal and differentiation. *Curr. Top Dev. Biol.* **107**, 293-331 (2014).
10. Kanda, S. et al. Sall1 maintains nephron progenitors and nascent nephrons by acting as both an activator and a repressor. *J. Am. Soc. Nephrol.* **25**, 2584-95 (2014).
11. Kalatzis, V., Sahly, I., El-Amraoui, A. & Petit, C. Eya1 expression in the developing ear and kidney: towards the understanding of the pathogenesis of Branchio-Oto-Renal (BOR) syndrome. *Dev. Dyn.* **213**, 486-99 (1998).

- 1 12. Xu, J., Liu, H., Park, J.S., Lan, Y. & Jiang, R. Osr1 acts downstream of and interacts  
2 synergistically with Six2 to maintain nephron progenitor cells during kidney organogenesis.  
3 *Development* **141**, 1442-52 (2014).
- 4 13. Torres, M. *et al.* Pax-2 controls multiple steps of urogenital development. *Development*  
5 **121**, 4057– 4065 (1995).
- 6 14. Kreidberg, J.A. *et al.* WT-1 is required for early kidney development. *Cell* **74**, 679–691  
7 (1993).
- 8 15. Barker, N. *et al.* Lgr5(+ve) stem/progenitor cells contribute to nephron formation during  
9 kidney development. *Cell Rep.* **2**, 540-52 (2012).
- 10 16. Fujimura, S., Jiang, Q., Kobayashi, C. & Nishinakamura, R. Notch2 activation in the  
11 embryonic kidney depletes nephron progenitors. *J. Am. Soc. Nephrol.* **21**, 803-10 (2010).
- 12 17. Dekel, B. *et al.* Multiple imprinted and stemness genes provide a link between normal and  
13 tumor progenitor cells of the developing human kidney. *Cancer Res.* **66**, 6040-9 (2006).
- 14 18. Metsuyanin, S. *et al.* Expression of stem cell markers in the human fetal kidney. *PLoS One*  
15 **4**, e6709 (2009).
- 16 19. Harari-Steinberg O, *et al.* Identification of human nephron progenitors capable of  
17 generation of kidney structures and functional repair of chronic renal disease. *EMBO Mol. Med.*  
18 **5**, 1556-68 (2013)
- 19 20. Kim, K. *et al.* Expression of stem cell marker CD133 in fetal and adult human kidneys and  
20 pauci-immune crescentic glomerulonephritis. *Histol. Histopathol.* **26**, 223–232 (2011).
- 21 21. Lazzeri, E. *et al.* Regenerative potential of embryonic renal multipotent progenitors in acute  
22 renal failure. *J. Am. Soc. Nephrol.* **18**, 3128-38 (2007).
- 23 22. Bussolati, B., Collino, F. & Camussi, G. CD133+ cells as a therapeutic target for kidney  
24 diseases. *Expert Opin. Ther. Targets.* **16**, 157-65 (2012).
- 25 23. Hartman, H.A., Lai, H.L. & Patterson, L.T. Cessation of renal morphogenesis in mice. *Dev.*  
26 *Biol.* **310**, 379–387 (2007).
- 27 24. Maeshima, A. *et al.* Involvement of Pax-2 in the action of activin A on tubular cell  
28 regeneration. *J. Am. Soc. Nephrol.* **13**, 2850–59 (2002).
- 29 25. Humphreys, B.D. *et al.* Intrinsic epithelial cells repair the kidney after injury. *Cell Stem Cell*  
30 **6**, 284-91 (2008).
- 31 26. Humphreys, B.D. *et al.* Repair of injured proximal tubule does not involve specialized  
32 progenitors. *Proc. Natl. Acad. Sci. USA* **108**, 9226-231 (2011).

- 1 27. Berger, K. *et al.* Origin of regenerating tubular cells after acute kidney injury. *Proc. Natl.*  
2 *Acad. Sci. USA* **111**, 1533-8 (2014).
- 3 28. Berger, K. *et al.* The regenerative potential of parietal epithelial cells in adult mice. *J. Am.*  
4 *Soc. Nephrol.* **25**, 693-705 (2014).
- 5 Berger, K, & Moeller, M.J. Mechanisms of epithelial repair and regeneration after acute kidney  
6 injury. *Semin. Nephrol.* **34**, 394-403 (2014).
- 7 29. Maeshima, A., Yamashita, S. & Nojima, Y. Identification of renal progenitor-like tubular cells  
8 that participate in the regeneration processes of the kidney. *J. Am. Soc. Nephrol.* **14**, 3138-46  
9 (2003).
- 10 30. Oliver, J.A. *et al.* The renal papilla is a niche for adult kidney stem cells. *J. Clin. Invest.* **114**,  
11 795-804 (2004).
- 12 31. Langworthy, M., Zhou, B., de Caestecker, M., Moeckel, G. & Baldwin, H.S. NFATc1  
13 identifies a population of proximal tubule cell progenitors. *J. Am. Soc. Nephrol.* **20**, 311-21  
14 (2009).
- 15 32. Rangel, E.B. *et al.* C-kit(+) cells isolated from developing kidneys are a novel population of  
16 stem cells with regenerative potential. *Stem Cells* **31**, 1644-56 (2013).
- 17 33. Yin, A.H. *et al.* AC133, a novel marker for human hematopoietic stem and progenitor cells.  
18 *Blood* **90**, 5002–5012 (1997).
- 19 34. Kemper, K. *et al.* The AC133 epitope, but not the CD133 protein, is lost upon cancer stem  
20 cell differentiation. *Cancer Res.* **70**, 719-29 (2010).
- 21 35. Sagrinati, C. *et al.* J. Isolation and characterization of multipotent progenitor cells from the  
22 Bowman's capsule of adult human kidneys. *J. Am. Soc. Nephrol.* **17**, 2443-56 (2006).
- 23 36. Bussolati, B. *et al.* Hypoxia modulates the undifferentiated phenotype of human renal inner  
24 medullary CD133+ progenitors through Oct4/miR-145 balance. *Am. J. Physiol. Renal Physiol.*  
25 **302**, F116-28 (2012)
- 26 37. Smeets, B. *et al.* Proximal tubular cells contain a phenotypically distinct, scattered cell  
27 population involved in tubular regeneration. *J. Pathol.* **229**, 645-59 (2013).
- 28 38. Ward, H.H. *et al.* Adult human CD133/1(+) kidney cells isolated from papilla integrate into  
29 developing kidney tubules. *Biochim. Biophys. Acta.* **1812**, 1344-57 (2011).
- 30 39. Lindgren, D. *et al.* Isolation and characterization of progenitor-like cells from human renal  
31 proximal tubules. *Am. J. Pathol.* **178**, 828-37 (2011).
- 32 40. Ronconi E. *et al.* Regeneration of glomerular podocytes by human renal progenitors. *J. Am.*

- 1 *Soc. Nephrol.* **20**, 322-32 (2009).
- 2 41. Angelotti, M.L. *et al.* Characterization of renal progenitors committed toward tubular lineage  
3 and their regenerative potential in renal tubular injury. *Stem Cells* **30**, 1714-25 (2012).
- 4 42. Bourseau-Guilmain, E., Griveau, A., Benoit, J.P. & Garcion, E. The importance of the stem  
5 cell marker prominin-1/CD133 in the uptake of transferrin and in iron metabolism in human  
6 colon cancer Caco-2 cells. *PLoS One* **6**, e25515 (2011).
- 7 43. Griguer, C.E. *et al.* CD133 is a marker of bioenergetic stress in human glioma. *PLoS One*  
8 **3**, e3655 (2008).
- 9 44. Loverre, A., Capobianco, C., Ditunno, P., Battaglia, M., Grandaliano, G. & Schena F.P.  
10 Increase of proliferating renal progenitor cells in acute tubular necrosis underlying delayed graft  
11 function. *Transplantation* **85**, 1112-9 (2008).
- 12 45. Ye, Y. *et al.* Proliferative capacity of stem/progenitor-like cells in the kidney may associate  
13 with the outcome of patients with acute tubular necrosis. *Hum. Pathol.* **42**, 1132-41 (2011).
- 14 46. Canaud, G. & Bonventre, J.V. Cell cycle arrest and the evolution of chronic kidney disease  
15 from acute kidney injury. *Nephrol. Dial. Transplant.* pii: gfu230. [Epub ahead of print] (2014)
- 16 Yang, L., Besschetnova, T.Y., Brooks, C.R., Shah, J.V. & Bonventre, J.V. Epithelial cell cycle  
17 arrest in G2/M mediates kidney fibrosis after injury. *Nat Med.* **16**, 535-43 (2010).
- 18 47. Smeets, B. *et al.* Renal progenitor cells contribute to hyperplastic lesions of  
19 podocytopathies and crescentic glomerulonephritis. *J. Am. Soc. Nephrol.* **20**, 2593-603 (2009)
- 20 48. Carvalhosa, R. *et al.* Cystogenic potential of CD133+ progenitor cells of human polycystic  
21 kidneys. *J. Pathol.* **225**, 129-41 (2011).
- 22 49. Kitamura, S. *et al.* Establishment and characterization of renal progenitor like cells from S3  
23 segment of nephron in rat adult kidney. *FASEB J.* **19**, 1789-97 (2005).
- 24 50. Maeshima, A., Sakurai, H. & Nigam S.K. Adult kidney tubular cell population showing  
25 phenotypic plasticity, tubulogenic capacity, and integration capability into developing kidney. *J*  
26 *Am Soc Nephrol.* **17**, 188-98 (2006).
- 27 51. Gupta, S. *et al.* Isolation and characterization of kidney-derived stem cells. *J. Am. Soc.*  
28 *Nephrol.* **17**, 3028-40 (2006).
- 29 52. Challen, G.A., Bertoncetto, I., Deane, J.A., Ricardo, S.D. & Little MH. Kidney side  
30 population reveals multilineage potential and renal functional capacity but also cellular  
31 heterogeneity. *J. Am. Soc. Nephrol.* **17**, 1896-912 (2006).
- 32 53. Dekel, B. *et al.* Isolation and characterization of nontubular sca-1+lin- multipotent



- 1 stem/progenitor cells from adult mouse kidney. *J. Am. Soc. Nephrol.* **17**, 3300-14 (2006).
- 2 54. Bussolati, B. *et al.* Isolation of renal progenitor cells from adult human kidney. *Am. J.*  
3 *Pathol.* **166**, 545-55 (2005).
- 4 55. Sallustio, F. *et al.* TLR2 plays a role in the activation of human resident renal  
5 stem/progenitor cells. *FASEB J.* **24**, 514-525 (2010).
- 6 56. Bussolati, B., Lauritano, C., Moggio, A., Collino, F., Mazzone, M. & Camussi, G. Renal  
7 CD133(+)/CD73(+) progenitors produce erythropoietin under hypoxia and prolyl hydroxylase  
8 inhibition. *J. Am. Soc. Nephrol.* **24**, 1234-41 (2013).
- 9 57. Buzhor, E. *et al.* Reactivation of NCAM1 defines a subpopulation of human adult kidney  
10 epithelial cells with clonogenic and stem/progenitor properties. *Am. J. Pathol.* **183**, 1621-33  
11 (2013).
- 12 58. Lazzeri, E. *et al.* Human Urine-Derived Renal Progenitors for Personalized Modeling of  
13 Genetic Kidney Disorders. *J. Am. Soc. Nephrol.* pii: ASN.2014010057. [Epub ahead ofprint]  
14 (2015).
- 15 59. Bombelli, S. *et al.* PKH(high) cells within clonal human nephrospheres provide a purified  
16 adult renal stem cell population. *Stem Cell Res.* **11**, 1163-77 (2013).
- 17 60. Abbate, M., Brown, D. & Bonventre, J.V. Expression of NCAM recapitulates tubulogenic  
18 development in kidneys recovering from acute ischemia. *Am. J. Physiol. Renal. Physiol.* **277**,  
19 F454–F463 (1999).
- 20 61. Crisan, M. *et al.* A perivascular origin for mesenchymal stem cells in multiple human  
21 organs. *Cell Stem Cell* **3**:301–313 (2008).
- 22 62. Wang, H. *et al.* Adult renal mesenchymal stem cell-like cells contribute to juxtaglomerular  
23 cell recruitment. *J. Am. Soc. Nephrol.* **24**, 1263-73 (2013).
- 24 63. Li, J. *et al.* Collecting duct-derived cells display mesenchymal stem cell properties and  
25 retain selective in vitro and in vivo epithelial capacity. *J. Am. Soc. Nephrol.* **26**, 81-94 (2015).
- 26 64. Pelekanos, R.A. *et al.* Comprehensive transcriptome and immunophenotype analysis of  
27 renal and cardiac MSC-like populations supports strong congruence with bone marrow MSC  
28 despite maintenance of distinct identities. *Stem Cell Res.* **8**, 58-73 (2012).
- 29 65. Bruno, S. *et al.* Isolation and characterization of resident mesenchymal stem cells in human  
30 glomeruli. *Stem Cells Dev.* **18**, 867-80 (2009).
- 31 66. Zhang, Y. *et al.* Urine derived cells are a potential source for urological tissue  
32 reconstruction. *J. Urol.* **180**, 2226-33 (2008).

- 1 67. Bharadwaj, S. *et al.* Multipotential differentiation of human urine-derived stem cells:  
2 potential for therapeutic applications in urology. *Stem Cells* **31**, 1840-56 (2013).
- 3 68. Da Sacco, S. *et al.* Human amniotic fluid as a potential new source of organ specific  
4 precursor cells for future regenerative medicine applications. *J. Urol.* **183**, 1193-200 (2010).
- 5 69. Kusaba, T. & Humphreys, B.D. Controversies on the origin of proliferating epithelial cells  
6 after kidney injury. *Pediatr. Nephrol.* **29**, 673-9 (2014).
- 7 70. Witzgall, R., Brown, D., Schwarz, C. & Bonventre, J.V. Localization of proliferating cell  
8 nuclear antigen, vimentin, c-Fos, and clusterin in the postischemic kidney. Evidence for a  
9 heterogenous genetic response among nephron segments, and a large pool of mitotically  
10 active and dedifferentiated cells. *J. Clin. Invest.* **93**, 2175-88 (1994).
- 11 71. Sallustio, F. *et al.* miR-1915 and miR-1225-5p regulate the expression of CD133, PAX2  
12 and TLR2 in adult renal progenitor cells. *PLoS One* **8**, e68296 (2013).
- 13 72. Morales, E.E. & Wingert, R.A. Renal stem cell reprogramming: Prospects in regenerative  
14 medicine. *World J. Stem Cells* **6**, 458-66 (2014).
- 15 Taguchi, A., *et al.* Redefining the in vivo origin of metanephric nephron progenitors enables  
16 generation of complex kidney structures from pluripotent stem cells. *Cell Stem Cell.* **14**, 53-67  
17 (2014).
- 18 Takasato, M., *et al.* Directing human embryonic stem cell differentiation towards a renal lineage  
19 generates a self-organizing kidney. *Nat. Cell Biol.* **16**, 118-26 (2014).
- 20 73. Araoka, T. *et al.* Efficient and rapid induction of human iPSCs/ESCs into nephrogenic  
21 intermediate mesoderm using small molecule-based differentiation methods. *PLoS One* **9**,  
22 e84881 (2014).
- 23 74. Lam, A.Q., Freedman, B.S., Morizane, R., Lerou, P.H., Valerius, M.T. & Bonventre, J.V.  
24 Rapid and efficient differentiation of human pluripotent stem cells into intermediate mesoderm  
25 that forms tubules expressing kidney proximal tubular markers. *J. Am. Soc. Nephrol.* **25**, 1211-  
26 25 (2014).
- 27 75. Hendry, C.E. *et al.* Direct transcriptional reprogramming of adult cells to embryonic nephron  
28 progenitors. *J. Am. Soc. Nephrol.* **24**, 1424-34 (2013).
- 29 76. Diep, C.Q. *et al.* Identification of adult nephron progenitors capable of kidney regeneration  
30 in zebrafish. *Nature* **470**, 95-100 (2011).
- 31 77. Kinomura, M. *et al.* Amelioration of cisplatin-induced acute renal injury by renal progenitor-  
32 like cells derived from the adult rat kidney. *Cell Transplant.* **172**, 143-58, (2008).

- 1 78. Grange, C., Moggio, A., Tapparo, M., Porta, S., Camussi, G. & Bussolati, B. Protective  
2 effect and localization by optical imaging of human renal CD133+ progenitor cells in an acute  
3 kidney injury model. *Physiol. Rep.* **2**, e12009 (2004).
- 4 79. De Chiara, L. *et al.* Renal cells from spermatogonial germline stem cells protect against  
5 kidney injury. *J. Am. Soc. Nephrol.* **25**, 316-28 (2014).
- 6 80. Bruno, S. *et al.* Mesenchymal stem cell-derived microvesicles protect against acute tubular  
7 injury. *J. Am. Soc. Nephrol.* **20**, 1053-67 (2009).
- 8 81. Giron-Michel, J. *et al.* Interleukin-15 is a major regulator of the cell-microenvironment  
9 interactions in human renal homeostasis. *Cytokine Growth Factor Rev.* **24**, 13-22 (2013).
- 10 82. Sallustio F. *et al.* Human renal stem/progenitor cells repair tubular epithelial cell injury  
11 through TLR2-driven inhibin-A and microvesicle-shuttled decorin. *Kidney Int.* **83**, 392-403  
12 (2013).
- 13 83. Chen, J. *et al.* Kidney-derived mesenchymal stem cells contribute to vasculogenesis,  
14 angiogenesis and endothelial repair. *Kidney Int.* **74**, 879-89 (2008).
- 15 84. Choi, H.Y. *et al.* Microparticles from kidney-derived mesenchymal stem cells act as carriers  
16 of proangiogenic signals and contribute to recovery from acute kidney injury. *PLoS One.* **9**,  
17 e87853 (2014).
- 18 85. Yamaleyeva, L.M. *et al.* Cell therapy with human renal cell cultures containing  
19 erythropoietin-positive cells improves chronic kidney injury. *Stem Cells Transl. Med.* **1**, 373-83  
20 (2012).
- 21 86. Kelley, R. *et al.* Tubular cell-enriched subpopulation of primary renal cells improves survival  
22 and augments kidney function in rodent model of chronic kidney disease. *Am. J. Physiol. Renal*  
23 *Physiol.* **299**, F1026-39 (2010).
- 24 87. Kelley, R. *et al.* A population of selected renal cells augments renal function and extends  
25 survival in the ZSF1 model of progressive diabetic nephropathy. *Cell Transplant.* **22**, 1023-39  
26 (2013).
- 27 88. Genheimer, C.W. *et al.* Molecular characterization of the regenerative response induced by  
28 intrarenal transplantation of selected renal cells in a rodent model of chronic kidney disease.  
29 *Cells Tissues Organs* **196**, 374-84 (2012).
- 30 89. Presnell, S.C. *et al.* Isolation, characterization, and expansion methods for defined primary  
31 renal cell populations from rodent, canine, and human normal and diseased kidneys. *Tissue*  
32 *Eng. Part C Methods* **17**, 261-73 (2011).

1

2 Prescott, L.F. The normal urinary excretion rates of renal tubular cells, leucocytes and red  
3 blood cells. *Clin. Sci.* **31** 425-435 (1966).

4 Imberti, B., *et al.* Renal progenitors derived from human iPSCs engraft and restore function in a  
5 mouse model of acute kidney injury. *Sci. Rep.* **5**, 8826 (2015).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

### Key points: RPC characterization

- Embryonic progenitors represent the prototype of the renal cell population, with nephrogenic potential towards all the different types of nephron epithelia.
- In the adult kidney, the nephrogenic mesenchymal progenitor population is absent.
- Adult RPCs appear as segment-restricted progenitors. Whether they are a fixed population or rather a transient functional state modulated by microenvironmental stimuli is still unclear.
- Human CD133+ cells with phenotypic and functional progenitor-like properties are present in all nephron segments as scattered distinct cells with a dedifferentiated phenotype, and survive and proliferate after damage.
- Adult RPC cultures have been obtained by different nephron compartments or by urine and show homogeneous characteristics *in vitro* although distinct integration potential in pre-clinical models, possibly due to segment commitment.
- All the cell lines of human RPCs possess a limited life span and undergo senescence over time.

### Key points: Acquired knowledge and limitations of RPC therapy

- **Pro:** Concerted efforts have been employed in developing the technology to generate metanephric mesenchyme progenitors by differentiation of pluripotent stem cells or by direct reprogramming of renal cells.  
**Con:** Maldifferentiation, oncogenic potential and expansion limits may impair their application.
  - **Pro:** Adult human RPCs show therapeutic potential in acute tubular and glomerular damage due to both paracrine mechanisms and differentiation.  
**Con:** No differences are observed in pre-clinical models with respect to MSCs, although these cells lack integration potential. MSCs may benefit from more readily-available sources and reduced immune compatibility issues.
  - **Pro:** Adult and foetal RPCs show long-term engraftment in glomeruli and tubules, in pre-clinical models of CKD and possibly a low risk of maldifferentiation. The generation of
- 41

1 patient-derived RPCs could have significant relevance in chronic settings as a possible  
2 autologous therapy that does not require immunosuppression.

3 **Con:** Knowledge of the differentiation, integration potential and persistence of the  
4 different cell types is still required.

- 5
- 6 • **Pro:** Adult sources of renal cells were shown to be beneficial in CKD and are currently  
7 undergoing a phase 1 clinical trial using an intra-parenchymal injection of autologous  
8 cells.

9 **Con:** Cells derived from CKD patients may lose regenerative ability, and additional  
10 reprogramming technology could be required. Mechanisms involved should be  
11 investigated more fully.

- 12
- 13 • The administration modalities for cell therapy with RPCs needs to be defined. As tubular  
14 localization requires the passage through glomerular or tubular basal membrane, a local  
15 administration appears more feasible.

16 **Pro.** Reduced cell number required, reduce extra-renal localization

17 **Con.** Possible local damage (necrosis or thrombosis), local rather than general diffusion  
18 of cells.

19

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

**Table I.** Identification of RPCs in renal rodent or human tissue using different markers or functional tools. Localization within nephron districts and evidences of *in vivo* function are indicated.

	Identification tool	Tissue location	Tissue function	<i>In vitro</i> culture	Refs
<b>Murine/rat tissue</b>	Scattered cells in PEC-rtTA mice	BC, PT, DT	Proliferation after injury	n.d.	27,28
	Sca-1	interstitium	-	multipotency	53
	BrdU retention (short chase)	PT, TL-HL, DT, CD	Proliferation after injury	epithelial diff.	29
	BrdU retention (long chase)	papilla	Proliferation/migration after injury	multipotency,	30
	Side population	PT	-	multipotency	52
	Lgr5	TL-HL and DT	-	n.d.	15
	nFatC1+	PT	apoptosis-resistance proliferation and repair	n.d.	27 31
	c-kit	TL-HL and DT		multipotency	32
<b>human tissue</b>	CD133+/CD24+	BC, PT, HL, DT	apoptosis-resistance proliferation	multipotency	3, 36-41

Abbreviations: BC: Bowmann’s capule, PT: proximal tubules, TL-HL: thick segment of the Henle’s loop, HL: Henle’s loop, DT: distal tubules, CD: connecting ducts.

1  
2  
3  
4  
5

**Table II.** Localization of CD133+ RPCs in different nephron districts, and marker co-expression.

Nephron district	Tissue location	Markers	Co-expression of scattered cell markers	Co-expression of segment markers	Re s
<b>Bowmann's capsule</b>	Tubular pole	CD133+/CD24+/ PDX -	Vimentin Collagen 7A Claudin-1	no	21 39 40
<b>Proximal Tubules</b>	Tubular plicae	CD133+/CD24+/C D106+	Vimentin CK7, CK19 Collagen 7A Claudin-1, 3 and 7 annexin 2, S100A6 CXCR4	Megalin Aquaporin-1	3 36 37 39 41
<b>Henle's loop</b>	Segments and loop	CD133+ Nestin+	-	no	36 38
<b>Distal tubules</b>	In contact with the glomerular vascular pole	CD133+/ CD24+	-	Thiazide-sensitive Na/Cl cotransporter	41

6 Abbreviations: PDX: podocalixin, CXCR4: C-X-C chemokine receptor type 4 , CK: cytokeratin.  
7



1  
2  
3  
4

**Table III.** Cell therapy using rodent renal-derived cells in pre-clinical models of AKI and CKD.

	Cell type	Model	administration	Benefit	Mechanism	ref
<b>adult RPCs</b>	S3-derived tubular RPCs	IRI cisplatin	subcapsule	Improvement of renal function	Tubular integration/ Paracrine	49, 75
	Multipotent RPCs	IRI	intra-arterially	Improvement of renal function	Tubular integration No paracrine	51
	Tubule-derived side population cells	adriamycin	I.V. and intra-parenchymal	Reduced albuminuria	Paracrine	52
	kidney-derived MSCs	IRI	I.V.	Improvement of renal function and of angiogenesis	Paracrine EV release	83, 84
	Sca-1 interstitial cells	IRI	Intra-parenchymal	Improvement of renal function	Tubular integration	53
<b>Adult renal cells</b>	Human EPO+ cells	IRI and gentamycin	Intra-parenchymal	Improvement of renal function- Reduction of fibrosis	Paracrine	85
	Rat renal cells	5/6 neph.	intra-parenchymal	Reduction of disease progression	Tubule Integration/ Paracrine	86
	Rat renal cells	Diabetes	intra-parenchymal	Reduction of disease progression	Tubule Integration/ Paracrine	87
<b>ESC-differentiated cells</b>	Tubular differentiated-spermatogonia I stem cells	IRI	Intra-parenchymal	Improvement of function- reduced development of fibrosis	Paracrine	79

5 Abbreviations: MSCs: mesenchymal stromal cells, IRI: ischemia-reperfusion injury, 5/6 neph.:  
6 5/6 nephrectomy, EPO: erythropoietin, I.V.: intravenously.  
7  
8  
9  
10

1 **Table IV.** Cell therapy using human renal-derived cells in pre-clinical models of AKI and CKD.

	Cell type	Model	adminis- tration	Benefit	Mechanism	ref
<b>Foetal RPCs</b>	Foetal CD133+/CD24+ RPCs	AKI (glycerol)	i.v.	Improvement of renal function	Tubular integration	21
	Fetal NCAM+ RPCs	CKD (5/6 neph.)	Intra-parenchymal	Prevention of disease progression	Tubular integration	19
<b>Adult RPCs</b>	Bowmann's capsule CD133+/CD24+/PDX- RPCs	FSGS (adryam.)	i.v.	Improvement of function-reduced fibrosis	Podocytic/Tubular integration	40
		AKI (glycerol)	i.v.	Improvement of function	Tubular integration	35
	Cortical CD133+/CD24+/CD106+ RPCs	FSGS (adryam.)	i.v.	Improvement of function	Tubular differentiation only	41
		AKI (glycerol)	i.v.			
	Cortical CD133+ RPCs	AKI (glycerol)	i.v.	Improvement of function	Tubular differentiation	54
	Medullary CD133+ RPCs	AKI (glycerol)	i.v.	Improvement of function	Tubular integration/paracrine	36,78
	Urine CD133+/CD24+ RPCs	FSGS (adryam.)	i.v.	Improvement of function	Podocytic/Tubular integration	58
NCAM+ RPCs	AKI (glycerol)	i.v.	Improvement of function	-	19	
<b>Adult renal cells</b>	Renal CD24-/CD133- cells	FSGS (adryam.)	i.v.	No improvement of renal function	-	35
		AKI (glycerol)	i.v.			41
	Bowmann's capsule CD133+/CD24+/PDX+ cells	FSGS (adryam.)	i.v.			40
	NCAM- cells from total epithelial cell culture	AKI (glycerol)	i.v.	Improvement of function	-	19
<b>iPS-derived RPCs</b>	iPS-derived RPCs	AKI (Cisplatin)	i.v.	Improvement of function	Tubular differentiation	ref

2 Abbreviations: PDX: podocalixin, FSGS: focal segmental glomerulosclerosis, induced by  
3 administration of adriamycin (adryam.). 5/6 nephrectomy: 5/6 neph. IRI: ischemia-reperfusion  
4 injury.  
5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**Legend for Figure**

Figure 1. Potential strategies of cell therapy for kidney regeneration. Abbreviations: GPSC, germline cell-derived pluripotent stem cells; iPS, induced pluripotent stem cells; ESC: embryonic stem cells; EPO: erythropoietin.

**Biographies:**

For each author please provide the correct current main job title and affiliation for the print version of your article, as well as a brief (just 50-100 words) biography detailing career, training and research interests for the online version.

**Review criteria:**

We searched for original articles in the PubMed database published between 1990 and 2015 containing the search terms: “renal regeneration”, “renal repair”, “kidney stem cells”, “kidney progenitors”, “iPS”, “kidney injury”, “cell therapy”, “preclinical models”, “CD133”, alone and in combination. All papers identified were English-language, full-text articles. Papers were prioritized on the basis of relevance and we apologize to authors of manuscripts that might have been missed.