AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Inflammation response and cytotoxic effects in human THP-1 cells of size-fractionated PM10 extracts in a polluted urban site.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1533689> since 2015-12-16T13:39:45Z

Published version:

DOI:10.1016/j.chemosphere.2015.11.074

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

**[Chemosphere, Vol. 145, February 2016, DOI:
10.1016/j.chemosphere.2015.11.074]**

<http://www.sciencedirect.com/science/article/pii/S0045653515303805>

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform]

Manuscript Number: CHEM37545R1

Title: Inflammation response and cytotoxic effects in human THP-1 cells of size-fractionated PM10 extracts in a polluted urban site.

Article Type: Research paper

Section/Category: Environmental Toxicology and Risk Assessment

Keywords: size-fractionated PM10; THP-1 cells; LDH; IL-8; TNFalpha; CYP1A1

Corresponding Author: Prof. Tiziana Schilirò,

Corresponding Author's Institution: University of Torino

First Author: Tiziana Schilirò

Order of Authors: Tiziana Schilirò; Luca Alessandria , PhD; Sara Bonetta, PhD; Elisabatta Carraro, Prof.; Giorgio Gilli, Prof.

Response to Reviewers: RESPONSE TO REVIEWER COMMENTS

Reviewer #1:

1. The title, abstract, introduction, and parts of the discussion are suitable for the study of size related effects of particles. In this study, in vitro assays were performed with extracts of PM, meaning without particles present. The title and text need to be rewritten to clearly state the hypothesis driving the study of extracts instead of PM, discuss the usefulness of the extracts and different extraction methods, and then in the end discuss how the differences in toxic activity of the extracts may in part explain size-related effects observed in PM-exposure studies.

- The text was modified as required to better explain the hypothesis driving the study of extracts instead of PM. Moreover the discussion was implemented to highlight the importance to use different extraction methods to evaluate the biological effects of different fractions (in particular the finest fractions).

2. Explanation of the in vitro tests statistical analysis in sufficient detail: type of controls used (is the description in LL 158-160 part of the controls?), solvent controls (especially important for potential interference with the substrates/read-out of the assays, RNA extraction), number of technical and biological replicates, outcome of the ANOVA (F, p, d values in all cases), were parts of the values considered as dependent (e.g. extracts of the same PM sample)? Explain which statistics were done on which datasets exactly. Why is Wilcoxon mentioned in the figure captions but no ANOVA results? Did you correct for multiple comparisons? Have you tried RDA on your data?

- A description of the type of controls used for each biological test (in particular QC filter; control cells and solvent control), the number of replicates made and the different outcome of the ANOVA were added. Moreover, the information about the statistics used and the different dataset used for ANOVA test were also added. In the figure only the Wilcoxon test was mentioned considering the results of biological test respect to control cells. On the contrary, we preferred to report the results of the ANOVA in the text. We don't have made correction for multiple comparison and we don't have tried the RDA on our data.

3. Explain motivation for the exposure scenario and concentration chosen

- 200 ug/mL is the concentration that always induced a significant effect with samples from Torino described in previous works and 48h is a sufficient exposure time to determine a significant extract effect (Alessandria et al., 2014; Schilirò et al., 2015).

4. The M&M section is not consistent regarding the use of „particle concentration“, „particle extracts“, and the mass concentration „200 ug/mL“. E.g. L 172 reads „particle concentration of 200 ug/mL“ - I understood that the exposure was done with extracts only?

- The text was modified to clarify this point. The cells were exposed to PM extracts corresponding to an equivalent particle concentration of 200 µg/mL.

5. Improvement of the figure captions: state the number of replicates, replace size-fractionated PM with the type of extract used, include enough details so that the figure is self-explanatory.

- The figure captions were modified and detailed as required.

6. Briefly discuss the limitations of using a model cell line (LL 166-167 „more sensitive“ than what?) regarding the conclusions for PM-regulation and the use of standard cell culture medium during exposure.

- Specific suggestion were added as required
The distribution patterns of cytotoxicity and/or inflammation among different cell lines (e.g. A549, THP1) after exposure to PM extracts are in some cases different confirming that the cell lines respond differently to PM components. In some works THP-1 cells tends to show more IL-8 production than other cell lines (e.g. A549 cells) (Corsini et al., 2013).

In toxicity testing it is accepted that cell lines offer the advantage of being more homogeneous and standardized than primary cell cultures. They are well characterized, easy to cultivate and reproducible results are easier to obtain. On the other hand, they may be quite different from the original tissue due to the fact that established cell lines have undergone a number of transformations implying some loss of differentiation. Moreover the use of standard cell culture media during exposure facilitate adaptation of cells to the culture and allow a better standardization of experimental conditions (Astashkina et al., Pharmacology & Therapeutics 134 (2012) 82-106; Ekwall et al., Chapter 7,

Short-term Toxicity Tests for Non-genotoxic Effects, 1990 SCOPE by John Wiley & Sons Ltd)

7. LL 409-412: A specific suggestion based on your observations would be more valuable than a general statement.

Specific suggestion were added as required

Reviewer #2:

General comments:

- In the "Material and Methods" paragraph (lines 119-120) the Authors say that the samplings were made during summer and winter periods but they don't specify the year of the study. The year of the study should be added.

The year of the study was added as required by the Reviewer.

- In the "Material and Methods" paragraph (line 151) "monthly strips" are reported. Why? Were the samples combined and considered monthly?

The strips were combined and considered seasonally. The text was corrected and this point was clarified.

- In the "Discussion and conclusions" paragraph (lines 386-387) the sentence "One possible explanation ...PM10 fraction" is not clear. The Authors should explain it better.

The sentence was clarified as required by the Reviewer.

The manuscript should be better checked. There are some typing errors. Some words should be uniformed both in text and in figure captions: IL-8 or IL8; THP1 or THP-1; PM10 or PM10; ml or ML. The words (IL-8, THP-1, PM10, mL) was uniformed in text and figure captions.

Some paragraph titles should be uniformed.

Paragraph title was uniformed.

References quoted in the text should be checked as to both style and punctuation.

References in the text was checked for style and punctuation.

The reference list should be checked.

The reference list was checked.

Specific comments:

Line 66: write full PAHs (as it is the first time they are mentioned) and insert a comma

The text was modified.

Line 71: "polycyclic aromatic hydrocarbons (PAH)" should be "PAHs" because already mentioned before (line 66)

The text was modified.

Line 72: the year of the reference is 2012 in the text but 2011 in the reference list

The year of the reference was modified.
 Line 122: "electronically controlled flow" instead of "flow electronically controlled"
 The text was modified.
 Line 124: "filters" instead of "filter"
 The text was modified.
 Line 138: "weighted" instead of "weighed"
 The text was modified.
 Line 156: "foetal calf serum (FCA)" should be "FCA" because already mentioned before (line 147)
 The text was modified.
 Line 224: Ni instead of Nichel (as other metals)
 The text was modified.
 Line 225: indeno(1,2,3-cd)pyrene instead of indeno(1,2,3-cd)pirene and delete the space before the parenthesis
 The text was modified.
 Line 258: delete the comma after and
 The text was modified.
 Line 309: northwest instead of northewest
 The text was modified.
 Line 312: "Torino PM concentration" instead of "The Torino PM concentration"
 The text was modified.
 Line 336: appeared instead of appear
 The text was modified.
 Line 351: researches instead of research
 The text was modified.
 Line 360: "especially the summer greater fractions" instead of "especially the by the summer greater fractions"
 The text was modified.
 Line 361: the reference Hetland et al., 2005 should be checked because 2005 is in the text and 2004 in the reference list
 The reference "Hetland et al., 2005" was checked and modified.
 Line 364: see line 361
 The reference "Hetland et al., 2005" was checked and modified.
 Line 372: see line 361
 The reference "Hetland et al., 2005" was checked and modified.
 Line 382: "compared to" instead of "in confront of"
 The text was modified.

There are some errors in the reference list and the reference list should be checked as to style and punctuation. In particular:
 Lines 427-429: the reference should be corrected (year and name of journal)
 The reference was corrected (year and name of journal).
 Line 440: the name of an author should be corrected (Gnauk instead of GnaUK)
 The name of the author was corrected.
 Lines 443-444: grp collaborativo EpiAir should be added
 grp collaborativo EpiAir was added.
 Line 486: see line 361
 The reference "Hetland et al., 2005" was checked.
 Line 509: IARC working group should be added
 IARC working group was added.
 Lines 546-548: this reference is not present in the text (Schins et al., 2004) and it should be deleted

The reference is present in the text (line 357 in the word text and line 363 in the pdf).

Table 1: indeno(1,2,3-cd)pyrene instead of indeno(1,2,3-cd)pirene

The text was modified.

Reviewer #3:

Key words:THP-1 cells;

The key word was modified.

Introduction

Line 73: Please, remove "exists" from the sentence.

Line 101: (Group 1)

The text was modified.

Material and Method

From environmental sampling you can obtain six particle size fractions, as explained at lines 126-134, but biological endpoints were only investigated on pooled three main size fractions. Why didn't you investigate the biological targets on each of the six particle size fractions? Clarify this point.

- The six particle fractions have been pooled in order to obtain a sufficient mass to PM to be analysed by biological tests.

You declare THP1 cell line as more sensitive in cytotoxicity test (line 166-167). You should better describe the reason of this choice.

- The distribution patterns of cytotoxicity and/or inflammation among different cell lines (e.g. A549, THP1) after exposure to PM extracts are in some cases different confirming that the cell lines respond differently to PM components. In some works THP-1 cells tends to show more IL-8 production than other cell lines (e.g. A549 cells) (Corsini et al., 2013).

Results / Discussion and Conclusion

There are many results, and many types of comparisons (organic vs aqueous extraction; summer vs winter; comparisons with negative controls...) resulting -sometimes- quite difficult to follow adequately the argumentation. For example, I think there is mismatch between the information from the Results concerning the inhibition cell proliferation in summer/winter (lines 272 - 274) and the Discussion at lines 350 - 352. Please clarify Results and/or Discussion and Conclusion consistently.

- Results and/or Discussion and Conclusion sections were checked and modified as required to enhance the text understanding

Line 288: Please, change "alpha" with " α ", since it was already made explicit in the Introduction as well in Mat & Met. Be consistent.

The text was modified.

Line 290: Please add a reference.

The reference was added

Line 385: PAH

The text was modified.

Ms. Ref. No.: CHEM37545

Dear Editor,

please find enclosed the revised manuscript "*Inflammation response and cytotoxic effects in human THP-1 cells of size-fractionated PM₁₀ extracts in a polluted urban site.*" by Tiziana Schilirò, Luca Alessandria, Sara Bonetta, Elisabetta Carraro and Giorgio Gilli.

We have answered to the Reviewers' comments, in particular relevant changes have been written in red all over the enclosed text.

Finally, our responses to each Reviewers comments have been reported as follows, highlighted in yellow.

Best regards,

Tiziana Schilirò and Co-authors.

RESPONSE TO REVIEWER COMMENTS

Reviewer #1:

1. The title, abstract, introduction, and parts of the discussion are suitable for the study of size related effects of particles. In this study, in vitro assays were performed with extracts of PM, meaning without particles present. The title and text need to be rewritten to clearly state the hypothesis driving the study of extracts instead of PM, discuss the usefulness of the extracts and different extraction methods, and then in the end discuss how the differences in toxic activity of the extracts may in part explain size-related effects observed in PM-exposure studies.

- *The text was modified as required to better explain the hypothesis driving the study of extracts instead of PM. Moreover the discussion was implemented to highlight the importance to use different extraction methods to evaluate the biological effects of different fractions (in particular the finest fractions).*

2. Explanation of the in vitro tests statistical analysis in sufficient detail: type of controls used (is the description in LL 158-160 part of the controls?), solvent controls (especially important for potential interference with the substrates/read-out of the assays, RNA extraction), number of technical and biological replicates, outcome of the ANOVA (F, p, d values in all cases), were parts of the values considered as dependent (e.g. extracts of the same PM sample)? Explain which statistics were done on which datasets exactly. Why is Wilcoxon mentioned in the figure captions but no ANOVA results? Did you correct for multiple comparisons? Have you tried RDA on your data?

- *A description of the type of controls used for each biological test (in particular QC filter; control cells and solvent control), the number of replicates made and the different outcome of the ANOVA were added. Moreover, the information about the statistics used and the different dataset used for ANOVA test were also added. In the figure only the Wilcoxon test was mentioned considering the results of biological test respect to control cells. On the contrary, we preferred to report the results of the ANOVA in the text. We don't have made correction for multiple comparison and we don't have tried the RDA on our data.*

3. Explain motivation for the exposure scenario and concentration chosen

- *200 ug/mL is the concentration that always induced a significant effect with samples from Torino described in previous works and 48h is a sufficient exposure time to determine a significant extract effect (Alessandria et al., 2014; Schilirò et al., 2015).*

4. The M&M section is not consistent regarding the use of „particle concentration“, „particle extracts“, and the mass concentration „200 ug/mL“. E.g. L 172 reads „particle concentration of 200 ug/mL“ - I understood that the exposure was done with extracts only?

- *The text was modified to clarify this point. The cells were exposed to PM extracts corresponding to an equivalent particle concentration of 200 µg/mL.*

5. Improvement of the figure captions: state the number of replicates, replace size-fractionated PM with the type of extract used, include enough details so that the figure is self-explanatory.

- *The figure captions were modified and detailed as required.*

6. Briefly discuss the limitations of using a model cell line (LL 166-167 „more sensitive“ than what?) regarding the conclusions for PM-regulation and the use of standard cell culture medium during exposure.

- *Specific suggestion were added as required
The distribution patterns of cytotoxicity and/or inflammation among different cell lines (e.g. A549, THP1) after exposure to PM extracts are in some cases different confirming that the cell lines respond differently to PM components. In some works THP-1 cells tends to show more IL-8 production than other cell lines (e.g. A549 cells) (Corsini et al., 2013).
In toxicity testing it is accepted that cell lines offer the advantage of being more homogeneous and standardized than primary cell cultures. They are well characterized, easy to cultivate and reproducible results are easier to obtain. On the other hand, they may be quite different from the original tissue due to the fact that established cell lines have undergone a number of transformations implying some loss of differentiation. Moreover the use of standard cell culture media during exposure facilitate adaptation of cells to the culture and allow a better standardization of experimental conditions (Astashkina et al., Pharmacology & Therapeutics 134 (2012) 82–106; Ekwall et al., Chapter 7, Short-term Toxicity Tests for Non-genotoxic Effects, 1990 SCOPE by John Wiley & Sons Ltd)*

7. LL 409-412: A specific suggestion based on your observations would be more valuable than a general statement.

Specific suggestion were added as required

Reviewer #2:

General comments:

- In the "Material and Methods" paragraph (lines 119-120) the Authors say that the samplings were made during summer and winter periods but they don't specify the year of the study. The year of the study should be added.

The year of the study was added as required by the Reviewer.

- In the "Material and Methods" paragraph (line 151) "monthly strips" are reported. Why? Were the samples combined and considered monthly?

The strips were combined and considered seasonally. The text was corrected and this point was clarified.

- In the "Discussion and conclusions" paragraph (lines 386-387) the sentence "One possible explanation ...PM10 fraction" is not clear. The Authors should explain it better.

The sentence was clarified as required by the Reviewer.

The manuscript should be better checked. There are some typing errors. Some words should be uniformed both in text and in figure captions: IL-8 or IL8; THP1 or THP-1; PM10 or PM10; ml or ML.

The words (IL-8, THP-1, PM₁₀, mL) was uniformed in text and figure captions.

Some paragraph titles should be uniformed.

Paragraph title was uniformed.

References quoted in the text should be checked as to both style and punctuation.

References in the text was checked for style and punctuation.

The reference list should be checked.

The reference list was checked.

Specific comments:

Line 66: write full PAHs (as it is the first time they are mentioned) and insert a comma

The text was modified.

Line 71: "polycyclic aromatic hydrocarbons (PAH)" should be "PAHs" because already mentioned before (line 66)

The text was modified.

Line 72: the year of the reference is 2012 in the text but 2011 in the reference list

The year of the reference was modified.

Line 122: "electronically controlled flow" instead of "flow electronically controlled"

The text was modified.

Line 124: "filters" instead of "filter"

The text was modified.

Line 138: "weighted" instead of "weighed"

The text was modified.

Line 156: "foetal calf serum (FCA)" should be "FCA" because already mentioned before (line 147)

The text was modified.

Line 224: Ni instead of Nichel (as other metals)

The text was modified.

Line 225: indeno(1,2,3-cd)pyrene instead of indeno(1,2,3-cd)pirene and delete the space before the parenthesis

The text was modified.

Line 258: delete the comma after and

The text was modified.

Line 309: northwest instead of northewest

The text was modified.

Line 312: "Torino PM concentration" instead of "The Torino PM concentration"

The text was modified.

Line 336: appeared instead of appear

The text was modified.

Line 351: researches instead of research

The text was modified.

Line 360: "especially the summer greater fractions" instead of "especially the by the summer greater fractions"

The text was modified.

Line 361: the reference Hetland et al., 2005 should be checked because 2005 is in the text and 2004 in the reference list

The reference "Hetland et al., 2005" was checked and modified.

Line 364: see line 361

The reference "Hetland et al., 2005" was checked and modified.

Line 372: see line 361

The reference "Hetland et al., 2005" was checked and modified.

Line 382: "compared to" instead of "in confront of"

The text was modified.

There are some errors in the reference list and the reference list should be checked as to style and punctuation. In particular:

Lines 427-429: the reference should be corrected (year and name of journal)

The reference was corrected (year and name of journal).

Line 440: the name of an author should be corrected (Gnauk instead of GnaUK)

The name of the author was corrected.

Lines 443-444: grp collaborativo EpiAir should be added

grp collaborativo EpiAir was added.

Line 486: see line 361

The reference "Hetland et al., 2005" was checked.

Line 509: IARC working group should be added

IARC working group was added.

Lines 546-548: this reference is not present in the text (Schins et al., 2004) and it should be deleted

The reference is present in the text (line 357 in the word text and line 363 in the pdf).

Table 1: indeno(1,2,3-cd)pyrene instead of indeno(1,2,3-cd)pirene

The text was modified.

Reviewer #3:

Key words:THP-1 cells;

The key word was modified.

Introduction

Line 73: Please, remove "exists" from the sentence.

Line 101: (Group 1)

The text was modified.

Material and Method

From environmental sampling you can obtain six particle size fractions, as explained at lines 126-134, but biological endpoints were only investigated on pooled three main size fractions. Why didn't you investigate the biological targets on each of the six particle size fractions? Clarify this point.

- *The six particle fractions have been pooled in order to obtain a sufficient mass to PM to be analysed by biological tests.*

You declare THP1 cell line as more sensitive in cytotoxicity test (line 166-167). You should better describe the reason of this choice.

- *The distribution patterns of cytotoxicity and/or inflammation among different cell lines (e.g. A549, THP1) after exposure to PM extracts are in some cases different confirming that the cell lines respond differently to PM components. In some works THP-1 cells tends to show more IL-8 production than other cell lines (e.g. A549 cells) (Corsini et al., 2013).*

Results / Discussion and Conclusion

There are many results, and many types of comparisons (organic vs aqueous extraction; summer vs winter; comparisons with negative controls....) resulting -sometimes- quite difficult to follow adequately the argumentation. For example, I think there is mismatch between the information from the Results concerning the inhibition cell proliferation in summer/winter (lines 272 - 274) and the Discussion at lines 350 - 352.

Please clarify Results and/or Discussion and Conclusion consistently.

- *Results and/or Discussion and Conclusion sections were checked and modified as required to enhance the text understanding*

Line 288: Please, change "alpha" with "<alpha>", since it was already made explicit in the Introduction as well in Mat & Met. Be consistent.

The text was modified.

Line 290: Please add a reference.

The reference was added

Line 385: PAH

The text was modified.

1 **Inflammation response and cytotoxic effects in human THP-1 cells of size-fractionated PM₁₀**
2 **extracts in a polluted urban site.**

4 T. Schilirò ^{*}^a, L. Alessandria ^a[^], S. Bonetta ^a[^], E. Carraro ^a, G. Gilli ^a

6 ^a *Department of Public Health and Pediatrics, University of Torino, Piazza Polonia, 94 - 10126,*
7 *Torino, Italy; tiziana.schiliro@unito.it; luca.al@hotmail.it; sara.bonetta@unito.it;*
8 *elisabetta.carraro@unito.it; giorgio.gilli@unito.it*

9 [^]These authors contributed equally to the work.

11 ***Corresponding author:**

12 **Tiziana Schilirò**

13 *Department of Public Health and Pediatrics,*
14 *University of Torino,*
15 *Piazza Polonia, 94 - 10126, Torino, Italy,*
16 Tel: +390116705820
17 e-mail address: tiziana.schiliro@unito.it

29 **ABSTRACT**

30 To contribute to a greater characterization of the airborne particulate matter's toxicity, size-
31 fractionated PM₁₀ was sampled during different seasons in a polluted urban site in Torino, a
32 northern Italian city. Three main size fractions (PM₁₀ - 3 µm; PM₃ - 0.95 µm; PM <0.95 µm)
33 extracts (organic and aqueous) were assayed with THP-1 cells to evaluate their effects on cell
34 proliferation, LDH activity, TNFα, IL-8 and CYP1A1 expression.

35 The mean PM₁₀ concentrations were statistically different in summer and in winter and the finest
36 fraction PM_{<0.95} was always higher than the others. Size-fractionated PM₁₀ extracts, sampled in an
37 urban traffic meteorological–chemical station produced size-related toxicological effects in relation
38 to season and particles extraction. The PM summer extracts induced a significant release of LDH
39 compared to winter and produced a size-related effect, with higher values measured with PM₁₀₋₃.
40 Exposure to size-fractionated PM₁₀ extracts did not induce significant expression of TNFα. IL-8
41 expression was influenced by exposure to size-fractionated PM₁₀ extracts and statistically
42 significant differences were found between kind of extracts for both seasons. The mean fold
43 increases in CYP1A1 expression were statistically different in summer and in winter; winter fraction
44 extracts produced a size-related effect, in particular for organic samples with higher values
45 measured with PM_{<0.95} extracts.

46 Our results confirm that the only measure of PM can be misleading for the assessment of air
47 quality moreover we support efforts toward identifying potential effect-based tools (e.g. *in vitro* test)
48 that could be used in the context of the different monitoring programs.

49

50 **Keywords:** size-fractionated PM₁₀; THP-1 cells; LDH; IL-8; TNFα; CYP1A1

51

52

53

54

55

56

57 1. INTRODUCTION

58 In the last ten years, a considerable number of new studies have been published, providing
59 evidence on the health effects of size fractions, components and sources of particulate matter
60 (PM). Health effects are observed with short-term and long-term exposures to airborne particles
61 indicating an increase in cardiac and respiratory morbidity and associated mortality (WHO-Europe,
62 2013).

63 An increasing body of epidemiological and toxicological evidences have shown PM mass (PM_{2.5}
64 and PM₁₀) comprises fractions with varying types and degrees of health effects, suggesting a role
65 for both the chemical composition (such as transition metals, polycyclic aromatic hydrocarbons -
66 PAHs), physical properties (size, particle number and surface area) and source (Brook et al., 2010;
67 Kroll et al., 2013; Krzyzanowski, 2008; Steenhof et al., 2011).

68 Airborne particles are generally heterogeneous mixtures consisting of inorganic components (e.g.,
69 transition metals), salts, carbonaceous material, volatile organic compounds (VOC), PAHs and
70 biological materials such as endotoxins, fungal spores and pollen (Anderson et al., 2012). Although
71 regulatory standards for PM₁₀ and PM_{2.5} as well as specific particle constituents exist, there is still
72 no clear **consensus among** researchers regarding the mechanisms of toxicity following exposure to
73 current levels of air pollution PM (Schwarze et al., 2007).

74 Many of the observed health endpoints may result, at least in part, from oxidative stress initiated by
75 the formation of reactive oxygen species (ROS) upon the interaction of PM with epithelial cells and
76 macrophages (Knaapen et al., 2004). Induction of cellular oxidative stress and resulting activation
77 of pro-inflammatory mediators are considered to play a central role in the development of airway
78 diseases (Lonkar and Dedon, 2011; Oh et al., 2011). Pulmonary inflammation is generally
79 characterized by local recruitment of proinflammatory cells such as neutrophils and macrophages,
80 which are involved in the up regulation of various signaling molecules, such as cytokines (TNF α ,
81 IL-6) and chemokines (IL-8) (Lauer et al., 2009; Michael et al., 2013). Based on these
82 mechanisms, increased production and release of inflammatory mediators by cells are relevant
83 and often used parameters suggesting PM-induced toxicity. These mediators in connection with an
84 increasing ROS formation play a key role in the development of several inflammatory diseases like

acute lung injury, COPD, chronic bronchitis and asthma (Valavanidis et al., 2008). Besides the ability to introduce inflammation, particles may also cause oxidative DNA damage because of their physicochemical properties. Within this so-called primary genotoxicity, surface associated free radicals and transition metals are considered to play a major role (Roig et al., 2013). Moreover particle-elicited inflammation and subsequent generation of ROS can lead to oxidative DNA damage, and this pathway is defined as secondary genotoxicity (Schins and Hei, 2006). In addition, organic compounds, such as polycyclic aromatic hydrocarbons (PAH) coated onto PM may induce DNA damage, mutations and cytotoxicity at a cellular level (Bonetta et al., 2009). Upon entering the organism, PAHs are first metabolized to transdihydrodiols by the activity of the cytochromes P450 (CYP) superfamily member CYP1A1 enzyme and epoxide hydrolase and then oxidized to reactive electrophiles by two pathways (Hanzalova et al., 2010; Spink et al., 2008). Intermediates thereafter interact with DNA target sites to produce adducts, mutation, DNA strand breaks and eventually tumour initiation (Schwarze et al., 2007).

In this regards, recently, outdoor air pollution and its major component, outdoor particulate matter, were classified as carcinogenic for humans (**Group1**) (Loomis et al., 2013). Consequently, reducing air pollution and particle matter to the lowest amount possible is becoming a marked priority. However, in order to define the most cost-effective policies, the most health-relevant PM fractions will have to be identified based on toxicological characteristics. With this purpose, to contribute to a greater characterization of the airborne particulate matter's toxicity, the biological effects of airborne size-fractionated PM₁₀ collected in a high polluted urban site during different seasons were examined. The effects of size fractionated PM₁₀ extracts (organic and aqueous) from winter and summer seasons were evaluated in human cell cultured *in vitro* (THP-1 monocytes) considering as endpoints: the inhibition of cell proliferation, lactate dehydrogenase (LDH) activity and tumour necrosis factor alpha (TNF α), interleukin 8 (IL-8) and CYP1A1 expressions.

110

111 2. MATERIALS AND METHODS

112 2.1 Size-fractionated PM₁₀ sampling and gravimetric analysis

113 The sampling site was located in Torino (northwest of the Italian Padana Plain), in a
114 meteorological–chemical station of the Environmental Protection Regional Agency (Piedmont
115 A.R.P.A.), it was representative of typical urban pollution, with high vehicle traffic impact
116 (coordinates utm, WGS84, X: 394836, Y: 4996153).

117 The samplings were made during two periods, summer (July/August 2011) and winter
118 (January/February 2012), PM₁₀ samples were collected using a Sierra-Andersen high-volume
119 cascade impactor (AirFlow PM₁₀-HVS sampler which a multi-stage cascade impactor, with pre-
120 selector complying with the EN-12341 norm by Analitica Strumenti) at an electronically controlled
121 flow at 1160 L/min. In each period, sampling durations were 24 hours and were repeated on
122 Tuesday and Friday for three weeks (6 samplings and 6 filters for each fraction, for a total of 36
123 filters in each period).

124 First, PM₁₀ was selected by a pre-selector, and then the multistage impactor determined the
125 division of different particle sizes of the sampled particles by the differentiation of the aerodynamic
126 diameter, which can identify the type of trajectory that particles take inside the suction flow related
127 to the three main aerodynamic factors of the particles themselves: dimension, shape and density
128 (Analitica Strumenti). Particles with sufficient inertia will impact on that particular stage collection
129 plate, whilst smaller particles will remain entrained in the air stream and pass to the next stage
130 where the process is repeated. The stages are assembled in a stack or row in order of decreasing
131 particle size. The particle size fractions are as follows: 10.0-7.2, 7.2-3.0, 3.0-1.5, 1.5-0.95, 0.95-
132 0.49, and <0.49 µm. Glass microfiber filters with splits (Type A/E; 8"x10"; Gelman Sciences, MI,
133 USA) were used to collect particles on each impactor plate; finally, glass microfiber filters (20.32 ×
134 25.40 cm; Pall Corporation, NY, USA) were present as back-up filters to collect the finest particles
135 (<0.49 µm). All of the filters were pre- and post-conditioned by placing them in a dry and dark
136 environment for 48 h, and then they were weighted in a room with a controlled temperature and
137 humidity. The PM concentrations in the air volume sampled was calculated as previously described
138 (Schilirò et al., 2010; Traversi et al., 2011), according to the European Committee for
139 Standardization (CEN, 1998).

140 **2.2 Particles extraction**

141 After gravimetric analyses, one strip was cut from each fractionated PM filters. The strips were
142 pooled into three main size fractions (PM_{10-3} : 10 - 3 μm ; $PM_{3-0.95}$: 3 – 0.95 μm ; and $PM_{<0.95}$: <0.95
143 μm) (two strips for each main size fraction) for a total of 12 strips for each main fraction both in
144 summer and in winter. This procedure was repeated two times to perform two different extractions
145 chosen for different abilities in the extraction of different compounds: acetone was selected as the
146 solvent for organic-extractable compounds (e.g., PAHs) (Claxton et al., 2004) and RPMI1640,
147 without foetal calf serum (FCS,) was chosen to extract water-soluble components (e.g., metals),
148 theoretically comparable to the extraction at the lung cells (Hetland et al., 2004). The 12 strips for
149 each main fraction both in summer and winter were cut into small pieces and were placed in 50 mL
150 polypropylene sterile tubes with 15 mL of each extraction media. The tubes were placed in an
151 ultrasonic water bath for 10 min, followed by 1 min of vortexing. This procedure was repeated 3
152 times (45 mL of extracts). The samples were centrifuged at 5000 x g for 10 min to remove the filter
153 material, and supernatant was collected. Acetone extract (organic) for biological tests was
154 evaporated with a rotary evaporator and suspended in dimethyl sulfoxide (DMSO) for the THP-1
155 cells tests. RPMI extracts (aqueous) were directly assayed for biological analysis (Alessandria et
156 al., 2014; Schilirò et al., 2010). Each extraction was also performed on a QC laboratory filter
157 (treated with the same method as the samples). Unless specified otherwise, all chemicals were
158 purchased from Sigma, St. Louis, MO, USA.

159 **2.3 Cell culture**

160 The human THP-1 monocytes from Interlab Cell Line Collection (Genova, IT) were used as
161 surrogates of lung monocytes and were grown, maintained and treated in RPMI1640
162 supplemented with 10 % (v/v) FCS, 2 % L-glutamine 200 mM and 1 % penicillin/streptomycin 10
163 mg/mL, at 37 °C in an humidified atmosphere containing 5% CO₂. Serum-free RPMI1640 without
164 phenol red constituted the experimental medium. The use of serum-free allow a better
165 standardization of experimental conditions. THP-1 cells are more sensitive in cytotoxicity tests
166 respect to other cell lines (e.g. A549 cells) (Corsini et al., 2013).

167 **2.4 Cell viability- MTT assay**

MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay was used to measure cell viability as previously described (Schilirò et al., 2015); briefly THP-1 cells were seeded, using experimental medium, in 24-well plates at a density of 4×10^4 cell/well and exposed to an equivalent particle concentration of 200 $\mu\text{g/mL}$. After treatment with PM extracts (72 h), the MTT (final concentration 0.5 mg/mL) was added to cells and incubated at 37°C for 4 h. Then, 1 mL 10 % SDS/0.01 M HCl was added in each well, and the cells were incubated overnight. Cell proliferation was determined by measuring absorbance at 570 nm with a micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc.). Each PM extract was tested in triplicate. QC laboratory filter extract was tested in the same manner of the PM extracts. The % of inhibition of cell proliferation was calculated comparing the absorbance of exposed cells with the absorbance of control cells (for aqueous extract non-exposed cells and for organic extracts solvent exposed cells).

2.5 Cytotoxicity-LDH assay

To evaluate PM extracts cytotoxicity, LDH activities from damaged cells were measured in cell-free culture supernatants, as previously described and modified for cells in suspension (Schilirò et al., 2015). Briefly, THP-1 cells were seeded in 6-well plates at a density of 1×10^6 cells/well with experimental medium and exposed to PM extracts containing an equivalent particle concentration of 200 $\mu\text{g/mL}$ (Alessandria et al., 2014). At 72 h, LDH activity was measured in the supernatant and cell lysate. LDH activity was calculated as the ratio of extracellular LDH (measured in the supernatant) and total LDH (expressed as sum of LDH measured in supernatant and cell lysate). To obtain cell lysates, cells were centrifuged to eliminate the supernatant and resuspended with 1 mL of TRAP (82.3 mM triethanolamine hydrochloride, pH 7.6) and sonicated for 10 s. Then, LDH was measured by adding 250 μL of a mix containing 0.25 mM NADH and 0.5 mM pyruvate, and consumption of NADH was measured as absorbance at 340 nm in a micro-plate reader (Benchmark Plus Microplate Reader, Biorad). Each PM extracts was tested in triplicate. QC laboratory filter extract was tested in the same manner of the PM extracts. QC laboratory filter extracts were tested in the same manner of PM₁₀ extracts samples. LDH activity of exposed cells is expressed as a percentage of control cells (for aqueous extract non-exposed cells and for organic extracts solvent exposed cells).

2.6 IL-8, TNF α and CYP1A1 gene expression

For quantitative Real Time PCR (qRT-PCR), THP-1 cells were seeded in 6-well culture plates and cultured overnight. Then, the cell culture medium was replaced by a fresh **experimental medium** and **PM extracts were tested at an equivalent particle concentration of** 200 μ g/mL for 48 h. Total RNA was isolated from treated and untreated THP-1 cells as previously described (Schilirò et al., 2015). Briefly total RNA (1 μ g) was used for the first-strand cDNA reaction with reverse transcriptase using an iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA). Relative gene expression was determined by real-time PCR with a Chromo 4 Real-Time System (Bio-Rad, Hercules, CA, USA) using 1 μ L of cDNA sample aliquot (10 ng total mRNA) as a template with Sso Fast EvaGreen SuperMix (Bio-Rad, Hercules, CA, USA).

Three genes were analysed, TNF α , IL-8, CYP1A1. The following primer sequences were used for relative gene expression analysis:

TNF α - f: 5'-ATGAGCACTGAAAGCATGATCCG-3',

TNF α - r: 5'- CAGGCTTGTCACCTCGGGGTTC-3';

IL-8 - f: 5'- TGCCAAGGAGTGCTAAAG-3',

IL-8 - r: 5'-CTCCACAACCCTCTGCAC-3';

CYP1A1 - f: sense 5'- GGCAGATCAACCATGACCAGAAG-3' ,

CYP1A1 - r: 5'-ACAGCAGGCATGCTTCATGGTTAG-3'.

Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and hypoxanthine-guanine phosphoribosyltransferase (HPRT1) gene expression were used as the internal control as they are housekeeping genes and were analysed in each experiment for normalisation using the following primers:

GAPDH - f: 5'-CCCTTCATTGACCTCAACTACATG-3',

GAPDH - r: 5'-TGGGATTTCCATTGATGACAAGC-3';

HPRT1- f: 5'-TGACACTGGCAAAACAATGCA-3',

HPRT1 - r: 5'-GGTCCTTTTCACCAGCAAGCT-3'.

Relative fold inductions were calculated using the Δ Ct formula (Scheffe et al., 2006). **QC laboratory filter extract was tested in the same manner of PM extracts.** All real-time RT-PCR assays for

relative gene expression were repeated at least three times in duplicates from independent total RNA samples for the same treatment conditions. non-exposed cells and solvent exposed cells were used as control cells for aqueous and organic extracts respectively.

2.7 Chemicals and meteo data

Chemical data (metals: As, Cd, Ni, Pb and PAHs: Benzo(a)anthracene, Benzo(b+j+k)fluorantene, Benzo(a)pyrene, Indeno(1,2,3-cd)pyrene) were extracted from a specialised database provided by the Regional System for the real-time monitoring of Air Quality, AriaWeb (ARPA Piemonte, 2015). The data were obtained for the same meteorological–chemical station and for the same days as our sampling. For example the metals data represent a mean of daily data collected using the standard monitoring method EN 14902:2005 «Standard method for measurement of Pb/Cd/As/Ni in the PM₁₀ fraction of suspended particulate matter» (2008/50/EC, annex VI, section B). All the adopted methods conform to the directive and were validated before being published in the AriaWeb database (ARPA Piemonte, 2015).

2.8 Statistical analysis

Statistical analyses were performed using the SPSS Package, version 21.0 for Windows. The Spearman rank order correlation coefficient was used to assess relationships between variables, a Wilcoxon test was used to compare means, and ANOVA was used for multivariate analysis in which we assumed an equal variance while using Tukey as post hoc multiple comparisons. The mean differences and correlations were considered significant at $p < 0.05$.

3. RESULTS

3.1 PM concentration and meteo-chemicals data

The mean PM₁₀ concentration was $23.9 \pm 8.4 \mu\text{g}/\text{m}^3$ in summer and $83.0 \pm 36.2 \mu\text{g}/\text{m}^3$ in winter, and this difference was statistically significant (Wilcoxon test, $p < 0.01$) (Figure 1).

The winter fractions concentrations compared with the corresponding summer ones showed significant differences (Wilcoxon test, $p < 0.01$).

ANOVA analysis performed assuming the gravimetric results as a dependent variable and the different PM₁₀ size fractions as the independent variable reported significant general results

(summer: $F = 15.091$, $p < 0.001$, $d=5$; and winter: $F = 4.904$, $p < 0.01$, $d=5$): the finest fraction $PM_{<0.49}$ was always significantly higher than the others in summer (Tukey test, $p < 0.01$) and represented about 37% of the total PM_{10} mass, while in winter was significantly higher only vs $PM_{10.0-7.2}$ and $PM_{3.0-1.5}$ (Tukey test, $p < 0.01$) and represented about 28% of the total PM_{10} mass; Figure 1 shows also the percentage size composition of PM_{10} .

Table 1 reports the descriptive analysis of the chemical and meteo-climatic parameters collected during the summer and winter samplings.

The mean temperature difference between sampling seasons was significant (Wilcoxon test, $p < 0.01$): the mean winter temperature was $1.2 \pm 4.2^{\circ}C$ and the mean summer temperature was $23.0 \pm 1.8^{\circ}C$. Also radiation was significantly different (Wilcoxon test, $p < 0.01$) in the two considered periods: $23 \pm 3 MJ/m^2$ in summer and $11 \pm 3 MJ/m^2$ in winter.

Neither the average humidity nor the wind speed were significantly different (Wilcoxon test, $p > 0.05$), with 82% humidity during the winter vs. 63% summer and a range wind speed in both seasons of 1.0-2.0 m/s.

ANOVA analysis performed assuming the gravimetric results as a dependent variable and the three main size fractions (analyzed two by two: PM_{10-3} : 10 - 3 μm ; $PM_{3-0.95}$: 3 - 0.95 μm ; and $PM_{<0.95}$: 0.95 - < 0.49 μm) as the independent variable reported a significant result in which the finest fraction ($PM_{<0.95}$) was significantly higher than the others both in summer and in winter (summer: $F = 11.170$, $p < 0.01$, $d=2$; and winter: $F = 5.790$, $p < 0.05$, $d=2$). Beyond this $PM_{<0.95}$ represented about 53% of PM_{10} mass in this urban site both in summer and in winter.

Referring to the annual limits of metals set by the European Directive 2008/50/EC, these are observed even if our sampling reflects only spot seasonal situations (3 weeks in each season) and the European limits refer to yearly averages. Cd and Pb winter concentrations showed significant differences (Wilcoxon test, $p < 0.05$) if compared with the corresponding summer concentrations.

Moreover both total PAHs and single PAHs concentrations were higher during the winter than summer (Wilcoxon test, $p < 0.05$).

3.2 Cell proliferation

Figure 2a shows effects of the different extracts on cell proliferation: inhibition of THP-1 cell proliferation observed *in vitro* was significant respect to control cells for all PM extracts although the observed effect was independent of the aerodynamic diameter of the particles and of kind of PM extract. Both aqueous and organic extracts inhibited cell proliferation with similar effects (Wilcoxon test, $p > 0.05$). The mean percentage proliferation inhibition was 30.8 ± 6.0 % for summer and 28.4 ± 6.9 % for winter fraction extracts. The difference between summer and winter samples was not significant both for organic and aqueous extracts (Wilcoxon test, $p > 0.05$).

3.3 LDH release

LDH assay indicates loss in cell membrane integrity and it was used for cytotoxicity determination. In general, the summer PM extracts induced a more significant release of LDH compared to winter (54.3 ± 36.5 % and 22.6 ± 17.7 % respectively; Wilcoxon test, $p < 0.001$). Significant differences in toxicity between the three different size fraction extracts were observed: summer organic extracts produced a marked size-related effect, with higher values measured with $PM_{10-3} > PM_{3-0.95} > PM_{<0.95}$; considering the winter extracts, LDH release over the negative control was lower than summer ones, except for the finest winter fraction $PM_{<0.95}$ of aqueous extract (Figure 2b).

3.4 IL-8, $TNF\alpha$ and CYP1A1 gene expression

The production of the pro-inflammatory cytokine $TNF\alpha$, as well as the chemokine IL-8 were determined in the cell culture supernatant after 48h of size-fractionated PM extracts exposure; these signaling molecules constitute a major parameter in the mediation of airway inflammation (Gualtieri et al., 2010). Exposure of THP-1 cells to size-fractionated PM_{10} extracts did not induce significant expression of $TNF\alpha$, except for the finest winter fraction both organic and aqueous samples (Figure 3a). On the other side, particles extracts had a stronger effect on IL-8 expression especially organic extracts in summer. Statistically significant differences were found between the two kind of extracts for both season (Wilcoxon test, $p < 0.05$) and comparing summer (4.13 ± 1.34 fold increase) and winter (2.78 ± 0.96 fold increase) extracts (Wilcoxon test, $p < 0.001$) (Figure 3b). No significant differences in toxicity between the extracts of the three different size fractions were observed.

306 The mean fold increase in CYP1A1 expression for summer **extracts** was 3.61 ± 1.18 and for winter
 307 was 3.01 ± 1.46 , and this difference was significant (Figure 3c). The average effect was higher in
 308 summer with the exception of the organic finest fraction **extracts**.
 309 In general in winter the effect was higher for organic extract, indeed there were statistically
 310 significant differences in the effects induced by the two type of extracts (**Wilcoxon test**, $p < 0.05$).
 311 PM **extracts** produced a significant marked size-related CYP1A1 **expression in winter**; ANOVA
 312 analysis performed assuming the **CYP1A1 expression as a dependent variable and the different**
 313 **size fractionated extracts as the independent variable showed a significant difference in the effects**
 314 **produced by winter extracts ($F = 8.756$, $p < 0.01$, $d=2$ organic; $F = 6.291$, $p < 0.01$ $d=2$ aqueous)** with
 315 higher values measured with $PM_{<0.95} > PM_{3-0.95} > PM_{10-3}$.

316

317 **4. DISCUSSION and CONCLUSIONS**

318 In our study size-fractionated PM_{10} was collected in a polluted urban site in Torino (**northwest**
 319 Italian city) during different seasons. In the north of Italy, in particular in Padana Plain, there is a
 320 widespread air pollution; the weak dispersion rate, observed during winter, represents a relevant
 321 factor and it is principally due to the conformation of the territory (Cadum et al., 2009). **Torino PM**
 322 **concentrations** are characterized by high levels of fine particles in winter: during high pollution
 323 periods the $PM_{2.5}/PM_{10}$ ratio increased up to 0.87; in this area the winter PM chemical composition
 324 is characterised by high levels of PAHs and metals (Romanazzi et al., 2014; Traversi et al., 2015)
 325 whereas in summer PM is rich of ions such sulphates (Schilirò et al., 2015). These seasonal
 326 differences could be explained as the winter contribution of domestic heating and of a lower
 327 atmospheric mixing layer also the low temperatures facilitate the condensation and absorption of
 328 volatile compounds on particle surfaces (Brüggemann et al., 2009); during summer, the
 329 photochemical reactions, associated with elevated solar radiation, modify the PM_{10} chemical
 330 constituents.

331 In the present study the observed PM_{10} winter levels are higher than both the WHO guidelines
 332 (Krzyzanowski, 2008) and the EU regulations 2008/50/CE. In addition, critical particle
 333 concentrations are present especially for the finest fraction $PM < 0.49 \mu m$ that represented a very

334 high proportion of PM₁₀: 37% in summer and 28% in winter. Various studies confirmed that the
335 finest fractions are the most abundant in the atmosphere because of the finest particulate pollution
336 is homogeneously diffused (Perez et al., 2010).

337 The high temperature and radiation observed in summer respect to winter attested more frequent
338 photochemical reactions in summer that promotes the formation of secondary particles as
339 highlighted by the percentage of fine fraction (PM_{<0.49}) in summer. In general the percentage of
340 secondary particles increases with decreasing particle size, this is in agreement with the literature
341 data according to which the components of the inorganic fraction of the secondary particles
342 (ammonium, nitrate and sulphate) are present mainly in the finer fraction (Pateraki et al., 2012;
343 Perrino et al., 2014)

344 Among the metals monitored in this study only Cd and Pb had a seasonal trend, As and Ni had
345 similar concentrations in winter and summer; in a previous work in this area, metals appeared to be
346 variously distributed among all the fractions, especially in winter season: As, Ni and Pb were
347 preferentially accumulated on the coarse fractions while Cd was equally distributed (Romanazzi et
348 al., 2014). Contrariwise PAHs showed a marked seasonal trend and the mean winter
349 benzo(a)pyrene concentration was higher both than the annual limit set by the European Directive
350 2008/50/EC (1 ng/m³) and the WHO guide line value of 0.12 ng/m³ (Krzyzanowski, 2008; WHO-
351 Europe, 2013) even if our sampling, as for metals, reflects only a spot seasonal situation. A higher
352 PAHs contamination was observed in the winter and this is probably due to the winter atmospheric
353 conditions that may promote an accumulation of primary pollutants and because of low
354 temperatures, the condensation of atmospheric pollutants in the particle phase (Ebi and McGregor,
355 2008); this seasonal trend was confirmed in other studies (Sisovic et al., 2008).

356 Size-fractionated PM₁₀ extracts (organic and aqueous) exerted a significant impact on cells both in
357 winter and in summer. The presented results indicated that different **PM₁₀ fraction extracts** lead to
358 clear changes in some toxicological responses.

359 Inhibition of proliferation showed a trend without significant differences among fraction **extracts** as
360 found in other **researches** (Longhin et al., 2013; Wessels et al., 2010), however it was significant
361 both in summer and in winter. **The particles size influenced the cytotoxic (LDH release) and pro-**

inflammatory potential (TNF α concentrations). Organic PM extracts induced a significant size-related LDH release, with greater release by the greater fractions. LDH release was significantly greater in summer than in winter with the exception of aqueous extracts of the finest fraction in which transition metals are mainly accumulated (Daher et al., 2014). Both organic and aqueous winter extracts of the finer fraction induced a significant TNF α expression that may be associated to the presence of PAHs and transition metals respectively in the finer fractions (de Kock et al., 2006).

Summer fraction extracts were able to trigger a pro-inflammatory response with the release of significant quantity of IL-8, especially the summer greater fractions. Actually various *in vitro* data showed the higher inflammatory potential of PM extracts of coarse fraction (Hetland et al., 2004; Jalava et al., 2007). Inflammatory effect of summer PM extracts could be related to the presence of biogenic compounds (Gualtieri et al., 2010; Schins et al., 2004); the role of endotoxins in leading the PM inflammatory effects has been well documented (Hetland et al., 2004; Traversi et al., 2010). Inflammatory effect of aqueous extracts of PM₁₀ fractions could also be related to the particles bound metals; the PM chemical composition describes the presence of crustal elements, ions and metals, variously distributed among all the fractions (Romanazzi et al., 2014), which have been showed to induce cell inflammation (Øvrevik et al., 2005; Veranth et al., 2006).

Apparently Torino summer PM₁₀ fraction extracts in a traffic site had a higher proinflammatory potential and induces stronger IL-8 responses in confront of a remote site (Schilirò et al., 2015), similar results have been presented by numerous authors (Camatini et al., 2012; Duvall et al., 2008; Gualtieri et al., 2010; Hetland et al., 2004).

PM extracts induced a significant size-related CYP1A1 expression, in particular for organic winter extracts. The importance of organic compounds, with regard to CYP1A1 expression, was confirmed as all the winter organic extracts induced significant size-related levels of this biological marker, with greater increase by the finer fraction extracts. This can easily be associated to the link between the majority of particle-bound PAHs and the fine and ultra-fine particles (Claxton et al., 2004; Di Filippo et al., 2010a). In this study, CYP1A1 expression was significant also for the summer extracts even if PAHs levels were lower highlighting the presence of other chemical

390 compounds capable to induce the CYP1A1 expression (e.g. dioxin like molecules) (Wenger et al.,
391 2009).

392 Torino PM₁₀ winter fractions in a traffic site showed a higher CYP1A1 expression compared to a
393 background urban site (Schilirò et al., 2015) especially the finer fractions extracts.

394 No statistically significant association were found between the THP-1 endpoints and chemical
395 (metals or PAHs) or meteo-climatic parameters collected during the summer and winter samplings.
396 One possible explanation for the lack of correlations could be that the chemical parameters were
397 quantified in total PM₁₀ samples instead of separated fractions.

398 Comparing results obtained in this work considering a trafficated Torino urban site with those
399 reported in a previous study conducted in a background Torino urban site in the same seasons
400 (Schilirò et al., 2015), the total PM₁₀ concentration was higher in the trafficated urban site in winter
401 but showed similar values respect to background site in summer. Moreover, the biological effects
402 were higher in the trafficated site mostly considering inhibition of cell proliferation and CYP1A1
403 expression in winter and LDH release and IL-8 expression in summer.

404 Our results show that biological responses significantly differ after exposure to equal mass
405 concentrations of urban traffic and background Torino urban site. These observations confirm the
406 hypothesis that particle composition, as well as source, constitute an important factor in PM
407 induced toxicity as underlined in other studies (Michael et al., 2013).

408 It is now widely accepted that air pollution and its major components have a marked seasonality,
409 and the toxic content in particulates can vary based upon the meteo-climatic conditions (Albinet et
410 al., 2008). However, biological effects underlined the potential importance of the non-regulated
411 pollutants to assess the air quality (Di Filippo et al., 2010a). This complexity is the reason for
412 difficulties in solving the question of PM toxicity.

413 In addition, our results confirm that the only measure of PM can be misleading for the assessment
414 of air quality (Di Filippo et al., 2010b) in fact PM₁₀ or PM_{2.5} monitoring itself is not informative
415 enough on toxic compounds bound to particles (Topinka et al., 2015). Evaluation of air pollution
416 should not be limited just to gravimetric assessment of PM but should consider also the size
417 spectrum of more dangerous PM fractions. Because of their large and irregular specific surface,

the finest fractions can favour the adsorption of mutagenic and carcinogenic species (such as PAHs, nitro-PAHs and metals) (Claxton et al., 2004; Topinka et al., 2015), moreover they can reach the deepest sites of the respiratory system (alveolar lung region). The use of different types of extract allows to evaluate the role of chemical composition on biological effects of different PM fraction extracts.

Moreover, the use of *in vitro* test, that integrate toxic potencies of samples including unidentified as well as identified chemicals, can give a correct estimation of the impact of a complex mixture.

Finally in this study we support efforts toward establishing more effective and source-specific regulations for mitigating PM toxicity (Daher et al., 2014) and identifying potential effect-based tools (e.g. *in vitro* test on specific cell line such as THP-1 or A549) that could be used in the context of the different monitoring programmes after being properly standardized. These kind of effect-based monitoring tools could also be valuable in toxicity identification evaluation (TIE) approaches to identify toxic fractions and provide guidance for the identification of causative agents.

431

432 ACKNOWLEDGMENTS

The authors kindly thank Drs M. Sacco, F. Lollobrigida and M. Grosa of the Environmental Protection Regional Agency (Piedmont A.R.P.A.). This study was financed by Torino University Local Research grant (ex-60%-2012).

436

437 REFERENCES

Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., Jaffrezo, J.L., 2008. Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys Part 1: concentrations, sources and gas/particle partitioning. *Atmos. Environ.* 42, 43-54.

Alessandria, L., Schilirò, T., Degan, R., Traversi, D., Gilli, G., 2014. Cytotoxic response in human lung epithelial cells and ion characteristics of urban-air particles from Torino, a northern Italian city. *Environ. Sci. Pollut. Res. Int.* 21, 5554-5564.

Anderson, J.O., Thundiyil, J.G., Stolbach, A., 2012. Clearing the air: review of the effects of particulate matter air pollution on human health. *J. Med. Toxicol.* 8, 166–175.

ARPA Piemonte, S.P., CSI Piemonte, 2015, ARIAWEB – Sistema Regionale Di Rilevamento Della Qualità Dell'aria Della Regione Piemonte.

Bonetta, S., Gianotti, V., Bonetta, S., Gosetti, F., Oddone, M., Gennaro, M.C., Carraro, E., 2009. DNA damage in A549 cells exposed to different extracts of PM_(2.5) from industrial, urban and highway sites. *Chemosphere* 77, 1030-1034.

Brook, R.D., Rajagopalan, S., Pope, C.A., Brook, J.R., Bhatnagar, A., Diez-Roux, A.V., Holguin Hong, Y.L., Luepker, R.V., Mittleman, M.A., Peters, A., Siscovick, D., Smith, S.C., Whitsel, L., Kaufman, J.D., Amer Heart Assoc Council E, Council Kidney Cardiovasc D, Council Nutr Phys Activity M., 2010. Particulate Matter Air Pollution and Cardiovascular Disease An Update to the Scientific Statement From the American Heart Association. *Circulation* 121, 2331-2378.

Brüggemann, E., Gerwig, H., Gnauk, T., Muller, K., Herrmann, H., 2009. Influence of seasons, air mass origin and day of the week on size-segregated chemical composition of aerosol particles at a kerbside. *Atmos. Environ.* 43, 2456-2463.

Cadum, E., Berti, G., Biggeri, A., Bisanti, L., Faustini, A., Forastiere, F., Grp Collaborativo EpiAir, 2009. The results of EpiAir and the national and international literature. *Epidemiol. Prev.* 33, 113-119.

Camatini, M., Corvaja, V., Pezzolato, E., Mantecca, P., Gualtieri, M., 2012. PM₁₀-biogenic fraction drives the seasonal variation of proinflammatory response in A549 cells. *Environ. Toxicol.* 27, 63-73.

CEN, 1998. Air quality - determination of the PM₁₀ fraction of suspended particulate matter – reference method and field test procedure to demonstrate reference equivalence of measurement methods. Brussels: European Committee for Standardization; (European Standard EN 12341).

Claxton, L.D., Matthews, P.P., Warren, S.H., 2004. The genotoxicity of ambient outdoor air, a review: Salmonella mutagenicity. *Mutat. Res.* 567, 347-399.

472 Corsini, E., Budello, S., Marabini, L., Galbiati, V., Piazzalunga, A., Barbieri, P., Cozzutto, S.,
 473 Marinovich, M., Pitea, D., Galli, C.L., 2013. Comparison of wood smoke PM_{2.5} obtained from the
 474 combustion of FIR and beech pellets on inflammation and DNA damage in A549 and THP-1
 475 human cell lines. *Arch. Toxicol.* 87, 2187-2199.

476 Daher, N., Saliba, N.A., Shihadeh, A.L., Jaafar, M., Baalbaki, R., Shafer, M.M., Schauer, J.J.,
 477 Sioutas, C., 2014. Oxidative potential and chemical speciation of size-resolved particulate
 478 matter (PM) at near-freeway and urban background sites in the greater Beirut area. *Sci.Total*
 479 *Environ.* 470-471, 417-426.

480 Danielsen, P.H., Møller, P., Jensen, K.A., Sharma, A.K., Wallin, H., Bossi, R., Autrup, H., Mølhave,
 481 L., Ravanat, J.L., Briedé, J.J., de Kok, T.M., Loft, S., 2011. Oxidative stress, DNA damage, and
 482 inflammation induced by ambient air and wood smoke particulate matter in human A549 and
 483 THP-1 cell lines. *Chem. Res. Toxicol.* 24,168-184.

484 de Kock, T.M., Driee, H.A.L., Hogervorst, J.G.F., Briede, J.J., 2006. Toxicological assessment of
 485 ambient and traffic-related ambient particulate matter: a review of recent studies. *Mutat. Res.*
 486 613, 103-122.

487 Di Filippo, P., Riccardi, C., Pomata, D., Buiarelli, F., 2010a. Concentrations of PAHs, and nitro- and
 488 methyl- derivatives associated with a size-segregated urban aerosol. *Atmos. Environ.* 44, 2742-
 489 2749.

490 Di Filippo, P., Riccardi, C., Pomata, D., Gariazzo, C., Buiarelli, F., 2010b. Seasonal Abundance of
 491 Particle-Phase Organic Pollutants in an Urban/Industrial Atmosphere. *Water Air Soil Pollut.*
 492 211, 231–250.

493 Duvall, R.M., Norris, G.A., Dailey, L.A., Burke, J.M., McGee, J.K., Gilmour, M.I., Gordon, T., Devlin,
 494 R.B., 2008. Source apportionment of particulate matter in the U.S. and associations with lung
 495 inflammatory markers. *Inhal. Toxicol.* 20, 671-683.

496 Ebi, K.L., McGregor, G., 2008. Climate change, tropospheric ozone and particulate matter, and
 497 health impacts. *Environ. Health Perspect.* 116, 1449-1455.

498 Gualtieri, M., Øvrevik, J., Holme, J.A., Perrone, M.G., Bolzacchini, E., Schwarze, P.E., Camatini,
 499 M., 2010. Differences in cytotoxicity versus pro-inflammatory potency of different PM fractions
 500 in human epithelial lung cells. *Toxicol. In Vitro* 24, 29-39.

501 Hanzalova, K., Rossner, Jr P., Sram, R.J., 2010. Oxidative damage induced by carcinogenic
 502 polycyclic aromatic hydrocarbons and organic extracts from urban air particulate matter. *Mutat.*
 503 *Res.* 696, 114–121.

504 Hetland, R.B., Cassee, F.R., Refsnes, M., Schwarze, P.E., Lag, M., Boere, A.J., Dybing, E., 2004.
 505 Release of inflammatory cytokines, cell toxicity and apoptosis in epithelial lung cells after
 506 exposure to ambient air particles of different size fractions. *Toxicol. In Vitro* 18, 203-212.

507 Jalava, P.I., Salonen, R.O., Pennanen, A.S., Sillanpää, M., Hälinen, A.I., Happonen, M.S., Hillamo, R.,
 508 Brunekreef, B., Katsouyanni, K., Sunyer, J., Hirvonen, M.R., 2007. Heterogeneities in
 509 inflammatory and cytotoxic responses of RAW 264.7 macrophage cell line to urban air coarse,
 510 fine, and ultrafine particles from six European sampling campaigns. *Inhal. Toxicol.* 19, 213-225.

511 Knaapen, A.M., Borm, P.J., Albrecht, C., Schins, R.P., 2004. Inhaled particles and lung cancer.
 512 Part A: Mechanisms. *Int. J. Cancer* 109, 799-809.

513 Kroll, A., Gietl, J.K., Wiesmüller, G.A., Günsel, A., Wohlleben, W., Schnekenburger, J., Klemm, O.,
 514 2013. In vitro toxicology of ambient particulate matter: correlation of cellular effects with particle
 515 size and components. *Environ. Toxicol.* 28, 76-86.

516 Krzyzanowski, M., 2008. WHO air quality guidelines for Europe. *J. Toxicol. Environ. Health A* 71,
 517 47–50.

518 Lauer, F.T., Mitchell, L.A., Bedrick, E., McDonald, J.D., Lee, W.Y., Li, W.W., Olvera, H., Amaya,
 519 M.A., Berwick, M., Gonzales, M., Currey, R., Pingitore, N.E., Jr., Burchiel, S.W., 2009.
 520 Temporal spatial analysis of U.S.-Mexico border environmental fine and coarse PM air sample
 521 extract activity in human bronchial epithelial cells. *Toxicol. Appl. Pharmacol.* 238, 1–10.

522 Longhin, E., Pezzolato, E., Mantecca, P., Holme, J.A., Franzetti, A., Camatini, M., Gualtieri, M.,
 523 2013. Season linked responses to fine and quasi-ultrafine Milan PM in cultured cells. *Toxicol. In*
 524 *Vitro* 27, 551-559.

525 Lonkar, P., Dedon, P.C., 2011. Reactive species and DNA damage in chronic inflammation:
 526 reconciling chemical mechanisms and biological fates. *Int. J. Cancer* 128, 1999-2009.

527 Loomis, D., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L.,
 528 Guha, N., Baan, R., Mattock, H., Straif, K., IARC working group, 2013. The carcinogenicity of
 529 outdoor air pollution. *Lancet Oncol.* 14, 1262–1263.

530 Michael, S., Montag, M., Dott, W., 2013. Pro-inflammatory effects and oxidative stress in lung
 531 macrophages and epithelial cells induced by ambient particulate matter. *Environ. Pollut.* 183,
 532 19-29.

533 Oh, S.M., Kim, H.R., Park, Y.J., Lee, S.Y., Chung, K.H., 2011. Organic extracts of urban air
 534 pollution particulate matter (PM_{2.5})-induced genotoxicity and oxidative stress in human lung
 535 bronchial epithelial cells (BEAS-2B cells). *Mutat. Res.* 723, 142-151.

536 Øvrevik, J., Myran, T., Refnes, M., Lag, M., Becher, R., Hetland, R.B., 2005. Mineral particles of
 537 varying composition induce differential chemokine release from epithelial lung cells: importance
 538 of physicochemical characteristics. *Ann. Occup. Hyg.* 49, 219-231.

539 Pateraki, S., Assimakopoulos, V.D., Bougiatioti, A., Kouvarakis, G., Mihalopoulos, N., Vasilakos,
 540 Ch., 2012. Carbonaceous and ionic compositional patterns of fine particles over an urban
 541 Mediterranean area. *Sci. Total Environ.* 424, 251-263.

542 Perez, N., Pey, J., Cusack, M., Reche, C., Querol, X., Alastuey, A., Viana, M., 2010. Variability of
 543 Particle Number, Black Carbon, and PM₁₀, PM_{2.5}, and PM₁ Levels and Speciation: Influence of
 544 Road Traffic Emissions on Urban Air Quality. *Aerosol Sci. Tech.* 44, 487-499.

545 Perrino, C., Catrambone, M., Dalla Torre, S., Rantica, E., Sargolini, T., Canepari, S., 2014.
 546 Seasonal variations in the chemical composition of particulate matter: a case study in the Po
 547 Valley. Part I: macro-components and mass closure. *Environ. Sci. Pollut. Res. Int.* 21, 3999-
 548 4009.

549 Roig, N., Sierra, J., Rovira, J., Schuhmacher, M., Domingo, J.L., Nadal, M., 2013. In vitro tests to
 550 assess toxic effects of airborne PM₍₁₀₎ samples. Correlation with metals and chlorinated dioxins
 551 and furans. *Sci. Total Environ.* 443, 791-797.

552 Romanazzi, V., Casazza, M., Malandrino, M., Maurino, V., Piano, A., Schilirò, T., Gilli, G., 2014.
 553 PM₁₀ size distribution of metals and environmental-sanitary risk analysis in the city of Torino.
 554 *Chemosphere* 112, 210-216.

555 Schefe, J.H., Lehmann, K.E., Buschmann, I.R., Unger, T., Funke-Kaiser, H., 2006. Quantitative
 556 real-time RT-PCR data analysis: current concepts and the novel "gene expression's C-T
 557 difference" formula. *J. Mol. Med.* 84, 901-910.

558 Schilirò, T., Alessandria, L., Degan, R., Traversi, D., Gilli, G., 2010. Chemical characterisation and
 559 cytotoxic effects in A549 cells of urban-air PM₁₀ collected in Torino, Italy. *Environ. Toxicol.*
 560 *Pharmacol.* 29, 150-157.

561 Schilirò, T., Bonetta, S., Alessandria, L., Gianotti, V., Carraro, E., Gilli, G., 2015. PM₁₀ in a
 562 background urban site: chemical characteristics and biological effects. *Environ. Toxicol.*
 563 *Pharmacol.* 39, 833-844.

564 Schins, R.P., Lightbody, J.H., Borm, P.J., Shi, T., Donaldson, K., Stone, V., 2004. Inflammatory
 565 effects of coarse and fine particulate matter in relation to chemical and biological constituents.
 566 *Toxicol. Appl. Pharmacol.* 195, 1-11.

567 Schins, R.P.F., Hei, T.K., 2006. Genotoxic effects of particles, in: Donaldson, K., Borm, P., (Eds.),
 568 *Particle Toxicology*. CRC Press/Taylor and Francis Group, Boca Raton, FL, pp 285-298.

569 Schwarze, P.E., Ovreivik, J., Hetland, R.B., Becher, R., Cassee, F.R., Lag, M., Lovik, M., Dybing,
 570 E., Refsnes, M., 2007. Importance of size and composition of particles for effects on cells in
 571 vitro. *Inhal. Toxicol.* 19, 17-22.

572 Sisovic, A., Beslic, I., Segar, K., Vadjic, V., 2008. PAH mass concentrations measured in PM₁₀
 573 particle fraction. *Environ. Int.* 34, 580-584.

574 Spink, D.C., Wu, S.J., Spink, B.C., Hussain, M.M., Vakharia, D.D., Pentecost, B.T., Kaminsky,
 575 L.S., 2008. Induction of CYP1A1 and CYP1B1 by benzo(k)fluoranthene and benzo(a)pyrene in

576 T-47D human breast cancer cells: roles of PAH interactions and PAH metabolites. *Toxicol.*
577 *Appl. Pharmacol.* 226, 213-24.

578 Steenhof, M., Gosens, I., Strak, M., Godri, K.J., Hoek, G., Cassee, F.R., Mudway, I.S., Kelly, F.J.,
579 Harrison, R.M., Lebret, E., Brunekreef, B., Janssen, N.A.H., Pieters, R.H.H., 2011. In vitro
580 toxicity of particulate matter (PM) collected at different sites in the Netherlands is associated
581 with PM composition, size fraction and oxidative potential e the RAPTES project. *Part. Fibre*
582 *Toxicol.* 8, 26.

583 Traversi, D., Alessandria, L., Schilirò, T., Chiadò Piat, S., Gilli, G., 2010. Meteo-climatic conditions
584 influence the contribution of endotoxins to PM₁₀ in an urban polluted environment. *J. Environ.*
585 *Monit.* 12, 484-490.

586 Traversi, D., Alessandria, L., Schilirò, T., Gilli, G., 2011. Size-fractionated PM₁₀ monitoring in
587 relation to the contribution of endotoxins in different polluted areas. *Atmos. Environ.* 45, 3515-
588 3521.

589 Traversi, D., Festa, E., Pignata, C., Gilli, G., 2015. Aero-dispersed mutagenicity attributed to
590 particulate and semi volatile phase in an urban environment. *Chemosphere.* 124, 163-169.

591 Valavanidis, A., Fiotakis, K., Vlachogianni, T., 2008. Airborne particulate matter and human health:
592 toxicological assessment and importance of size and composition of particles for oxidative
593 damage and carcinogenic mechanisms. *J. Environ. Sci. Health C* 26, 339-362.

594 Veranth, J.M., Moss, T.A., Chow, J.C., Labban, R., Nichols, W.K., Walton, J.C., Watson, J.G.,
595 Yost, G.S., 2006. Correlation of in vitro cytokine responses with the chemical composition of
596 soil-derived particulate matter. *Environ. Health Perspect.* 114, 314-349.

597 Wenger, D., Gerecke, A.C., Heeb, N.V., Hueglin, C., Seiler, C., Haag, R., Naegeli, H., Zenobi, R.,
598 2009. Aryl hydrocarbon receptor-mediated activity of atmospheric particulate matter from an
599 urban and a rural site in Switzerland. *Atmos. Environ.* 43, 3556-3562.

600 Wessels, A., Birmili, W., Albrecht, C., Hellack, B., Jeremann, E., Wick, G., Harrison, R.M., Schins,
601 R.P.F., 2010. Oxidant Generation and Toxicity of Size-Fractionated Ambient Particles in
602 Human Lung Epithelial Cells. *Environ. Sci. Technol.* 44, 3539-3545.

603 WHO-Europe, 2013. Review of Evidence On Health Aspects Of Air Pollution – Technical Report –
604 REVIHAAP Project.

HIGHLIGHTS

The PM₁₀ finest fraction PM < 0.49 µm was the highest one both in winter and in summer

The particles size influenced both LDH release and IL-8 concentrations

Size-fractionated PM₁₀ extracts induced a significant size-related CYP1A1 expression

Air pollution evaluation should consider the size spectrum of dangerous PM fractions

Inflammation response and cytotoxic effects in human THP-1 cells of size-fractionated PM₁₀ extracts in a polluted urban site.

T. Schilirò^{*a}, L. Alessandria^{a^}, S. Bonetta^{a^}, E. Carraro^a, G. Gilli^a

^a Department of Public Health and Pediatrics, University of Torino, Piazza Polonia, 94 - 10126, Torino, Italy; tiziana.schiliro@unito.it; luca.al@hotmail.it; sara.bonetta@unito.it; elisabetta.carraro@unito.it; giorgio.gilli@unito.it

[^]These authors contributed equally to the work.

***Corresponding author:**

Tiziana Schilirò

Department of Public Health and Pediatrics,
University of Torino,
Piazza Polonia, 94 - 10126, Torino, Italy,
Tel: +390116705820
e-mail address: tiziana.schiliro@unito.it

ABSTRACT

To contribute to a greater characterization of the airborne particulate matter's toxicity, size-fractionated PM₁₀ was sampled during different seasons in a polluted urban site in Torino, a northern Italian city. Three main size fractions (PM₁₀₋₃ - 3 µm; PM_{3-0.95} - 0.95 µm; PM_{<0.95} - <0.95 µm) extracts (organic and aqueous) were assayed with THP-1 cells to evaluate their effects on cell proliferation, LDH activity, TNFα, IL-8 and CYP1A1 expression.

The mean PM₁₀ concentrations were statistically different in summer and in winter and the finest fraction PM_{<0.95} was always higher than the others. Size-fractionated PM₁₀ extracts, sampled in an urban traffic meteorological–chemical station produced size-related toxicological effects in relation to season and particles extraction. The PM summer extracts induced a significant release of LDH compared to winter and produced a size-related effect, with higher values measured with PM₁₀₋₃. Exposure to size-fractionated PM₁₀ extracts did not induce significant expression of TNFα. IL-8 expression was influenced by exposure to size-fractionated PM₁₀ extracts and statistically significant differences were found between kind of extracts for both seasons. The mean fold increases in CYP1A1 expression were statistically different in summer and in winter; winter fraction extracts produced a size-related effect, in particular for organic samples with higher values measured with PM_{<0.95} extracts.

Our results confirm that the only measure of PM can be misleading for the assessment of air quality moreover we support efforts toward identifying potential effect-based tools (e.g. *in vitro* test) that could be used in the context of the different monitoring programs.

Keywords: size-fractionated PM₁₀; THP-1 cells; LDH; IL-8; TNFα; CYP1A1

57 1. INTRODUCTION

58 In the last ten years, a considerable number of new studies have been published, providing
59 evidence on the health effects of size fractions, components and sources of particulate matter
60 (PM). Health effects are observed with short-term and long-term exposures to airborne particles
61 indicating an increase in cardiac and respiratory morbidity and associated mortality (WHO-Europe,
62 2013).

63 An increasing body of epidemiological and toxicological evidences have shown PM mass (PM_{2.5}
64 and PM₁₀) comprises fractions with varying types and degrees of health effects, suggesting a role
65 for both the chemical composition (such as transition metals, polycyclic aromatic hydrocarbons -
66 PAHs), physical properties (size, particle number and surface area) and source (Brook et al., 2010;
67 Kroll et al., 2013; Krzyzanowski, 2008; Steenhof et al., 2011).

68 Airborne particles are generally heterogeneous mixtures consisting of inorganic components (e.g.,
69 transition metals), salts, carbonaceous material, volatile organic compounds (VOC), PAHs and
70 biological materials such as endotoxins, fungal spores and pollen (Anderson et al., 2012). Although
71 regulatory standards for PM₁₀ and PM_{2.5} as well as specific particle constituents exist, there is still
72 no clear consensus among researchers regarding the mechanisms of toxicity following exposure to
73 current levels of air pollution PM (Schwarze et al., 2007).

74 Many of the observed health endpoints may result, at least in part, from oxidative stress initiated by
75 the formation of reactive oxygen species (ROS) upon the interaction of PM with epithelial cells and
76 macrophages (Knaapen et al., 2004). Induction of cellular oxidative stress and resulting activation
77 of pro-inflammatory mediators are considered to play a central role in the development of airway
78 diseases (Lonkar and Dedon, 2011; Oh et al., 2011). Pulmonary inflammation is generally
79 characterized by local recruitment of proinflammatory cells such as neutrophils and macrophages,
80 which are involved in the up regulation of various signaling molecules, such as cytokines (TNF α ,
81 IL-6) and chemokines (IL-8) (Lauer et al., 2009; Michael et al., 2013). Based on these
82 mechanisms, increased production and release of inflammatory mediators by cells are relevant
83 and often used parameters suggesting PM-induced toxicity. These mediators in connection with an
84 increasing ROS formation play a key role in the development of several inflammatory diseases like

acute lung injury, COPD, chronic bronchitis and asthma (Valavanidis et al., 2008). Besides the ability to introduce inflammation, particles may also cause oxidative DNA damage because of their physicochemical properties. Within this so-called primary genotoxicity, surface associated free radicals and transition metals are considered to play a major role (Roig et al., 2013). Moreover particle-elicited inflammation and subsequent generation of ROS can lead to oxidative DNA damage, and this pathway is defined as secondary genotoxicity (Schins and Hei, 2006). In addition, organic compounds, such as polycyclic aromatic hydrocarbons (PAH) coated onto PM may induce DNA damage, mutations and cytotoxicity at a cellular level (Bonetta et al., 2009). Upon entering the organism, PAHs are first metabolized to transdihydrodiols by the activity of the cytochromes P450 (CYP) superfamily member CYP1A1 enzyme and epoxide hydrolase and then oxidized to reactive electrophiles by two pathways (Hanzalova et al., 2010; Spink et al., 2008). Intermediates thereafter interact with DNA target sites to produce adducts, mutation, DNA strand breaks and eventually tumour initiation (Schwarze et al., 2007).

In this regards, recently, outdoor air pollution and its major component, outdoor particulate matter, were classified as carcinogenic for humans (Group1) (Loomis et al., 2013). Consequently, reducing air pollution and particle matter to the lowest amount possible is becoming a marked priority. However, in order to define the most cost-effective policies, the most health-relevant PM fractions will have to be identified based on toxicological characteristics. With this purpose, to contribute to a greater characterization of the airborne particulate matter's toxicity, the biological effects of airborne size-fractionated PM₁₀ collected in a high polluted urban site during different seasons were examined. The effects of size fractionated PM₁₀ extracts (organic and aqueous) from winter and summer seasons were evaluated in human cell cultured *in vitro* (THP-1 monocytes) considering as endpoints: the inhibition of cell proliferation, lactate dehydrogenase (LDH) activity and tumour necrosis factor alpha (TNF α), interleukin 8 (IL-8) and CYP1A1 expressions.

110

111 2. MATERIALS AND METHODS

112 2.1 Size-fractionated PM₁₀ sampling and gravimetric analysis

113 The sampling site was located in Torino (northwest of the Italian Padana Plain), in a
114 meteorological–chemical station of the Environmental Protection Regional Agency (Piedmont
115 A.R.P.A.), it was representative of typical urban pollution, with high vehicle traffic impact
116 (coordinates utm, WGS84, X: 394836, Y: 4996153).

117 The samplings were made during two periods, summer (July/August 2011) and winter
118 (January/February 2012), PM₁₀ samples were collected using a Sierra-Andersen high-volume
119 cascade impactor (AirFlow PM₁₀-HVS sampler which a multi-stage cascade impactor, with pre-
120 selector complying with the EN-12341 norm by Analitica Strumenti) at an electronically controlled
121 flow at 1160 L/min. In each period, sampling durations were 24 hours and were repeated on
122 Tuesday and Friday for three weeks (6 samplings and 6 filters for each fraction, for a total of 36
123 filters in each period).

124 First, PM₁₀ was selected by a pre-selector, and then the multistage impactor determined the
125 division of different particle sizes of the sampled particles by the differentiation of the aerodynamic
126 diameter, which can identify the type of trajectory that particles take inside the suction flow related
127 to the three main aerodynamic factors of the particles themselves: dimension, shape and density
128 (Analitica Strumenti). Particles with sufficient inertia will impact on that particular stage collection
129 plate, whilst smaller particles will remain entrained in the air stream and pass to the next stage
130 where the process is repeated. The stages are assembled in a stack or row in order of decreasing
131 particle size. The particle size fractions are as follows: 10.0-7.2, 7.2-3.0, 3.0-1.5, 1.5-0.95, 0.95-
132 0.49, and <0.49 µm. Glass microfiber filters with splits (Type A/E; 8"x10"; Gelman Sciences, MI,
133 USA) were used to collect particles on each impactor plate; finally, glass microfiber filters (20.32 ×
134 25.40 cm; Pall Corporation, NY, USA) were present as back-up filters to collect the finest particles
135 (<0.49 µm). All of the filters were pre- and post-conditioned by placing them in a dry and dark
136 environment for 48 h, and then they were weighted in a room with a controlled temperature and
137 humidity. The PM concentrations in the air volume sampled was calculated as previously described
138 (Schilirò et al., 2010; Traversi et al., 2011), according to the European Committee for
139 Standardization (CEN, 1998).

140 **2.2 Particles extraction**

141 After gravimetric analyses, one strip was cut from each fractionated PM filters. The strips were
142 pooled into three main size fractions (PM_{10-3} : 10 - 3 μm ; $PM_{3-0.95}$: 3 – 0.95 μm ; and $PM_{<0.95}$: <0.95
143 μm) (two strips for each main size fraction) for a total of 12 strips for each main fraction both in
144 summer and in winter. This procedure was repeated two times to perform two different extractions
145 chosen for different abilities in the extraction of different compounds: acetone was selected as the
146 solvent for organic-extractable compounds (e.g., PAHs) (Claxton et al., 2004) and RPMI1640,
147 without foetal calf serum (FCS,) was chosen to extract water-soluble components (e.g., metals),
148 theoretically comparable to the extraction at the lung cells (Hetland et al., 2004). The 12 strips for
149 each main fraction both in summer and winter were cut into small pieces and were placed in 50 mL
150 polypropylene sterile tubes with 15 mL of each extraction media. The tubes were placed in an
151 ultrasonic water bath for 10 min, followed by 1 min of vortexing. This procedure was repeated 3
152 times (45 mL of extracts). The samples were centrifuged at 5000 x g for 10 min to remove the filter
153 material, and supernatant was collected. Acetone extract (organic) for biological tests was
154 evaporated with a rotary evaporator and suspended in dimethyl sulfoxide (DMSO) for the THP-1
155 cells tests. RPMI extracts (aqueous) were directly assayed for biological analysis (Alessandria et
156 al., 2014; Schilirò et al., 2010). Each extraction was also performed on a QC laboratory filter
157 (treated with the same method as the samples). Unless specified otherwise, all chemicals were
158 purchased from Sigma, St. Louis, MO, USA.

159 **2.3 Cell culture**

160 The human THP-1 monocytes from Interlab Cell Line Collection (Genova, IT) were used as
161 surrogates of lung monocytes and were grown, maintained and treated in RPMI1640
162 supplemented with 10 % (v/v) FCS, 2 % L-glutamine 200 mM and 1 % penicillin/streptomycin 10
163 mg/mL, at 37 °C in an humidified atmosphere containing 5% CO₂. Serum-free RPMI1640 without
164 phenol red constituted the experimental medium. The use of serum-free allow a better
165 standardization of experimental conditions. THP-1 cells are more sensitive in cytotoxicity tests
166 respect to other cell lines (e.g. A549 cells) (Corsini et al., 2013).

167 **2.4 Cell viability- MTT assay**

MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay was used to measure cell viability as previously described (Schilirò et al., 2015); briefly THP-1 cells were seeded, using experimental medium, in 24-well plates at a density of 4×10^4 cell/well and exposed to an equivalent particle concentration of 200 $\mu\text{g/mL}$. After treatment with PM extracts (72 h), the MTT (final concentration 0.5 mg/mL) was added to cells and incubated at 37°C for 4 h. Then, 1 mL 10 % SDS/0.01 M HCl was added in each well, and the cells were incubated overnight. Cell proliferation was determined by measuring absorbance at 570 nm with a micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc.). Each PM extract was tested in triplicate. QC laboratory filter extract was tested in the same manner of the PM extracts. The % of inhibition of cell proliferation was calculated comparing the absorbance of exposed cells with the absorbance of control cells (for aqueous extract non-exposed cells and for organic extracts solvent exposed cells).

2.5 Cytotoxicity-LDH assay

To evaluate PM extracts cytotoxicity, LDH activities from damaged cells were measured in cell-free culture supernatants, as previously described and modified for cells in suspension (Schilirò et al., 2015). Briefly, THP-1 cells were seeded in 6-well plates at a density of 1×10^6 cells/well with experimental medium and exposed to PM extracts containing an equivalent particle concentration of 200 $\mu\text{g/mL}$ (Alessandria et al., 2014). At 72 h, LDH activity was measured in the supernatant and cell lysate. LDH activity was calculated as the ratio of extracellular LDH (measured in the supernatant) and total LDH (expressed as sum of LDH measured in supernatant and cell lysate). To obtain cell lysates, cells were centrifuged to eliminate the supernatant and resuspended with 1 mL of TRAP (82.3 mM triethanolamine hydrochloride, pH 7.6) and sonicated for 10 s. Then, LDH was measured by adding 250 μL of a mix containing 0.25 mM NADH and 0.5 mM pyruvate, and consumption of NADH was measured as absorbance at 340 nm in a micro-plate reader (Benchmark Plus Microplate Reader, Biorad). Each PM extracts was tested in triplicate. QC laboratory filter extract was tested in the same manner of the PM extracts. QC laboratory filter extracts were tested in the same manner of PM₁₀ extracts samples. LDH activity of exposed cells is expressed as a percentage of control cells (for aqueous extract non-exposed cells and for organic extracts solvent exposed cells).

2.6 IL-8, TNF α and CYP1A1 gene expression

For quantitative Real Time PCR (qRT-PCR), THP-1 cells were seeded in 6-well culture plates and cultured overnight. Then, the cell culture medium was replaced by a fresh experimental medium and PM extracts were tested at an equivalent particle concentration of 200 μ g/mL for 48 h. Total RNA was isolated from treated and untreated THP-1 cells as previously described (Schilirò et al., 2015). Briefly total RNA (1 μ g) was used for the first-strand cDNA reaction with reverse transcriptase using an iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA). Relative gene expression was determined by real-time PCR with a Chromo 4 Real-Time System (Bio-Rad, Hercules, CA, USA) using 1 μ L of cDNA sample aliquot (10 ng total mRNA) as a template with Sso Fast EvaGreen SuperMix (Bio-Rad, Hercules, CA, USA).

Three genes were analysed, TNF α , IL-8, CYP1A1. The following primer sequences were used for relative gene expression analysis:

TNF α - f: 5'-ATGAGCACTGAAAGCATGATCCG-3',

TNF α - r: 5'-CAGGCTTGTCACCTCGGGGTTC-3';

IL-8 - f: 5'-TGCCAAGGAGTGCTAAAG-3',

IL-8 - r: 5'-CTCCACAACCCTCTGCAC-3';

CYP1A1 - f: sense 5'-GGCAGATCAACCATGACCAGAAG-3',

CYP1A1 - r: 5'-ACAGCAGGCATGCTTCATGGTTAG-3'.

Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and hypoxanthine-guanine phosphoribosyltransferase (HPRT1) gene expression were used as the internal control as they are housekeeping genes and were analysed in each experiment for normalisation using the following primers:

GAPDH - f: 5'-CCCTTCATTGACCTCAACTACATG-3',

GAPDH - r: 5'-TGGGATTTCCATTGATGACAAGC-3';

HPRT1- f: 5'-TGACACTGGCAAAACAATGCA-3',

HPRT1 - r: 5'-GGTCCTTTTCACCAGCAAGCT-3'.

Relative fold inductions were calculated using the Δ Ct formula (Scheffe et al., 2006). QC laboratory filter extract was tested in the same manner of PM extracts. All real-time RT-PCR assays for

relative gene expression were repeated at least three times in duplicates from independent total RNA samples for the same treatment conditions. non-exposed cells and solvent exposed cells were used as control cells for aqueous and organic extracts respectively.

2.7 Chemicals and meteo data

Chemical data (metals: As, Cd, Ni, Pb and PAHs: Benzo(a)anthracene, Benzo(b+j+k)fluorantene, Benzo(a)pyrene, Indeno(1,2,3-cd)pyrene) were extracted from a specialised database provided by the Regional System for the real-time monitoring of Air Quality, AriaWeb (ARPA Piemonte, 2015). The data were obtained for the same meteorological–chemical station and for the same days as our sampling. For example the metals data represent a mean of daily data collected using the standard monitoring method EN 14902:2005 «Standard method for measurement of Pb/Cd/As/Ni in the PM₁₀ fraction of suspended particulate matter» (2008/50/EC, annex VI, section B). All the adopted methods conform to the directive and were validated before being published in the AriaWeb database (ARPA Piemonte, 2015).

2.8 Statistical analysis

Statistical analyses were performed using the SPSS Package, version 21.0 for Windows. The Spearman rank order correlation coefficient was used to assess relationships between variables, a Wilcoxon test was used to compare means, and ANOVA was used for multivariate analysis in which we assumed an equal variance while using Tukey as post hoc multiple comparisons. The mean differences and correlations were considered significant at $p < 0.05$.

3. RESULTS

3.1 PM concentration and meteo-chemicals data

The mean PM₁₀ concentration was $23.9 \pm 8.4 \mu\text{g}/\text{m}^3$ in summer and $83.0 \pm 36.2 \mu\text{g}/\text{m}^3$ in winter, and this difference was statistically significant (Wilcoxon test, $p < 0.01$) (Figure 1).

The winter fractions concentrations compared with the corresponding summer ones showed significant differences (Wilcoxon test, $p < 0.01$).

ANOVA analysis performed assuming the gravimetric results as a dependent variable and the different PM₁₀ size fractions as the independent variable reported significant general results

(summer: $F = 15.091$, $p < 0.001$, $d=5$; and winter: $F = 4.904$, $p < 0.01$, $d=5$): the finest fraction $PM_{<0.49}$ was always significantly higher than the others in summer (Tukey test, $p < 0.01$) and represented about 37% of the total PM_{10} mass, while in winter was significantly higher only vs $PM_{10.0-7.2}$ and $PM_{3.0-1.5}$ (Tukey test, $p < 0.01$) and represented about 28% of the total PM_{10} mass; Figure 1 shows also the percentage size composition of PM_{10} .

Table 1 reports the descriptive analysis of the chemical and meteo-climatic parameters collected during the summer and winter samplings.

The mean temperature difference between sampling seasons was significant (Wilcoxon test, $p < 0.01$): the mean winter temperature was $1.2 \pm 4.2^{\circ}C$ and the mean summer temperature was $23.0 \pm 1.8^{\circ}C$. Also radiation was significantly different (Wilcoxon test, $p < 0.01$) in the two considered periods: $23 \pm 3 MJ/m^2$ in summer and $11 \pm 3 MJ/m^2$ in winter.

Neither the average humidity nor the wind speed were significantly different (Wilcoxon test, $p > 0.05$), with 82% humidity during the winter vs. 63% summer and a range wind speed in both seasons of 1.0-2.0 m/s.

ANOVA analysis performed assuming the gravimetric results as a dependent variable and the three main size fractions (analyzed two by two: PM_{10-3} : 10 - 3 μm ; $PM_{3-0.95}$: 3 - 0.95 μm ; and $PM_{<0.95}$: 0.95 - < 0.49 μm) as the independent variable reported a significant result in which the finest fraction ($PM_{<0.95}$) was significantly higher than the others both in summer and in winter (summer: $F = 11.170$, $p < 0.01$, $d=2$; and winter: $F = 5.790$, $p < 0.05$, $d=2$). Beyond this $PM_{<0.95}$ represented about 53% of PM_{10} mass in this urban site both in summer and in winter.

Referring to the annual limits of metals set by the European Directive 2008/50/EC, these are observed even if our sampling reflects only spot seasonal situations (3 weeks in each season) and the European limits refer to yearly averages. Cd and Pb winter concentrations showed significant differences (Wilcoxon test, $p < 0.05$) if compared with the corresponding summer concentrations.

Moreover both total PAHs and single PAHs concentrations were higher during the winter than summer (Wilcoxon test, $p < 0.05$).

3.2 Cell proliferation

279 Figure 2a shows effects of the different extracts on cell proliferation: inhibition of THP-1 cell
280 proliferation observed *in vitro* was significant respect to control cells for all PM extracts although
281 the observed effect was independent of the aerodynamic diameter of the particles and of kind of
282 PM extract. Both aqueous and organic extracts inhibited cell proliferation with similar effects
283 (Wilcoxon test, $p > 0.05$). The mean percentage proliferation inhibition was 30.8 ± 6.0 % for
284 summer and 28.4 ± 6.9 % for winter fraction extracts. The difference between summer and winter
285 samples was not significant both for organic and aqueous extracts (Wilcoxon test, $p > 0.05$).

286 **3.3 LDH release**

287 LDH assay indicates loss in cell membrane integrity and it was used for cytotoxicity determination.
288 In general, the summer PM extracts induced a more significant release of LDH compared to winter
289 (54.3 ± 36.5 % and 22.6 ± 17.7 % respectively; Wilcoxon test, $p < 0.001$). Significant differences in
290 toxicity between the three different size fraction extracts were observed: summer organic extracts
291 produced a marked size-related effect, with higher values measured with $PM_{10-3} > PM_{3-0.95} >$
292 $PM_{<0.95}$; considering the winter extracts, LDH release over the negative control was lower than
293 summer ones, except for the finest winter fraction $PM_{<0.95}$ of aqueous extract (Figure 2b).

294 **3.4 IL-8, $TNF\alpha$ and CYP1A1 gene expression**

295 The production of the pro-inflammatory cytokine $TNF\alpha$, as well as the chemokine IL-8 were
296 determined in the cell culture supernatant after 48h of size-fractionated PM extracts exposure;
297 these signaling molecules constitute a major parameter in the mediation of airway inflammation
298 (Gualtieri et al., 2010). Exposure of THP-1 cells to size-fractionated PM_{10} extracts did not induce
299 significant expression of $TNF\alpha$, except for the finest winter fraction both organic and aqueous
300 samples (Figure 3a). On the other side, particles extracts had a stronger effect on IL-8 expression
301 especially organic extracts in summer. Statistically significant differences were found between the
302 two kind of extracts for both season (Wilcoxon test, $p < 0.05$) and comparing summer (4.13 ± 1.34
303 fold increase) and winter (2.78 ± 0.96 fold increase) extracts (Wilcoxon test, $p < 0.001$) (Figure 3b).
304 No significant differences in toxicity between the extracts of the three different size fractions were
305 observed.

306 The mean fold increase in CYP1A1 expression for summer extracts was 3.61 ± 1.18 and for winter
307 was 3.01 ± 1.46 , and this difference was significant (Figure 3c). The average effect was higher in
308 summer with the exception of the organic finest fraction extracts.

309 In general in winter the effect was higher for organic extract, indeed there were statistically
310 significant differences in the effects induced by the two type of extracts (Wilcoxon test, $p < 0.05$).

311 PM extracts produced a significant marked size-related CYP1A1 expression in winter; ANOVA
312 analysis performed assuming the CYP1A1 expression as a dependent variable and the different
313 size fractionated extracts as the independent variable showed a significant difference in the effects
314 produced by winter extracts ($F = 8.756$, $p < 0.01$, $d=2$ organic; $F = 6.291$, $p < 0.01$ $d=2$ aqueous) with
315 higher values measured with $PM_{<0.95} > PM_{3-0.95} > PM_{10-3}$.

316

317 **4. DISCUSSION and CONCLUSIONS**

318 In our study size-fractionated PM_{10} was collected in a polluted urban site in Torino (northwest
319 Italian city) during different seasons. In the north of Italy, in particular in Padana Plain, there is a
320 widespread air pollution; the weak dispersion rate, observed during winter, represents a relevant
321 factor and it is principally due to the conformation of the territory (Cadum et al., 2009). Torino PM
322 concentrations are characterized by high levels of fine particles in winter: during high pollution
323 periods the $PM_{2.5}/PM_{10}$ ratio increased up to 0.87; in this area the winter PM chemical composition
324 is characterised by high levels of PAHs and metals (Romanazzi et al., 2014; Traversi et al., 2015)
325 whereas in summer PM is rich of ions such sulphates (Schilirò et al., 2015). These seasonal
326 differences could be explained as the winter contribution of domestic heating and of a lower
327 atmospheric mixing layer also the low temperatures facilitate the condensation and absorption of
328 volatile compounds on particle surfaces (Brüggemann et al., 2009); during summer, the
329 photochemical reactions, associated with elevated solar radiation, modify the PM_{10} chemical
330 constituents.

331 In the present study the observed PM_{10} winter levels are higher than both the WHO guidelines
332 (Krzyzanowski, 2008) and the EU regulations 2008/50/CE. In addition, critical particle
333 concentrations are present especially for the finest fraction $PM < 0.49 \mu m$ that represented a very

high proportion of PM₁₀: 37% in summer and 28% in winter. Various studies confirmed that the finest fractions are the most abundant in the atmosphere because of the finest particulate pollution is homogeneously diffused (Perez et al., 2010).

The high temperature and radiation observed in summer respect to winter attested more frequent photochemical reactions in summer that promotes the formation of secondary particles as highlighted by the percentage of fine fraction (PM_{<0.49}) in summer. In general the percentage of secondary particles increases with decreasing particle size, this is in agreement with the literature data according to which the components of the inorganic fraction of the secondary particles (ammonium, nitrate and sulphate) are present mainly in the finer fraction (Pateraki et al., 2012; Perrino et al., 2014)

Among the metals monitored in this study only Cd and Pb had a seasonal trend, As and Ni had similar concentrations in winter and summer; in a previous work in this area, metals appeared to be variously distributed among all the fractions, especially in winter season: As, Ni and Pb were preferentially accumulated on the coarse fractions while Cd was equally distributed (Romanazzi et al., 2014). Contrariwise PAHs showed a marked seasonal trend and the mean winter benzo(a)pyrene concentration was higher both than the annual limit set by the European Directive 2008/50/EC (1 ng/m³) and the WHO guide line value of 0.12 ng/m³ (Krzyzanowski, 2008; WHO-Europe, 2013) even if our sampling, as for metals, reflects only a spot seasonal situation. A higher PAHs contamination was observed in the winter and this is probably due to the winter atmospheric conditions that may promote an accumulation of primary pollutants and because of low temperatures, the condensation of atmospheric pollutants in the particle phase (Ebi and McGregor, 2008); this seasonal trend was confirmed in other studies (Sisovic et al., 2008).

Size-fractionated PM₁₀ extracts (organic and aqueous) exerted a significant impact on cells both in winter and in summer. The presented results indicated that different PM₁₀ fraction extracts lead to clear changes in some toxicological responses.

Inhibition of proliferation showed a trend without significant differences among fraction extracts as found in other researches (Longhin et al., 2013; Wessels et al., 2010), however it was significant both in summer and in winter. The particles size influenced the cytotoxic (LDH release) and pro-

inflammatory potential (TNF α concentrations). Organic PM extracts induced a significant size-related LDH release, with greater release by the greater fractions. LDH release was significantly greater in summer than in winter with the exception of aqueous extracts of the finest fraction in which transition metals are mainly accumulated (Daher et al., 2014). Both organic and aqueous winter extracts of the finer fraction induced a significant TNF α expression that may be associated to the presence of PAHs and transition metals respectively in the finer fractions (de Kock et al., 2006).

Summer fraction extracts were able to trigger a pro-inflammatory response with the release of significant quantity of IL-8, especially the summer greater fractions. Actually various *in vitro* data showed the higher inflammatory potential of PM extracts of coarse fraction (Hetland et al., 2004; Jalava et al., 2007). Inflammatory effect of summer PM extracts could be related to the presence of biogenic compounds (Gualtieri et al., 2010; Schins et al., 2004); the role of endotoxins in leading the PM inflammatory effects has been well documented (Hetland et al., 2004; Traversi et al., 2010). Inflammatory effect of aqueous extracts of PM₁₀ fractions could also be related to the particles bound metals; the PM chemical composition describes the presence of crustal elements, ions and metals, variously distributed among all the fractions (Romanazzi et al., 2014), which have been showed to induce cell inflammation (Øvrevik et al., 2005; Veranth et al., 2006).

Apparently Torino summer PM₁₀ fraction extracts in a traffic site had a higher proinflammatory potential and induces stronger IL-8 responses in confront of a remote site (Schilirò et al., 2015), similar results have been presented by numerous authors (Camatini et al., 2012; Duvall et al., 2008; Gualtieri et al., 2010; Hetland et al., 2004).

PM extracts induced a significant size-related CYP1A1 expression, in particular for organic winter extracts. The importance of organic compounds, with regard to CYP1A1 expression, was confirmed as all the winter organic extracts induced significant size-related levels of this biological marker, with greater increase by the finer fraction extracts. This can easily be associated to the link between the majority of particle-bound PAHs and the fine and ultra-fine particles (Claxton et al., 2004; Di Filippo et al., 2010a). In this study, CYP1A1 expression was significant also for the summer extracts even if PAHs levels were lower highlighting the presence of other chemical

390 compounds capable to induce the CYP1A1 expression (e.g. dioxin like molecules) (Wenger et al.,
391 2009).

392 Torino PM₁₀ winter fractions in a traffic site showed a higher CYP1A1 expression compared to a
393 background urban site (Schilirò et al., 2015) especially the finer fractions extracts.

394 No statistically significant association were found between the THP-1 endpoints and chemical
395 (metals or PAHs) or meteo-climatic parameters collected during the summer and winter samplings.
396 One possible explanation for the lack of correlations could be that the chemical parameters were
397 quantified in total PM₁₀ samples instead of separated fractions.

398 Comparing results obtained in this work considering a trafficated Torino urban site with those
399 reported in a previous study conducted in a background Torino urban site in the same seasons
400 (Schilirò et al., 2015), the total PM₁₀ concentration was higher in the trafficated urban site in winter
401 but showed similar values respect to background site in summer. Moreover, the biological effects
402 were higher in the trafficated site mostly considering inhibition of cell proliferation and CYP1A1
403 expression in winter and LDH release and IL-8 expression in summer.

404 Our results show that biological responses significantly differ after exposure to equal mass
405 concentrations of urban traffic and background Torino urban site. These observations confirm the
406 hypothesis that particle composition, as well as source, constitute an important factor in PM
407 induced toxicity as underlined in other studies (Michael et al., 2013).

408 It is now widely accepted that air pollution and its major components have a marked seasonality,
409 and the toxic content in particulates can vary based upon the meteo-climatic conditions (Albinet et
410 al., 2008). However, biological effects underlined the potential importance of the non-regulated
411 pollutants to assess the air quality (Di Filippo et al., 2010a). This complexity is the reason for
412 difficulties in solving the question of PM toxicity.

413 In addition, our results confirm that the only measure of PM can be misleading for the assessment
414 of air quality (Di Filippo et al., 2010b) in fact PM₁₀ or PM_{2.5} monitoring itself is not informative
415 enough on toxic compounds bound to particles (Topinka et al., 2015). Evaluation of air pollution
416 should not be limited just to gravimetric assessment of PM but should consider also the size
417 spectrum of more dangerous PM fractions. Because of their large and irregular specific surface,

418 the finest fractions can favour the adsorption of mutagenic and carcinogenic species (such as
419 PAHs, nitro-PAHs and metals) (Claxton et al., 2004; Topinka et al., 2015), moreover they can
420 reach the deepest sites of the respiratory system (alveolar lung region). The use of different types
421 of extract allows to evaluate the role of chemical composition on biological effects of different PM
422 fraction extracts.

423 Moreover, the use of *in vitro* test, that integrate toxic potencies of samples including unidentified as
424 well as identified chemicals, can give a correct estimation of the impact of a complex mixture.

425 Finally in this study we support efforts toward establishing more effective and source-specific
426 regulations for mitigating PM toxicity (Daher et al., 2014) and identifying potential effect-based
427 tools (e.g. *in vitro* test on specific cell line such as THP-1 or A549) that could be used in the context
428 of the different monitoring programmes after being properly standardized. These kind of effect-
429 based monitoring tools could also be valuable in toxicity identification evaluation (TIE) approaches
430 to identify toxic fractions and provide guidance for the identification of causative agents.

431

432 **ACKNOWLEDGMENTS**

433 The authors kindly thank Drs M. Sacco, F. Lollobrigida and M. Grosa of the Environmental
434 Protection Regional Agency (Piedmont A.R.P.A.). This study was financed by Torino University
435 Local Research grant (ex-60%-2012).

436

437 **REFERENCES**

438 Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., Jaffrezo, J.L., 2008. Nitrated and
439 oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French
440 alpine valleys Part 1: concentrations, sources and gas/particle partitioning. Atmos. Environ. 42,
441 43-54.

442 Alessandria, L., Schilirò, T., Degan, R., Traversi, D., Gilli, G., 2014. Cytotoxic response in human
443 lung epithelial cells and ion characteristics of urban-air particles from Torino, a northern Italian
444 city. Environ. Sci. Pollut. Res. Int. 21, 5554-5564.

Anderson, J.O., Thundiyil, J.G., Stolbach, A., 2012. Clearing the air: review of the effects of particulate matter air pollution on human health. *J. Med. Toxicol.* 8, 166–175.

ARPA Piemonte, S.P., CSI Piemonte, 2015, ARIAWEB – Sistema Regionale Di Rilevamento Della Qualità Dell'aria Della Regione Piemonte.

Bonetta, S., Gianotti, V., Bonetta, S., Gosetti, F., Oddone, M., Gennaro, M.C., Carraro, E., 2009. DNA damage in A549 cells exposed to different extracts of PM_(2.5) from industrial, urban and highway sites. *Chemosphere* 77, 1030-1034.

Brook, R.D., Rajagopalan, S., Pope, C.A., Brook, J.R., Bhatnagar, A., Diez-Roux, A.V., Holguin Hong, Y.L., Luepker, R.V., Mittleman, M.A., Peters, A., Siscovick, D., Smith, S.C., Whitsel, L., Kaufman, J.D., Amer Heart Assoc Council E, Council Kidney Cardiovasc D, Council Nutr Phys Activity M., 2010. Particulate Matter Air Pollution and Cardiovascular Disease An Update to the Scientific Statement From the American Heart Association. *Circulation* 121, 2331-2378.

Brüggemann, E., Gerwig, H., Gnauk, T., Muller, K., Herrmann, H., 2009. Influence of seasons, air mass origin and day of the week on size-segregated chemical composition of aerosol particles at a kerbside. *Atmos. Environ.* 43, 2456-2463.

Cadum, E., Berti, G., Biggeri, A., Bisanti, L., Faustini, A., Forastiere, F., Grp Collaborativo EpiAir, 2009. The results of EpiAir and the national and international literature. *Epidemiol. Prev.* 33, 113-119.

Camatini, M., Corvaja, V., Pezzolato, E., Mantecca, P., Gualtieri, M., 2012. PM₁₀-biogenic fraction drives the seasonal variation of proinflammatory response in A549 cells. *Environ. Toxicol.* 27, 63-73.

CEN, 1998. Air quality - determination of the PM₁₀ fraction of suspended particulate matter – reference method and field test procedure to demonstrate reference equivalence of measurement methods. Brussels: European Committee for Standardization; (European Standard EN 12341).

Claxton, L.D., Matthews, P.P., Warren, S.H., 2004. The genotoxicity of ambient outdoor air, a review: Salmonella mutagenicity. *Mutat. Res.* 567, 347-399.

472 Corsini, E., Budello, S., Marabini, L., Galbiati, V., Piazzalunga, A., Barbieri, P., Cozzutto, S.,
 473 Marinovich, M., Pitea, D., Galli, C.L., 2013. Comparison of wood smoke PM_{2.5} obtained from the
 474 combustion of FIR and beech pellets on inflammation and DNA damage in A549 and THP-1
 475 human cell lines. *Arch. Toxicol.* 87, 2187-2199.

476 Daher, N., Saliba, N.A., Shihadeh, A.L., Jaafar, M., Baalbaki, R., Shafer, M.M., Schauer, J.J.,
 477 Sioutas, C., 2014. Oxidative potential and chemical speciation of size-resolved particulate
 478 matter (PM) at near-freeway and urban background sites in the greater Beirut area. *Sci.Total*
 479 *Environ.* 470-471, 417-426.

480 Danielsen, P.H., Møller, P., Jensen, K.A., Sharma, A.K., Wallin, H., Bossi, R., Autrup, H., Mølhave,
 481 L., Ravanat, J.L., Briedé, J.J., de Kok, T.M., Loft, S., 2011. Oxidative stress, DNA damage, and
 482 inflammation induced by ambient air and wood smoke particulate matter in human A549 and
 483 THP-1 cell lines. *Chem. Res. Toxicol.* 24,168-184.

484 de Kock, T.M., Driee, H.A.L., Hogervorst, J.G.F., Briede, J.J., 2006. Toxicological assessment of
 485 ambient and traffic-related ambient particulate matter: a review of recent studies. *Mutat. Res.*
 486 613, 103-122.

487 Di Filippo, P., Riccardi, C., Pomata, D., Buiarelli, F., 2010a. Concentrations of PAHs, and nitro- and
 488 methyl- derivatives associated with a size-segregated urban aerosol. *Atmos. Environ.* 44, 2742-
 489 2749.

490 Di Filippo, P., Riccardi, C., Pomata, D., Gariazzo, C., Buiarelli, F., 2010b. Seasonal Abundance of
 491 Particle-Phase Organic Pollutants in an Urban/Industrial Atmosphere. *Water Air Soil Pollut.*
 492 211, 231–250.

493 Duvall, R.M., Norris, G.A., Dailey, L.A., Burke, J.M., McGee, J.K., Gilmour, M.I., Gordon, T., Devlin,
 494 R.B., 2008. Source apportionment of particulate matter in the U.S. and associations with lung
 495 inflammatory markers. *Inhal. Toxicol.* 20, 671-683.

496 Ebi, K.L., McGregor, G., 2008. Climate change, tropospheric ozone and particulate matter, and
 497 health impacts. *Environ. Health Perspect.* 116, 1449-1455.

498 Gualtieri, M., Øvrevik, J., Holme, J.A., Perrone, M.G., Bolzacchini, E., Schwarze, P.E., Camatini,
 499 M., 2010. Differences in cytotoxicity versus pro-inflammatory potency of different PM fractions
 500 in human epithelial lung cells. *Toxicol. In Vitro* 24, 29-39.

501 Hanzalova, K., Rossner, Jr P., Sram, R.J., 2010. Oxidative damage induced by carcinogenic
 502 polycyclic aromatic hydrocarbons and organic extracts from urban air particulate matter. *Mutat.*
 503 *Res.* 696, 114–121.

504 Hetland, R.B., Cassee, F.R., Refsnes, M., Schwarze, P.E., Lag, M., Boere, A.J., Dybing, E., 2004.
 505 Release of inflammatory cytokines, cell toxicity and apoptosis in epithelial lung cells after
 506 exposure to ambient air particles of different size fractions. *Toxicol. In Vitro* 18, 203-212.

507 Jalava, P.I., Salonen, R.O., Pennanen, A.S., Sillanpää, M., Hälinen, A.I., Happonen, M.S., Hillamo, R.,
 508 Brunekreef, B., Katsouyanni, K., Sunyer, J., Hirvonen, M.R., 2007. Heterogeneities in
 509 inflammatory and cytotoxic responses of RAW 264.7 macrophage cell line to urban air coarse,
 510 fine, and ultrafine particles from six European sampling campaigns. *Inhal. Toxicol.* 19, 213-225.

511 Knaapen, A.M., Borm, P.J., Albrecht, C., Schins, R.P., 2004. Inhaled particles and lung cancer.
 512 Part A: Mechanisms. *Int. J. Cancer* 109, 799-809.

513 Kroll, A., Gietl, J.K., Wiesmüller, G.A., Günsel, A., Wohlleben, W., Schnekenburger, J., Klemm, O.,
 514 2013. In vitro toxicology of ambient particulate matter: correlation of cellular effects with particle
 515 size and components. *Environ. Toxicol.* 28, 76-86.

516 Krzyzanowski, M., 2008. WHO air quality guidelines for Europe. *J. Toxicol. Environ. Health A* 71,
 517 47–50.

518 Lauer, F.T., Mitchell, L.A., Bedrick, E., McDonald, J.D., Lee, W.Y., Li, W.W., Olvera, H., Amaya,
 519 M.A., Berwick, M., Gonzales, M., Currey, R., Pingitore, N.E., Jr., Burchiel, S.W., 2009.
 520 Temporal spatial analysis of U.S.-Mexico border environmental fine and coarse PM air sample
 521 extract activity in human bronchial epithelial cells. *Toxicol. Appl. Pharmacol.* 238, 1–10.

522 Longhin, E., Pezzolato, E., Mantecca, P., Holme, J.A., Franzetti, A., Camatini, M., Gualtieri, M.,
 523 2013. Season linked responses to fine and quasi-ultrafine Milan PM in cultured cells. *Toxicol. In*
 524 *Vitro* 27, 551-559.

525 Lonkar, P., Dedon, P.C., 2011. Reactive species and DNA damage in chronic inflammation:
 526 reconciling chemical mechanisms and biological fates. *Int. J. Cancer* 128, 1999-2009.

527 Loomis, D., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L.,
 528 Guha, N., Baan, R., Mattock, H., Straif, K., IARC working group, 2013. The carcinogenicity of
 529 outdoor air pollution. *Lancet Oncol.* 14, 1262–1263.

530 Michael, S., Montag, M., Dott, W., 2013. Pro-inflammatory effects and oxidative stress in lung
 531 macrophages and epithelial cells induced by ambient particulate matter. *Environ. Pollut.* 183,
 532 19-29.

533 Oh, S.M., Kim, H.R., Park, Y.J., Lee, S.Y., Chung, K.H., 2011. Organic extracts of urban air
 534 pollution particulate matter (PM_{2.5})-induced genotoxicity and oxidative stress in human lung
 535 bronchial epithelial cells (BEAS-2B cells). *Mutat. Res.* 723, 142-151.

536 Øvrevik, J., Myran, T., Refnes, M., Lag, M., Becher, R., Hetland, R.B., 2005. Mineral particles of
 537 varying composition induce differential chemokine release from epithelial lung cells: importance
 538 of physicochemical characteristics. *Ann. Occup. Hyg.* 49, 219-231.

539 Pateraki, S., Assimakopoulos, V.D., Bougiatioti, A., Kouvarakis, G., Mihalopoulos, N., Vasilakos,
 540 Ch., 2012. Carbonaceous and ionic compositional patterns of fine particles over an urban
 541 Mediterranean area. *Sci. Total Environ.* 424, 251-263.

542 Perez, N., Pey, J., Cusack, M., Reche, C., Querol, X., Alastuey, A., Viana, M., 2010. Variability of
 543 Particle Number, Black Carbon, and PM₁₀, PM_{2.5}, and PM₁ Levels and Speciation: Influence of
 544 Road Traffic Emissions on Urban Air Quality. *Aerosol Sci. Tech.* 44, 487-499.

545 Perrino, C., Catrambone, M., Dalla Torre, S., Rantica, E., Sargolini, T., Canepari, S., 2014.
 546 Seasonal variations in the chemical composition of particulate matter: a case study in the Po
 547 Valley. Part I: macro-components and mass closure. *Environ. Sci. Pollut. Res. Int.* 21, 3999-
 548 4009.

549 Roig, N., Sierra, J., Rovira, J., Schuhmacher, M., Domingo, J.L., Nadal, M., 2013. In vitro tests to
 550 assess toxic effects of airborne PM₍₁₀₎ samples. Correlation with metals and chlorinated dioxins
 551 and furans. *Sci. Total Environ.* 443, 791-797.

552 Romanazzi, V., Casazza, M., Malandrino, M., Maurino, V., Piano, A., Schilirò, T., Gilli, G., 2014.
 553 PM₁₀ size distribution of metals and environmental-sanitary risk analysis in the city of Torino.
 554 *Chemosphere* 112, 210-216.

555 Schefe, J.H., Lehmann, K.E., Buschmann, I.R., Unger, T., Funke-Kaiser, H., 2006. Quantitative
 556 real-time RT-PCR data analysis: current concepts and the novel "gene expression's C-T
 557 difference" formula. *J. Mol. Med.* 84, 901-910.

558 Schilirò, T., Alessandria, L., Degan, R., Traversi, D., Gilli, G., 2010. Chemical characterisation and
 559 cytotoxic effects in A549 cells of urban-air PM₁₀ collected in Torino, Italy. *Environ. Toxicol.*
 560 *Pharmacol.* 29, 150-157.

561 Schilirò, T., Bonetta, S., Alessandria, L., Gianotti, V., Carraro, E., Gilli, G., 2015. PM₁₀ in a
 562 background urban site: chemical characteristics and biological effects. *Environ. Toxicol.*
 563 *Pharmacol.* 39, 833-844.

564 Schins, R.P., Lightbody, J.H., Borm, P.J., Shi, T., Donaldson, K., Stone, V., 2004. Inflammatory
 565 effects of coarse and fine particulate matter in relation to chemical and biological constituents.
 566 *Toxicol. Appl. Pharmacol.* 195, 1-11.

567 Schins, R.P.F., Hei, T.K., 2006. Genotoxic effects of particles, in: Donaldson, K., Borm, P., (Eds.),
 568 *Particle Toxicology*. CRC Press/Taylor and Francis Group, Boca Raton, FL, pp 285-298.

569 Schwarze, P.E., Ovreivik, J., Hetland, R.B., Becher, R., Cassee, F.R., Lag, M., Lovik, M., Dybing,
 570 E., Refsnes, M., 2007. Importance of size and composition of particles for effects on cells in
 571 vitro. *Inhal. Toxicol.* 19, 17-22.

572 Sisovic, A., Beslic, I., Segar, K., Vadjic, V., 2008. PAH mass concentrations measured in PM₁₀
 573 particle fraction. *Environ. Int.* 34, 580-584.

574 Spink, D.C., Wu, S.J., Spink, B.C., Hussain, M.M., Vakharia, D.D., Pentecost, B.T., Kaminsky,
 575 L.S., 2008. Induction of CYP1A1 and CYP1B1 by benzo(k)fluoranthene and benzo(a)pyrene in

576 T-47D human breast cancer cells: roles of PAH interactions and PAH metabolites. *Toxicol.*
577 *Appl. Pharmacol.* 226, 213-24.

578 Steenhof, M., Gosens, I., Strak, M., Godri, K.J., Hoek, G., Cassee, F.R., Mudway, I.S., Kelly, F.J.,
579 Harrison, R.M., Lebret, E., Brunekreef, B., Janssen, N.A.H., Pieters, R.H.H., 2011. In vitro
580 toxicity of particulate matter (PM) collected at different sites in the Netherlands is associated
581 with PM composition, size fraction and oxidative potential e the RAPTES project. *Part. Fibre*
582 *Toxicol.* 8, 26.

583 Traversi, D., Alessandria, L., Schilirò, T., Chiadò Piat, S., Gilli, G., 2010. Meteo-climatic conditions
584 influence the contribution of endotoxins to PM₁₀ in an urban polluted environment. *J. Environ.*
585 *Monit.* 12, 484-490.

586 Traversi, D., Alessandria, L., Schilirò, T., Gilli, G., 2011. Size-fractionated PM₁₀ monitoring in
587 relation to the contribution of endotoxins in different polluted areas. *Atmos. Environ.* 45, 3515-
588 3521.

589 Traversi, D., Festa, E., Pignata, C., Gilli, G., 2015. Aero-dispersed mutagenicity attributed to
590 particulate and semi volatile phase in an urban environment. *Chemosphere.* 124, 163-169.

591 Valavanidis, A., Fiotakis, K., Vlachogianni, T., 2008. Airborne particulate matter and human health:
592 toxicological assessment and importance of size and composition of particles for oxidative
593 damage and carcinogenic mechanisms. *J. Environ. Sci. Health C* 26, 339-362.

594 Veranth, J.M., Moss, T.A., Chow, J.C., Labban, R., Nichols, W.K., Walton, J.C., Watson, J.G.,
595 Yost, G.S., 2006. Correlation of in vitro cytokine responses with the chemical composition of
596 soil-derived particulate matter. *Environ. Health Perspect.* 114, 314-349.

597 Wenger, D., Gerecke, A.C., Heeb, N.V., Hueglin, C., Seiler, C., Haag, R., Naegeli, H., Zenobi, R.,
598 2009. Aryl hydrocarbon receptor-mediated activity of atmospheric particulate matter from an
599 urban and a rural site in Switzerland. *Atmos. Environ.* 43, 3556-3562.

600 Wessels, A., Birmili, W., Albrecht, C., Hellack, B., Jermann, E., Wick, G., Harrison, R.M., Schins,
601 R.P.F., 2010. Oxidant Generation and Toxicity of Size-Fractionated Ambient Particles in
602 Human Lung Epithelial Cells. *Environ. Sci. Technol.* 44, 3539-3545.

603 WHO-Europe, 2013. Review of Evidence On Health Aspects Of Air Pollution – Technical Report –
604 REVIHAAP Project.

1 **Table 1.** Descriptive analysis for chemical and meteo-climatic parameters in summer and winter
2 samplings (means with standard deviations, minimum and maximum values) in the urban
3 meteorological–chemical station. nd: not detected

Parameters	Mean (sd)		Min		Max	
	summer	winter	summer	winter	summer	winter
Temperature (°C)	23.0 (1.8)	1.2 (4.2)	13.7	- 5.5	32.3	16.7
Wind (m/s)	1.7 (0.3)	1.6 (0.4)	1.1	1.0	2.0	2.0
Precipitations (mm)	0.2 (sum)	0.0 (sum)	0.0	0.0	0.2	0.0
Humidity (%)	63 (20)	82 (21)	16	27	96	100
Radiation (MJ/m ²)	23 (3)	11 (3)	19	6	28	12
PM 10-3 µm (µg/m ³)	6.6 (1.0)	18.0 (2.3)	4.3	10.1	10.8	23.9
PM 3 – 0.95 µm (µg/m ³)	4.5 (0.6)	20.9 (5.2)	2.9	9.2	6.6	45.0
PM < 0.95 µm (µg/m ³)	12.7 (1.9)	44.1 (8.7)	6.8	24.1	19.5	85.2
As (ng/m ³)	0.7 (0.3)	0.7 (0.3)	nd	nd	0.8	0.7
Cd (ng/m ³)	0.2 (0.1)	0.5 (0.2)	0.1	nd	0.2	0.5
Ni (ng/m ³)	5.9 (2.5)	6.5 (2.6)	nd	nd	6.3	7.6
Pb (ng/m ³)	6.2 (2.5)	15.5 (1.2)	nd	13	7.0	16
Benzo(a)anthracene	0.04 (0.01)	2.08 (0.93)	nd	nd	0.04	3.16
Benzo(b+j+k)fluorantene	0.20 (0.08)	4.04 (1.81)	nd	nd	0.25	6.38
Benzo(a)pyrene	0.08 (0.05)	2.20 (0.98)	nd	nd	0.10	3.30
Indeno(1,2,3-cd)pyrene	0.06 (0.02)	2.18 (0.97)	nd	nd	0.06	3.12

4

5

FIGURE CAPTIONS

Figure 1. PM₁₀ fractions concentration and percentage size composition in winter (a) and in summer (b), sampled in a polluted urban meteorological–chemical station.

Figure 2. Inhibition of proliferation (a) and LDH release (b) of THP-1 cells exposed (48h, 200µg/mL) to organic (black bars) and aqueous (grey bars) PM fraction extracts of summer and winter season. Bars represent the mean value from three replicates; error bars represent standard error of mean. *p< 0.05 vs control (control level is at 0%) according to Wilcoxon test.

Figure 3. TNF α (a), IL-8 (b) and CYP1A1 (c) expressions of THP-1 cells exposed (48h, 200µg/mL) to organic (black bars) and aqueous (grey bars) PM fraction extracts of summer and winter season. Bars represent the mean value from three analysis in duplicate; error bars represent standard error of mean. *p< 0.05 vs control cells (control level is at 1 fold increase) according to Wilcoxon test.

Figure

[Click here to download high resolution image](#)

Figure
[Click here to download high resolution image](#)

Figure
[Click here to download high resolution image](#)