AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Ten years of subproteome investigations in lactic acid bacteria: A key for food starter and probiotic typing

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1557624> since 2016-03-14T11:05:11Z

Published version:

DOI:10.1016/j.jprot.2015.04.028

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Journal of Proteomics, Volume 127, Part B, 2015, DOI: 10.1016/j.jprot.2015.04.028]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[\[http://www.sciencedirect.com/science/article/pii/S1874391915002092\]](http://www.sciencedirect.com/science/article/pii/S1874391915002092)

Abstract

The definition of safety and efficacy of food-employed bacteria as well as probiotic strains is a continuous, often unattended, challenge. Proteomic techniques such as 2DE, DIGE and LC/LC-MS/MS are suitable and powerful tools to reveal new aspects (positive and negative) of “known” and “unknown” strains that can be employed in food making and as nutraceutical supplements for human health. Unfortunately, these techniques are not used as extensively as it should be wise. The present report describes the most significant results obtained by our research group in 10 years of study on subproteomes in bacteria, chiefly lactic acid bacteria. Production of desired and undesired metabolites, differences between strains belonging to same species but isolated from different ecological niches, the effect of cryoprotectants on survival to lyophilization as well as the adhesive capability of strains, were elucidated by analysis of cytosolic, membrane-enriched, surface and extracellular proteomes. The present review opens a window on a yet largely underexplored field and highlights the huge potential of subproteome investigations for

more rational choice of microbial strains as food starters, probiotics and for production of nutraceuticals. These analyses will hopefully contribute to manufacturing safer and healthier food and food supplements in the near future.

This article is part of a Special Issue entitled: HUPO 2014.

Graphical abstract

[Figure options](#)

Abbreviations

- EF, elongation factor;
- GAPDH, glyceraldehyde 3-phosphate dehydrogenase;
- PGK, phosphoglycerate kinase

Keywords

- 2DE;
- Subproteomes;
- Moonlighting proteins;
- Food bacteria;
- Adhesion;
- Biogenic amines

1. Introduction

Safety of bacterial starters employed in food fermentations as well as efficacy of probiotic strains used as food additives or nutraceutical supplements is an urgent concern. In spite of the improved analytical controls, food-borne syndromes other than infections are still a reality in western countries [1], [2] and [3]. Probiotics have been receiving growing attention in recent years for their ability to modulate physiological functions such as

nutrition and metabolism [4] and [5], immunity [6] and [7] and gut-brain axis [8] and [9]. They play significant roles in the fight against pathological phenomena such as infections [10], cancer [11], inflammation, allergies, autoimmune diseases [7], oxidative stress [12] and [13], cardiovascular problems [5] and psychiatric disorders [14] which affect health and life quality. However, several biological activities supposed on the basis of analyses of microbial genomes still need to be proved by functional investigations. Resistance to technological treatments or phenotypic modifications occurring during lyophilization, processing and storage, are also important issues to ascertain that the viability and the beneficial properties of a particular strain (employed as starter or as bio-control agent) are conserved [15], [16] and [17].

Current genetic typing of a strain is far to be able to provide an accurate picture of bacterial physiology and phenotypes. Actually, a huge number of epigenetic modulations are known to occur in complex ecosystems such as food matrices or animal intestine where most food and probiotic bacteria live. For instance, in lactic acid bacteria (LAB), the most employed probiotics and starters, a number of metabolic pathways are controlled by pH, such as some routes activated only by acidic pH conditions and generating toxic compounds (e.g., biogenic amine production or arginine deiminase pathway) [18]. The attitude to exclude *a priori* a certain starter strain for such genetic traits is incorrect if the food matrix pH is neutral or alkaline [18]. Furthermore, in the complex gut ecological niche, reciprocal interactions between bacteria and host occur: it has been demonstrated that transit through the mouse intestine changes the expression level of several bacterial genes involved in the induction of specific cross-talking signals [19] and [20], and that exposure to probiotic strains also affect colonocyte transcriptome [21]. On the other hand, it is worth reminding that gene function annotation by genomic analyses is based only on sequence homologies with genes present in databases and final validation should always be obtained by functional studies.

All these considerations underline the importance of studying bacterial protein profiles in different experimental conditions. Comparative proteomic analysis is among the most suitable tools to evaluate the phenotypic responses of a food starter or a probiotic strain to a certain environment. For instance, lactose starvation stress and the effect of the surfactant Tween-80 as a protective agent, have been evaluated in *Lactobacillus casei* GCR163 by a comparative study based on nanoLC-MS/MS [Al Naseri 22]. A gel-free proteomic approach was used to elucidate the acidic stress response in *Lactobacillus plantarum* 423 [Heunis 23]. As far as proteomics is concerned, 2DE in-gel techniques proved to be of particular interest allowing us also to detect enzyme isoforms. For instance, it is possible to discriminate a differential biosynthesis of isoenzymes involved in either glycolytic or gluconeogenic direction of Embden-Meyerhof-Parnas route [24]. Since the beginning of the proteomic era, several protocols have been set up to recover proteins from different cell districts allowing to separately analyze different subproteomes expressed by microorganisms (Fig. 1). The analysis of subproteomes represents a

valuable tool to obtain additional information on specific problems: membrane protein patterns can reveal induction of transporters [24] and, since the first adaptive modifications to hostile environments occur at envelope level, stress tolerance pathways [25]. On the other hand, surface and extracellular proteomes help typing probiotic bacteria by elucidating what happens in the bacterium-host interface. In this context, a very recent cutting-edge paper [Le Marechal 26] describes three methods to recover key surface proteins involved in immune-modulation in *Propionibacterium freudenreichii*.

Fig. 1.

Flow chart illustrating sample preparation strategies for the analysis of the different subproteomes.

[Figure options](#)

For all these reasons we are willing to contribute with a review reporting examples in which the analysis of LAB subproteomes proved to be essential for reconstructing metabolic pathways, obtaining information about bacterial physiology and behavior and characterizing starters and probiotic bacteria in view of industrial or *in vivo* applications.

1.1. Biogenic amine risk: tracking the history of an undesired metabolic pathway by 2-DE

LAB are quite acid tolerant. Because of their inability to biosynthesize heme, they cannot get functional respiratory chains and have evolved a substrate-level phosphorylation mainly based on lactic acid fermentation. Apart from being a low-energy gain route, this pathway causes pH-lowering and LAB have to face acidification problems [18]. Some LAB strains display a natural ability to tolerate the harsh wine environment (pH = 3) as well as

the very acidic pH of stomach. This phenotype is based on several strategies to neutralizing pH lowering and/or to adapting to acidic environments and is highly appreciated in starter and probiotic LAB. Unfortunately, some of these metabolic strategies generate undesired molecules such as spoilage amines, like putrescine and cadaverine that alter the organoleptic properties of food [27]. Also bio-active amines acting on the Central Nervous System (CNS) and the vascular apparatus like histamine, an hypotensive molecule involved in allergies, and tyramine, an hypertensive metabolite sporadically involved in cerebral hemorrhage [28], [29] and [30], can be produced.

Some years ago, our research group used proteomic approach to evaluate the potential of LAB strains, either isolated from food or present in our microbial collection, to produce biogenic amines in different environmental conditions [24], [31], [32] and [33]. Once established that certain strains possess the genetic determinants for amino acid decarboxylation to amines, it still remained to elucidate if amines were actually produced and if their production was constitutive, modulated by environmental factors through regulation of gene expression or rather controlled at catalytic level.

In toto comparative proteomes of wine- and collection-isolated lactobacilli (harboring gene determinants for histamine production) grown with or without the precursor amino acid were analyzed [24] and [32]. The results obtained demonstrated that histidine decarboxylase (HDC) biosynthesis is significantly affected by the presence of histidine in the culture medium [24]. These preliminary data indicated that: i) histamine production, apart from being controlled through catalytic modulation of HDC by several effectors [34] and [35], was also regulated at biosynthetic level by histidine; ii) starter strains possessing the genetic determinants for histamine production could be safely employed in wine for malo-lactic fermentation since the content of histidine in wine is generally very low and the only amino acid present in a significant amount after alcoholic fermentation and separation of dead yeast cells is arginine [36]. However, since induction by the precursor amino acid is not the only factor triggering histamine biosynthesis, caution should be applied and other inducers should be tested, as reported in the next paragraphs. As far as tyrosine decarboxylation is concerned, soluble protein analysis failed to reveal differential production of tyrosine decarboxylase (TDC) in a tyramine-producing *Enterococcus faecalis* DISAV1022 isolated from a goat cheese. For this reason, we addressed our interest to membrane protein profiles. Membrane-enriched comparative proteome, obtained by an extensive ultracentrifugation followed by extraction with thrifluoroethanol/chloroform mixture [37] and [38], enabled us to identify a spot corresponding to tyrosine decarboxylase (TDC) which was detectable only in cultures supplemented with 1 g/L tyrosine [31]. This study demonstrated that also the tyrosine decarboxylation pathway was induced at biosynthetic level by the precursor amino acid. Unfortunately, tyrosine content in cheese is very high, suggesting the exclusion of this *E. faecalis* strain as starter for cheese making.

On the other hand, our proteomic approaches revealed that the risk of undesired bio-active amine production can also be limited by alternative strategies based on induction of pathways competing with amino acid decarboxylation. It is known that similar alkalization and energy gain can be obtained through either malo-lactic fermentation (MLF) or arginine deiminase pathway (ADI) [18]. The former route is very similar to amino acid decarboxylation and consists in the conversion of the dicarboxylic malic acid into the monocarboxylic lactic acid. This reaction contributes to both pH buffering and generation of metabolic energy (PMF). ADI pathway produces ATP and significant alkalization as well [18]. On the basis of these considerations possible synergies/competitions among the three pathways (amino acid decarboxylation, MLF and ADI) were explored [32]. Comparative proteomic analyses of soluble extracts of *Lactobacillus hilgardii* ISE 5211 grown on high amounts of malic acid and/or arginine and/or histidine demonstrated that histidine negatively controls ADI pathway enzyme biosynthesis and that reciprocally arginine down-regulates HDC production. Although these findings offer an interesting paradigm of reciprocal regulation between these pathways, they cannot be exploited from an applicative standpoint since the ADI alkaline end-product is ammonia, a metabolite which is more toxic and dangerous than histamine for human health. Unfortunately, MLF did not show any significant competition against histidine decarboxylation or ADI in *L. hilgardii* [32]. However, recent results by Laroute and co-workers (Laroute et al., unpublished data) indicated negative modulation of amine production by malate in a *Lactococcus lactis* strain, extensively employed in the dairy industry. In such a case, malate supplementation could be considered an industrial fermentation strategy to re-direct microbial metabolism towards lactic acid instead of amine production.

A particular case of bio-active amine produced by LAB is gamma-amino butyrate (GABA) originating from glutamate decarboxylation. GABA is an appreciated molecule since it acts as a relaxing agent on the smooth muscles of the intestine [39] and also exerts positive effects on mood promoting tranquility [40], [41] and [42]. GABA-producing LAB strains show improved survival to gastrointestinal transit and have been added to several fermented foods [43] with the aim of releasing GABA directly in the gut. A combined fermentation-transcriptomic-proteomic investigation, performed on *L. lactis* NCDO2118, revealed that: i) glutamate-fortified culture medium can enhance GABA biosynthesis, and also glutamine can serve as a precursor in absence of glutamate thus allowing the recovery of a significant amount of GABA; ii) glutamate supplementation down-regulated the biosynthesis of ADI pathway enzymes [33]. As illustrated in Fig. 2 both pathways contribute to energy gain and pH lowering. These results suggest that, in strains bearing both the genes for ADI and glutamate decarboxylation, an excess of glutamate or glutamine in the food matrix allows the preventing of ammonia accumulation.

Fig. 2.

Competing routes for arginine deimination (ADI) and glutamate decarboxylation pathway which contribute to acid resistance and generation of metabolic energy in lactic acid bacteria. *ADI*, arginine deiminase; Arg, arginine; *CK*, carbamate kinase; GABA, γ -aminobutyric acid; *GAD*, glutamate decarboxylase; Glu, glutamate; Orn, ornithine; *OTC*, ornithine transcarbamylase; PFM, proton motive force.

[Figure options](#)

The overall results we obtained by comparative proteomic experiments performed on biogenic amine-producing LAB are summarized in [Table 1](#). The presence of the precursor amino acid: i) induces a cytosolic overproduction of histidine decarboxylase (in *Lactobacillus w53*, *Lactobacillus* 30a and *L. hilgardii* ISE 5211); ii) triggers an overexpression of membrane-bound tyrosine decarboxylase in *E. faecalis* DISAV1022; and iii) does not induce differential biosynthesis of glutamate decarboxylase in *L. lactis* NCDO2218 as shown by both cytosol and membrane protein analysis.

Table 1.

Summary of the results obtained from comparative proteomic analyses of different biogenic amine producing LAB. NP, not performed.

Bacterial strain	Stimulated condition	Biogenic amine	Involved enzyme	Cytosol proteome	Membrane-enriched proteome	Reference
<i>Lactobacillus w53</i>	10 g/L histidine	Histamine	Histidine decarboxylase	Enzyme up-regulated	NP	[24]
<i>Lactobacillus</i> 30a	10 g/L histidine			Enzyme up-regulated	NP	[24]
<i>Lactobacillus hilgardii</i> ISE5211	4 g/L histidine			Enzyme up-regulated	NP	[32]
<i>Enterococcus faecalis</i> DISAV1022	1 g/L tyrosine	Tyramine	Tyrosine decarboxylase	Enzyme not up-regulated	Enzyme up-regulated	[31]
<i>Lactococcus lactis</i> NCDO2118	5 g/L glutamate	GABA	Glutamate decarboxylase	Enzyme not up-regulated	Enzyme not up-regulated	[33]

[Table options](#)

1.2. Surviving lyophilization without losing efficacy: what DIGE proteomics can add

The use of optimized and stable bacterial cultures both as starters for industrial processes and as probiotics is a well established practice [\[44\]](#) and [\[45\]](#). Furthermore, the use of selected LAB (such as bacteriocin-producing strains) as bio-control agents to counteract food-borne infections and food spoilage has been expanding in the last decade [\[46\]](#). Nowadays, few highly performant strains are available and used throughout the world

(e.g., by food fermentation and pharmaceutical plants) for each of these applications. Hence, these bacterial strains need to be sent without damage by starter-producing companies. This can be achieved by lyophilization procedures. The same treatment is also used for probiotic strains that are not resistant to gastrointestinal transit and therefore need to be encapsulated before administration. The main problem related to the use of these standardized procedure is represented by cell death during the freeze-drying process. Several cryoprotectants have been suggested to enhance the survival rate: carbohydrates such as polyfructose, trehalose and raffinose display a protective effect on membrane and a stabilizing activity on proteins [47] and [48]. Trehalose proved to be one of the most performant cryoprotectants [49], so we decided to evaluate its effect on the cheese-isolated *L. lactis* strain 11D. The growth of this strain in the presence of trehalose as carbon source resulted in a higher growth rate than in the presence of glucose as carbon source [50]. To understand if this biomass increase corresponds to significant physiological modifications (and, notably, to potentially harmful phenotype) a 2D-fluorescence difference gel electrophoresis (DIGE) experiment was performed to highlight differences in protein profiles between cultures grown on trehalose- and glucose-containing medium. The main goal was to exclude induction of potential virulence factors by trehalose, and also to ascertain if the main probiotic characters of *L. lactis* were conserved during growth on this sugar. Inhibition of the biosynthesis of two stress proteins (GroEL and proline dipeptidase) by trehalose was observed. Such attenuation of the stress response is probably due to the protective effect of trehalose itself. Furthermore, trehalose up-regulated the production of a chitinase, suggesting that growth on this sugar could enhance the potential of this bacterial strain against fungi [50]. It has to be mentioned that in some LAB strains, chitinase gene(s) are found within an operon also encoding a chitin-binding protein involved in adhesion to chitin but also to mucins since both chitins and mucins contain N-acetylglucosamine units [51]. Hence, these data suggest that the gut adhesive potential of *L. lactis* is increased by trehalose. The biosynthesis of a cell-wall renewal enzyme was also stimulated by trehalose, confirming that this sugar can improve the growth rate of this strain.

1.3. Controversial strain typing: combining in-gel and gel-free proteomics to overcome the concept of species-pathogenicity

Some bacterial species, such as *E. faecalis*, have long been employed in food-making. *E. faecalis* is especially used in the Mediterranean area for cheese production and it is appreciated for its capability to confer specific texture and organoleptic properties to food due to its proteolytic and esterolytic activity and diacetyl production [52], [53] and [54]. *Enterococcus* strains have also been proposed as probiotics, mainly for their ability to produce bacteriocins called enterocins [1], [55] and [56]. However, because of their resistance to a wide range of pHs and temperatures, enterococci are ubiquitous and able to colonize heterogeneous niches (soils, food and the

gastro-intestinal tract of humans and animals). It has recently been demonstrated that human enterococci display the same antibiotypes and toxinogenic profiles than pig-, cattle- and sheep-isolated enterococci and probably the strains hosted in the human gut are acquired from food [57]. An increasing number of multidrug resistant (MDR) enterococci has been reported in the literature: they can become the dominant flora under antibiotic pressure, predisposing immune-compromized patients to severe hospital infections including sepsis with a high mortality degree [58]. Apart from antibiotic resistance, virulence factors such as proteolytic enzymes have been also demonstrated to be present in enterococci [59] and [60]. Therefore the safe use of this bacterial species in food is questionable [61].

To shed light on the potential pathogenicity characters of *E. faecalis*, a proteomic investigation was carried out to compare food-derived and hospital-isolated strains [62]. A combined in-gel (exoproteome) and gel-free (*in toto* proteome by nano LC-MALDI-TOF/TOF MS) approach was used and *E. faecalis* strains of different origin proved to be significantly different. First of all, the exoproteome of the hospital-isolate contained several virulence factors, like gelatinase, extracellular serine protease and superoxide dismutase which were absent in the food-isolate protein extracts. The-gel-free approach complemented these results by revealing an increased production of additional proteins potentially involved in pathogenicity in the hospital strain. Among the latter, the endocarditis-specific antigen, a chitin-binding protein, an adhesion lipoprotein and a two-component regulatory system for alkaline phosphatase activation were identified. On the other hand, the food-isolated strain expressed proteins related to bacteria–bacteria communication suggesting a good adaptation to the complex multi-organism food matrix [62]. All these results highlight the need of detailed typing of strains to be employed in food fermentation or as probiotics and proteomics proved to be an excellent tool for this purpose.

1.4. Exploring exoproteomes for evaluating probiotic persistence, safety and efficacy

Probiotic bacteria are defined as live microorganisms that, when administered in adequate amounts, confer a health benefit to the host [63]. Probiotics are used either as food-additives or as (lyophilized or encapsulated) pharmaceutical preparations. They can be employed to counteract intestinal infections in the newborns, traveler's diarrhea, post-antibiotic unbalance of the gut microbiota and also as immune-modulators controlling inflammation, allergies, and autoimmune diseases [64]. Apart from these ascertained functions, they have received growing attention as cell cycle regulators since they can produce short chain fatty acids (SCFA), controlling and preventing colon cancer, and conjugated linoleic acids (CLA), acting as modulators on both lipid and sugar metabolism [65], [66] and [67]. Nevertheless, the latter functions still have to be validated and controversial results are reported in the literature [68] and [69]. The best recognized probiotics belong to the genera *Lactobacillus* and *Bifidobacterium*. Recently, also

Propionibacteria and some yeasts, like *Saccharomyces boulardii*, have started to be considered suitable for probiotic use [70], [71] and [72].

To be recognized as probiotics, bacteria need to satisfy the following safety and efficacy requirements: 1. to be GRAS (generally recognized as safe), even in immunocompromized hosts; 2. to possess a desirable antibiotic sensitivity profile (lacking antibiotic resistance traits); 3. to be able to survive to the gastro-intestinal transit, resisting to acidic pH, gastric enzymes and bile salts; and 4. to be able to adhere to gut cells [73].

Some of the tests needed to proving these requisites are quite simple, while more sophisticated analyses are necessary to the establishment of adhesion capacity and immune stimulating properties. The ability to permanently adhere to the gut mucosa is the key for probiotics persistence in the human gut. Adhesion is based on complex and sequential events including: i) chemotactic movements towards mucus; ii) weak interactions with mucus and mucosa; and iii) production of specific adhesive molecules interacting with both gut cell surface and extracellular matrix proteins (EMP) [64]. If this process is correctly achieved, probiotics can exclude or displace other bacteria thus permanently colonizing human gut [74]. Specific adhesive molecules involved in this mechanism are present at the cell surface of probiotic strains: teichoic and lipoteichoic acids, exopolysaccharides (EPS) and, above all, proteins. In this context the study of the cell surface proteome allows to establish which adhesive proteins are involved in this phenomenon for each probiotic strain. It has to be underlined that most proteins displayed on the bacterial surface are weakly bound to the cell wall, and hence they can be recovered from the culture medium. This weak association depends on the fact that these proteins are anchored to the cell wall through non-covalent interactions [75], being “easily” released during cell-wall turnover [76], or as a result of pH changes [77] in the extracellular environment where they exert different functions. This is an interesting phenomenon allowing the use of a single protein for multiple functions. Actually, most adhesins belong to the class of moonlighting proteins, displaying several different roles in relation to the cellular compartment in which they are [78]. Roles of different moonlighting proteins are summarized in Table 2. All these considerations underline that the analysis of the exoproteome is the best method to establish how many different proteins are involved in the adhesion process. Of course, to fully elucidate the overall adhesion potential of a strain, exoproteome findings have to be complemented and validated by phenotypic tests using cell culture methods (such as adhesion tests to model gut cells) that unequivocally demonstrate adhesive capabilities.

Table 2.

Established role of different moonlighting proteins.

Moonlighting protein	Role	References
α -enolase	Adhesion, plasminogen binding	[77] and [79]
Glyceraldehyde 3-phosphate dehydrogenase	Adhesion	[77]
Phosphoglycerate mutase	Plasminogen binding	[79]
EF-Tu	Adhesion	[80] and [81]

Moonlighting protein	Role	References
GroEL	Adhesion	[50]
DnaK	Plasminogen binding	[79]
Ornithine transcarbamylase	Immunogenic surface protein	[82]

[Table options](#)

In our recent researches, the exoproteome and the adhesion attitude of three *Lactobacillus* strains (*Lactobacillus reuteri* Lb2 BM DSM 16143, *L. plantarum* S11T3E and *Lactobacillus pentosus* S3T60C) were investigated to correlate adhesion potential to specific proteins [\[83\]](#) and [\[84\]](#). The adhesive ability of all these strains had been previously proved, by using Caco-2 cells as a model of gut mucosa. All the exoproteomes were analyzed during the middle exponential phase to avoid cytosolic contaminations due to cell lysis during the stationary growth phase. *L. reuteri* exoproteome revealed the presence of several moonlighting proteins (PGK, GAPDH, EF-Ts, EF-Tu and trigger factor). Some of them (EF-Tu, GAPDH and trigger factor) have been previously referred to as adhesins. The fact that most of these proteins are enzymes involved in central metabolism or protein biosynthesis, strongly underlines the importance of subproteome evaluation. Currently available bioinformatic tools can hardly predict if the product of a gene will be secreted or not, since some proteins are known to be translocated across the cell membrane even if they do not possess a “traditional” signal peptide for secretion [\[85\]](#). As a consequence, gene-function predictive software can hardly help us to ensure if a gene product with a proved intracellular function (involvement in central energy metabolism) also possesses a more sophisticated and interactive role in the cell wall or extracellular environment. It is interesting to underline that some moonlighting proteins found in *L. reuteri* exoproteome (like EF-Tu) can have immunomodulating functions [\[86\]](#) and this is a valuable information about a probiotic strain. Actually, probiotic and gut bacteria can act on dendritic cells by means of interactions with Toll-like receptors and direct them towards specific differentiation [\[87\]](#). In this way, different immune functions (e.g., production of pro-inflammatory or anti-inflammatory cytokines, T helper/T regulatory cell ratio) can be controlled.

L. plantarum S11T3E exoproteome revealed the presence of several proteins with adhesive functions (adherence protein with chitin-binding domain, GAPDH, M23 family peptidase) confirming the hypothesis that its adhesive ability mainly depends on this family of macromolecules. On the contrary, only one potential adhesive protein (M23 family peptidase) was present in *L. pentosus* S3T60C exoproteome, indicating that its adhesive capability could be mediated chiefly by macromolecules different from proteins. Notably, exopolysaccharides could be responsible for this property as confirmed by its ability to produce high amounts of these compounds [\[84\]](#).

1.5. Subproteomes for reconstructing metabolic pathways: *L. reuteri* story of selenium fixing and releasing

L. reuteri is a probiotic species possessing both acidic pH and bile salt resistance [\[88\]](#) and [\[89\]](#) and appreciated for its ability to produce microbial interference

molecules such as reuterin, reutericin and reutericycline useful to counteract infections [90],[91] and [92]. It is also able to compete with pathogenic bacteria for adhesion [93] and to attenuate toxin production in both bacteria [94] and fungi [95]. *L. reuteri* Lb2 BM DSM 16143 is able to grow on sodium selenite, recovering it from the culture medium. Part of this selenite is reduced to elemental selenium and accumulated on the cell surface as nanoparticles [12], but a significant part follows another way. In order to clarify the metabolic destiny of selenium (Se) into the cell, a comparative proteomic investigation of the *in toto* fraction by 2DE was set up by growing *L. reuteri* Lb2 BM DSM 16143 with and without sodium selenite [12]. An enhanced biosynthesis of selenocysteine lyase was detected in stimulated condition, suggesting, for the first time, a possible ability of this strain to fix selenium into selenocysteines. Selenium is usually inserted into two amino acids: selenomethionine (SeMet) and selenocysteine (SeCys). While SeMet is formed by a simple replacement of sulfur with selenium in methionine, SeCys is defined as the 21st amino acid and it is genetically encoded by the UGA codon[96]. Further investigations on this metabolic ability were performed by laser ablation inductively coupled plasma mass spectrometry imaging (LA-ICP MSI) of Se-containing proteins in 2D gel electrophoresis, followed by their identification by capillary HPLC-electro-spray Orbitrap MS/MS assisted by the quantitative control of selenium elution by ICP MS [97]. This approach provided a highly reliable demonstration that this *L. reuteri* strain fixes Se exclusively into SeCys, whereas Met/SeMet substitution did not occur. This study also allowed the identification of the proteins in which Se is stored as SeCys (phosphoketolase, pyruvate kinase, arginine deiminase, 6-phosphogluconate dehydrogenase, GAPDH, ornithine carbamoyltransferase, ribonucleoside hydrolase RihC), as well as the exact position in which SeCys is inserted within the protein primary structure (Fig. 3). This work was a further demonstration of the power of different and combined proteomic approaches in the study and characterization of properties and abilities of probiotic bacteria.

Fig. 3.

Incorporation of selenium as selenocysteine in specific proteins of *L. reuteri* LB2 BM DSM16143 as revealed by a 2D LA-ICP MSI approach.

[Figure options](#)

The study of the exoproteome of *L. reuteri* Lb2 BM DSM 16143 grown in the presence of sodium selenite and the manual interpretation of the raw data of the LC-MS/MS, revealed the presence of two peptides containing a SeCys and belonging to GAPDH and phosphoketolase, respectively [13]. This experimental evidence has strengthened the probiotic potential of this strain since it demonstrated its ability to secrete organic bio-available forms of Se (Se-containing proteins). Furthermore, this is also a very appreciated feature in the perspective of using the strain as nutraceutical supplement to treat Se deficiencies, or as an antioxidant to prevent viral infections and aging-related diseases [98].

2. Conclusions

Ten years of investigations on probiotic and food bacteria proteomes by 2DE, DIGE and gel-free approaches have demonstrated that proteomic techniques can offer highly valuable and efficient help in typing strains to be employed as food starters, probiotics and nutraceutical supplements. The main results discussed in this review are summarized in [Table 3](#). However, such a challenge is still at its beginnings. Nowadays, the probiotic/starter potential is still largely underexplored and underexploited, and further investigations of the protein profiles can add consistence to claims and open new relevant perspectives in the fields of food industry and human health.

Table 3.

Contributions concerning proteomic analyses revised in this manuscript.

Physiological Function Studied	Method	Reference
Lactose starvation	LC-MS/MS	[22]
Acid stress response	LC-MS/MS	[23]
Transporter induction	Total in-gel proteome	[24]
Envelope stress response	High isoelectric point subproteome	[25]
Acclimation before MLF	Total in-gel proteome	[17]
Se-fixing pathway	Total in-gel proteome	[12]
Moonlighting proteins	Envelope-enriched and exoproteome by in-gel proteome	[13]
Extracellular protein modulation by human gut factors	SDS-PAGE and MALDI MS/MS	[20]
Tyrosine and phenylalanine decarboxylase enzymes	Membrane in-gel proteome	[31]
ADI and histidine decarboxylation reciprocal competition	Total in-gel proteome	[32]
Membrane proteins solubilization	Membrane-enriched in-gel proteome	[37]
GABA production	In-gel proteome and transcripts	[33]
Separation of membrane proteins	Membrane in-gel proteome	[38]
Lyophilization resistance	DIGE in-gel proteomic	[50]
Virulence factors in food and hospital Enterococci	In-gel exoproteome and gel-free total proteome	[62]
Exported proteins	In-gel exoproteome	[76]
Adhesion	In-gel exoproteome	[83]
Bile salts response	Total in-gel proteome	[88]
Adhesion	In-gel exoproteome	[84]
Selenocysteine-containing proteins	LA-ICP MSI	[97]

[Table options](#)

Conflict of interests

The authors declare no conflict of interests.

References

1.
 - [\[1\]](#)

- L.M. Perin, R.O. Miranda, S.D. Todorov, B.D. Franco, L.A. Nero
- **Virulence, antibiotic resistance and biogenic amines of bacteriocinogenic lactococci and enterococci isolated from goat milk**

○ Int J Food Microbiol, 185 (2014), pp. 121–126

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(7\)](#)

2.

- [\[2\]](#)
- B.K. Schindler, S. Bruns, G. Lach
- **Biogenic amines — a possible source for nicotine in mushrooms? A discussion of published literature data**

○ Food Chem, 171 (2015), pp. 379–381

○ [Article](#)

|

○ [Purchase PDF](#)

3.

- [\[3\]](#)
- P. Fernandez de Palencia, M. Fernandez, M.L. Mohedano, V. Ladero, C. Quevedo, M.A. Alvarez, *et al.*
- **Role of tyramine synthesis by food-borne *Enterococcus durans* in adaptation to the gastrointestinal tract environment**

○ Appl Environ Microbiol, 77 (2011), pp. 699–702

○ [Full Text via CrossRef](#)

4.

- [\[4\]](#)
- H. Meisel
- **Overview on milk protein-derived peptides**

○ Int Dairy J, 8 (1998), pp. 363–373

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(235\)](#)

5.

- [\[5\]](#)
- R.M. Pigeon, E.P. Cuesta, S.E. Gililliand
- **Binding of free bile acids by cells of yogurt starter culture bacteria**
- J Dairy Sci, 85 (2002), pp. 2705–2710
- [Article](#)
- |
- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(50\)](#)

6.

- [\[6\]](#)
- T. Pessi, Y. Sütas, M. Hurme, E. Isolauri
- **Interleukin-10 generation in atopic children following oral *Lactobacillus rhamnosus* GG**
- Clin Exp Allergy, 30 (2000), pp. 1804–1808
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)

|

[Citing articles \(271\)](#)

7.

- [\[7\]](#)
- K. Karimi, M.D. Inman, J. Bienenstock, P. Forsythe
- ***Lactobacillus reuteri*-induced regulatory T cells protect against an allergic airway response in mice**
- Am J Respir Crit Care Med, 179 (2009), pp. 186–193
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)

|

[Citing articles \(133\)](#)

8.

- [\[8\]](#)
- Y.R. Cho, J.Y. Chang, H.C. Chang
- **Production of gamma-aminobutyric acid (GABA) by *Lactobacillus buchneri* isolated from kimchi and its neuroprotective effect on neuronal cells**
- J Microbiol Biotechnol, 17 (2007), pp. 104–109
- [View Record in Scopus](#)
- |
- [Citing articles \(84\)](#)

9.

- [\[9\]](#)
- R. Mazzoli
- **Neuro-active compounds produced by probiotics**
- E. Pessione (Ed.), Interactive Probiotics, CRC press, Boca Raton (2014), pp. 148–176
- [View Record in Scopus](#)
- |
- [Citing articles \(2\)](#)

10.

- [\[10\]](#)
- C. Lamberti, F. Genovese, J.D. Coisson, G. Lo Bianco, L.S. Coccolin, E. Pessione
- **Anti-*S. aureus* and anti-*L. monocytogenes* molecules produced by cheese-isolated lab**
- Czech J Food Sci, 32 (2014), pp. 54–60
- [View Record in Scopus](#)
- |
- [Citing articles \(2\)](#)

11.

- [\[11\]](#)
- M. Raman, P. Ambalam, K.K. Kondepudi, S. Pithva, C. Kothari, A.T. Patel, *et al.*
- **Potential of probiotics, prebiotics and synbiotics for management of colorectal cancer**
- Gut Microbes, 4 (3) (2013), pp. 181–192
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(17\)](#)

12.

- [\[12\]](#)
- C. Lamberti, E. Mangiapane, A. Pessione, R. Mazzoli, C. Giunta, E. Pessione

- **Proteomic characterization of a selenium-metabolizing probiotic *Lactobacillus reuteri* Lb2 BM for nutraceutical applications**

- Proteomics, 11 (2011), pp. 2212–2221

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(13\)](#)

13.

- [\[13\]](#)

- E. Mangiapane, C. Lamberti, A. Pessione, E. Galano, A. Amoresano, E. Pessione

- **Selenium effects on the metabolism of a Se-metabolizing *Lactobacillus reuteri*: analysis of envelope-enriched and extracellular proteomes**

- Mol Biosyst, 10 (6) (2014), pp. 1272–1280

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(3\)](#)

14.

- [\[14\]](#)

- R. Wall, J.F. Cryan, R.P. Ross, G.F. Fitzgerald, T.G. Dinan, C. Stanton

- **Bacterial neuroactive compounds produced by psychobiotics**

- Adv Exp Med Biol, 817 (2014), pp. 221–239

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(12\)](#)

15.

- [\[15\]](#)

- S. Song, D.W. Bae, K. Lim, M.W. Griffiths, S. Oh

- **Cold stress improves the ability of *Lactobacillus plantarum* L67 to survive to freezing**

- Int J Food Microbiol, 191C (2014), pp. 135–143

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

Citing articles (5)

16.

- [\[16\]](#)
- H. Velly, M. Bouix, S. Passot, C. Penicaud, H. Beinstainer, S. Ghorbal, *et al.*
- **Cyclopropanation of unsaturated fatty acids and membrane rigidification improve the freeze-drying resistance of *Lactococcus lactis* subsp. *lactis* TOMSC161**
- Appl Microbiol Biotechnol, 99 (2015), pp. 907–918
- [Full Text via CrossRef](#)

17.

- [\[17\]](#)
- D. Cecconi, A. Milli, S. Rinalducci, L. Zolla, G. Zapparoli
- **Proteomic analysis of *Oenococcus oeni* freeze-dried cultures in order to assess the importance of cell acclimatation to conduct malo-lactic fermentation in wine**
- Electrophoresis, 30 (2009), pp. 2988–2995
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)

|

Citing articles (18)

18.

- [\[18\]](#)
- A. Pessione, C. Lamberti, E. Pessione
- **Proteomics as a tool to study biogenic amine production**
- Mol Biosyst, 6 (8) (2010), pp. 1419–1430
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)

|

Citing articles (11)

19.

- [\[19\]](#)
- M. O'Connel Motherway, A. Zomer, S.C. Leahy, J. Reunanen, F. Bottacini, M.J. Claesson, *et al.*
- **Functional genome analysis of *Bifidobacterium breve* UCC2003 reveals type IVb tight adherence (Tad) pili as an essential and conserved host-colonization factor**
- PNAS, 108 (27) (2011), pp. 11217–11222
-

20.

- [\[20\]](#)
- B. Sanchez, L. Ruiz, A. Suarez, C.G. de los Reyes-Gavilan, A. Margolles
- **Human cecum content modulates production of extracellular proteins by food and probiotic bacteria**
- FEMS Microbiol Lett, 324 (2011), pp. 189–194
- [View Record in Scopus](#)
|
[Full Text via CrossRef](#)
|
[Citing articles \(2\)](#)

1.

- [\[21\]](#)
- S. O'Flaherty
- **Probiotic-mediated modulation of host gene expression**
- E. Pessione (Ed.), Interactive Probiotics, CRC press, Boca Raton (2014), pp. 81–98
- [View Record in Scopus](#)
|
[Citing articles \(1\)](#)

2.

- [\[22\]](#)
- A. Al Naseri, J.P. Bowman, R. Wilson, R.E. Nilsson, M.L. Britz
- **Impact of lactose starvation on the physiology of *Lactobacillus casei* GCRL163 in the presence or absence of Tween 80**
- J Proteome Res, 12 (2013), pp. 5313–5322
- [View Record in Scopus](#)
|
[Full Text via CrossRef](#)
|
[Citing articles \(6\)](#)

3.

- [\[23\]](#)
- T. Heunis, S. Deane, S. Smit, L.M. Dicks
- **Proteomic profiling of the acid stress response in *Lactobacillus plantarum* 423**
- J Proteome Res, 13 (2014), pp. 4028–4039
- [View Record in Scopus](#)
|

[Full Text via CrossRef](#)

|

[Citing articles \(6\)](#)

4.

- [\[24\]](#)
- E. Pessione, R. Mazzoli, M.G. Giuffrida, C. Lamberti, E. Garcia-Moruno, C. Barello, *et al.*
- **A proteomic approach to studying biogenic amine producing lactic acid bacteria**
- Proteomics, 5 (3) (2005), pp. 687–698

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(46\)](#)

5.

- [\[25\]](#)
- R. Mazzoli, P. Fattori, C. Lamberti, M.G. Giuffrida, M. Zapponi, C. Giunta, *et al.*
- **High isoelectric point sub-proteomes analysis of *Acinetobacter radioresistens* S13 reveals envelope stress responses induced by aromatic compounds**
- Mol Biosyst, 7 (3) (2011), pp. 598–607

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(8\)](#)

6.

- [\[26\]](#)
- C. Le Marechal, V. Peton, C. Plé, C. Vroland, J. Jardin, V. Briard-Bion, *et al.*
- **Surface proteins of *Propionibacterium freudenreichii* are involved in its anti-inflammatory properties**
- J Proteomics, 113 (2015), pp. 447–461

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(9\)](#)

7.

- [\[27\]](#)
- A. Costantini, R. Pietroniro, F. Doria, E. Pessione, E. Garcia-Moruno
- **Putrescine production from different amino acid precursors by lactic acid bacteria from wine and cider**

○ Int J Food Microbiol, 165 (2013), pp. 11–17

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(8\)](#)

8.

- [\[28\]](#)
- M.V. Moreno-Arribas, M.C. Polo, F. Jorganes, R. Muñoz
- **Screening of biogenic amine production by lactic acid bacteria isolated from grape must and wine**

○ Int J Food Microbiol, 84 (1) (2003), pp. 117–123

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(140\)](#)

9.

- [\[29\]](#)
- J.G. Millichap, M.M. Yee
- **The diet factor in pediatric and adolescent migraine**

○ Pediatr Neurol, 28 (2003), pp. 9–15

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(109\)](#)

10.

- [\[30\]](#)
- W.N. Konings
- **Microbial transport: adaptation to natural environments**
- Antonie Van Leeuwenhoek, 90 (4) (2006), pp. 325–342
- [View Record in Scopus](#)
- |
- [Citing articles \(43\)](#)

11.

- [\[31\]](#)
- E. Pessione, A. Pessione, C. Lamberti, J.D. Coisson, K. Riedel, R. Mazzoli, *et al.*
- **First evidence of a membrane-bound, tyramine and β -phenylethylamine producing, tyrosine decarboxylase in *Enterococcus faecalis*: a two-dimensional electrophoresis proteomic study**
- Proteomics, 9 (10) (2009), pp. 2695–2710
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(24\)](#)

12.

- [\[32\]](#)
- C. Lamberti, M. Purrotti, R. Mazzoli, P. Fattori, C. Barello, J.D. Coisson, *et al.*
- **ADI pathway and histidine decarboxylation are reciprocally regulated in *Lactobacillus hilgardii* SE 5211: proteomic evidence**
- Amino Acids, 41 (2) (2011), pp. 517–527
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(2\)](#)

13.

- [\[33\]](#)
- R. Mazzoli, E. Pessione, M. Dufour, V. Laroute, M.G. Giuffrida, C. Giunta, *et al.*
- **Glutamate-induced metabolic changes in *Lactococcus lactis* NCDO 2118 during GABA production: combined transcriptomic and proteomic analysis**
- Amino Acids, 39 (3) (2010), pp. 727–737
- [View Record in Scopus](#)
- |

[Full Text via CrossRef](#)

|

[Citing articles \(13\)](#)

14.

- [\[34\]](#)
- P.A. Recsei, E.E. Snell
- **Pyruvoyl-dependent histidine decarboxylases. Mechanism of cleavage of the proenzyme from *Lactobacillus buchneri***
- J Biol Chem, 260 (5) (1985), pp. 2804–2806

- [View Record in Scopus](#)

|

[Citing articles \(14\)](#)

15.

- [\[35\]](#)
- C. Rodriguez-Caso, D. Rodriguez-Agudo, A.A. Moya-Garcia, I. Fajardo, M.A. Medina, V. Subramaniam, *et al.*
- **Local changes in the catalytic site of mammalian histidine decarboxylase can affect its global conformation and stability**
- Eur J Biochem, 270 (21) (2003), pp. 4376–4387

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(26\)](#)

16.

- [\[36\]](#)
- J.P. Bourdineaud
- **Both arginine and fructose stimulate pH-independent resistance in the wine bacteria *Oenococcus oeni***
- Int J Food Microbiol, 107 (3) (2006), pp. 274–280

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(12\)](#)

17.

- [\[37\]](#)
- K. Zuobi-Hasona, P.J. Crowley, A. Hasona, A.S. Bleiweis, L.J. Brady
- **Solubilization of cellular membrane proteins from *Streptococcus mutans* for two-dimensional electrophoresis**
- Electrophoresis, 26 (2005), pp. 1200–1205
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(37\)](#)

18.

- [\[38\]](#)
- J.M.P. Deshusses, J.A. Burgess, A. Scherl, Y. Wenger, N. Walter, V. Converset, *et al.*
- **Exploitation of specific properties of trifluoroethanol for extraction and separation of membrane proteins**
- Proteomics, 3 (8) (2003), pp. 1418–1424
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(64\)](#)

19.

- [\[39\]](#)
- G. Clarke, J.F. Cryan, T.G. Dinan, E.M. Quigley
- **Review article: probiotics for the treatment of irritable bowel syndrome-focus on lactic acid bacteria**
- Aliment Pharmacol Ther, 35 (2012), pp. 403–413
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(77\)](#)

20.

- [\[40\]](#)
- A.C. Logan, M. Katzman
- **Major depressive disorder: probiotics may be an adjuvant therapy**
- Med Hypotheses, 64 (3) (2005), pp. 533–538

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

Citing articles (64)

1.

- [\[41\]](#)

- J.A. Bravo, P. Forsythe, M.V. Chew, E. Escaravage, H.M. Savignac, T.G. Dinan, *et al.*

- **Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse *via* the vagus nerve**

- PNAS, 108 (38) (2011), pp. 16050–16055

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

Citing articles (355)

2.

- [\[42\]](#)

- X. Ma, Y.K. Mao, B. Wang, J.D. Huizinga, J. Bienenstock, W. Kunze

- ***Lactobacillus reuteri* ingestion prevents hyperexcitability of colonic DRG neurons induced by noxious stimuli**

- Am J Physiol Gastrointest Liver Physiol, 296 (2009), pp. G868–G875

- [Full Text via CrossRef](#)

3.

- [\[43\]](#)

- N.Y. Kim, G.E. Ji

- **Characterization of the production of biogenic amines and gamma-aminobutyric acid in the soybean pastes fermented by *Aspergillus orizae* and *Lactobacillus brevis***

- J Microbiol Biotechnol (2014) <http://dx.doi.org/10.4014/jmb.1409.09081>

-

4.

- [\[44\]](#)

- J.C. Nielsen, C. Prah, A. Lonvaud-Funel

- **Malolactic fermentation in wine by direct inoculation with freeze-dried *Leuconostoc oenos* cultures**

- Am J Enol Vitic, 47 (1996), pp. 42–48

- [View Record in Scopus](#)

|
[Citing articles \(76\)](#)

5.

- [\[45\]](#)
- A. Saadatzadeh, M.R. Fazeli, H. Jamalifar, R. Dinarvand
- **Probiotic properties of lyophilized cell free extract of *Lactobacillus casei***
- Jundishapur J Nat Pharm Prod, 8 (3) (2013), pp. 131–137
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(2\)](#)

6.

- [\[46\]](#)
- N. Benkerroum, A. Daoudi, T. Hamraoui, H. Ghalfi, M. Duroy, P. Evrart, *et al.*
- **Lyophilized preparations of bacteriocinogenic *Lactobacillus curvatus* and *Lactococcus lactis*ssp. *lactis* as potential protective adjuncts to control *Listeria monocytogenes* in dry-fermented sausages**
- J Appl Microbiol, 98 (2005), pp. 56–63
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(48\)](#)

7.

- [\[47\]](#)
- G. Moiset, C.A. Lopez, R. Bartelds, L. Syga, E. Rijpkema, A. Cukkemane, *et al.*
- **Disaccharides impact the lateral organization of lipid membranes**
- J Am Chem Soc, 136 (2014), pp. 16167–16175
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(7\)](#)

8.

- [\[48\]](#)
- E.E. Tymczyszyn, A. Gomez-Zavaglia, E.A. Disalvo

- **Effects of sugars and growth media on the dehydration of *Lactobacillus delbrueckii ssp.bulgaricus***

- J Appl Microbiol, 102 (2007), pp. 845–851

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(22\)](#)

9.

- [\[49\]](#)

- A. Jofré, T. Aymerich, M. Garriga

- **Impact of different cryoprotectants on the survival of freeze-dried *Lactobacillus rhamnosus* and *Lactobacillus casei/paracasei* during long term storage**

- Benefic Microbes, 6 (2014), pp. 1–6

- [View Record in Scopus](#)

|

[Citing articles \(1\)](#)

10.

- [\[50\]](#)

- F. Genovese, J.D. Coisson, A. Majumder, A. Pessione, B. Svensson, S. Jacobsen, *et al.*

- **An exoproteome approach to monitor safety of a cheese-isolated *Lactococcus lactis***

- Food Res Int, 54 (1) (2013), pp. 1072–1079

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(4\)](#)

11.

- [\[51\]](#)

- B. Sanchez, C. Gonzales-Tejedo, P. Ruas-Madiedo, M. Urdaci, A. Margolles

- ***L. plantarum* extracellular chitin-binding protein and its role in the interaction between CaCo2 cells and mucin**

- Appl Environ Microbiol, 77 (2011), pp. 1123–1126

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(16\)](#)

12.

- [\[52\]](#)
- J.A. Centeno, S. Menendez, M. Hermida, J.L. Rodriguez-Otero
- **Effects of the addition of *Enterococcus faecalis* in Cebreiro cheese manufacture**
- Int J Food Microbiol, 48 (2) (1999), pp. 97–111

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(107\)](#)

13.

- [\[53\]](#)
- G. Giraffa
- **Functionality of enterococci in dairy products**
- Int J Food Microbiol, 88 (2003), pp. 215–222

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(310\)](#)

14.

- [\[54\]](#)
- C.M.A.P. Franz, M.E. Stiles, K.H. Schleifer, W.H. Holzapfel
- **Enterococci in foods, a conundrum for food safety**
- Int J Food Microbiol, 88 (2003), pp. 105–122

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(288\)](#)

15.

- [\[55\]](#)
- E. Balla, J.M.T. Dicks, M. Du Toit, M.J. van der Merwe, W.H. Holzapfel
- **Characterization and cloning of the genes encoding enterocin 1071a and enterocin 1071b, two antimicrobial peptides produced by *Enterococcus faecalis* BFE1071**
- Appl Environ Microbiol, 66 (2000), pp. 1298–1304
- [View Record in Scopus](#)
|
[Full Text via CrossRef](#)
|
[Citing articles \(70\)](#)

16.

- [\[56\]](#)
- C.M.A.P. Franz, R.W. Worobo, L.E.N. Quadri, U. Schillinger, W.H. Holzapfel, J.C. Vederas, *et al.*
- **Atypical genetic locus associated with constitutive production of enterocin B by *Enterococcus faecium* BFE 900**
- Appl Environ Microbiol, 65 (5) (1999), pp. 2170–2178
- [View Record in Scopus](#)
|
[Citing articles \(75\)](#)

17.

- [\[57\]](#)
- S. Ramos, G. Igrejas, J. Rodrigues, J.L. Capelo-Martinez, P. Poeta
- **Genetic characterization of antibiotic resistance and virulence factors in vanA-containing enterococci from cattle, sheep and pigs subsequent to the discontinuation of the use of avoparcin**
- Vet J, 193 (1) (2012), pp. 301–303
- [Article](#)
|
○ [Purchase PDF](#)
|
[View Record in Scopus](#)
|
[Citing articles \(6\)](#)

18.

- [\[58\]](#)
- W.R. Miller, J.M. Munita, C.A. Arias
- **Mechanisms of antibiotic resistance in enterococci**

- Expert Rev Anti Infect Ther, 12 (10) (2014), pp. 1221–1236

- [View Record in Scopus](#)

|

- [Full Text via CrossRef](#)

|

- [Citing articles \(4\)](#)

19.

- [\[59\]](#)

- N. Shankar, A.S. Baghdayan, M.S. Gilmore

- **Modulation of virulence within a pathogenicity island in vancomycin-resistant *Enterococcus faecalis***

- Nature, 417 (2002), pp. 746–750

- [View Record in Scopus](#)

|

- [Full Text via CrossRef](#)

|

- [Citing articles \(224\)](#)

20.

- [\[60\]](#)

- T. Semedo, M.A. Santos, M.F.S. Lopes, J.J.F. Marques, M.T.B. Crespo, R. Tenreiro

- **Virulence factors in food, clinical and reference enterococci: a common trait in the genus?**

- Syst Appl Microbiol, 26 (2003), pp. 13–22

- [Article](#)

|

- [Purchase PDF](#)

|

- [View Record in Scopus](#)

|

- [Full Text via CrossRef](#)

|

- [Citing articles \(107\)](#)

1.

- [\[61\]](#)

- F. Pérez-Martin, S. Seseña, P.M. Izquierdo, M.L. Palop

- **Are *Enterococcus* populations present during malolactic fermentation of red wine safe?**

- Food Microbiol, 42 (2014), pp. 95–101

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

Citing articles (5)

2.

- [\[62\]](#)
- A. Pessione, C. Lamberti, L. Cocolin, S. Campolongo, A. Grunau, S. Giubergia, *et al.*
- **Different protein expression profiles in cheese and clinical isolates of *Enterococcus faecalis* revealed by proteomic analysis**
- Proteomics, 12 (3) (2012), pp. 431–447

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

Citing articles (10)

3.

- [\[63\]](#)
- Anonymous
- **Guidelines for the evaluation of probiotics in food**
- Joint FAO/WHO Working Group Report on Drafting Guidelines for the Evaluation of Probiotics in Food, FAO, London, Ontario (2002), p. 11

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

Citing articles (1)

4.

- [\[64\]](#)
- E. Pessione
- **Lactic acid bacteria contribution to gut complexity: lights and shadows**
- Front Cell Infect Microbiol, 2 (2012), p. 86

- [View Record in Scopus](#)

|

Citing articles (23)

5.

- [\[65\]](#)
- J. Sauer, K.K. Richter, B.L. Pool-Zobel

- **Physiological concentration of butyrate favorably modulate genes of oxidative and metabolic stress in primary human colon cells**

- J Nutr Biochem, 18 (2007), pp. 736–745

- [Article](#)

|

- [Purchase PDF](#)

|

[View Record in Scopus](#)

|

[Citing articles \(56\)](#)

6.

- [\[66\]](#)

- C. Pellizzaro, D. Coradini, M.G. Daidone

- **Modulation of angiogenesis-related proteins synthesis by sodium butyrate in colon cancer cell line HT29**

- Carcinogenesis, 23 (2002), pp. 735–740

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(33\)](#)

7.

- [\[67\]](#)

- S. Minucci, P.G. Pelicci

- **Histone deacetylase inhibitors and the promise of epigenetic (and more) treatments for cancer**

- Nat Rev Cancer, 6 (2006), pp. 38–51

- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

[Citing articles \(1368\)](#)

8.

- [\[68\]](#)

- K.W. Wahle, S.D. Heys, D. Rotondo

- **Conjugated linoleic acids: are they beneficial or detrimental to health?**

- Prog Lipid Res, 43 (2004), pp. 553–587

- [Article](#)

|
○ [Purchase PDF](#)

|
[View Record in Scopus](#)

|
[Citing articles \(347\)](#)

- 9.
- [\[69\]](#)
 - J.R. Lupton
 - **Microbial degradation products influence colon cancer risk: the butyrate controversy**
 - J Nutr, 134 (2004), pp. 479–482
 - [View Record in Scopus](#)
- |
[Citing articles \(133\)](#)

- 10.
- [\[70\]](#)
 - S.-M. Deutsch, J. Gwenaël
 - **New probiotics. Dairy propionibacteria and the modulation of gut microbiota and physiology**
 - E. Pessione (Ed.), Interactive Probiotics, CRC press, Boca Raton (2014), pp. 122–147
 - [View Record in Scopus](#)
- |
[Citing articles \(1\)](#)

- 11.
- [\[71\]](#)
 - I.M. Smith, J.E. Christensen, N. Arneborg, L. Jespersen
 - **Yeast modulation of human dendritic cell cytokine secretion: an *in vitro* study**
 - PLoS One, 9 (5) (2014), p. e96595
 - [Full Text via CrossRef](#)

- 12.
- [\[72\]](#)
 - R. Pontier-Bres, P. Munro, L. Boyer, R. Anty, V. Imbert, C. Terciolo, *et al.*
 - ***Saccharomyces boulardii* modifies *Salmonella Typhimurium* traffic and host immune responses along the intestinal tract**
 - PLoS One, 9 (8) (2014), p. e103069
 - [Full Text via CrossRef](#)

- 13.
- [\[73\]](#)

- J.K. Collins, G. Thornton, O.G. Sullivan
- **Selection of probiotic strains for human applications**

○ Int Dairy J, 8 (1998), pp. 487–490

○ [Article](#)

|

○ [Purchase PDF](#)

|

[View Record in Scopus](#)

|

Citing articles (159)

14.

- [\[74\]](#)
- S. Even, D. Bouchard, Y. Le Loir
- **Lactic acid bacteria to modulate virulence expression in pathogenic bacteria**
- E. Pessione (Ed.), Interactive Probiotics, CRC press, Boca Raton (2014), pp. 52–80
- [View Record in Scopus](#)

|

Citing articles (2)

15.

- [\[75\]](#)
- M. Desvaux, E. Dumas, I. Chafsey, M. Hébraud
- **Protein cell surface display in gram-positive bacteria: from single protein to macromolecular protein structure**
- FEMS Microbiol Lett, 256 (2006), pp. 1–15
- [View Record in Scopus](#)

|

[Full Text via CrossRef](#)

|

Citing articles (101)

16.

- [\[76\]](#)
- B. Sanchez, M.C. Champomier-Verges, P. Anglade, F. Baraige, C.G. de los Reyes-Gavilan, A. Margolles, *et al.*
- **A preliminary analysis of *Bifidobacterium longum* exported proteins by two-dimensional electrophoresis**
- J Mol Microbiol Biotechnol, 14 (2008), pp. 74–79
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(17\)](#)

17.

- [\[77\]](#)
- J. Antikainen, V. Kuparinen, K. Lahteenmaaki, T.K. Kohronen
- **pH-Dependent association of enolase and glyceraldehyde 3-phosphate dehydrogenase of *Lactobacillus crispatus* with the cell wall and lipoteichoic acids**
- J Bacteriol, 189 (2007), pp. 4539–4543
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(69\)](#)

18.

- [\[78\]](#)
- C.J. Jeffery
- **Moonlighting proteins: complications and implications for proteomic research**
- Drug Discov Today TARGETS, 3 (2004), pp. 71–78
- [Article](#)
- [Purchase PDF](#)

|
[View Record in Scopus](#)

|
[Citing articles \(15\)](#)

19.

- [\[79\]](#)
- M. Candela, S. Bergmann, M. Vici, B. Vitali, S. Turrone, B.J. Eikmanns, *et al.*
- **Binding of human plasminogen to *Bifidobacterium***
- J Bacteriol, 189 (16) (2007), pp. 5929–5936
- [View Record in Scopus](#)

|
[Full Text via CrossRef](#)

|
[Citing articles \(49\)](#)

20.

- [\[80\]](#)
- D. Granato, G.E. Bergonzelli, R.D. Pridmore, L. Marvin, M. Rouvet, I. Corthesy-Theulaz
- **Cell surface-associated elongation factor Tu mediates the attachment of *Lactobacillus johnsonii* NCC533 (La1) to human intestinal cells and mucin**
- Infect Immun, 72 (2004), pp. 2160–2169
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(195\)](#)

1.

- [\[81\]](#)
- O. Gilad, B. Svensson, A.H. Viborg, B. Stuer-Lauridsen, S. Jacobsen
- **The extracellular proteome of *Bifidobacterium animalis* subsp. *lactis* BB-12 reveals proteins with putative roles in probiotic effects**
- Proteomics, 12 (12) (2011), pp. 2503–2514
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(26\)](#)

2.

- [\[82\]](#)
- S.I. Alam, S. Bansod, R.B. Kumar, N. Sengupta, L. Singh
- **Differential proteomic analysis of *Clostridium perfringens* ATCC13124; identification of dominant, surface and structure associated proteins**
- BMC Microbiol, 9 (2009), p. 162
- [Full Text via CrossRef](#)

3.

- [\[83\]](#)
- E. Mangiapane, C. Lamberti, A. Pessione, P. Ceruti, E. Galano, F. Novelli, *et al.*
- **Integrating proteomics and physiological tests to understand adhesion mechanism of the probiotic *Lactobacillus reuteri* Lb2 BM DSM 16341**
- Jiomics, 3 (2013), pp. 145–156
- [View Record in Scopus](#)
- |
- [Citing articles \(4\)](#)

4.

- [\[84\]](#)
- Pessione A, Lo Bianco G, Mangiapane E, Cirrincione S, Pessione E. Characterization of potentially probiotic lactic acid bacteria isolated from olives: evaluation of short chain fatty acids production and analysis of the extracellular proteome. Food Res Int 201;67:247-254.

5.

- [\[85\]](#)
- S. Yan, G. Wu
- **Signal peptide of cellulase**
- Appl Microbiol Biotechnol, 98 (2014), pp. 5329–5362
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(4\)](#)

6.

- [\[86\]](#)
- D. Granato, G.E. Bergonzelli, R.D. Pridmore, L. Marvin, M. Rouvet, I. Corthesy-Theulaz
- **Cell surface-associated elongation factor Tu mediates the attachment of *Lactobacillus johnsonii*NCC533 (La1) to human intestinal cells and mucin**
- Infect Immun, 72 (2004), pp. 2160–2169
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(195\)](#)

7.

- [\[87\]](#)
- B. Sanchez, M.C. Urdaci, A. Margolles
- **Extracellular proteins secreted by probiotic bacteria as mediators of effects that promote mucosa-bacteria interactions**
- Microbiology, 156 (2010), pp. 3232–3242
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(55\)](#)

8.

- [\[88\]](#)
- K. Lee, H.G. Lee, Y.J. Choi
- **Proteomic analysis of the effects of bile salts on the intestinal and probiotic bacterium *Lactobacillus reuteri***
- J Biotechnol, 137 (2008), pp. 14–19
- [Article](#)
- |
- [Purchase PDF](#)
- |

[View Record in Scopus](#)

|

Citing articles (27)

9.

- [\[89\]](#)
- K. Lee, K. Pi
- **Effect of transient acid stress on the proteome of intestinal probiotic *Lactobacillus reuteri***
- Biochemistry, 75 (2010), pp. 460–465
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- Citing articles (9)

10.

- [\[90\]](#)
- L.T. Axelsson, T.C. Chung, W.J. Dobrogosz, S.E. Lindgren
- **Production of a broad spectrum antimicrobial substance by *Lactobacillus reuteri***
- Microb Ecol Health Dis, 2 (1989), pp. 131–136
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- Citing articles (179)

11.

- [\[91\]](#)
- T. Kabuchi, T. Saito, Y. Kawai, J. Uemura, T. Itoh
- **Production, purification and characterization of reutericin 6, a bacteriocin with lytic activity produced by *Lactobacillus reuteri* LA6**
- Int J Food Microbiol, 34 (1997), pp. 145–156

-
- 12.
 - [\[92\]](#)
 - A. Holtzel, M.G. Ganzle, G.J. Nicholson, W.P. Hammes, G. Jung
 - **The first low molecular weight antibiotic from lactic acid bacteria: reutericyclin, a new tetrameric acid**
 - Angew Chem Int Ed, 39 (2000), pp. 2766–2768
 - [View Record in Scopus](#)
 - |
 - [Full Text via CrossRef](#)
 - |
 - [Citing articles \(76\)](#)
- 13.
 - [\[93\]](#)
 - K. Vielfort, H. Sjolinder, S. Roos, H. Jonsson, H. Aro
 - **Adherence of clinically isolated lactobacilli to human cervical cells in competition with *Neisseria gonorrhoeae***
 - Microbes Infect, 10 (2008), pp. 1325–1334
 - [Article](#)
 - |
 - [Purchase PDF](#)
 - |
 - [View Record in Scopus](#)
 - |
 - [Citing articles \(11\)](#)
- 14.
 - [\[94\]](#)
 - J. Li, W. Wang, S.X. Xu, N.A. Margarvey, J.K. McCormick
 - **Lactobacillus reuteri-produced cyclic dipeptides quench agr-mediated expression of toxic shock syndrome toxin-1 in staphylococci**
 - PNAS, 108 (2011), pp. 3360–3365
 - [View Record in Scopus](#)
 - |
 - [Full Text via CrossRef](#)
 - |
 - [Citing articles \(68\)](#)
- 15.
 - [\[95\]](#)

- A. Hernandez-Mendoza, A.F. Gonzalez-Cordova, B. Vallejo-Cordoba, H.S. Garcia
- **Effect of oral supplementation of *Lactobacillus reuteri* in reduction of intestinal absorption of aflatoxin B(1) in rats**
- J Basic Microbiol, 51 (2011), pp. 263–268
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(7\)](#)

16.

- [\[96\]](#)
- S. Gromer, J.K. Eubel, B.L. Lee, J. Jacob
- **Human selenoproteins at a glance**
- Cell Mol Life Sci, 62 (2005), pp. 2414–2437
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(136\)](#)

17.

- [\[97\]](#)
- E. Galano, E. Mangiapane, J. Bianga, A. Palmese, E. Pessione, J. Szpunar, *et al.*
- **Privileged incorporation of selenium as selenocysteine in *Lactobacillus reuteri* proteins demonstrated by selenium-specific imaging and proteomics**
- Mol Cell Proteomics, 12 (8) (2013), pp. 2196–2204
- [View Record in Scopus](#)
- |
- [Full Text via CrossRef](#)
- |
- [Citing articles \(10\)](#)

18.

- [\[98\]](#)
- E. Mangiapane, A. Pessione, E. Pessione
- **Selenium and selenoproteins: an overview on different biological systems**
- Curr Protein Pept Sci, 15 (6) (2014), pp. 598–607
- [View Record in Scopus](#)
- |

[Full Text via CrossRef](#)

|

[Citing articles \(5\)](#)

This article is part of a Special Issue entitled: HUPO 2014.