AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Allogeneic transplantation for multiple myeloma: Yes, no or maybe?

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1560262> since 2016-04-27T15:16:38Z

Published version:

DOI:10.1038/bmt.2016.12

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Bone Marrow Transplantation (2016) 51, 506–507; doi:10.1038/bmt.2016.12;.]

ovvero [Bruno B.]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://www.nature.com.offcampus.dam.unito.it/bmt/journal/v51/n4/full/bmt201612a.html>]

Allogeneic transplantation for multiple myeloma: yes, no or maybe?

B Bruno^{1,2}

1. ¹Department of Oncology, Division of Hematology, AOU Città della Salute e della Scienza di Torino, University of Torino, Torino, Italy
2. ²Department of Molecular Biotechnology and Health Sciences, University of Torino, Torino, Italy

Correspondence: Dr B Bruno, Department of Molecular Biotechnology and Health Sciences, University of Torino, Via Genova 3, Torino 10126, Italy. E-mail: benedetto.bruno@unito.it

Received 22 December 2015;
Advance online publication 22 February 2016

Accepted 7 January 2016

In this edition of *Bone Marrow Transplantation*, Ahmad *et al.*¹ report on a cohort of 92 newly diagnosed myeloma patients treated at a single institution from 2001 through 2010, with a planned standard autograft followed by an allograft from a HLA-identical sibling. At induction, 75/92 (82%) of patients were treated with VCR-adriamycin-dexamethasone and only a minority, 18/92 (18%), with bortezomib-based therapies. The conditioning regimen before the allograft consisted of IV fludarabine (30 mg/m²) and IV cyclophosphamide (300 mg/m²) for 5 days, followed by G-CSF-mobilised PBSC. The allograft was entirely delivered as an outpatient procedure. No maintenance or consolidation therapy with ‘new drugs’ was allowed. After a remarkable median follow-up of 8.8 years, probability of 10-year overall survival and PFS were 62% and 41%, respectively. Of note, the cumulative incidence of treatment-related mortality was 10% at 10 years. The cumulative incidence of extensive chronic GvHD was 79%. The majority of long-term survivors were, however, off immunosuppressive drugs at 5 years post transplant.

The results of this well-sized study reported by Ahmad *et al.* allow the following observations. First, despite the fact that the trial predated the introduction of so-called new drugs with a potent anti-myeloma activity, over 40% of patients became long-term survivors in continuous CR and at least a subset may have reached disease eradication due to graft-versus-myeloma. Second, transplant-related mortality was low and this finding clearly shows that the high transplant mortality rates observed decades ago have successfully been overcome. Third, long-term disease control and relapse mortality have become the major causes of treatment failure also in the setting of allografting. Fourth, although the incidence of chronic GvHD was high, most patients were off immunosuppression at year 5 post transplant and this undoubtedly allowed improvement in the quality of life.

In the light of the study by Ahmad *et al.* and of those by other groups, the abandonment of allografting, as some have suggested, appears rather premature even in newly diagnosed myeloma patients. Conversely, the role of allografting should become a matter of sound scientific debate in the myeloma community. Autologous transplantation undoubtedly remains the standard of care for younger medically fit myeloma patients. Three recent large studies that included an autograft followed by maintenance therapy with lenalidomide reported overall survivals of 85% at 3 years, and 73% and 81% at 4 years, respectively.^{2, 3, 4} However, there is no clear evidence of cure and in a subset of high-risk patients prognosis remains dismal.

In the early 2000s, as in the study by Ahmad *et al.*, a number of trials employed the frontline tandem approach with an autograft followed by reduced-intensity or non-myeloablative allografts.⁵

6, 7, 8, 9, 10, 11, 12 Some prospective trials also compared allografting with autografting using a biological assignment based on the availability of a HLA-identical donor. Substantial differences in inclusion criteria and treatment schemas partly contributed to conflicting outcomes. One meta-analysis, which included six clinical trials containing 1192 newly diagnosed patients who received tandem auto–auto and 630 who received a tandem auto-non-myeloablative allo, showed that the rates of CR were higher in the auto–allo group.¹³ However, this did not translate into a survival advantage in the first 3 years or after 3 years from assignment given the higher non-relapse mortality in the auto–allo group. Follow-up duration greatly varied among studies. Of note, the survival advantage in the auto–allo group, reported in two comparative studies, became statistically significant after a follow-up of at least 3 years. Importantly, a new meta-analysis on the same trials with updated long-term clinical outcomes has been planned (Costa L., personal communication).

All these studies were, however, designed well before potent anti-myeloma agents such as bortezomib, carfilzomib or lenalidomide and pomalidomide became readily available. Although the annual reports of the European Society of Blood and Marrow Transplantation show an important transplant activity in plasma cell disorders in recent years, the real role of the combination of new drugs with graft-versus-myeloma has never been fully explored in this incurable disease. This may partly be attributed to the current limited interest on cell therapy strategies in myeloma. New drugs and graft-versus-myeloma are not mutually exclusive, and their synergy has clearly been shown in relapsed patients.¹⁴

Patient-risk stratification and depth of response have become important factors to evaluate prognosis and guide treatment decisions in hematological malignancies. In multiple myeloma, the recently Revised International Staging System by the International Myeloma Working Group showed that there is a subset of patients where overall and PFSs are very poor even in the era of new drugs.¹⁵ In this patient subset, new effective treatments should be sought. The negative prognostic impact of high-risk cytogenetics appeared to be partly neutralized by graft-versus-myeloma in two recent studies.^{12, 16}

Depth of response is another key factor to predict clinical outcomes in myeloma. Thanks to advanced molecular technology, PCR-based minimal residual disease (MRD) detection currently represents a powerful outcome predictor. In the GIMEMA-VEL-03-096 trial, 39 patients who received consolidation with a combination bortezomib-thalidomide-dexamethasone (VTD) after an autograft were monitored for MRD by qualitative nested-PCR and quantitative PCR (qPCR). After a median follow-up of 8 years, overall survival was 72% for patients in major MRD response versus 48% for those with MRD persistence ($P=0.041$).¹⁷ This was a hallmark study that clearly showed that molecular remissions could be obtained with new drug-based consolidation treatments. However, the same group recently reported on the long-term outcomes of molecular monitoring after a tandem ‘auto-non-myeloablative allo’ approach.¹⁸ Twenty-six patients were prospectively evaluated by both nested-PCR and qPCR. Specific markers were generated in 19/26. At the median follow-up of 12.1, overall survival for the whole-patient cohort was not reached and event-free survival was 4 years from the allograft. Rate of nested-PCR negativity increased up to 47% at 1 year after the allograft, whereas the molecular response rate was 63% at 2 years by qPCR. At a remarkable median follow-up of 12.1, median overall survival and event-free survival were not reached in patients who achieved nested-PCR negativity while they were 3.3 and 1.5 years, respectively, in the remaining patients. Graft-versus-myeloma appeared to determine prolonged molecular remission rates, higher than those reported after autografting and VTD. Although the number of patients studied was small, one may provocatively infer that, had these findings been obtained with other treatment modalities, they would have been emphasized more in the scientific community.

In summary, despite the recent dramatic improvement in survival, the overwhelming majority of myeloma patients invariably relapse. Given the potentially curative effect of graft-versus-myeloma, it may become ethical to evaluate its combination with potent anti-myeloma agents in young high-risk and/or early-relapsed patients where life expectancy is currently very poor.

Conflict of interest

The author declare no conflict of interest.

References

1. Ahmad I, LeBlanc R, Cohen S, Lachance S, Kiss T, Sauvageau G *et al.* Favorable long-term outcome of patients with multiple myeloma using a frontline tandem approach with autologous and non-myeloablative allogeneic transplantation. *Bone Marrow Transplant* (e-pub ahead of print 21 December 2015; doi:10.1038/bmt.2015.319).
2. McCarthy PL, Owzar K, Hofmeister CC, Hurd DD, Hassoun H, Richardson PG *et al.* Lenalidomide after stem-cell transplantation for multiple myeloma. *N Engl J Med* 2012; 366: 1770–1781.
3. Attal M, Lauwers-Cances V, Marit G, Caillot D, Moreau P, Facon T *et al.* Lenalidomide maintenance after stem-cell transplantation for multiple myeloma. *N Engl J Med* 2012; 366: 1782–1791.
4. Palumbo A, Cavallo F, Gay F, Di Raimondo F, Ben Yehuda D, Petrucci MT *et al.* Autologous transplantation and maintenance therapy in multiple myeloma. *N Engl J Med* 2014; 371: 895–905.
5. Bruno B, Rotta M, Patriarca F, Rotta M, Sorasio R, Allione B *et al.* A comparison of allografting with autografting for newly-diagnosed myeloma. *N Engl J Med* 2007; 356: 1110–1120.
6. Moreau P, Garban F, Attal M, Michallet M, Marit G, Hulin C *et al.* Long-term follow-up results of IFM99-03 and IFM99-04 trials comparing nonmyeloablative allotransplantation with autologous transplantation in high-risk de novo multiple myeloma. *Blood* 2008; 112: 3914–3915.
7. Bruno B, Rotta M, Patriarca F, Mattei D, Allione B, Carnevale-Schianca F *et al.* Non-myeloablative allografting for newly diagnosed multiple myeloma: the experience of the Gruppo Italiano Trapianti di Midollo. *Blood* 2009; 113: 3375–3382.
8. Rotta M, Storer BE, Sahebi F, Shizuru JA, Bruno B, Lange T *et al.* Long-term outcome of patients with multiple myeloma after autologous hematopoietic cell transplantation and nonmyeloablative allografting. *Blood* 2009; 113: 3383–3391.
9. Giaccone L, Storer B, Patriarca F, Rotta M, Sorasio R, Allione B *et al.* Long-term follow-up of a comparison of nonmyeloablative allografting with autografting for newly diagnosed myeloma. *Blood* 2011; 117: 6721–6727.
10. Krishnan A, Pasquini MC, Logan B, Stadtmauer EA, Vesole DH, Alyea E 3rd *et al.* Tandem autologous versus single autologous transplantation followed by allogeneic hematopoietic cell transplantation for patients with multiple myeloma: results from the blood and marrow transplant clinical trials network (BMT CTN) 0102 trial. *Lancet Oncol* 2011; 12: 1195–2003.
11. Bjorkstrand B, Iacobelli S, Hegenbart U, Gruber A, Greinix H, Volin L *et al.* Tandem autologous/reduced-intensity conditioning allogeneic stem-cell transplantation versus autologous transplantation in myeloma: long-term follow-up. *J Clin Oncol* 2011; 29: 3016–3022.

12. Gahrton G, Iacobelli S, Björkstrand B, Hegenbart U, Gruber A, Greinix H *et al.* Autologous/reduced-intensity conditioning allogeneic stem cell transplantation versus autologous transplantation in multiple myeloma: long term results of the EBMT-NMAM2000 study. *Blood* 2013; 121: 5055–5063.
13. Armeson KE, Hill EG, Costa LJ. Tandem autologous vs autologous plus reduced intensity allogeneic transplantation in the upfront management of multiple myeloma: meta-analysis of trials with biological assignment. *Bone Marrow Transplant* 2013; 48: 562–567.
14. Palumbo A, Avet-Loiseau H, Oliva S, Lokhorst HM, Goldschmidt H, Rosinol L *et al.* Revised International Staging System for Multiple Myeloma: A Report From International Myeloma Working Group. *J Clin Oncol* 2015; 33: 2863–2869. |
15. Festuccia M, Martino M, Ferrando F, Messica G, Moscato T, Fedele R *et al.* Allogeneic stemcell transplantation in multiple myeloma: immunotherapy and new drugs. *Expert Opin Biol Ther* 2015; 15: 857–872.
16. Kröger N, Badbaran A, Zabelina T, Ayuk F, Wolschke C, Alchalby H *et al.* Impact of high-risk cytogenetics and achievement of molecular remission on long-term freedom from disease after autologous-allogeneic tandem transplantation in patients with multiple myeloma. *Biol Blood Marrow Transplant* 2013; 19: 398–404.
17. Ferrero S, Ladetto M, Drandi D, Cavallo F, Genuardi E, Urbano M *et al.* Long-term results of the GIMEMA VEL-03-096 trial in MM patients receiving VTD consolidation after ASCT: MRD kinetics' impact on survival. *Leukemia* 2015; 29: 689–695.
18. Ladetto M, Ferrero S, Drandi D, Festuccia M, Patriarca F, Mordini N *et al.* Prospective molecular monitoring of minimal residual disease after non-myceloablative allografting in newly diagnosed multiple myeloma. *Leukemia* (e-pub ahead of print 6 October 2015;