

Intrathecal or intraventricular colistin: a review

Olivia Bargiacchi¹, Francesco Giuseppe De Rosa²

¹Infectious Diseases Section, "Maggiore della Carità" Hospital, Novara, Italy;

²Department of Medical Sciences, University of Turin. Infectious Diseases Section at Amedeo di Savoia Hospital, Turin, Italy

SUMMARY

Central Nervous System (CNS) infections related to external ventricular derivation are a major complication of patients undergoing neurosurgical procedures. Antimicrobial treatment of CNS infections should be based not only on the susceptibility of the isolated microorganism, but also on the treatment's pharmacokinetic properties demonstrating the passage of the molecule through the blood-brain barrier. When CNS infections

are caused by multi-drug resistant Gram-negative bacteria, intrathecal colistin is considered an effective and safe option. We review the literature of intrathecal/intraventricular use of colistin, comprehensive of both pharmacokinetic data and clinical experiences.

Keywords: colistin, central nervous system infection, intrathecal, intraventricular, pharmacokinetics.

INTRODUCTION

Meningitis and or ventriculitis may be a significant complication in patients undergoing craniotomy and the incidence is up to 8% in patients with external catheter derivation (EVD) [1, 2]. Coagulase-negative staphylococci are predominant amongst Gram-positive and, amongst Gram-negative rods, *Acinetobacter baumannii* has an incidence of 3.6-11% [3]. Risk factors for infection are represented by leakage of the cerebrospinal fluid (CSF), concomitant infection at the time of incision, prolonged or repeated surgery and prolonged ventricular drainage, surgery through the sinuses as well as severity of underlying condition [4, 5]. Aetiology by pseudomonas and carbapenemase-producing Enterobacteriaceae is even more complex and difficult to treat [6-9].

The choice and kinetics of antimicrobials in central nervous system (CNS) infections is of utmost importance and the blood-brain barrier (BBB) has specific features that limit the passage of mol-

ecules on several parameters such as size, lipophilia, plasma protein binding and transporter affinity [10]. In patients with acute bacterial meningitis the CNS transfer of drugs may be increased due to opening of intracellular tight junctions, increased CSF outflow resistance, low pH and also by transporters stimulated by proinflammatory cytokines. There is a poor CSF penetration of beta-lactams ($AUC_{CSF}/AUC_{serum} < 0.15$) that increases with inflammation, without requiring intraventricular administration. Other antibiotics such as aminoglycosides, glycopeptides and polymyxins have a very low therapeutic index, with low CSF concentrations and therefore require lumbar, intrathecal or intraventricular (ITH/IVT) administration; only the latter ensure distribution in the whole CSF space.

Pharmacokinetic of colistin

Colistimethate (CSM) is eliminated by the kidneys and has a half-life of 2.7-4.5 hours; after a standard intravenous (IV) administration of 5 mg/Kg/day, the average plasma concentration is 8.1 mg/mL. The few data available on the pharmacokinetics of polymyxins in CSF reported a low diffusion through the CNS, with unclear rate of penetration and a possible role for meningeal inflammation

Corresponding author

Francesco Giuseppe De Rosa

E-mail: francescogiuseppe.derosa@unito.it

[11]. When colistin concentrations were measured by microbiological assay, the low CNS concentrations were not enhanced by meningeal inflammation [12]. Methodological issues in studies before 2009 may be responsible for the low concentration of both colistin and CMS, in a context in heterogeneous patient populations composed by children, adults, neurosurgical patients with EVD and variable amount of daily CSF drainage, with or without meningeal inflammation. Moreover, standard dosage of IV administration of colistin was not universally applied, as we did in a series of three patients [13, 14].

Jimenez Mejitas et al. reported CSF concentrations as high as 25% of serum concentration, with an IV infusion of 1,000,000 IU every 6 hours by a broth-dilution assay of polymyxins, using *Escherichia coli* strain ATCC 25922, whilst Markantonis et al. used high-performance liquid chromatography (HPLC) in five non-consecutive critically ill adult patients receiving intravenous CMS at variable dosages from 150 mg daily to 225 mg three times a day (1,875,000-2,812,500 IU): three patients had meningitis without considerable meningeal inflammation [11, 15, 16]. Serum and CSF samples were collected after two days of treatment by lumbar puncture in four patients and from EVD in one patient, with a ratio of the area under the concentration-time curve (AUC) for CSF-serum of 0.051. It is interesting that the AUC of colistin in CSF were parallel with simultaneous peaks and similar same half-life. According to these results, the conclusion was that the CSF penetration was only of 5%, consistently with previous studies performed by microbiological assays, suggesting a low possibility of eradication of infections by Gram-negative bacteria [12]. Antachopoulos et al. used liquid chromatography tandem mass spectrometry (LC-MS-MS) to study the CMS concentrations in three paediatric patients with EVD [17]. Intravenous colistin had low CSF concentrations below 0.2 mcg/mL and higher (34% to 67% of sera concentrations) levels of colistin were observed with meningeal inflammation.

According to these studies, IV colistin alone does not provide CSF concentrations high enough to reach a MIC of 2 mcg/mL for MDR Gram-negative rods and therefore topical administration is needed to support the treatment. Only two studies described the PK of ITH/IVT colistin: Imberti et al. studied by HPLC and by one-compartment

model colistin CSF in nine patients only treated with IVT colistin at variable dosages, from 2.61 to 10.44 mg (32,000-130,000 IU) [18]. The *C-trough* concentration ranged between 2,0 and 9,7 mcg/mL and the CSF concentration were always >2 mcg/mL when colistin was administered at a dose >65,200 IU/day, with a great inter-patients variability, perhaps due to the unknown fractions of CMS metabolized to colistin, the diffusion from the CSF to the systemic circulation and the cerebral tissue and the elimination through the external efflux of CSF, in addition to the unpredictable amount of the CSF spontaneously drained through the EVD by the fluctuation of intracranial pressure. Therefore it is not surprising that the ITH/IVT dose of 125,000 IU/day suggested by the IDSA guidelines seems to be more appropriate [19].

Twelve patients treated with IVT-IV colistin (3,000,000 IU every 8 hours and IVT dose of 125,000 IU daily) and EVD were studied by HPLC by Ziaka, of which five were treated only IV, three only by IVT administration and four patients by IVT-IV. The CSF concentrations in the IV group were very low, with a ratio of 0.07 [20]. There was no accumulation of colistin concentration over time, the mean CSF/serum concentration was higher in IVT-IV combination than in the other groups and the mean AUC CSF/AUC serum ratios were 60% higher in patients with ventriculitis than in control patients. Perhaps a greater penetration could be achieved with meningeal inflammation but a median CSF concentration >0.5 mcg/mL was only observed in the combination group, suggesting that IV colistin could increase CSF concentrations of ITH/IVT colistin; the conclusion was that combined treatment had a better chance to eradicate severe CNS infections [20].

Clinical data

Intrathecal/Intraventricular (IVT/ITH) colistin administration was associated with better outcomes compared to IV colistin alone [21-23]. There are guidelines available such as those by the Infectious Diseases Society of America (IDSA) in 2004 that suggested a daily ITH dosage of 10 mg (125,000 IU) [19]. However, in clinical practice, the dose is often chosen empirically, ranging between 1.6 and 40 mg (20,000-500,000 IU) either as a single dose or in divided doses. We synthesized in Table 1 the main data collected from all

Table 1 - Reported cases of Gram-negative CNS infections in adults treated with intrathecal or intraventricular colistin. Dosages reported in the original paper are converted in International Units (1 mg of colistin equals 30,000 IU of colistin and 1 mg of colistimethate equals 12.500 IU of colistin). *ITH*: intrathecal, *IVT*: intraventricular, *IV*: intravenous, *NR*: not reported, *qd*: once daily, *bid*: twice a day.

| Author, year, reference | Case no. | Microorganisms | ITH, IVT Colistin dose (IU), timing, (days of treatment) | IV Colistin | Other antibiotics | Infection outcome |
|--------------------------------|----------|---|--|-------------|----------------------------|-------------------|
| Price, 1972 [24] | 5 | <i>K. aerogenes</i> | IVT 50,000 bid → qd (NR) | Yes | gentamicin | 2/5 Cure |
| Kaplan, 1990 [25] | 1 | <i>A. baumannii</i> | ITH NR (20) | No | cefotaxime, aminoglycoside | Cure |
| Fernandez-Viladrich, 1999 [26] | 2 | <i>A. baumannii</i> | ITH 62,000-125,000 bid (17-19) | No | tobramycin | Cure |
| Vasen, 2000 [27] | 1 | <i>A. baumannii</i> | ITH 62,500→125,000 qd (21) | No | No | Cure |
| Benifla, 2004 [28] | 1 | <i>A. baumannii</i> | IVT 40,000 qd (17) | No | ampicillin/sulbactam | Cure |
| Quinn, 2005 [29] | 2 | <i>P. aeruginosa</i> | ITH 125,000 qd (14-16) | No | No | Cure |
| Suarez Fernandez, 2005 [30] | 7 | <i>A. baumannii</i> | IVT ITH 300,000 bid (8-21) | Yes | No | 5/7 Cure |
| Gump, 2005 [31] | 1 | <i>P. aeruginosa</i> | ITH 250,000-125,000 qd (10) | Yes | No | Cure |
| Kasiakou, 2005 [32] | 1 | <i>A. baumannii</i> | IVT 20,000 qd (21)→ 40,000 bid (42) | Yes | amikacin | Relapse → Cure |
| Bukhary, 2005 [33] | 1 | <i>A. baumannii</i> | IVT 125,000 bid (21) | No | No | Cure |
| Sueke, 2005 [34] | 1 | <i>A. baumannii</i> | ITH 75,000 bid (14) | No | No | Cure |
| Berlana, 2005 [35] | 2 | <i>A. baumannii</i> , <i>P. aeruginosa</i> | ITH 125,000-250,000 qd (8-10) | Yes | No | Cure |
| Charra, 2005 [36] | 1 | <i>A. baumannii</i> | ITH 62,500-125,000 qd (21) | No | No | Cure |
| Karakitsos, 2006 [37] | 6 | <i>A. baumannii</i> | IVT 125,000-250,000 qd (16-20) | No | No | Cure |
| Yagmur, 2006 [38] | 1 | <i>P. aeruginosa</i> | ITH 62,500 qd (21) | No | No | Cure |
| Ng, 2006 [39] | 5 | <i>A. baumannii</i> | ITH IVT 62,500-125,000 qd (3-24) | Yes | amikacin | 4/5 Cure |
| Al Shirawi, 2006 [40] | 1 | <i>A. baumannii</i> | IVT 40,000 qd (28) | No | No | Cure |
| Motaouakkil, 2006 [41] | 1 | <i>A. baumannii</i> | ITH 62,000-125,000 qd (21) | No | rifampin | Cure |
| Schina, 2006 [42] | 1 | <i>P. aeruginosa</i> | ITH 20,000-60,000 qd (26) | Yes | ceftazidime, amikacin | Cure |
| Metan, 2007 [43] | 1 | <i>A. baumannii</i> | ITH 125,000 qd (21) | No | meropenem | Death |
| Enfanto, 2007 [44] | 1 | <i>K. pneumoniae</i> | ITH 125,000 qd (7) | No | vancomycin, meropenem | Cure |
| Ho, 2007 [45] | 1 | <i>A. baumannii</i> | ITH 80,000 qd (28) | Yes | No | Cure |
| Paramythiotou, 2007 [46] | 1 | <i>A. baumannii</i> | IVT 125,000 qd (20) | Yes | No | Cure |
| Lee, 2008 [47] | 1 | <i>A. baumannii</i> | IVT 62,500 qd (3) | Yes | meropenem, sulbactam | Cure |
| Rodriguez Guardado, 2008 [48] | 8 | <i>A. baumannii</i> | IVT 125,000 bid (21) | Yes | No | Cure |
| Hachimi, 2008 [49] | 1 | <i>A. baumannii</i> | IVT 62,000-125,000 qd (22) | No | amikacin | Cure |
| Rodriguez, 2009 [50] | 1 | <i>A. baumannii</i> | ITH 125,000 qd (NR) | No | rifampin | Cure |

Continues >>>

Continued from previous page >>>

| Author, year, reference | Case no. | Microorganisms | ITH, IVT Colistin dose (IU), timing, (days of treatment) | IV Colistin | Other antibiotics | Infection outcome |
|----------------------------|-----------------------|---|--|-------------|--|--------------------|
| Lopez-alvarez, 2009 [51] | 3 | <i>A. baumannii</i> | IVT 62,500-100,000 bid (9-29) | Yes | No | Cure |
| Dalgic, 2009 [52] | 1 | <i>A. baumannii</i> | IVT 12,500→125,000 qd (20) | Yes | No | Cure |
| Baiocchi, 2010 [53] | 1 | <i>P. aeruginosa</i> | ITH 125,000 qd (NR) | No | aztreonam, amikacin | Cure |
| De Pascale, 2010 [54] | 1 | <i>A. baumannii</i> | ITH 75,000-150,000 qd (22) | Yes | No | Cure |
| Khawcharoenporn, 2010 [55] | 7 | <i>A. baumannii</i> | ITH 300,000-500,000 qd (2-21) | No | imipenem, sulbactam, ciprofloxacin | 5/7 Cure |
| Cascio, 2010 [56] | 1 | <i>A. baumannii</i> | ITH 125,000 qd (10) | No | No | Cure |
| Kim, 2010 [57] | 1 | <i>A. baumannii</i> | ITH 62,500 bid (17) | Yes | rifampin | Cure |
| Ozdemir, 2010 [58] | 1 | <i>A. baumannii</i> | ITH 125,000 qd (35) | Yes | meropenem, sulbactam, amikacin | Cure |
| Razmkon, 2011 [59] | 6 | <i>A. baumannii</i> | IVT 125,000 qd (NR) | Yes | No | 3/6 Cure |
| Patel, 2011 [60] | 1 | <i>A. baumannii</i> | IVT 125,000 qd (36) | Yes | No | Cure |
| Saleem, 2011 [61] | 5 | <i>A. baumannii</i> | ITH 4,800/kg-12,000/kg qd (5-21) | Yes | No | 4/5 Cure |
| Alaoui, 2011 [62] | 1 | <i>A. baumannii</i> | ITH 48,000 qd (NR) | Yes | No | Cure |
| Wang, 2012 [63] | 4 | <i>A. baumannii</i> | IVT 25,000-200,000 bid or qd (7-27) | Yes | meropenem, sulbactam, imipenem, | 3/4 Cure |
| Imberti, 2012 [18] | 9 | <i>A. baumannii</i> (2) <i>K. pneumoniae</i> (6) <i>P. aeruginosa</i> (1) | IVT 31,250-62,500 qd – bid (11-36) | Yes (5/9) | rifampin, meropenem, linezolid, sulbactam, amikacin, ceftazidime, fosfomicin, vancomycin | 8/9 Cure |
| Ziaka, 2013 [20] | 7 | <i>A. baumannii</i> (3) <i>K. pneumoniae</i> (2) Negative cultures (2) | IVT 125,000 qd (NR) | 4/7 Yes | NR | 4/5 Cure |
| Karaiskos, 2013 [64] | 6 | <i>A.baumannii</i> | IVT 500,000→125,000 qd→ every 48h (12-21) | Yes | No | Cure |
| Bargiacchi, 2014 [14] | 3 | <i>A. baumannii</i> (1) <i>K. pneumoniae</i> (1) <i>P. aeruginosa</i> (1) | IVT 250,000 qd (14-20) | Yes | rifampin, tigecycline | Cure |
| De Bonis, 2015 [23] | 9 | <i>A. baumannii</i> | IVT 125,000-250,000 qd (14-40) | Yes | NR | Cure |
| Shofty, 2016 [22] | 22 (+15 ITH amikacin) | <i>A. baumannii</i> (51% in whole) | ITH/IVT 50,000-250,000 qd (median duration 9 days) | Yes | NR | 5/37 in whole Cure |
| Fotakopoulos, 2016 [21] | 23 | <i>A. baumannii</i> (15) <i>K. pneumoniae</i> (2) <i>P. aeruginosa</i> (8) <i>E. cloacae</i> (1) | 150,000-200,000 qd (8-24) | Yes | NR | 20/23 Cure |

the published cases and case series with IVT/ITH colistin [14, 18, 20-23, 24-64].

In a recent paper, Karaiskos and coll. proposed an interesting schedule of IVT administration with a loading dose of 500,000 IU followed by a dose of 125,000-250,000 IU every 24h, followed by a q48h ITH administration after the CSF culture becomes negative, to reduce the risk of chemical meningitis or ventriculitis and the numbers of manipulations of the EVD [64]. However, in published studies, where ITH/IVT colistin was mainly used for infections by *A. baumannii*, there was no difference in the cure rate and the time to CSF sterilization with daily ITH/IVT dosages higher or lower than 125,000 IU. Moreover, this dosage should be high

enough to treat even *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. Figure 1 shows that a cumulative dose above 1,750,000 IU of ITH/IVT colistin (which correspond to 125,000 IU once daily for 14 days) is associated to better outcomes.

The duration of therapy described is quite variable, from one to four weeks. Shorter treatments (less than a week) seem to correlate with a higher mortality, although in critical patients death may occur for other underlying conditions before the conclusion of the scheduled treatment (Figure 2). By converse, higher dosages may be administered in more severe patients, with a consistent bias toward higher and early mortality in those patients (Figures 3 and 4). An important issue influencing the duration of

Figure 1 - Cure rate according to cumulative dose of intrathecal/intraventricular colistin. The cumulative dose of 1.750.000 correspond to 125.000 IU of colistin once daily (dose suggested by IDSA Guidelines) for 14 days.





Figure 2 - Cure rates according to duration of treatment. In brackets the number of patients, whose duration of treatment was defined.






Figure 3 - Cure rates according to duration of treatment and dose of intrathecal colistin for those patients whose duration and dose of treatment was defined.




Figure 4 - Cure rates according to dose of intrathecal/intraventricular colistin. In brackets the number of patients, whose duration of treatment was defined.

treatment is the time to the first negative culture suggesting that a fast clearance of the infection may result in a better outcome with an earlier chance to change an infected catheter or to replace the external catheter, an essential factor to destroy the bacterial biofilm adhering to the device.

It is unclear which role IV colistin may play on the outcome of the CNS infection, when ITH/IVT colistin is used. Therefore, if the source of infection is strictly related to the colonization of the EVD and the infection is limited to the CNS, ITH/IVT treatment may be sufficient to sterilize CSF, but if ventriculitis is the result of a bacteraemia or there are other sites infected, IV colistin should be added to ITH/IVT colistin and perhaps a second

systemic antimicrobial, chosen according to the susceptibility and synergy tests [65].

Combination antimicrobial treatment is reported in many cases choosing antibiotics according to the susceptibility tests. Rifampin has optimal CNS diffusion and may have a synergistic effect with colistin against Gram negative bacteria, such as *A. baumannii*, *P. aeruginosa* and, perhaps, KPC [18, 41, 50, 57, 66-69].

Side effects are reported: seizures, controlled by anticonvulsant therapy (1 case), chemical meningitis (5 cases), and *cauda equina* syndrome (1 case), with complete resolution after ITH administration discontinuation [39, 51, 57, 64]. No nephrotoxicity is reported with ITH/IVT administration.

■ CONCLUSIONS

PK studies and clinical data suggest that intravenous colistin alone is not sufficient to treat meningitis and ventriculitis caused by Gram-negative multidrug resistant bacteria. Intrathecal or intraventricular colistin has proved to be effective and safe. The suggested dose of 125,000 IU by IDSA Guidelines is still valid if administered once daily for at least 14 days. IV and IVT colistin (at least 125,000 IU/die) for at least 10-14 days are probably required in most patients with CNS infections.

■ REFERENCES

- [1] Laxmi S., Tunkel A.R. Healthcare-associated bacterial meningitis. *Curr. Infect. Dis. Rep.* 13, 367-373, 2011.
- [2] Lozier A.P., Sciacca R.R., Romagnoli M.F., Connolly E.S. Jr. Ventriculostomy-related infections: a critical review of the literature. *Neurosurgery* 62, 688-700, 2008.
- [3] Kim B.N., Peleg A.Y., Lodise T.P., et al., Management of meningitis due to antibiotic-resistant *Acinetobacter* species. *Lancet Infect. Dis.* 9, 245-255, 2009.
- [4] Kim J.H., Desai N.S., Ricci J. et al. Factors contributing to ventriculostomy infection. *World Neurosurg.* 77, 135-140, 2012.
- [5] Gutiérrez-González R., Boto G.R., Pérez-Zamarrón A. Cerebrospinal fluid diversion devices and infection. A comprehensive review. *Eur. J. Clin. Microbiol. Infect. Dis.* 31, 889-897, 2012.
- [6] Ece G., Samlioglu P., Atalay S., Kose S. Evaluation of the in vitro colistin susceptibility of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* strains at a tertiary care centre in Western Turkey. *Infez. Med.* 1, 36-40, 2014.
- [7] Tascini C., Menichetti F., Stefanelli A., Loni C., Lambellet P. Clinical efficacy of intravenous colistin therapy in combination with ceftazidime in severe MDR *P. aeruginosa* systemic infections in two haematological patients. *Infez. Med.* 1, 41-44, 2006.
- [8] Bassetti M., Righi E., Esposito S., Petrosillo N., Nicolini L. Drug treatment for multidrug-resistant *Acinetobacter baumannii* infections. *Future Microbiology.* 3, 6, 649-660, 2008.
- [9] Esposito S., Esposito I., Leone S. Considerations of antibiotic therapy duration in community- and hospital-acquired bacterial infections. *J. Antimicrob. Chemother.* 67, 11, 2570-2575, 2012.
- [10] Nau R., Sörgel F., Eiffert H. Penetration of drugs through the blood-cerebrospinal fluid/blood-brain barrier for treatment of central nervous system infections. *Clin. Microbiol. Rev.* 23, 858-883, 2010.
- [11] Jiménez-Mejías M.E., Pichardo-Guerrero C., Márquez-Rivas F.J., Martín-Lozano D., Prados T., Pachón J. Cerebrospinal fluid penetration and pharmacokinetic/pharmacodynamic parameters of intravenously administered colistin in a case of multidrug-resistant *Acinetobacter baumannii* meningitis. *Eur. J. Clin. Microbiol. Infect. Dis.* 21, 212-214, 2002.
- [12] Everett E.D., Strausbaugh L.J. Antimicrobial agents and the central nervous system. *Neurosurgery.* 6, 691-714, 1980.
- [13] Plachouras D., Karvanen M., Friberg L.E. Population pharmacokinetic analysis of colistin methanesulfonate and colistin after intravenous administration in critically ill patients with infections caused by gram-negative bacteria. *Antimicrob. Agents Chemother.* 53, 3430-3436, 2009.
- [14] Bargiacchi O., Rossati A., Car P., et al. Intrathecal/intraventricular colistin in external ventricular device-related infections by multi-drug resistant Gram negative bacteria: case reports and review. *Infection.* 42, 801-809, 2014.
- [15] Jiménez-Mejías M.E., Becerril B., Márquez-Rivas F.J., Pichardo C., Cuberos L., Pachón J. Successful treatment of multidrug-resistant *Acinetobacter baumannii* meningitis with intravenous colistin sulfomethate sodium. *Eur. J. Clin. Microbiol. Infect. Dis.* 19, 970-971, 2000.
- [16] Markantonis S.L., Markou N., Fousteri M., et al. Penetration of colistin into cerebrospinal fluid. *Antimicrob. Agents Chemother.* 53, 4907-4910, 2009.
- [17] Antachopoulos C., Karvanen M., Iosifidis E., et al. Serum and cerebrospinal fluid levels of colistin in pediatric patients. *Antimicrob. Agents Chemother.* 54, 3985-3987, 2010.
- [18] Imberti R., Cusato M., Accetta G., et al. Pharmacokinetics of colistin in cerebrospinal fluid after intraventricular administration of colistin methanesulfonate. *Antimicrob. Agents Chemother.* 56, 4416-4421, 2012.
- [19] Tunkel A.R., Hartman B.J., Kaplan S.L., et al. Practice guidelines for the management of bacterial meningitis. *Clin. Infect. Dis.* 39, 1267-1284, 2004.
- [20] Ziaka M., Markantonis S.L., Fousteri M., et al. Combined intravenous and intraventricular administration of colistin methanesulfonate in critically ill patients with central nervous system infection. *Antimicrob. Agents Chemother.* 57, 1938-1940, 2013.
- [21] Fotakopoulos G., Makris D., Chatzi M., Tsimitrea E., Zakyntinos E., Fountas K. Outcomes in meningitis/ventriculitis treated with intravenous or intraventricular plus intravenous colistin. *Acta Neurochir. (Wien).* 158: 603-610, 2016.
- [22] Shofty B., Neuberger A., Naffaa M.E., et al. Intrathecal or intraventricular therapy for post-neurosurgical Gram-negative meningitis: matched cohort study. *Clin. Microbiol. Infect.* 22, 66-70, 2016.
- [23] De Bonis P., Lofrese G., Scoppettuolo G., et al. Intraventricular versus intravenous colistin for the treatment of extensively drug resistant *Acinetobacter baumannii* meningitis. *Eur. J. Neurol.* 23, 68-75, 2016.

- [24] Price D.J., Sleight J.D. Klebsiella meningitis-report of nine cases. *J. Neurol. Neurosurg. Psychiatry.* 35, 903-908, 1972.
- [25] Kaplan S.L., Patrick C.C. Cefotaxime and aminoglycoside treatment of meningitis caused by gram-negative enteric organisms. *Pediatr. Infect. Dis. J.* 9, 810-814, 1990.
- [26] Fernandez-Viladrich P., Corbella X., Corral L., Tubau F., Mateu A. Successful treatment of ventriculitis due to carbapenem-resistant *Acinetobacter baumannii* with intraventricular colistin sulfomethate sodium. *Clin. Infect. Dis.* 28(4), 916-917, 1999.
- [27] Vasen W., Desmery P., Ilutovich S., Di Martino A., Intrathecal use of colistin. *J. Clin. Microbiol.* 38, 3523, 2000.
- [28] Benifla M., Zucker G., Cohen A., Alkan M. Successful treatment of *Acinetobacter meningitis* with intrathecal polymyxin E. *J. Antimicrob. Chemother.* 54, 290-292, 2004.
- [29] Quinn A.L., Parada J.P., Belmares J., O'Keefe J.P. Intrathecal colistin and sterilization of resistant *Pseudomonas aeruginosa* shunt infection. *Ann. Pharmacother.* 39, 949-952, 2005.
- [30] Suárez Fernández C., Fernández-Viladrich P., Tubau F., Corral L., Mateu A. Intrathecal colistin treatment of central nervous system infections due to multi-resistant *Acinetobacter baumannii*. In *Program and Abstracts of 15th European Congress of Clinical Microbiology and Infectious Diseases Copenhagen, Denmark, April 2-5, 2005*, Abstract number: 1134_02_277.
- [31] Gump W.C., Walsh J.W. Intrathecal colistin for treatment of highly resistant *Pseudomonas ventriculitis*. Case report and review of the literature. *J. Neurosurg.* 102, 915, 2005.
- [32] Kasiakou S.K., Rafailidis P.I., Liaropoulos K., Falagas M.E. Cure of post-traumatic recurrent multiresistant Gram-negative rod meningitis with intraventricular colistin. *J. Infect.* 50, 348-352, 2005.
- [33] Bukhary Z., Mahmood W., Al-Khani A., Al-Abdeley H.M. Treatment of nosocomial meningitis due to a multidrug resistant *Acinetobacter baumannii* with intraventricular colistin. *Saudi Med. J.* 26(4), 656-658, 2005.
- [34] Sueke H., Marsh H., Dhital A., Using intrathecal colistin for multidrug resistant shunt infection. *Br. J. Neurosurg.* 19, 51-52, 2005.
- [35] Berlana D., Llop J.M., Fort E., Badia M.B., Jódar R. Use of colistin in the treatment of multiple-drug-resistant gram-negative infections. *Am. J. Health Syst. Pharm.* 62, 39-47, 2005.
- [36] Charra Charra B., Hachimi A., Benslama A., Motaouakkil S. Intrathecal use of colistin. *Ann. Fr. Anesth. Reanim.* 25, 215, 2006.
- [37] Karakitsos D., Paramythiotou E., Samonis G., Karabinis A. Is intraventricular colistin an effective and safe treatment for post-surgical ventriculitis in the intensive care unit? *Acta Anaesthesiol. Scand.* 50, 1309-1310, 2006.
- [38] Yagmur R., Esen F. Intrathecal colistin for treatment of *Pseudomonas aeruginosa ventriculitis*: report of a case with successful outcome. *Crit. Care.* 10, 428, 2006.
- [39] Ng J., Gosbell I.B., Kelly J.A., Boyle M.J., Ferguson J.K. Cure of multiresistant *Acinetobacter baumannii* central nervous system infections with intraventricular or intrathecal colistin: case series and literature review. *J. Antimicrob. Chemother.* 58, 1078-1081, 2006.
- [40] Al Shirawi N., Memish Z.A., Cherfan A., Al Shimermeri A. Post-neurosurgical meningitis due to multidrug-resistant *Acinetobacter baumannii* treated with intrathecal colistin: case report and review of the literature. *J. Chemother.* 18, 554-558, 2006.
- [41] Motaouakkil S., Charra B., Hachimi A., et al. Colistin and rifampin in the treatment of nosocomial infections from multiresistant *Acinetobacter baumannii*. *J. Infect.* 53, 274-278, 2006.
- [42] Schina M., Spyridi E., Daoudakis M., Mertzanos E., Korfiatis S. Successful treatment of multidrug-resistant *Pseudomonas aeruginosa meningitis* with intravenous and intrathecal colistin. *Int. J. Infect. Dis.* 10, 178-179, 2006.
- [43] Metan G., Alp E., Aygen B., Sumerkan B., *Acinetobacter baumannii meningitis* in post-neurosurgical patients: clinical outcome and impact of carbapenem resistance. *J. Antimicrob. Chemother.* 60, 197-199, 2007.
- [44] Enfanto C.M., Shin S. Treatment with Intrathecal Colistin of a Resistant Extended-Spectrum beta-Lactamase-Producing *Klebsiella pneumoniae* in the Cerebrospinal Fluid. *Infect. Dis. Clin. Pract.* 15, 274-277, 2007.
- [45] Ho Y.H., Wang L.S., Chao H.J., Chang K.C., Su C.F. Successful treatment of meningitis caused by multidrug-resistant *Acinetobacter baumannii* with intravenous and intrathecal colistin. *J. Microbiol. Immunol. Infect.* 40, 537-540, 2007.
- [46] Paramythiotou E., Karakitsos D., Aggelopoulou H., Sioutos P., Samonis G., Karabinis A. Post-surgical meningitis due to multiresistant *Acinetobacter baumannii*. Effective treatment with intravenous and/or intraventricular colistin and therapeutic dilemmas. *Med. Mal. Infect.* 37, 124-125, 2007.
- [47] Lee C.H., Tang Y.F., Su L.H., Chien C.C., Liu J.W. Antimicrobial effects of varied combinations of meropenem, sulbactam, and colistin on a multidrug-resistant *Acinetobacter baumannii* isolate that caused meningitis and bacteremia. *Microb. Drug Resist.* 14, 233-237, 2008.
- [48] Rodríguez Guardado A., Blanco A., Asensi V., et al. Multidrug-resistant *Acinetobacter meningitis* in neurosurgical patients with intraventricular catheters: assessment of different treatments. *J. Antimicrob. Chemother.* 61, 908-913, 2008.
- [49] Hachimi A., Charra B., Benslama A., Motaouakkil S. Intrathecal use of amikacin and colistin. *Ann. Fr. Anesth. Reanim.* 27, 113-114, 2008.
- [50] Rodríguez C.H., Bombicino K., Granados G., Nas-

- tro M., Vay C., Famiglietti A. Selection of colistin-resistant *Acinetobacter baumannii* isolates in postneurosurgical meningitis in an intensive care unit with high presence of heteroresistance to colistin. *Diagn. Microbiol. Infect. Dis.* 65, 188-191, 2009.
- [51] López-Alvarez B., Martín-Láez R., Fariñas M.C., Paternina-Vidal B., García-Palomo J.D., Vázquez-Barquero A. Multidrug-resistant *Acinetobacter baumannii* ventriculitis: successful treatment with intraventricular colistin. *Acta Neurochir. (Wien)*. 51, 1465-1472, 2009.
- [52] Dalgic N., Ceylan Y., Sancar M. et al. Successful treatment of multidrug-resistant *Acinetobacter baumannii* ventriculitis with intravenous and intraventricular colistin. *Ann. Trop. Paediatr.* 29, 141-147, 2009.
- [53] Baiocchi M., Catena V., Zago S., Badolati L., Baccharin M. Intrathecal colistin for treatment of multidrug resistant (MDR) *Pseudomonas aeruginosa* after neurosurgical ventriculitis. *Infez. Med.* 18, 182-186, 2010.
- [54] De Pascale G., Pompucci A., Maviglia R., et al. Successful treatment of multidrug-resistant *Acinetobacter baumannii* ventriculitis with intrathecal and intravenous colistin. *Minerva Anestesiol.* 76, 957-960, 2010.
- [55] Khawcharoenporn T., Apisarnthanarak A., Mundy L.M. Intrathecal colistin for drug-resistant *Acinetobacter baumannii* central nervous system infection: a case series and systematic review. *Clin. Microbiol. Infect.* 16, 888-894, 2010.
- [56] Cascio A., Conti A., Sinardi L., et al. Post-neurosurgical multidrug-resistant *Acinetobacter baumannii* meningitis successfully treated with intrathecal colistin. A new case and a systematic review of the literature. *Int. J. Infect. Dis.* 14, e572-579, 2010.
- [57] Kim K., Kang H.S., Sohn C.H., Oh B.M. Cauda equina syndrome misdiagnosed as aggravated hydrocephalus: neurological complication of intrathecal colistin in post-surgical meningitis. *Acta Neurochir. (Wien)*. 153, 425-427, 2011.
- [58] Ozdemir H., Tapisiz A., Ciftçi E., et al. Successful treatment of three children with post-neurosurgical multidrug-resistant *Acinetobacter baumannii* meningitis. *Infection.* 38, 241-244, 2010.
- [59] Razmkon A., Mehrafshan A., Bakhtzad A. Management of post-neurosurgical *Acinetobacter* infections: experiences obtained during an outbreak. *Acta Neurochir. (Wien)*. 153, 435-436, 2011.
- [60] Patel J.A., Pacheco S.M., Postelnick M., Sutton S. Prolonged triple therapy for persistent multidrug-resistant *Acinetobacter baumannii* ventriculitis. *Am. J. Health Syst. Pharm.* 68, 1527-1531, 2011.
- [61] Saleem A.F., Shah M.S., Shaikh A.S., Mir F., Zaidi A.K. *Acinetobacter* species meningitis in children: a case series from Karachi, Pakistan. *J. Infect. Dev. Ctries.* 5, 809-814, 2011.
- [62] Alaoui S.Y., Nejmi S.E., Chakir A.A., Hmamouchi B., Chlilek A. Intraventricular colistin use in neonatal meningitis caused by *Acinetobacter baumannii*. *Ann. Fr. Anesth. Reanim.* 30, 854-855, 2011.
- [63] Wang J.H., Lin P.C., Chou C.H., et al. Intraventricular antimicrobial therapy in postneurosurgical Gram-negative bacillary meningitis or ventriculitis: A hospital-based retrospective study. *J. Microbiol. Immunol. Infect.* 47, 204-210, 2014.
- [64] Karaiskos I., Galani L., Baziaka F., et al. Successful treatment of extensively drug-resistant *Acinetobacter baumannii* ventriculitis and meningitis with intraventricular colistin after application of a loading dose: a case series. *Int. J. Antimicrob. Agents.* 41, 480-483, 2013.
- [65] Tumbarello M., Viale P., Viscoli C., et al. Predictors of mortality in bloodstream infections caused by *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae*: importance of combination therapy. *Clin. Infect. Dis.* 55, 943-950, 2012.
- [66] Liang W., Liu X.F., Huang J., Zhu D.M., Li J., Zhang J. Activities of colistin- and minocycline-based combinations against extensive drug resistant *Acinetobacter baumannii* isolates from intensive care unit patients. *BMC Infect. Dis.* 27, 11, 109, 2011.
- [67] Tan T.Y., Lim T.P., Lee W.H., Sasikala S., Hsu L.Y., Kwa A.L. In vitro antibiotic synergy in extensively drug-resistant *Acinetobacter baumannii*: the effect of testing by time-kill, checkerboard, and Etest methods. *Antimicrob. Agents Chemother.* 55, 436-438, 2011.
- [68] Tascini C., Gemignani G., Ferranti S., et al. Microbiological activity and clinical efficacy of a colistin and rifampin combination in multidrug-resistant *Pseudomonas aeruginosa* infections. *J. Chemother.* 16, 282-287, 2004.
- [69] Tascini C., Tagliaferri E., Giani T., et al. Synergistic activity of colistin plus rifampin against colistin-resistant KPC-producing *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 57, 3990-3993, 2013.