

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: A MINDMAPS study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1567891> since 2016-06-19T10:05:28Z

Published version:

DOI:10.1097/PSY.000000000000174

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Psychosomatic medicine, 77(4), 2015, doi: 10.1097/PSY.000000000000174

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://journals.lww.com/psychosomaticmedicine/pages/articleviewer.aspx?year=2015&issue=05000&article=00009&type=abstract>

Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study

Running head: Sex differences in depression post-MI

Frank Doyle PhD¹, Hannah McGee PhD², R Conroy DSc³, Elisabeth H. Bos PhD⁴, Anna Meijer MSc⁵, Richard Steeds MD⁶, Hiroshi Sato PhD⁷, Donna Stewart MD⁸, Kapil Parakh PhD⁹, Robert Carney PhD¹⁰, Kenneth Freedland PhD¹¹, Matteo Anselmino PhD¹², Roxanne Pelletier PhD¹³, Henk Jan Conradi PhD¹⁴, Peter de Jonge PhD¹⁵.

Manuscript word count: 6752 (entire paper), including 4 Tables and 1 Figure

Affiliations

1. Frank Doyle, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
2. Hannah McGee, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
3. R Conroy, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

4. Elisabeth H. Bos, University Medical Center Groningen: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
5. Anna Meijer MSc, University Medical Center Groningen: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
6. Richard Steeds, Queen Elizabeth Hospital: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
7. Hiroshi Sato, Kwansai Gakuin University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
8. Donna Stewart, University of Toronto: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
9. Kapil Parakh, Johns Hopkins University School of Medicine: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
10. Robert Carney, Washington University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

11. Kenneth Freedland, Washington University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
12. Matteo Anselmino, University of Turin: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
13. Roxanne Pelletier, McGill University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
14. Henk Jan Conradi, University of Amsterdam: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
15. Peter de Jonge, University Medical Center Groningen: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

Corresponding author:

Frank Doyle, PhD

Division of Population Health Sciences (Psychology),

Royal College of Surgeons in Ireland,

123 St Stephen's Green,

Dublin 2,

Ireland

Email: fdoyl4@csi.ie, Tel: + 353 1 4022718, Fax: + 353 1 4022764

Funding: A.M., H.J.C., and P. de J. have support from a Vidi grant from the Dutch Medical Research Council (grant no. 016.086.397)

Conflict of interest disclosures

Disclosures and grant support

Frank Doyle, PhD (No conflicts of interest or disclosures)

Hannah McGee PhD (No conflicts of interest or disclosures)

R Conroy DSc (No conflicts of interest or disclosures)

Elske H. Bos PhD (No conflicts of interest or disclosures)

Anna Meijer MSc (No conflicts of interest or disclosures)

Richard Steeds MD (No conflicts of interest or disclosures)

Hiroshi Sato PhD (supported by a grant-in-aid for University and Society Collaboration (15590743) from the Japanese Ministry of Education, Culture, Sports, Science, and Technology (Tokyo, Japan) and by a research grant from the Japan Arteriosclerosis Prevention Fund (Tokyo, Japan))

Donna Stewart MD (Potential conflicts are a grant from Canadian Institute of Health Research on Gender, Mental Health and Violence, and membership on the Duloxetine and Pregnancy Registry Scientific Advisory Board of Eli Lilly Inc. I also received travel support from Ranbaxy Pharma for a psychiatric meeting in Thailand)

Kapil Parakh PhD (No conflicts of interest or disclosures)

Robert Carney PhD (Dr. Carney or a member of his family own stock in Pfizer Inc and Forest Laboratories.)

Kenneth Freedland PhD (No conflicts of interest or disclosures)

Matteo Anselmino PhD (No conflicts of interest or disclosures)

Roxanne Pelletier PhD (no conflict of interest to disclose and I am funded by the Canadian Institutes of Health Research)

Henk Jan Conradi PhD (No conflicts of interest or disclosures)

Peter de Jonge PhD (No conflicts of interest or disclosures)

Keywords

*Keywords*ó Depression; myocardial infarction; gender differences; prognosis; individual patient data meta-analysis

The MIND-MAPS Group:

E.H. Bos*, H.J. Conradi**, P. de Jonge*, A. Meijer*, J. Ormel*

* Interdisciplinary Center Psychopathology and Emotion regulation, University Medical Center Groningen, 9713 GZ, Groningen, The Netherlands ** Department of Clinical Psychology, University of Amsterdam, 1018 XA, Amsterdam, The Netherlands

S. Bergerone, M. Anselmino

Division of Cardiology, Department of Medical Sciences, University of Turin, Città della Salute e della Scienza Hospital, Turin, Italy

L.F. Berkman*, R. Carney**, K.E. Freedland**

* Centre for Population and Development Studies, Harvard University, Boston, Massachusetts 02138, USA, ** Department of Psychiatry, Washington University, St. Louis, Missouri 63108, USA

J. Denollet

Centre of Research on Psychology in Somatic Diseases, Tilburg University, 5000 LE, Tilburg, The Netherlands

F. Doyle*, H.M. McGee*, Ronán Conroy**

* Division of Population Health Sciences (Psychology), Royal College of Surgeons in Ireland, Dublin 2, Ireland; ** Division of Population Health Sciences (Epidemiology & Public Health Medicine), Royal College of Surgeons in Ireland, Dublin 2, Ireland

S.L. Grace

Women's Health Program, University Health Network; Faculty of Medicine, University of Toronto; School of Kinesiology and Health Science, York University, Toronto, Ontario M3J1P3, Canada

S.H. Hosseini

Psychiatry and Science Behavior Research Centre, Department of Psychiatry, Mazandaran University of Medical Sciences, Sari, 48171-19164, Iran

D.A. Lane

University of Birmingham Centre for Cardiovascular Sciences, City Hospital, Birmingham B18 7QH, UK

L. Pilote

Divisions of Epidemiology and Cardiology, McGill University Health Centre, Montréal H3A1A1, Canada

K. Parakh, R.C. Ziegelstein

Department of Medicine, Johns Hopkins University School of Medicine, Johns Hopkins Bayview Medical Centre, Baltimore, Maryland 21224, USA

C. Rafanelli

Department of Psychology, University of Bologna, 40127 Bologna, Italy

H. Sato

School of Human Welfare Studies, Kwansei Gakuin University, Nishinomiya, 662-8501 Japan

R.P. Steeds

Department of Cardiology, Queen Elizabeth Hospital, Birmingham B15 2TH, UK

B.D. Thombs

Lady Davis Institute for Medical Research, Jewish General Hospital and McGill University, Montréal H3T 1E4, Canada

C. Welin

College of Health Sciences, Department of Nursing, Göteborg University, SE-405 30 Göteborg, Sweden

Abstract

Objective Using combined individual patient data (IPD) from prospective studies, we explored sex differences in depression and prognosis post-myocardial infarction (MI), and determined whether disease indices could account for found differences.

Methods Meta-analysis of IPD from 10,175 MI patients who completed diagnostic interviews or depression questionnaires from 16 prospective studies of MI patients, identified by systematic review for the MINDMAPS study. Multilevel logistic and Cox regression models were used to determine sex differences in prevalence of depression and sex-specific effects of depression on prognosis.

Results Combined interview and questionnaire data from observational studies showed that 36% (635/1760) of women and 29% (1575/5526) of men reported elevated levels of depression (age-adjusted OR=0.68, 95% CI 0.60 to 0.77, $p<0.001$). The risk for all-cause mortality associated with depression was higher in men (HR=1.38, 95% CI 1.30 to 1.47, $p<0.001$) than in women (HR=1.22, 95% CI 1.14 to 1.31, $p<0.001$). Low left ventricular ejection fraction (LVEF) was associated with higher depression scores in men only (sex*LVEF interaction B=0.294, 95% CI 0.090 to 0.498, $p=0.005$), which attenuated the sex difference in the association between depression and prognosis.

Conclusions The prevalence of depression post-MI was higher in women than men, but the association between depression and cardiac prognosis was worse for men. LVEF was associated with depression in men only, and accounted for the increased risk of all-cause mortality in depressed men versus women, suggesting that depression in men post-MI may in part reflect cardiovascular disease severity.

Keywords: Depression; myocardial infarction; gender; prognosis; individual data meta-analysis

Acronyms

ACE = angiotensin converting enzyme

BMI ó body mass index

CABG = coronary artery bypass graft

CI = confidence interval

CVE ó cardiovascular events

HF = heart failure

HR = Hazard ratio

IPD = Individual patient data

LVEF = Left ventricular ejection fraction

MI = myocardial infarction

OR = odds ratio

PTCA = percutaneous transluminal coronary intervention

SD = standard deviation

Introduction

Depression in people with myocardial infarction (MI) has been the source of considerable study. While both clinically diagnosed depression and questionnaire-assessed depressive symptoms have been consistently associated with poorer cardiovascular prognosis in MI patients (1), the treatment of depression has not been demonstrated to reduce morbidity or mortality in randomized trials (2, 3). Recent estimates suggest a prevalence of diagnosed major depressive disorder in the US of 2.1% to 4.9% in adults aged 45 and over (4), with women having a higher prevalence (4.8%) than men (3.3%). Similarly, in adults aged 45 years and above, estimates of elevated depressive symptoms are also more prevalent among women (7-12%) than men (5-7%) (5). Similar findings are available for European countries (6, 7), although the differing methodologies adopted by different countries make overall estimates for major depression more difficult. According to a systematic review of the literature, the prevalence of depression is significantly higher in patients with MI, with 20% having major depression and up to 31% reporting elevated depressive symptoms, although sex differences were not reported (8).

The etiology of depression post-MI has been the subject of considerable debate. While depressive symptoms and trajectories of these symptoms post-MI appear to have similar risk factors to depression in the general population (e.g. stressful life events, personality etc.) (9-13), sex is not always a predictor of depression in these studies. This suggests that there may be a potentially different etiology for depressive symptoms in men and women with MI. Indeed, other researchers have demonstrated that elevated levels of depression, and its subsequent association with cardiovascular prognosis, may be a consequence of

cardiovascular disease severity (14, 15). For example, poor left ventricular ejection fraction (LVEF) has been demonstrated to not only correlate with depression, but also to predict the onset of major depression and depressive symptoms in patients with MI (14). In addition, this study found that poorer LVEF was a stronger predictor of onset of major depression for men than for women (14), again suggesting potentially different etiology for depression between men and women post-MI.

We aimed to establish whether the prevalence of diagnosed depression and elevated depressive symptomatology is similar in men and women with MI, and whether the diagnosed depression or depressive symptoms had a greater impact on prognosis for men than for women. We also investigated the role of LVEF in the etiology of depression and prognosis according to patients' sex. These questions have not been subject to systematic investigation and previous studies were not large enough to detect potential moderating effects. We therefore used data from the MINDMAPS individual patient meta-analytic database of 16 studies (16) to determine the following:

- 1 ó whether cardiovascular disease severity indices account for differences in post-MI prevalence of depressive symptoms between men and women; and
- 2 ó whether the association between depression and prognosis is stronger in men than in women, and whether disease severity indices (e.g. LVEF) account for these differences.

Methods

MINDMAPS

Details of MINDMAPS are available elsewhere (16). In brief, eligibility for inclusion in the MINDMAPS dataset was determined by a systematic review of published studies on post-MI depression (within 3 months post-hospitalisation for MI, assessed by validated questionnaire or diagnostic interview) and two prognostic endpoints (all-cause mortality, or major cardiovascular events ó CVE: recurrent MI, unstable angina, CABG) (1).

Authors of all selected studies were contacted and original data was requested (covering demographics, depression, disease severity [*LVEF*<40%, *Killip class (I vs. II, III, IV), and history of MI*], comorbidities, diabetes, smoking, body mass index [BMI], medication use, and outcomes). Of 30 eligible studies, 16 provided data (16).

Statistical analysis

Stata 11 was used for the main analyses (Statacorp LP, TX, USA). Multilevel logistic regression, adjusting for age, was used to obtain odds ratios (ORs) for differences in depression status between men (1) and women (0), using a binary depression variable (based on interview or recommended scale thresholds). Hazard ratios (HRs) were calculated using multilevel Cox proportional hazards regression analysis, to predict time-to-event (all-cause mortality or CVE), using depression, sex, and the interaction between sex and depression as predictors, adjusting for age. A continuous depression variable was used in these Cox analyses (z-transformed total scale scores). As different studies used different methods and are likely to represent populations with different characteristics (such as date of study, differences in therapies and assessments used, etc.), observations

within studies are unlikely to be fully independent. Therefore, a random intercept for study was included, resulting in a multilevel model. When calculating the prevalence estimates we omitted data from the ENRICHHD randomized trial and ancillary studies (2, 17) (which were combined in the original study (16)), as these were enrolled under strict inclusion criteria whereas observational studies would be more inclusive. To demonstrate the random effect variance in respect to the depression prevalence, we displayed the intraclass correlation coefficient, which shows what part of the total variance in prevalence is between studies. To investigate whether cardiovascular disease indices and other risk factors accounted for any sex differences in depression, a linear mixed model with random intercept was used, with depression z-scores as the dependent variable, and sex, disease indices and their interaction as predictors, adjusting for age. For each disease index a separate model was built. A contrast coding of -0.5/0.5 for dichotomous variables was used in the Cox regression analyses to ensure between-study variances between groups were equal (e.g. female vs. male) (18). Separate models were built for each disease and risk factor index. Also, two more multivariate models were built including all disease and risk factors, but omitting LVEF, together as covariates, and then a final model including all disease and risk factors. To increase confidence interval robustness, and in view of the non-normal distribution of the depression z-scores, bootstrapping with 1000 replications was used in all analyses (19). The handling of the variables used for adjustment and the applied outcomes was identical to that in the main MINDMAPS study (16). All tests were two-tailed.

Results

Sample description

Of 30 eligible studies (16), 16 provided individual patient data. These studies were from 9 different countries. The description of the combined sample (N=10,175), stratified by sex, is displayed in Table 1:

Insert Table 1 here

Overall, the mean age was 61 years (range 20-97), 71% of participants were men, and mean follow-up time per study was 3.15 years (range 0.96 to 6.65).

Prevalence of depression

Data from four diagnostic interview studies (omitting the ENRICHD data, n=900) showed that 20% (47/240) of women and 12% (76/660) of men had major depression. The age-adjusted multilevel logistic regression showed this to be a significant difference (OR=0.42, 95% CI 0.27 to 0.65, p=0.001). There was little heterogeneity between these interview studies in depression prevalence: the intraclass correlation coefficient was 0.01 (Chi² = 1.4; p = 0.115).

When using the questionnaire studies, excluding data from the ENRICHD and ancillary studies, the sex differences in prevalence remained significant (age-adjusted OR=0.69,

95% CI 0.61 to 0.77, $p < 0.001$), with a higher prevalence of depressive symptoms in women (38%, 657/1741) than men (30%, 1622/5461). There was some heterogeneity between these questionnaire studies in depression prevalence: the intraclass correlation coefficient was 0.10 ($\text{Chi}^2(1) = 436.3$; $p < 0.001$).

Combining the interview and questionnaire studies, but omitting the ENRICHD data, resulted in 36% (635/1760) of women and 29% (1575/5526) of men reporting elevated depressive symptoms (age-adjusted OR=0.68, 95% CI 0.60 to 0.77, $p < 0.001$). The intraclass correlation coefficient was 0.14 ($\text{Chi}^2(1) = 547.7$; $p > 0.001$), indicating significant, though not large, heterogeneity between studies in depression prevalence.

We also checked whether the sex difference in depression prevalence varied with age. We did so by including an interaction term sex*age in the above analysis. The interaction was not significant (OR=1.00, 95% CI 0.99 to 1.01, $p = 0.91$). So, the sex difference in depression prevalence did not vary with age.

Disease severity

Various disease indices and risk factors were analysed to determine if these accounted for sex differences in depression scores. Interactions between sex and history of MI, Killip class, diabetes, smoking or BMI were not associated with depression z-scores (data not shown). However, an interaction between sex and left ventricular ejection fraction was observed (see the upper panel of Table 2):

Insert Table 2 here

This interaction remained significant when adjusting for all other disease factors and risk indices (see the lower panel of Table 2). The results suggest that for men only, low LVEF was associated with higher levels of depression.

Moderation effects of sex in the association between depression and prognosis

Survival curves for all-cause mortality and depression status, and cardiovascular events and depression status are shown in Fig 1:

Insert Fig 1 here

The survival curves show that depression was related to poorer prognosis, in both men and women, and that the effect is more important in men. This is further investigated in Table 3:

Insert Table 3 here

The Cox regression analyses for all-cause mortality showed that the interaction between sex and depression was statistically significant (HR for interaction=1.12; 95% CI: 1.05 to 1.19), suggesting that the association between depression and mortality was indeed stronger for men than for women. The HR associated with depression was 12% higher in men compared to women (men, HR=1.38, 95% CI 1.30 to 1.47, $p<0.001$; women, HR=1.22, 95% CI 1.14 to 1.31, $p<0.001$). Omitting data from the ENRICHHD studies had little effect on the interaction term (HR for interaction=1.10, 95% CI 1.02 to 1.19, $p=0.014$). The interaction effect for CVE pointed into the same direction, but did not reach significance.

We also checked whether the sex difference in the impact of depression on cardiovascular outcomes varied with age. We did so by including an interaction term $\text{sex*age*depression}$ in the above analyses. The interaction was not significant (all-cause mortality HR =1.00. 95% CI 0.99 to 1.01, $p = 0.92$; CVE: HR =1.00. 95% CI 0.99 to 1.00, $p = 0.59$). So, the sex difference in the impact of depression on cardiovascular outcomes did not vary with age. Finally, we determined if the moderating effect of sex in the association between depression and all-cause mortality could be explained by the cardiovascular disease indices or risk factors. The same multilevel Cox regression analyses were used, predicting all-cause mortality by age, depression, sex, and sex*depression . Then, the size of the change in the estimate for the interaction sex*depression was determined when each of the disease variables was included in these separate models, and in two other multivariate models. Results are shown in Table 4:

Insert Table 4 here

In the separate models, LVEF accounted for the largest reduction in the size of the sex*depression interaction, and inclusion of this variable resulted in a non-significant interaction. History of MI also accounted for a substantial reduction in the interaction, but most other factors increased instead of reduced the size of the interaction. These differences could be the result of differences between studies or sample sizes, therefore, the analyses were repeated with only those participants who had data on all six cardiovascular disease variables and risk indices (History of MI, LVEF, and Killip class, diabetes, smoking, BMI). The last rows of Table 4 show the results of the multivariate models. While adjusting for all covariate except LVEF (5172 participants in 5 studies) resulted in an 8% increase in the sex*depression interaction, inclusion of all six covariates in this subset (2226 participants in 3 studies) resulted in a 32.5% reduction of the sex*depression interaction effect, and resulted in a non-significant interaction. Together, these results suggest that the moderating effect of sex on the association between depression and all-cause mortality can partially be explained by differences in LVEF and history of MI among depressed men and women, and that LVEF explained the largest part.

Furthermore, despite there being no statistically significant sex*depression interaction for predicting CVE, the percentage reduction was again largest for LVEF, showing a consistency across outcomes (results not shown).

Discussion

In this largest study to date, we found a number of results that are important in interpreting the association between post-MI depression and prognosis. Firstly, low LVEF was associated with depressive symptoms, but only in men. Secondly, the association between depression and prognosis was stronger in men than in women, which could partly be explained by depressive symptoms being a reflection of low LVEF more often for men than for women. LVEF attenuated the interaction effect between depression and sex in the prediction of prognosis.

In general, the prevalence of depression by diagnostic interview and questionnaire studies are similar to that previously reported (8), with higher proportions of women having depression according to both diagnostic interview and questionnaire studies. This higher prevalence of depression in women, added to the fact that depression was associated with disease severity in men only, demonstrates how depression in this population mimics depression in the general population. Other studies have also demonstrated how psychosocial risk factors for depression are predictors not only of elevated post-MI depressive symptomatology, but also of trajectories of depression post-discharge (9-13, 20), suggesting that depressive symptoms cannot be attributed solely to coronary disease

severity. However, the higher prevalence of depression reported using scale thresholds rather than interview techniques again provides evidence that psychometric scales overestimate the prevalence of depression in this population, with our findings demonstrating 2.5- to 3-fold higher prevalence, as found previously (8). This higher prevalence by questionnaire probably represents confounding between somatic symptoms of depression and symptoms of coronary disease (14, 15, 21), which cannot easily be thoroughly investigated as may happen during a diagnostic interview. Conversely, it has been demonstrated that even low levels of depression predict cardiovascular prognosis (22), thus relying on diagnostic interviews may actually underestimate meaningful levels of depression. Furthermore, while it has previously been suggested that the higher prevalence of depression in women might lead to poorer cardiovascular prognosis (23), we have found no evidence for this. Indeed, depression was more strongly related to a worse cardiac prognosis in men than in women. Single studies may not have had the power to detect such effects.

Depression was associated with a poor LVEF in men but not in women. Poor LVEF has been associated with depression in patients with coronary disease (24, 25). Previously, van Melle et al. (14) have shown that in 1989 post-MI patients LVEF was inversely correlated with depression status (from diagnostic interviews) and depression scale scores at several time points in the year post-MI, even in those who did not demonstrate depression at baseline. Furthermore, men demonstrated a trend towards a stronger increase in rates of depression across more severe LVEF categories, which was confirmed in the present findings. The data from that sample were included in the present

analyses. However, that was a single study, while the present analyses are from 16 studies. That depression was more strongly related to prognosis for men may be explained by the differential nature of MI in women and men. Women with MI tend to be older by approximately 10 years, have more diffuse coronary disease, have non-ST-elevation MI thereby not being eligible for emergency primary percutaneous coronary intervention (26) which is protective of subsequent lowering of LVEF (27-29), and therefore have worse hospital outcomes (26, 27). Men tend to have more focal lesions that result in ST-elevation MI. In the current data, women were less likely to receive percutaneous coronary intervention. As a result, women are more likely to end up with heart failure with preserved ejection fraction while men end up with heart failure with reduced ejection fraction (27-29). Thus, the differential attenuation by LVEF of the association of depression with outcomes may be partly explained by the fact that women are less likely to have the type of MI that results in a low LVEF. Further research is needed to clarify this possibility. Alternatively, the etiology of depression in post-MI patients may differ in women and men. Men may be more likely to have depression as a result of coronary disease severity, while women may more often have pre-existing depression (30). These sex differences are perhaps unlikely to be mediated by other biological (e.g. reduced cardiac autonomic tone, increased platelet aggregation, increased inflammatory processes) (31, 32) or behavioral (e.g. smoking, physical inactivity etc.) (32-34) mechanisms linking depression to MI, as no consistent links with men or women have been reported. Recent research also suggests that fetal exposure may have increased the risk of both depression and cardiovascular disease, and that this effect is stronger in women (35), although this requires further investigation. It is unclear why there was not a

significant moderating effect of sex in the association between depression and CVE. One reason may be that CVE is a less well-defined endpoint across the studies, and this may lead to measurement error, or that there were different numbers and types of studies and participants analysed for this outcome.

Limitations and strengths

The heterogeneity of study populations, depression, cardiovascular risk factor measurements, and endpoint assessments may be problematic, although the inclusion of the random intercept accounts for study differences. Measures of distress have been shown to demonstrate measurement invariance across sex and ethnic groups in a large cross-sectional study (36). All-cause mortality includes non-cardiac deaths, and may bias the results. The heterogeneity of cardiovascular risk factors reported in individual studies meant that full adjustment could not be made in the overall sample. Inclusion of variables such as New York Heart Association categorisation, diastolic function, blood pressure, and cholesterol were not included in the dataset, but might have changed the results. Similarly, psychiatric history was unavailable, as was alcohol abuse, and other non-cardiac co-morbidities or their treatments may also impact on the findings. Not all relevant studies were available for analysis (16), which may have biased the results, although most did not report sex differences or did not find any differences in prognosis .

Another major limitation of the present research, and indeed other studies in this area, is the lack of a plausible model that could explain the sex differences seen in this

manuscript. Other researchers have rightly called for more clarity in the analysis of observational studies which investigate risk factors in disease etiology (37, 38). To establish any single variable as a true independent risk factor, confounder or mediator for a given outcome, knowledge of temporal precedence of variables is required (37, 38). This is extremely difficult in the case of cardiovascular disease and depression. In the present analyses we cannot, for example, assume that depression precedes LVEF, or vice versa, as this data simply was not available to us. We have therefore avoided the terms *mediate* or *confound* etc. Further research with repeated measures of important variables is required before true risk factors and mediators of sex differences in depression and prognosis post-MI are elucidated.

The major strength of the study is the use of individual patient data, which allows for time-to-event analysis, and adjustment for potentially confounding variables systematically across studies, which cannot be done in summary meta-analyses, resulting in more accurate estimates of effect sizes, increased statistical power, generalizability and reliability of results (39). The inclusion of the random intercept term allows important between-study differences to be accounted for statistically, such as changing acute treatments (percutaneous coronary intervention has superseded thrombolysis in the timeframe spanned by the present studies) (27, 28), or differential assessment of LVEF δ which may impact on its prognostic ability (40, 41). The ability to analyse both interview and questionnaire studies is a strength, and the results demonstrated consistent patterns across these methods.

Conclusions

Using IPD from a large sample, the present study revealed significant differences in the prevalence, possible etiology and consequences of depression post-MI in men versus women. The prevalence of depression post-MI was higher in women than men, but the association between depression and cardiac prognosis was worse for men. LVEF was associated with depression in men only, and accounted for the increased risk of all-cause mortality in depressed men, suggesting that depression in men post-MI may in part reflect cardiovascular disease severity. Future studies, including intervention studies, should account for these sex differences.

References

1. Meijer A, Conradi HJ, Bos EH, Thombs BD, van Melle JP, de Jonge P. Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis of 25 years of research. *Gen Hosp Psychiatry*. 2011;33:203-16.
2. The ENRICHD Investigators. Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. *JAMA*. 2003;289:3106-16.
3. van Melle JP, de Jonge P, Honig A, Schene AH, Kuyper AM, Crijns HJ, Schins A, Tulner D, van den Berg MP, Ormel J. Effects of antidepressant treatment following myocardial infarction. *Br J Psychiat*. 2007;190:460-6.
4. Centers for Disease Control and Prevention. Morbidity and Mortality Weekly Report: Erratum. 2010 [cited 2012 08/06/2012]; Available from: http://www.cdc.gov/Features/dsDepression/Revised_Table_Estimates_for_Depression_MMWR_Erratum_Feb%202011.pdf.
5. Centers for Disease Control and Prevention. QuickStats: Prevalence of Current Depression* Among Persons Aged ≥ 12 Years, by Age Group and Sex—United States, National Health and Nutrition Examination Survey, 2007-2010. National Center for Health Statistics, 2012 Jan 6.
6. Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jonsson B, Olesen J, Allgulander C, Alonso J, Faravelli C, Fratiglioni L, Jennum P, Lieb R,

Maercker A, van Os J, Preisig M, Salvador-Carulla L, Simon R, Steinhausen HC. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharm* 2011;21:655-79.

7. Van de Velde S, Bracke P, Levecque K. Gender differences in depression in 23 European countries. Cross-national variation in the gender gap in depression. *Soc Sci Med*. 2010;71:305-13.

8. Thombs BD, Bass EB, Ford DE, Stewart KJ, Tsilidis KK, Patel U, Fauerbach JA, Bush DE, Ziegelstein RC. Prevalence of depression in survivors of acute myocardial infarction. *J Gen Intern Med*. 2006;21:30-8.

9. Rieckmann N, Burg MM, Gerin W, Chaplin WF, Clemow L, Davidson KW. Depression vulnerabilities in patients with different levels of depressive symptoms after acute coronary syndromes. *Psychotherap Psychosom*. 2006;75:353-61.

10. Doyle F, McGee H, Conroy R, Delaney M. What predicts depression in cardiac patients: Sociodemographic factors, disease severity or theoretical vulnerabilities? *Psychol Health*. 2011;26:619-34.

11. Doyle F, McGee HM, Delaney M, Motterlini N, Conroy RM. Depressive vulnerabilities predict depression status and trajectories of depression over one year in persons with acute coronary syndrome. *Gen Hosp Psychiat*. 2011;33:224-31.

12. Kaptein KI, de Jonge P, van den Brink RH, Korf J. Course of depressive symptoms after myocardial infarction and cardiac prognosis: a latent class analysis. *Psychosom Med*. 2006;68:662-8.

13. Martens EJ, Smith OR, Winter J, Denollet J, Pedersen SS. Cardiac history, prior depression and personality predict course of depressive symptoms after myocardial infarction. *Psychol Med.* 2008;38:257-64.
14. van Melle JP, de Jonge P, Ormel J, Crijns HJ, van Veldhuisen DJ, Honig A, Schene AH, van den Berg MP. Relationship between left ventricular dysfunction and depression following myocardial infarction: data from the MIND-IT. *Eur Heart J.* 2005;26:2650-6.
15. Nicholson A, Kuper H, Hemingway H. Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J.* 2006;27:2763-74.
16. Meijer A, Conradi HJ, Bos EH, Anselmino M, Carney R, Denollet J, Doyle F, Freedland KE, Grace SL, Hosseini SH, Lane DA, Pilote L, Parakh K, Rafanelli C, Sato H, Steeds RP, Welin C, de Jonge P. Adjusted prognostic association of post-myocardial infarction depression with mortality and cardiovascular events: an individual patient data meta-analysis. *Br J Psychiat.* 2013;203:90-102.
17. Carney RM, Blumenthal JA, Stein PK, Watkins L, Catellier D, Berkman LF, Czajkowski SM, O'Connor C, Stone PH, Freedland KE. Depression, heart rate variability, and acute myocardial infarction. *Circulation.* 2001;104:2024-8.
18. Smith CT, Williamson PR, Marson AG. An overview of methods and empirical comparison of aggregate data and individual patient data results for investigating heterogeneity in meta-analysis of time-to-event outcomes. *J Eval Clin Pract.* 2005;11:468-78.

19. Sauerbrei W, Schumacher M. A bootstrap resampling procedure for model building: application to the Cox regression model. *Stat Med.* 1992;11:2093-109.
20. Sorensen C, Brandes A, Hendricks O, Thrane J, Friis-Hasche E, Haghfelt T, Bech P. Psychosocial predictors of depression in patients with acute coronary syndrome. *Acta Psychiatr Scand.* 2005;111:116-24.
21. Doyle F, Conroy R, McGee H. Challenges in reducing depression-related mortality in cardiac populations: cognition, emotion, fatigue or personality? *Health Psychol Rev.* 2007;1:137-72.
22. Bush DE, Ziegelstein RC, Tayback M, Richter D, Stevens S, Zahalsky H, Fauerbach JA. Even minimal symptoms of depression increase mortality risk after acute myocardial infarction. *Am J Cardiol.* 2001;88:337-41.
23. Naqvi TZ, Naqvi SS, Merz CN. Gender differences in the link between depression and cardiovascular disease. *Psychosom Med.* 2005;67 Suppl 1:S15-8.
24. Lehto S, Koukkunen H, Hintikka J, Viinamaki H, Laakso M, Pyorala K. Depression after coronary heart disease events. *Scandinav Cardiovasc J.* 2000;34:580-3.
25. Freedland KE, Rich MW, Skala JA, Carney RM, Davila-Roman VG, Jaffe AS. Prevalence of depression in hospitalized patients with congestive heart failure. *Psychosom Med.* 2003;65:119-28.
26. Milcent C, Dormont B, Durand-Zaleski I, Steg PG. Gender differences in hospital mortality and use of percutaneous coronary intervention in acute

myocardial infarction: microsimulation analysis of the 1999 nationwide French hospitals database. *Circulation*. 2007;115:833-9.

27. Poon S, Goodman SG, Yan RT, Bugiardini R, Bierman AS, Eagle KA, Johnston N, Huynh T, Grondin FR, Schenck-Gustafsson K, Yan AT. Bridging the gender gap: Insights from a contemporary analysis of sex-related differences in the treatment and outcomes of patients with acute coronary syndromes. *Am Heart J*. 2012;163:66-73.

28. Gulati M, Shaw LJ, Bairey Merz CN. Myocardial ischemia in women: lessons from the NHLBI WISE study. *Clin Cardiol*. 2012;35:141-8.

29. Shaw LJ, Bugiardini R, Merz CN. Women and ischemic heart disease: evolving knowledge. *J Am Coll Cardiol*. 2009;54:1561-75.

30. Kendler KS, Gardner CO, Fiske A, Gatz M. Major depression and coronary artery disease in the Swedish twin registry: phenotypic, genetic, and environmental sources of comorbidity. *Arch Gen Psychiat*. 2009;66:857-63.

31. Carney RM, Freedland KE, Miller GE, Jaffe AS. Depression as a risk factor for cardiac mortality and morbidity: a review of potential mechanisms. *J Psychosom Res*. 2002;53:897-902.

32. Ormel J, de Jonge P. Unipolar depression and the progression of coronary artery disease: toward an integrative model. *Psychother Psychosom*. 2011;80:264-74.

33. Ziegelstein RC, Fauerbach JA, Stevens SS, Romanelli J, Richter DP, Bush DE. Patients with depression are less likely to follow recommendations to

reduce cardiac risk during recovery from a myocardial infarction. *Arch Intern Med.* 2000;160:1818-23.

34. Doyle F, Rohde D, Kowalczyk A, Morgan K, Cousins G, McGee H. Systematic review and meta-analysis of the impact of depression on subsequent smoking cessation in patients with coronary heart disease: 1990–2013. *Psychosom Med.* 2014;76:44-57.

35. Goldstein JM, Cherkerzian S, Buka SL, Fitzmaurice G, Hornig M, Gillman M, O'Toole S, Sloan RP. Sex-specific impact of maternal-fetal risk factors on depression and cardiovascular risk 40 years later. *J Develop Origin Health Disease.* 2011;2:353-64.

36. Clark CJ, Henderson KM, de Leon CF, Guo H, Lunos S, Evans DA, Everson-Rose SA. Latent constructs in psychosocial factors associated with cardiovascular disease: an examination by race and sex. *Front Psychiat.* 2012;3:5.

37. Christenfeld NJ, Sloan RP, Carroll D, Greenland S. Risk factors, confounding, and the illusion of statistical control. *Psychosom Med.* 2004;66:868-75.

38. Kraemer HC, Stice E, Kazdin A, Offord D, Kupfer D. How do risk factors work together? Mediators, moderators, and independent, overlapping, and proxy risk factors. *Am J Psychiat.* 2001;158:848-56.

39. Stewart LA, Tierney JF. To IPD or not to IPD? Advantages and disadvantages of systematic reviews using individual patient data. *Eval Health Prof.* 2002;25:76-97.

40. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelisa A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocard.* 2009;10:165-93.

41. Schepis T, Gaemperli O, Koepfli P, Valenta I, Strobel K, Brunner A, Leschka S, Desbiolles L, Husmann L, Alkadhi H, Kaufmann PA. Comparison of 64-slice CT with gated SPECT for evaluation of left ventricular function. *J Nuclear Med.* 2006;47:1288-94.

Table 1: Baseline characteristics by sex

	Men (n=7200)	Women (n=2975)	p-value	N measured
Demographic characteristics				
Age (mean, (SD))	59.7 (11.5)	64.1 (12.2)	<0.001***	10,171
Employment status (% employed)	53.6	27.0	<0.001***	6,528
Partner status (% with partner)	77.5	51.4	<0.001***	6,412
Cardiac disease severity				
History of MI (% yes)	19.2	18.0	<0.001***	9,646
LVEF (% of patients <40%)	22.6	22.4	0.565	3,505
Killip class (% poor)	16.1	22.9	<0.001***	7,532
PTCA (%)	47.2	42.1	<0.001***	7,679
History of PTCA (%)	11.4	11.0	0.113	4,830
CABG (%)	9.8	9.3	<0.001***	8,139
History of CABG (%)	10.8	7.9	<0.001***	4,849
Thrombolysis (%)	33.8	34.0	0.061	8,065
Congestive HF (%)	17.2	24.3	0.013*	6,104
Other risk factors				
Diabetes (%)	18.0	29.8	<0.001***	10,060
Smoking (%)	49.5	32.8	<0.001***	9,942
BMI (mean, (SD))	26.8 (4.5)	27.8 (6.0)	0.002**	7,188
Hyperlipidemia / hypercholesterolemia (%)	45.3	50.2	<0.001***	8,405
Hypertension (%)	31.6	38.6	<0.001***	8,301

History of hypertension (%)	43.1	62.2	<0.001***	5,348
Medication use				
Hypolipidemics (%)	43.5	44.1	0.291	4,004
Beta-blockers (%)	72.3	71.2	0.003**	8,833
Aspirin (%)	87.2	87.2	0.490	7,561
Calcium-channel blockers / antagonists (%)	14.5	21.3	<0.001***	7,056
ACE-inhibitors (%)	48.7	49.8	0.258	8,550
Antidepressant use (%)	5.8	8.1	0.013*	5,507
Study type				
Diagnostic interview (n)	660	240		900
Questionnaire (n)	7108	2942		10,050

ACE: angiotensin converting-enzyme; BMI: body mass index; CABG: coronary artery bypass graft; HF: heart failure;

LVEF: left ventricular ejection fraction; MI: myocardial infarction; PTCA: percutaneous transluminal coronary angioplasty; SD: standard deviation

***p < 0.001; **p < 0.01; *p < 0.05

Table 2: Linear mixed model with random intercept for study, predicting depression z-score (n=3115, 5 studies)

	B	Bootstrapped 95% CI	p
Adjusted for age (n=3115, 5 studies)			
Age	-0.001	-0.004 ó 0.002	0.571
LVEF	-0.015	-0.191 ó 0.161	0.869
Sex	-0.371	-0.476 ó -0.265	<0.001***
Sex*LVEF	0.294	0.090 ó 0.498	0.005**
Adjusted for all confounders (n=2226, 3 studies)			
Age	-0.002	-0.008 ó 0.003	0.363
History of MI	0.148	0.064 ó 0.233	0.001**
Killip class	0.132	0.029 ó 0.235	0.012*
Diabetes	0.145	0.059 ó 0.231	0.001**
Smoking	0.057	0.024 ó 0.090	0.001**
BMI	-0.011	-0.028 ó 0.006	0.207
LVEF	-0.019	-0.089 ó 0.052	0.605
Sex	-0.348	-0.426 ó -0.269	<0.001***
Sex*LVEF	0.274	0.061 ó 0.488	0.012*

***p < 0.001; **p < 0.01; *p < 0.05. Sex (0=female, 1=male)

Table 3: Multilevel Cox proportional hazards regression assessing sex differences in the association between depression and cardiovascular prognosis

	All-cause mortality (n=7628, 10 studies)			Cardiovascular events (CVE) (n=6556, 7 studies)		
	HR	Bootstrapped 95% CI	P	HR	Bootstrapped 95% CI	p
Age	1.07	1.06 ó 1.08	< 0.001***	1.01	1.01 ó 1.02	< 0.001***
Depression z-score	1.30	1.24 ó 1.37	< 0.001***	1.17	1.13 ó 1.22	< 0.001***
Sex	1.07	0.90 ó 1.28	0.441	0.93	0.78 ó 1.11	0.420
Sex * depression z-score	1.12	1.05 ó 1.19	< 0.001***	1.07	0.98 ó 1.17	0.118

***p < 0.001; **p < 0.01; *p < 0.05. Sex (-0.5=female, 0.5=male)

Table 4: Percentage change in sex*depression interaction effects in the prediction of all-cause mortality by additional disease indices or risk factors

Factor	HR for interaction sex*depression	Adjusted HR for interaction sex*depression	% change	N/n studies
History of MI	1.097***	1.088**	-8.9%	7543/10
LVEF	1.126*	1.086	-30.5%	3115/5
Killip class	1.137***	1.162***	+16.9%	5924/6
Diabetes	1.107***	1.130***	+20.2%	7587/10
Smoking	1.118**	1.119**	+0.8%	7485/10
BMI	1.128**	1.125**	-2.2%	6133/7
Adjusting for all covariates except LVEF	1.140***	1.152**	+8.0%	5172/5
Adjusting for all covariates	1.107**	1.071	-32.5%	2226/3

Adjusted for age, depression z-score, and sex, with study as a random factor

Note: The percentage change was estimated by using the log HR for the sex*depression interaction effect.

***p < 0.001; **p < 0.01; *p < 0.05

Figure Legends

Figure 1: Survival curves for all-cause mortality (ACM) and cardiovascular events (CVE), by depression category and sex (adjusted for age). depr = depression. Depression categories are based on a split-half of the continuous depression z-scores on self-report questionnaires. (Note: In all Cox regression analyses, the continuous depression z-score is used.)

