AperTO - Archivio Istituzionale Open Access dell'Università di Torino

LIVER DAMAGE CAN BE ASSOCIATED WITH DEREGULATION OF THE DE NOVO LIPOGENESIS PATHWAY IN SUBJECTS WITH NON ALCOHOLIC FATTY LIVER DISEASE

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/157807> since 2017-02-06T17:15:15Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

Digestive and Liver Disease, Volume 47, Supplement 1, 2015:e5

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://www.dldjournalonline.com/issue/S1590-8658%2814%29X0020-4>

Liver damage can be associated with deregulation of the de novo lipogenesis pathway in subjects with Non Alcoholic Fatty Liver Disease

C. Rosso¹, C. Saponaro^{2,3}, L. Mezzabotta¹, E. Vanni¹, R. Gambino¹, F. Saba¹, R. Ibrahim Kamal Jouness¹, M. Gaggini^{2,4}, E. Buzzigoli², GP Caviglia¹, ML Abate¹, F. Salomone⁵, A. Smedile¹, M. Rizzetto¹, M. Cassader¹, A. Gastaldelli², E. Bugianesi¹

¹*Department of Medical Sciences, University of Torino, Turin,*

²*Cardiometabolic Risk Unit, CNR-Institute of Clinical Physiology, Pisa, Italy*

³*Dottorato Pegaso Regione Toscana in Biochimica e Biologia Molecolare, University of Siena, Siena, Italy*

⁴*Dipartimento di Patologia Chirurgica, Molecolare e di Area Critica, University of Pisa, Pisa, Italy*

⁵*U.O.C. of Gastroenterology, Azienda Sanitaria Provinciale di Catania, Catania, Italy*

Introduction. Adipose tissue insulin resistance (IR) and elevated adipose free fatty acid (FFA) flux are prominent features in NAFLD. De novo lipogenesis (DNL) is up to 3-fold higher in subjects with fatty liver and is not suppressed on fasting. When DNL is stimulated, the production of saturated FAs is increased and the oxidation of FAs of any source is reduced. Both mechanisms can favor inflammation and IR.

Aim. We measured FFA flux/composition and used a surrogate index of DNL (DNLi) to evaluate their relationship with histological features in a group of non-diabetic subjects with biopsy-proven NAFLD.

Materials and Methods. Hepatic and adipose tissue-IR indices were derived from [2H5]glycerol and [2H2] glucose kinetics in a group of non-diabetic NAFLD patients in the basal state (n=40) and after a 4h oral glucose load test (n=20). Gas chromatography mass spectrometry was used to assess FFAs composition. DNLi was derived as the ratio palmitic/linoleic acid.

Results: Fasting plasma glucose/insulin, lipid profile, hepatic/adipose tissue IR indices and DNLi were similar in the two groups. Fasting DNLi was associated with triglycerides (TG) and FFAs levels and with adipose tissue IR ($r=0.597$, $r=0.330$ and $r=0.394$, respectively). Among histological features, fasting DNLi significantly correlated with steatosis ($r=0.364$, $P=0.02$) and NAS score ($r=0.306$, $P=0.05$). After the glucose load, TG levels initially increased despite elevated insulin levels, suggesting a significant contribution of DNL. Accordingly, DNLi consensually increased with TG levels ($r=0.749$, $P<0.005$) and was significantly related to the degree of fibrosis ($r=0.514$, $P=0.02$).

Conclusions. Oral glucose load is associated with changes in DNL and hepatic triglyceride synthesis that can favor liver fibrosis in patients with NAFLD.

Funded by FP7/2007-2013 under grant agreement n° HEALTH-F2-2009-241762 for the project FLIP and by PRIN 2009ARYX4T.