



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Forest dynamics and disturbance regimes in the Italian Apennines

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1614155 since 2017-05-19T12:41:15Z
Published version:
DOI:10.1016/j.foreco.2016.10.033
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Forest dynamics and disturbance regimes in the Italian Apennines

- 2 Giorgio Vacchiano^{*a}, Matteo Garbarino^b, Emanuele Lingua^c, Renzo Motta^d
- 3
- ^a Università degli Studi di Torino, DISAFA. Largo Braccini 2, 10095 Grugliasco (TO) Italy. giorgio.vacchiano@unito.it
 ^b Università Politecnica delle Marche, D3A. Via Brecce Bianche, 60131 AN Italy. matteo.garbarino@univpm.it
 ^c Università degli Studi di Padova, TESAF. Viale dell'Università 16, 35020 Legnaro (PD) Italy. emanuele.lingua@unipd.it
 ^d Università degli Studi di Torino, DISAFA. Largo Braccini 2, 10095 Grugliasco (TO) Italy. renzo.motta@unito.it
- 9 * corresponding author: <u>giorgio.vacchiano@unito.it</u>
- 10

11 Abstract

Forests of the Apennines are characterised by high canopy cover and high tree species diversity (being at the interface between two major climatic zones of Europe), and provide important ecosystem functions to millions of people. They exemplify cutting-edge themes such as forest ecology in warmer climates, consequences of heavy land use, and resilience at the trailing edge of the distribution of many European forest species (Silver fir, Norway spruce, Beech, Black pine, Birch).

17 We introduce the setting under the geological and climatological point of view and review the literature 18 on the interactions between these long-term drivers and the specific, structural, and genetic diversity of 19 these forest communities (e.g., effects of glacial refugia or tectonic/volcanic activity), followed by a 20 brief outline of what little is known about natural disturbance regimes and their range of variability. 21 Anthropogenic disturbances (fire, grazing) and land use changes (abandonment of cropland and pasture) 22 have been by far the main drivers of forest dynamics at least for the last two millennia, determining for 23 examples overageing of coppices, treeline advances, forest encroachment on former agricultural land. We suggest considerations about the interplay between these land use changes and disturbance drivers 24 25 (e.g. fuel continuity), summarise comparisons between managed and unmanaged forests (e.g., increase 26 in tree size, deadwood, biodiversity indicators), and elaborate on current proposals for climate-adapted

27 management, highlighting specific and genetic diversity as an important source of resilience and28 adaptive potential.

29

30 Keywords

- 31 Forest dynamics; Italy; Land use change; Mediterranean mountains; Natural disturbances;
- 32 Palaeoecology
- 33

34 Highlights

- Forests of the Apenninnes have been poorly explored in the ecological literature
- Anthropogenic disturbances and land use changes are the main drivers of forest dynamics
- Regimes of natural disturbances (avalanches, fires, wind, insects) are masked by human impacts
- Specific and genetic diversity (e.g., trailing edges) are an important source of resilience and
 adaptive potential
- 40

42 **1. Introduction**

Italy has two main mountain systems: the Alps and the Apennines (Figure 1). The alpine area represents
approximately one-quarter of the total Italian land surface (75,000 km²), while the Apennine region
accounts for approximately two-fifths (120,000 km²). Together they make up approximately 35% of the
total area of the country.

The Apennines include a series of mountain ranges (approximately 38-45° N and 8-17° E) bordered by 47 48 narrow coastlands that form the physical backbone of peninsular Italy. Their total length is 49 approximately 1,400 kilometres, and their width ranges from 40 to 200 km. Mount Corno Grande (2,912 50 m a.s.l.) is the highest point of the Apennines proper on the peninsula, while the stratovolcano Mount 51 Etna (3,323 m) in the island of Sicily is the highest peak, and the highest active volcano in Europe. 52 The Apennines are characterised by high forest cover and high tree species diversity (being at the 53 interface between the temperate and Mediterranean biomes of Europe). Apennine forests provide 54 important ecosystem services for millions of people, e.g., timber and energy wood, non-wood forest 55 products, water, biodiversity, and recreation (Vizzarri et al. 2015). Their functioning is driven by similar 56 macroecological factor as those at work in the Alps (Bebi et al. 2016), but the local peculiarities of 57 geological history, climate, and human influence have shaped a very different situation in terms of forest 58 composition, structure, and landscape mosaic. However, the Apennines have been much less explored 59 than the Alps in the literature in terms of disturbance regimes (both natural and anthropogenic) and their 60 effects on ecosystem processes.

The aim of this review is to summarize the main drivers of the composition, cover, structure, and
dynamics of Apennine forests, including both macroecological constraints (geology and climate), human
influence, and natural disturbance agents. In order to do so, we searched the existing scientific literature
(Google Scholar database) using the keywords "Apennine" and "forest", plus each of the following:
"pollen OR charcoal", "land use change", "(stand OR forest) structure", and "disturbances" (all words
used in either English or Italian languages). The search produced 9860 unique records. After filtering out

67 irrelevant and inaccessible papers, the total number of reviewed studies was 170 (i.e., 74, 42, 28, and 24
68 papers for each of the last four keywords, respectively) (Appendix 1).

Although focused on a specific geographic area, the history and dynamics of forests illustrated herein
are similar to those occurred in other heavily anthropized mountain regions of the world, and our
conclusions could be relevant in such areas as well.

72

73 2. Macroecological drivers

74 2.1 Geology

75 The Apennine orogeny began in the middle Miocene (about 20 million years ago, i.e., millions of years 76 later than the Alpine orogeny) (Carminati et al. 2012) as a consequence of the subduction of the African-77 Adriatic plate below the European plate (Carminati et al. 2003), and still continues today (Devoti et al. 78 2008). At the same time, large faults developed along the western side of the Apennines, connected to a 79 crustal thinning that resulted in the opening of the Tyrrhenian sea (Rosenbaum and Lister 2004). 80 Therefore, these young mountains of Italy are of paradoxical provenience, deriving from both 81 compression (folded systems) and extension (fault-block systems). This is responsible for the great 82 variety of rock types and the rugged appearance of the range today. In the north, sandstones, marls, and 83 greenstones occur. In the central Apennines clay, sandstone, and limestones are common. In the 84 southern Apennines large calcareous rock outcrops are separated by lowland areas of shale and 85 sandstone, and interrupted by extensive argillaceous rock types, which originate frequent erosion of the 86 "badlands" type.

87

88 2.2 Historical climate

89 Italy is placed at the boundary between the Mediterranean and the temperate zone of the boreal

90 hemisphere (Blasi et al. 2007). The Mediterranean Sea, which almost surrounds the country, is a

91 reservoir of heat and humidity. The Apennines intercept Atlantic perturbations and, with their rough

92 morphology, experience sharp temperature variations over short distances, frequent rainfalls for 93 adiabatic cooling and stau wind currents, and winter fog stagnation inside the several internal closed 94 basins and narrow valleys. The climate is of the mountain variety of the Mediterranean type, with dry 95 summers and rainy (and snowy) winters. Mean temperature ranges from 0 to 11 °C in January and from 24 to 28 °C in July. In the upper zone (over 2,000 m) there is snow 180-190 days per year. Annual 96 97 precipitation ranges from 600 to 4500 mm. Because the Adriatic Sea, rather thin and shallow, exerts a 98 less pronounced effect than the Tyrrhenian Sea, most of the precipitation falls on the western slopes of 99 the mountains (> 3,000 mm per year in Liguria). Precipitation seasonality is another typical trait of 100 Mediterranean climates; summer cyclones bring in torrential rains and may cause severe flooding. The 101 aridity index (ratio between precipitation and ET_0 : Wang et al. 2012) is often < 0.65 (sub-humid), 102 especially in Sicily, Apulia, Sardinia and Basilicata.

103

104 2.3 Climate change

105 The Mediterranean basin is very sensitive to climate variation (Lionello et al. 2006). In the last century, 106 Italian temperatures registered a warming (+1°C on average between 1865 and 2003) comparable to 107 Europe-wide trends (Brunetti et al. 2006). Annual precipitation and the number of rainfall days 108 decreased (811 to 723 mm in 1961-2000, and -10% in 1866-1996, respectively), but the intensity of 109 individual events increased (Brunetti et al. 2001). Hot waves increased in frequency and duration (66 to 110 187 hot days in 1951-2000) (Lionello et al. 2010). These trends are expected to worsen in the next 111 decades (Giorgi and Lionello 2008). Conditions simulated by HadCM3 forecast a remarkable increase of 112 maximum July temperatures over the whole country (+7.6 °C and +5.6°C in 2080 under A2 and B2 113 scenarios, respectively). The highest increases are concentrated in the northern Apennines, the lowest in 114 the South. Minimum January temperature is also expected to increase, albeit moderately (+1.8 to +2.8°C 115 in 2080). An overall reduction in annual precipitation is projected for year 2080, more pronounced under 116 the A2 (-18.1%) than the B2 scenario (-5.9%), mainly concentrated along the central Apennines (-22.8%)

and -9.3% under A2 and B2 scenarios, respectively). The highest reductions are expected in summer (48.5% and -41.5%) (Dibari et al. 2015). The potential consequences of the northward extension of the
Mediterranean subtropical climatic region in Italy include a decline in soil organic carbon, and a
reduction in snow cover associated with warming. In turn, a shallower, ephemeral snowpack will
promote soil freezing, with important consequences on soil nutrient dynamics (e.g., higher N losses)
(Edwards et al. 2007).

123

124 **3. Vegetation history**

The flora of the Apennines is the result of several "floristic streams" that have reached Italy since the 125 126 Tertiary (Valva 1992; Uzunov et al. 2005). Most species migrated from the balkanic-illiric province 127 during the Messinian salinity crisis in the high Miocene – a flow that originated most of the 128 Mediterranean flora (Bocquet et al. 1978) and which explains many disjunct distributions, e.g., Pinus 129 heldreichii Christ. (Piotti et al. 2014). Another wave of plant migration originated in the Iberian 130 peninsula and northern Africa, again during the Messinian, and enriched the Apennine flora with several 131 drought-tolerant elements (Biondi et al. 2015). During the later cold periods of the Pliocene, some alpine 132 and boreal species migrated from central and northern Europe (Pedrotti and Gafta 2003), while an 133 eastward migration, linked with plate movements, extended the distribution of some Tyrrhenian 134 elements to the southern tip of the peninsula (e.g., Alnus cordata (Loisel) Desf. and Pinus nigra Arnold 135 in the Calabrian Apennine) (Blasi et al. 2007). 136 After the Pleistocenic glacial (150,000-130,000 years Before Present), at least 24 transitions between

137 glacial (stadial) and warm (interstadial) climate occurred. Several tree species started to expand out of

their glacial refugia during each interstadial (e.g., oak in Sicily: Rossignol-Strick and Planchais 1989),

139 only to retreat again when cold and dry conditions returned (Figure 2). However, glacial refugia during

140 this time granted that some of the vegetational complexity that had originated in the Pliocene-

Pleistocene period could be maintained in part until the modern era (1800- first half of 1900), especiallyin the remote areas of the central and southern Apennines.

143 At the Last Glacial Maximum (22,000 to 14,000 years BP) summer temperatures were 6 to 8°C cooler 144 than at present, and sea level 120-140 m lower. Both paleoecological and simulation studies suggest 145 very little closed woody vegetation in the Mediterranean (Ray and Adams 2001; Di Rita et al. 2013), 146 with scattered trees or small pockets of open woodland from about 500 m above [present] sea level in 147 the Apennines and in the other Southern European peninsulas (Tzedakis et al. 1995). Since then, post-148 glacial migration followed four main routes: (1) northward and upward range expansion, e.g., beech and 149 fir (Bradshaw et al. 2010); (2) continuous persistence with increasing frequencies, e.g., oaks (Petit et al. 150 2002); (3) discontinuous persistence at low frequencies, e.g., birch (Plini and Tondi 1989); (4) failed 151 migration and persistence in localized refugia, e.g., spruce (Picea abies (L.) Karst.) (Ravazzi 2002). 152 Following the Younger Dryas cold interval (10,800-10,000 years BP), while beech (Fagus sylvatica L.) 153 was gaining dominance over much of temperate Europe (Bradshaw et al. 2010), the Mediterranean 154 region showed moister-than-present conditions. Near the coasts forests might have been dominated by 155 deciduous oaks (Lippi et al. 2007), while mountain areas developed two forest belts- the upper 156 dominated by fir (Abies alba Mill.), and the lower with mixed deciduous forests dominated by oak and 157 including maple, ash, linden and elm (Watson 1996).

During the Holocene thermal maximum (7,000 to 4,000 years BP), paleoecological records show a rapid and fairly simultaneous decline of elm throughout Europe (Huntley et al. 1989), and a strong increase in beech, which successfully invaded the fir woodland to form a mixed fir-beech complex (Bradshaw et al. 2010). Evergreen mediterranean vegetation replaced the deciduous forest on limestone soils (Di Rita and Magri 2009).

In the last 4,000 years, climate has been rather stable (except for short-lived fluctuations such as the 550
AD and the Little Ice Age cool periods). In the absence of land use by man, forests would dominate the
potential vegetation of most of the Apennine region, with an elevational separation between

166 Mediterranean evergreen, deciduous broadleaves, and conifers resulting from differences in cold 167 tolerance of each species (De Philippis 1937). Mountain conifers may descend in the broadleaves belt, as 168 deciduous broadleaves do in the evergreen mediterranean vegetation belt, due to the fact that high 169 temperature extremes produce less distinct zonations (Pignatti 2011). Besides temperature, water 170 availability is a major factor influencing potential forest vegetation: forest cover is generally higher in 171 the regions with temperate climate compared to those with Mediterranean climate, but also in regions 172 with high orographic precipitation (e.g., the Tyrrhenian side of the northern Apennines) with respect to 173 more continental and drier areas (Magri et al. 2015). The superposition of climatic gradients, post-glacial 174 migration routes, and complex orography has produced for most tree species in the Apennines a 175 fragmented distribution, which has promoted reproductive isolation and high genetic diversity (Magri et 176 al. 2006, Leonardi et al. 2012).

177

178 4. Human history and impacts on vegetation

Anthropogenic land use has had a profound effect on forests of the Apennine region and, in most cases,outweighted by far the effects of macroecological constraints (Brown et al. 2013).

181 Even before the advent of agriculture, the presence of humans has severely influenced the composition 182 and distribution of forests. This is exemplified by the late Holocene dynamics of fir and beech. During 183 the early Holocene (9000–6000 years BP), fir became abundant from the sea level to the mountains 184 (Montanari 1989). After 6,000 years BP, however, its presence declined in many sites (Magri et al. 185 2015), leading to the current reduced distributions. The reasons for such decline, and the relive 186 importance of climate, competition, and anthropogenic impact, are debated. On one hand, fir has 187 retreated into more or less the same areas where it was found at the beginning of the present interglacial 188 (Muller et al. 2007), supporting the hypothesis that its Holocene dynamics are driven by long-term 189 climatic and edaphic patterns (Joannin et al. 2012). On the other hand, multiple palaeoecological 190 evidence shows a synchrony between the reduction of fir at 6,000-3,000 years BP and an increase in

191 human activity, especially from pastoralism and fire (Henne et al. 2013). These dynamics would imply a 192 much wider climatic niche than previously thought for fir, which could have been part of sub-193 Mediterranean oak forests, as suggested also from Pleistocenic pollen evidence (Tinner et al. 2013). 194 At the same time, beech, which had started expanding into southern and central Italy already 18,000 195 years BP (Joannin et al. 2012), reached its maximum spread between 8,000 and 4,000 years BP (Branch 196 and Marini 2014), i.e. simultaneously to the decline of fir, leading to a mixed Abies-Fagus association in 197 the upper belt between from 5,200 yBP and then to a dominance of beech from around 2,900 years BP 198 (Watson 1996). The expansion of beech in the Northern Apennines may have been favored by a 199 decrease in summer insolation (Berger 1978) and a smoothing of seasonal climate extremes (Huntley et 200 al. 1989; Watson 1996), but also facilitated by human activities. Since beech seedlings require moderate 201 light intensity for development (Ellenberg 1986), it is reasonable that clearing of the pre-existing dense 202 fir forest has determined a shift in dominance between the two species (Valsecchi et al. 2008). 203 The transition of the first human communities out of hunting-gathering occurred around the start of the 204 second millennium BC. From this time, semi-nomadic agriculture and herding were accompanied by 205 widespread slash-and-burn deforestation (Watson 1996), a higher hydrogeologic instability (Cremaschi 206 et al. 2008) and an increase in soil erosion, which favored some pioneer tree species, e.g., Calabrian pine (*Pinus nigra* subsp. *laricio*) (Nicolaci et al. 2014). Starting from the 8th century BC, under the Roman 207 208 influence, Apennine forests were used for timber, fuelwood, and cleared for agriculture and pastures. 209 Forests in the vicinity of the sea or major rivers were harvested and floated to the nearest ports as civil 210 and naval timber (*silva incaedua*); the others were coppiced (*silva caedua*), harvested by hand or 211 animals, and used for tools and small crafts, or as energy wood. This system is believed to be 212 responsible for the degradation of forests along coasts (Calò et al. 2012), and on major river basins 213 (Mercuri and Sadori 2013). However, recent evidence (Sadori et al. 2015) shows that at least beech-fir 214 forests were more widely distributed than today, despite the fact that fir was the most desirable species for ship hulls, construction timber, and furniture (Allevato et al. 2010). Another source of forest 215

216 degradation was transhumant sheep herding, which was favored in southern Italy by the territorial unification brought about by the Romans, and was practiced up to the beginning of the 20th century. At 217 218 its apex (5.5 million sheep in year 1604: Venanzoni et al. 1993), the newtwork of seasonal transhumance 219 "highways" had a cumulative length of more than 3000 km. Some Apennine forests still show the legacy 220 of their past use as wooded pastures, e.g., simplified vertical structure (Mancini et al. 2016). 221 The demographic decline that followed the fall of the Roman Empire favored a partial recovery of forest 222 vegetation. In the year 410, emperor Constantine entrusted to the catholic Church all forests that 223 surrounded former sacred temples. Among the general forest abandonment of the early Middle Ages, 224 religious orders (Benedectins, Camaldolese, Carthusians, Vallombrosans) were the only subjects that 225 carried out forest management and conservation (Piccioli 1923), with long-lasting effects on forest 226 structure and composition, e.g., in most present-day pure fir forests of the Apennines (Costantini et al. 227 2010). The population increased again starting from the Late Middle Ages, with the exception of the 228 period 1350-1450 AD (i.e., a period of both climatic, economic, and demographic crisis: Sadori et al. 229 2016, when forests expanded again: Mensing et al. 2013), and clearing was resumed in community 230 forests (Guido et al. 2013), while those belonging to the noblety were restricted to all uses (foris stare – 231 "stay outside" edicts may have originated the word *forest*) and preserved as hunting grounds. In 232 response to increasing forest clearing, some states such as the Republic of Venice, the Papal States, and 233 the Granduchy of Tuscany started preserving public forests as a strategic resource for ship and 234 household building (Di Filippo et al. 2007).

Around year 1500, most oak high forests had been replaced by chestnut (*Castanea sativa* Mill.)
orchards, an important source of food, whose cultivation started during the Roman Age (Di Pasquale et
al. 2010) and rapidly spread outside the ecological niche of the species starting from the early Middle
Ages (Conedera et al. 2004). The introduction of chestnut coincided with a radical change in local use of
land. Fire was no longer used systematically to clear open spaces in forests. Instead, many wooded areas

were actively managed as chestnut groves and managed in coppices for pole production (Tinner andConedera 1995).

242

243 5. Land use changes in the last 200 years

The decline of forest cover continued in the 17-19th centuries and was finally exacerbated by the needs 244 245 of the newborn railway network (Zanotti Cavazzoni 1907). The recognition of the link between 246 deforestation and hydrogeologic instability resulted in measures to restore forest cover; between national 247 unification (1861) and 1950, 200,000 hectares of forests were planted (Patrone 1953), and a further 248 560,000 hectares between 1950 and 1980 as a consequence of occupational policies (Romano 1987). At 249 the same time, the abandonment of rural and mountain settlements (Munafò et al. 2015) caused a strong 250 increase in forest cover, ranging from +0.1% and +27% per decade (average: +10.3%) (Fig. 3). The rate 251 of change seems more dependent on local socio-economic changes than on the period, location, or 252 topographic features: forest and shrubland expansion were inversely related to population density 253 (Falcucci et al. 2007; Corona et al. 2008) and occurred mainly at elevations below 1,000 m, on moderate 254 slope gradients (< 40%), and near forest edges and roads (\leq 500 m) (Cimini et al. 2013). In most cases, 255 the process has been related to abandonment of traditional farming or grazing activities (e.g., De Sillo et 256 al. 2012). The period needed for tree canopy closure on former unforested land ranged from 25 to 50 257 years (e.g., Bracchetti et al. 2012). The mode of natural afforestation depended on previous landscape 258 configuration, species composition, and topography. In former agricultural areas, closure of open spaces 259 resulted in homogeneization of the landscape mosaic, while in former pastures complexity increased 260 with incoming forest cover (e.g., Corso et al. 2005; De Sillo et al. 2012). The density and complexity of 261 edges, however, has often increased due to non-homogenous colonization patterns of the incoming 262 forest.

Nine studies have addressed treeline shift, a much less researched issue than e.g. on the Alps (GehrigFasel et al. 2007; Leonelli et al. 2011; Bebi et al. 2016). Even though on the Apennines the treeline had

265 been depressed by human activity (fire, grazing) since the Late Holocene (Compostella et al. 2014), 266 recent change was less clearcut. The only reported figures were an increase of 1m per year upwards and 267 3m downwards in dwarf pine (Pinus mugo Turra) (Palombo et al. 2013) and an expansion of 1% per 268 year at the beech treeline (van Gils et al. 2008). The relative importance of recent climate warming and 269 land use change was debated, with the first factor playing out at regional level (e.g., simultaneous 270 increases in black pine treeline in the last 30-40 years in the Central Apennines: Piermattei et al. 2016) 271 and the second at local scale (Palombo et al. 2014). Where grazing is still active, it may prevent 272 establishment of shrubs that could subsequently act as nurse sites for beech seedlings (Catorci et al. 273 2012), such that the treeline is actually stable (Pezzi et al. 2007).

274

275 6. Changes in forest structure

276 Abandonment of forest management and rural landscapes has also determined a widespread ageing of all 277 Apennine forests (Tellini Florenzano 2004) and an increase of living and dead biomass (Marchetti et al. 278 2010; Motta et al. 2013). Many coppices have been subject to active conversion into high forest, while 279 forest regulations have been issued in many regions imposing a maximum coppice age above which 280 vegetative regeneration was forbidden in order to avoid exhaustion of stumps and fertility declines 281 (usually at 35-40 years, e.g., Regione Toscana 2003). Abandoned coppices of many species transitioned 282 spontaneously to high forest, e.g. in beech (Nocentini et al. 2009), birch (Bagnato et al. 2014), and 283 chestnut stands, which have undergone succession by more shade-tolerant broadleaves such as beech, 284 hop-hornbeam, and downy oak (Pezzi et al. 2011). More intensive silvicultural systems, e.g., uneven-285 aged coppicing, are all but lost (Coppini and Hermanin 2007). Re-naturalization dynamics and late-seral 286 succession were observed also in former black pine plantations (Tonon et al. 2005). 287 Land abandonment and decrease of forest harvesting has also raised the interest for forests with old-288

growth characteristics (Marchetti and Blasi 2010; Chirici and Nocentini 2013). These stands, prevalently

289 dominated by beech, house some of the tallest and oldest trees in Europe; they have a significantly

higher deadwood volume than ordinarily managed forests with similar composition (Lombardi et al.
2013), and a living biomass comparable or higher than central European old-growth forests, in both pure
beech (302-1383 m³ ha⁻¹) and mixed beech-fir stands (570-1189) (Calamini et al. 2011). Differences
between these old growth stands and managed one are very evident also in terms of genetic diversity
(Paffetti et al. 2012), lichen flora (Brunialti et al. 2010), forest dynamics (Travaglini et al. 2012), and
structural diversity (Alessandrini et al. 2011). All these factors have a preminent importance as
indicators of habitat and biodiversity (Schulz et al. 2014).

297

298 **7. Disturbance regimes**

299 Besides macroecological and anthropic drivers, natural disturbance regimes are one of the main factors 300 shaping the composition, structure, and patterns of forest ecosystems both at the local and at the regional 301 level (Kulakowski et al. 2016). However, the natural disturbance regimes of Southern European forests 302 have been long since masked by human modification to forest cover, composition, structure, and 303 continuity (Bengtsson et al. 2000). Whereas actual disturbance regimes have started to be addressed by 304 ecological research in the Italian Alps (Valese et al. 2014, Vacchiano et al. 2016), data and studies from 305 Apennine forests are conspicuously lacking, so that the properties of disturbance regimes and its 306 historical range of variability (Kulakowski et al. 2016) are still largely unknown.

Fire is the most common disturbance agent in peninsular Italy. In the period between 1980 and 2012 the average burned area in Italy was 113,496 ha per year (EC 2013), with a mean yearly frequency of 9,736 fires, i.e., 3.2 fires per km² (Spain: 3, Greece: 1.2). With the exception of climatically unfavorable years (e.g., 227,729 ha burned in 2007), the five-year average burned area has decreased steadily (from 46,800 ha in 1990-1995 to 36,800 ha in 2005-2010) due to improvements in fire policies and prevention (FAO 2010). Long-term data are not available at a regional detail; in year 2015, Apennine regions (i.e., excluding the Alpine space, Puglia, and islands) included 61% of all forest fires and 70% of total

forested burned area of the country (Corpo Forestale dello Stato 2016). The percent of wooded area

burned yearly is 1.14% nationally, i.e., the highest in Mediterranean Europe after Portugal (FAO 2010).
Mediterranean pine forests have a disproportionately high fire occurrence relative to their area, followed
by Mediterranean montane forests and evergreen broadleaves forests (Corona et al. 2014). Most fires are
anthropogenic: negligent motives are common, especially agricultural (stubbe burning, land cleaning
after harvesting) and pastoral burning (Lovreglio et al. 2010).

320 The National Forest Inventory of 2005 reports the following surfaces for other forest disturbance agents 321 in the Apennine regions: biotic agents 508,803 ha, browsing and grazing 160,965 ha, weather extremes 322 220,223 ha, i.e., respectively 11%, 3%, and 5% of total forest cover (MIPAAF 2007). Abiotic 323 disturbances have been analyzed mostly in relation to the ecology of beech forests; a common feature of 324 such studies is that, contrary to expectations, beech preserves its dominance over competitor species 325 even after low- and medium severity disturbance (van Gils et al. 2010). The high structural complexity 326 of overmature beech forests in the central Apennines has been related to frequent low-severity gap-327 forming events (Ziaco et al. 2012) (Fig. 4), similarly to central European old-growth beech forests 328 (Westphal et al. 2006). However, recent large stand-replacing windthrows in fir forests (Bottalico et al. 329 2015) suggest that the wind disturbance regime might be composed also by rarer, more intense events. 330 A much common form of disturbance is landslides, which occur in 9% of all Apennine areas (Triglia 331 and Iadanza 2014), with somewhat positive consequences for forest diversity (e.g., exposure of mineral 332 soil) but catastrophic results for society (e.g., the mudflow events in Sarno in 1998 or Genova in 2015). 333 No systematic assessment exists on the extent and severity of biotic disturbances in Apennine forests. 334 Historical evidence (e.g., chestnut blight) and case studies highlight the potential for high-severity 335 events from both native (Puddu et al. 2003; Luchi et al. 2014) and alien pathogens (Vettraino et al. 2005; 336 Luchi et al. 2016). Evidence of insect outbreaks is also limited, mostly to artificial tree plantations 337 (Vignali et al. 2015). However, it has been hypothesized that climate warming might exacerbate tree 338 drought stress, which is already inducing tree decline in sensitive populations (Piovesan et al. 2008; Di

Filippo et al. 2010), and hence facilitate the outbreaks of secondary insects and pathogens, especially insouthern, low-elevation beech stands (Luchi et al. 2015).

341

342 8. Management implications

343 Land use change is currently the dominant driver of forest dynamics in the Apennine, relative to both 344 climate and natural disturbances. If the urbanization trend and the abandonment of marginal lands 345 persist, forests in mountain areas will keep on accumulating live and dead biomass, and the connectivity 346 between old-growth patches will likely increase with ongoing secondary succession in fallow lands. 347 While this might lead to increased provision of some important forest ecosystem services, e.g., timber, 348 carbon stocking, avalanche and rockfall mitigation, or even carnivore habitat (Fabbri et al. 2007, Ciucci 349 and Boitani 2008), some other forest functions are expected decrease, e.g., recreational use, mechanical 350 stability, landscape diversity, and habitat for open-areas plant and animal species (Casanova et al. 2005). 351 Forest planning therefore needs to embrace a comprehensive zonation approach, building on science-352 based forecasts of forest structure and composition (Vacchiano et al. 2012b), and assigning spatially-353 explicit priorities to management alternatives needed to maximize each of the desired ecosystem service. 354 We also argue that the ongoing shift from segregation (i.e., each forest accomplishes a single 355 management objectives) to integration (multiple management objectives are sought in the same forest 356 area) should be strengthened, especially for what concerns biodiversity-oriented forest management 357 (Schulz et al. 2014).

So far, the pressure posed by climate warming on forest ecosystems has been buffered or even countered by forest expansion following land abandonment. However, while large uncertainties still exist regarding the climate resilience of several important tree species (see below), adaptive management strategies such as intensification of thinning regimes or assisted migration (Temperli et al. 2012) need to be further explored to provide means to counter the negative effects of climate change, e.g. forest decline in drought-prone conifer forests (Vacchiano et al. 2012a). However, the actual outcomes of

364 climate change, and the adaptive management actions needed, will be determined by its interactions with past and future land use, changes in disturbance regimes (e.g., van Gils et al. 2010) and legacies 365 366 (Vacchiano et al. 2014), endogenous stand dynamics (Long and Vacchiano 2014), resilience drivers 367 (e.g., reproduction dynamics: Ascoli et al. 2015), and novel communities (e.g., Benesperi et al. 2012). 368 In particular, the interplay of climate warming and forest expansion will exacerbate damages by extreme 369 events such as windthrow, forest fires, and insect outbreaks (Seidl et al. 2014). This is particularly 370 troubling where forests play important social functions, such as protection of infrastructure from rockfall 371 or debris flows. Apennine forest seem to have experienced low- to moderate-severity disturbance 372 regimes in the recent past, but managers should be prepared to plan and carry out forest restoration 373 measures on larger scales following the likely increase in frequency of stand-replacing events, as well as 374 measures to increase the resistance and resilience of forest ecosystems to such events. 375 The abundance of coppices due to the prevalent use of small, private forest lots as a source of energy 376 wood, and their subsequent abandonment, raise many question about the best silvicultural practices to 377 preserve the provision of such cultural landscape and its functions, especially soil protection and 378 biodiversity. Current forest regulations forbid clearcutting in overmature coppices; variable retention 379 systems that have recently been proposed for Alpine beech forests could be adopted to foster structural 380 differentiation and avoid vertical and horizontal simplification, which might lead to loss of habitat and 381 productivity (Negro et al. 2015).

The reduction of open meadows has also modified the feeding behaviour of wild ungulates, which have increasingly browsed forest regeneration. As repeated and selective browsing threatens the regeneration of forest stands and triggers a loss of ecosystem resilience (Cutini et al. 2011), forest management and wildlife management must be planned together, with an increased communication between stakeholders from the two sectors.

387

388 9. Directions for research

389 This review illustrates how cultural factors had a profound effect on forests of the Apennine region and, 390 in most cases, outweighed by far the effects of macroecological constraints. The millennia-long uses of 391 the forest by man, as well as more recent socio-economic changes, seem to play a more important role 392 for current forest structure and dynamics than natural processes driven by geology and climate. 393 Disturbance regimes have also been masked by the pervasive human influence, and their range of 394 variability still represent a large knowledge gap. Finally, the climate plasticity of some important forest 395 species has still to be completely understood (e.g., fir); the high specific and genetic diversity of 396 Apennine forests will have to be leveraged as an important source of resilience and adaptive potential. 397 The future extent, composition and functioning of Apennine forests is therefore dependent on the 398 changes in land use, disturbance regimes, and climate, but only the latter have been subject to rigorous 399 study or simulation. Prediction of future forest will need to incorporate future land use and disturbance 400 scenarios and the complex and spatially-explicit feedbacks between them. 401

402

403 **References**

404

Alessandrini A, Biondi F, Di Filippo A, Ziaco E, Piovesan G. Tree size distribution at increasing spatial
scales converges to the rotated sigmoid curve in two old-growth beech stands of the Italian Apennines.
Forest Ecology and Management. 2011 Dec 1;262(11):1950-62.

Allevato E, Ermolli ER, Boetto G, Di Pasquale G. Pollen-wood analysis at the Neapolis harbour site
(1st–3rd century AD, southern Italy) and its archaeobotanical implications. Journal of Archaeological
Science. 2010 Sep 30;37(9):2365-75.

411 Ascoli D, Vacchiano G, Maringer J, Bovio G, Conedera M. The synchronicity of masting and

412 intermediate severity fire effects favors beech recruitment. Forest Ecology and Management. 2015 Oct

413 1;353:126-35.

- 414 Bagnato S, La Piana V, Mercurio R, Merlino A, Scarfò F, Sciascia N, Solano F, Spampinato G.
- 415 Dinamiche evolutive in boschi cedui di betulla (Betula aetnensis Rafin) nel Monte Etna (Sicilia).
- 416 Forest@. 2014 Apr 18;11(51):52-64.
- 417 Bebi P, Seidl R, Motta R, Fuhr M, Firm D, Krumm F, Conedera M, Ginzler C, Wohlgemuth T,
- 418 Kulakowski D. Changes of forest cover and disturbance regimes in the mountain forests of the Alps.
- 419 Forest Ecology and Management. 2016; in press.
- 420 Benesperi R, Giuliani C, Zanetti S, Gennai M, Lippi MM, Guidi T, Nascimbene J, Foggi B. Forest plant
- 421 diversity is threatened by *Robinia pseudoacacia* (black-locust) invasion. Biodiversity and Conservation.
- 422 2012 Dec 1;21(14):3555-68.
- 423 Berger A. Long-term variations of daily insolation and Quaternary climatic changes. Journal of the
- 424 Atmospheric Sciences. 1978 Dec;35(12):2362-7.
- 425 Bertoldi R. Le vicende vegetazionali e climatiche nella sequenza paleobotanica wurmiana e post-
- 426 wurmiana di Ladgei (Appennino Settentrionale). Ateneo Parm. Acta Nat. 1981;16:147-75.
- 427 Biondi E, Allegrezza M, Casavecchia S, Galdenzi D, Gasparri R, Pesaresi S, Poldini L, Sburlino G,
- 428 Vagge I, Venanzoni R. New syntaxonomic contribution to the Vegetation Prodrome of Italy. Plant
- 429 Biosystems. 2015 May 4;149(3):603-15.
- 430 Blasi CA, Filibeck GO, Burrascano SA, Copiz RI, Celesti-Grapow L, Di Pietro R, Ercole S, Lattanzi E,
- 431 Rosati L, Tilia A. Primi risultati per una nuova regionalizzazione fitogeografica del territorio italiano.
- 432 Biogeographia sn. 2007;28:9-23.
- 433 Bocquet G, Widler B, Kiefer H. The Messinian model-A new outlook for the floristics and systematics
- 434 of the Mediterranean area. Candollea. 1978;33(2):269-87.
- 435 Bottalico F, Bottacci A, Galipò G, Nocentini S, Torrini L, Travaglini D, Ciancio O. Formazione dei gap
- 436 causati dal vento in soprassuoli coetanei di abete bianco (*Abies alba* Mill.). Un caso di studio nella

- 437 montagna appenninica (Italia centrale). In: O. Ciancio (ed) Atti del Secondo Congresso Internazionale di
- 438 Selvicoltura Proceedings of the Second International Congress of Silviculture. Accademia Italiana di
- 439 Scienze Forestali, Firenze, 2015;1: 257-62.
- 440 Bracchetti L, Carotenuto L, Catorci A. Land-cover changes in a remote area of central Apennines (Italy)
- and management directions. Landscape and Urban Planning. 2012 Feb 29;104(2):157-70.
- 442 Bradshaw RH, Kito N, Giesecke T. Factors influencing the Holocene history of Fagus. Forest Ecology
 443 and Management. 2010 May 15;259(11):2204-12.
- 444 Branch NP, Marini NA. Mid-Late Holocene environmental change and human activities in the northern
- 445 Apennines, Italy. Quaternary International. 2014 Dec 5;353:34-51.
- 446 Brown AG, Hatton J, Selby KA, Leng MJ, Christie N. Multi-proxy study of Holocene environmental
- change and human activity in the Central Apennine Mountains, Italy. Journal of Quaternary Science.
 2013 Jan 1;28(1):71-82.
- Brunetti M, Colacino M, Maugeri M, Nanni T. Trends in the daily intensity of precipitation in Italy from
 1951 to 1996. International Journal of Climatology. 2001 Mar 15;21(3):299-316.
- 451 Brunetti M, Maugeri M, Monti F, Nanni T. Temperature and precipitation variability in Italy in the last
- 452 two centuries from homogenised instrumental time series. International Journal of Climatology. 2006
 453 Mar 15;26(3):345-81.
- 454 Brunialti G, Frati L, Aleffi M, Marignani M, Rosati L, Burrascano S, Ravera S. Lichens and bryophytes
- 455 as indicators of old-growth features in Mediterranean forests. Plant Biosystems. 2010;144(1):221-33.
- 456 Calamini G, Maltoni A, Travaglini D, Iovino F, Nicolaci A, Menguzzato G, Corona P, Ferrari B, Di
- 457 Santo D, Chirici G, Lombardi F. Stand structure attributes in potential old-growth forests in the
- 458 Apennines, Italy. L'Italia Forestale e Montana. 2011;66:365–81.

- 459 Calò C, Henne PD, Curry B, Magny M, Vescovi E, La Mantia T, Pasta S, Vannière B, Tinner W.
- 460 Spatio-temporal patterns of Holocene environmental change in southern Sicily. Palaeogeography,
- 461 Palaeoclimatology, Palaeoecology. 2012 Mar 15;323:110-22.
- 462 Carminati E, Doglioni C, Scrocca D. Apennines subduction-related subsidence of Venice (Italy).
- 463 Geophysical Research Letters. 2003 Jul 1;30(13):1717.
- 464 Carminati E, Lustrino M, Doglioni C. Geodynamic evolution of the central and western Mediterranean:
- 465 Tectonics vs. igneous petrology constraints. Tectonophysics. 2012 Dec 5;579:173-92.
- 466 Casanova P, Memoli A, Pini L. La fauna a uccelli e mammiferi della foresta di Vallombrosa: problemi
- 467 di Habitat, di alimentazione e di gestione dell'ecosistema. L'Italia Forestale e Montana. 2015;60(2):213-
- 468 16.
- 469 Catorci A, Scapin W, Tardella MF, Vitanzi A. Seedling survival and dynamics of upper timberline in
- 470 Central Apennines. Polish Journal of Ecology. 2012 Jan 1;60(1):79-94.
- 471 Chirici G, Nocentini S. Old-growth forests in Italy: recent research developments and future
- 472 perspectives. L'Italia Forestale e Montana. 2013 May 16;65(5):475-80.
- 473 Cimini D, Tomao A, Mattioli W, Barbati A, Corona P. Assessing impact of forest cover change
- 474 dynamics on high nature value farmland in Mediterranean mountain landscape. Annals of Silvicultural
 475 Research 37 (1): 29-37
- 476 Ciucci P, Boitani L. The Apennine brown bear: a critical review of its status and conservation problems.
- 477 Ursus. 2008;19(2):130-45.
- 478 Compostella C, Mariani GS, Trombino L. Holocene environmental history at the treeline in the Northern
- 479 Apennines, Italy: A micromorphological approach. The Holocene. 2014 Feb 6;24(4):393–404.

- 480 Conedera M, Krebs P, Tinner W, Pradella M, Torriani D. The cultivation of *Castanea sativa* (Mill.) in
- 481 Europe, from its origin to its diffusion on a continental scale. Vegetation History and Archaeobotany.
 482 2004 Aug 1;13(3):161-79.
- 483 Coppini M, Hermanin L. Restoration of selective beech coppices: a case study in the Apennines (Italy).
- 484 Forest Ecology and Management. 2007 Sep 25;249(1):18-27.
- 485 Corona P, Calvani P, Mugnozza GS, Pompei E. Modelling natural forest expansion on a landscape level
- 486 by multinomial logistic regression. Plant Biosystems. 2008 Nov 1;142(3):509-17.
- 487 Corona P, Ferrari B, Cartisano R, Barbati A. Calibration assessment of forest flammability potential in
- 488 Italy. iForest, 2014;7:300–5.
- 489 Corpo Forestale dello Stato. Gli incendi boschivi nel 2015. MIPAAF. 2016;[online resource] URL:
- 490 http://www.corpoforestale.it/flex/cm/pages/ServeAttachment.php/L/IT/D/5%252F5%252Fe%252FD.87
- 491 <u>c1ebbd8f53c3aaada6/P/BLOB%3AID%3D11941/E/pdf</u>. Last accessed 12th October 2016.
- 492 Corso G, Carranza ML, Giancola C, Stanisci A. Trasformazioni del paesaggio altomontano
- 493 dell'Appennino Centrale negli ultimi 50 anni. Il caso del pSIC "La Gallinola-M. Miletto-Monti del
- 494 Matese". In: Atti 9° Conferenza Asita. 2015;9:799-804
- 495 Costantini EA, Fantappié M, L'Abate G. Climate and pedoclimate of Italy. The Soils of Italy. 2013:19496 37.
- 497 Costantini M, Di Pietro F, Marongiu S, Fossa U, Frigerio S, Socci F, Urbinati C, De Santis F. Le radici
- 498 della sostenibilità: la regola della vita eremitica, ovvero Le Constitutiones Camaldulenses: primo
- 499 volume. Sviluppo rurale. INEA, Roma. 2010.
- 500 Cremaschi M, Nicosia C, Salvioni MA. L'uso del suolo nell'Eneolitico e nel Bronzo antico, nuovi dati
- 501 dalla Pianura Padana centrale. In: . L'età del Rame in Italia. Atti della XLIII Riunione Scientifica
- 502 dell'Istituto Italiano di Preistoria e Protostoria (2008). Istituto italiano di Storia e Protostoria, Firenze.
- 503 2011; 225-31.

- Cutini A, Bongi P, Chianucci F, Pagon N, Grignolio S, Amorini E, Apollonio M. Roe deer (*Capreolus capreolus* L.) browsing effects and use of chestnut and Turkey oak coppiced areas. Annals of Forest
 Science. 2011;68(4):667-74.
- 507 De Philippis A. Classificazioni ed indici del clima in rapporto alla vegetazione forestale italiana.
- 508 Tipografia Mariano Ricci, Firenze. 1937.
- 509 De Sillo R, De Sanctis M, Bruno F, Attorre F. Vegetation and landscape of the Simbruini mountains
- 510 (Central Apennines). Plant Sociology. 2012;49(Suppl.1):3-64.
- 511 Devoti R, Riguzzi F, Cuffaro M, Doglioni C. New GPS constraints on the kinematics of the Apennines
- subduction. Earth and Planetary Science Letters. 2008 Aug 30;273(1):163-74.
- 513 Di Filippo A, Alessandrini A, Biondi F, Blasi S, Portoghesi L, Piovesan G. Climate change and oak
- 514 growth decline: Dendroecology and stand productivity of a Turkey oak (Quercus cerris L.) old stored
- 515 coppice in Central Italy. Annals of Forest Science. 2010 Oct 1;67(7):706.
- 516 Di Filippo A, Biondi F, Čufar K, De Luis M, Grabner M, Maugeri M, Presutti Saba E, Schirone B,
- 517 Piovesan G. Bioclimatology of beech (Fagus sylvatica L.) in the Eastern Alps: spatial and altitudinal
- 518 climatic signals identified through a tree-ring network. Journal of Biogeography. 2007 Nov
- 519 1;34(11):1873-92.
- 520 Di Pasquale G, Allevato E, Russo Ermolli E, Coubray S, Lubritto C, Marzaioli F, Yoneda M, Takeuchi
- 521 K, Kano Y, Matsuyama S, De Simone GF. Reworking the idea of chestnut (*Castanea sativa* Mill.)
- 522 cultivation in Roman times: new data from ancient Campania. Plant Biosystems. 2010 Dec
- **523** 1;144(4):865-73.
- 524 Di Rita F, Anzidei AP, Magri D. A Lateglacial and early Holocene pollen record from Valle di
- 525 Castiglione (Rome): vegetation dynamics and climate implications. Quaternary International. 2013 Mar
 526 4;288:73-80.

- 527 Di Rita F, Magri D. Holocene drought, deforestation and evergreen vegetation development in the
- 528 central Mediterranean: a 5500 year record from Lago Alimini Piccolo, Apulia, southeast Italy. The
- 529 Holocene. 2009 Mar 1;19(2):295-306.
- 530 Dibari C, Argenti G, Catolfi F, Moriondo M, Staglianò N, Bindi M. Pastoral suitability driven by future
- climate change along the Apennines. Italian Journal of Agronomy. 2015 Sep 18;10(3):109-16.
- 532 EC. Forest Fires in Europe, Middle East and North Africa 2012. Publications Office of the European
 533 Union, Luxembourg. 2013.
- 534 Edwards AC, Scalenghe R, Freppaz M. Changes in the seasonal snow cover of alpine regions and its
- effect on soil processes: a review. Quaternary international. 2007 Mar 31;162:172-81.
- 536 Ellenberg H. Vegetation Mitteleuropas mit den Alpen in okologischer Sicht. Fourth edition. Verlag
- 537 Eugen Ulmer, Stuttgart, Germany. 1986.
- 538 Falcucci A, Maiorano L, Boitani L. Changes in land-use/land-cover patterns in Italy and their
- implications for biodiversity conservation. Landscape Ecology. 2007 Apr 1;22(4):617-31.
- 540 FAO. Global forest resource assessment. UN Food and Agriculture Organisation, Rome. 2010.
- 541 Gehrig-Fasel J, Guisan A, Zimmermann NE. Tree line shifts in the Swiss Alps: Climate change or land
- abandonment?. Journal of Vegetation Science. 2007 Aug 1;18(4):571-82.
- 543 Giorgi F, Lionello P. Climate change projections for the Mediterranean region. Global and Planetary
- 544 Change. 2008 Sep 30;63(2):90-104.
- 545 Guido MA, Menozzi BI, Bellini C, Placereani S, Montanari C. A palynological contribution to the
- 546 environmental archaeology of a Mediterranean mountain wetland (North West Apennines, Italy). The
- 547 Holocene. 2013 Nov 1;23(11):1517-27.

- 548 Henne PD, Elkin C, Colombaroli D, Samartin S, Bugmann H, Heiri O, Tinner W. Impacts of changing
- 549 climate and land use on vegetation dynamics in a Mediterranean ecosystem: insights from paleoecology
- and dynamic modeling. Landscape Ecology. 2013 May 1;28(5):819-33.
- 551 Huntley B, Bartlein PJ, Prentice IC. Climatic control of the distribution and abundance of beech (Fagus
- L.) in Europe and North America. Journal of Biogeography. 1989 Nov 1;16(6):551-60.
- Joannin S, Brugiapaglia E, Beaulieu JL, Bernardo L, Magny M, Peyron O, Goring S, Vannière B.
- 554 Pollen-based reconstruction of Holocene vegetation and climate in southern Italy: the case of Lago
- 555 Trifoglietti. Climate of the Past. 2012 Dec 7;8(6):1973-96.
- 556 Kulakowski D, Seidl R, Holeksa J, Kuuluvainen T, Nagel T, Panayotov M, Svoboda M, Thorn S,
- 557 Vacchiano G, Whitlock C, Wohlgemuth T, Bebi P. A walk on the wild side: Disturbance dynamics and
- the conservation and management of European mountain forest ecosystems. Forest Ecology and
- 559 Management. 2016; in press.
- Leonardi S, Piovani P, Scalfi M, Piotti A, Giannini R, Menozzi P. Effect of habitat fragmentation on the
 genetic diversity and structure of peripheral populations of beech in Central Italy. Journal of Heredity.
 2012 May 1;103(3):408-17.
- 563 Leonelli G, Pelfini M, di Cella UM, Garavaglia V. Climate warming and the recent treeline shift in the
- 564 European Alps: the role of geomorphological factors in high-altitude sites. Ambio. 2011 May
 565 1;40(3):264-73.
- 566 Lionello P, Baldi M, Brunetti M, Cacciamani C, Maugeri M, Nanni T, Pavan V, Tomozeiu R. Eventi
- 567 climatici estremi: tendenze attuali e clima futuro sull'Italia. In: Castellari S, Artale V (eds), I
- 568 cambiamenti climatici in Italia: evidenze, vulnerabilità e impatti. Bologna University Press, Bologna.
 569 2010;81-106.
- 570 Lionello P, Malanotte-Rizzoli P, Boscolo R (eds). Mediterranean climate variability. Elsevier,
- 571 Amsterdam. 2006.

- 572 Lippi MM, Guido M, Menozzi BI, Bellini C, Montanari C. The Massaciuccoli Holocene pollen
- 573 sequence and the vegetation history of the coastal plains by the Mar Ligure (Tuscany and Liguria, Italy).
- 574 Vegetation History and Archaeobotany. 2007 May 1;16(4):267-77.
- 575 Lombardi F, Chirici G, Marchetti M, Tognetti R, Lasserre B, Corona P, Barbati A, Ferrari B, Di Paolo S,
- 576 Giuliarelli D, Mason F, Iovino F, Nicolaci A, Bianchi L, Maltoni A, Travaglini D. Deadwood in forest
- 577 stands close to old-growthness under Mediterranean conditions in the Italian Peninsula. L'Italia Forestale
- 578 e Montana. 2013; 65(5):481-504.
- 579 Long JN, Vacchiano G. A comprehensive framework of forest stand property–density relationships:
- 580 perspectives for plant population ecology and forest management. Annals of Forest Science. 2014 May
- 581 1;71(3):325-35.
- Lovreglio R, Leone V, Giaquinto P, Notarnicola A. Wildfire cause analysis: four case-studies in
 southern Italy. iForest. 2010 Jan 22;3(1):8-15.
- 584 Luchi N, Capretti P, Feducci M, Vannini A, Ceccarelli B, Vettraino AM. Latent infection of
- 585 *Biscogniauxia nummularia* in *Fagus sylvatica*: a possible bioindicator of beech health conditions.
- 586 iForest. 2015;9:49-54.
- 587 Luchi N, Ghelardini L, Santini A, Migliorini D, Capretti P. First record of ash dieback caused by
- *Hymenoscyphus fraxineus* on *Fraxinus excelsior* in the Apennines (Tuscany, Italy). Plant Disease. 2016
 Jan 5:PDIS-09.
- Luchi N, Longa O, Danti R, Capretti P, Maresi G. *Diplodia sapinea*: the main fungal species involved in
 the colonization of pine shoots in Italy. Forest Pathology. 2014 Oct 1;44(5):372-81.
- 592 Magri D, Agrillo E, Di Rita F, Furlanetto G, Pini R, Ravazzi C, Spada F. Holocene dynamics of tree
- taxa populations in Italy. Rev Palaeobot Palynol. 2015; 218:267–284.

Mancini NM, Mancini GM, Travaglini D, Nocentini S, Giannini R. First results on the structure and
seed production of beech stands at the timberline in the Monti della Laga (Gran Sasso and Monti della

Laga National Park). Italian Journal of Forest and Mountain Environments. 2016 May 10;71(1):31-47.

- Marchetti M, Blasi C. Old-growth forests in Italy: towards a first network. L'Italia Forestale e Montana.
 2010 Dec 30;65(6):679-98.
- 599 Marchetti M, Tognetti R, Lombardi F, Chiavetta U, Palumbo G, Sellitto M, Colombo C, Iovino P,
- 600 Alfani A, Baldantoni D, Barbati A. Ecological portrayal of old-growth forests and persistent woodlands
- in the Cilento and Vallo di Diano National Park (southern Italy). Plant Biosystems. 2010 Mar
 1;144(1):130-47.
- Mensing S, Tunno I, Cifani G, Florindo F, Noble P, Sagnotti L, Piovesan G. Effects of human impacts
 and climate variation on forests: the Rieti basin since medieval time. Annali di Botanica. 2013 Mar
 19;3:121-6.
- 606 Mercuri AM, Sadori L. L'uso dei microresti vegetali per le ricostruzioni paleoambientali e per la
- valutazione degli effetti dell'attività antropica: l'esempio dei laghi vulcanici laziali. Miscellanea INGV,
 2013;18:128-33.
- MIPAAF. Inventario Nazionale delle Foreste e dei Serbatoi di Carbonio INFC Le stime di superficie
 (Prima Parte). MIPAAF Corpo Forestale dello Stato, ISAFA, Trento. 2007.
- Montanari C. Recent pollen spectra in two small mountain basins of the Ligurian Apennines (northern
 Italy). Grana. 1989 Dec 1;28(4):305-15.
- 613 Motta R, Berretti R, Borchi S, Bresciani A, Garbarino M, Trucchi D. Stand structure and coarse woody
- 614 debris profile of «La Verna» forest (Arezzo, Italy). L'Italia Forestale e Montana. 2013 May
- 615 16;65(5):591-605.

- 616 Muller SD, Nakagawa T, De Beaulieu JL, Court-Picon M, Carcaillet C, Miramont C, Roiron P,
- 617 Boutterin C, Ali AA, Bruneton H. Post-glacial migration of silver fir (Abies alba Mill.) in the
- 618 south-western Alps. Journal of Biogeography. 2007 May 1;34(5):876-99.
- 619 Munafò M, Riitano N, Fasio F, Bakudila Mbuta A, Sallustio L, Marchetti M. I comuni e le comunità
- appenninici: evoluzione del territorio. Stati generali delle comunità dell'Appennino. 2015;[online
- 621 reource] URL: <u>http://www.slowfood.it/wp-content/uploads/2015/10/Studio-Appennini-2015.pdf</u>. Çast
- accessed 12th October, 2016.
- 623 Negro M, Vacchiano G, Berretti R, Chamberlain DE, Palestrini C, Motta R, Rolando A. Effects of forest
- 624 management on ground beetle diversity in alpine beech (Fagus sylvatica L.) stands. Forest Ecology and
- 625 Managemenet 2015;328:300-9.
- 626 Nicolaci A, Travaglini D, Menguzzato G, Nocentini S, Veltri A, Iovino F. Ecological and anthropogenic
- drivers of Calabrian pine (*Pinus nigra* JF Arn. ssp. *laricio* (Poiret) Maire) distribution in the Sila
- 628 mountain range. iForest. 2014;8(4):497-508.
- Nocentini S. Structure and management of beech (*Fagus sylvatica* L.) forests in Italy. iForest. 2009 Jun
 10;2(3):105-13.
- 631 Paffetti D, Travaglini D, Buonamici A, Nocentini S, Vendramin GG, Giannini R, Vettori C. The
- 632 influence of forest management on beech (*Fagus sylvatica* L.) stand structure and genetic diversity.
- 633 Forest Ecology and Management 2012; 284:34-44.
- 634 Palombo C, Battipaglia G, Cherubini P, Chirici G, Garfi V, Lasserre B, Lombardi F, Marchetti M,
- 635 Tognetti R. Warming-related growth responses at the southern limit distribution of mountain pine (Pinus
- 636 *mugo* Turra subsp. *mugo*). Journal of Vegetation Science. 2014 Mar 1;25(2):571-83.
- 637 Palombo C, Chirici G, Tognetti R, Marchetti M. Is land abandonment affecting forest dynamics at high
- elevation in Mediterranean mountains more than climate change? Plant Biosystems. 2013;147 (1):1-11.

- Patrone G. Il contributo dello Stato e degli Enti alle sistemazioni montane ed al miglioramento ed
 ampliamento dei boschi e dei pascoli montani dal 1867 al 1950. Coppini, Firenze. 1953.
- 641 Pedrotti F, Gafta D. The high mountain flora and vegetation of the Apennines and the Italian Alps.
- 642 Ecological Studies. 2003 Jan 1;1(167):73-84.
- 643 Petit RJ, Brewer S, Bordács S, Burg K, Cheddadi R, Coart E, Cottrell J, Csaikl UM, van Dam B, Deans
- 544 JD, Espinel S. Identification of refugia and post-glacial colonisation routes of European white oaks
- based on chloroplast DNA and fossil pollen evidence. Forest ecology and management. 2002 Feb1;156(1):49-74.
- 647 Pezzi G, Bitelli G, Ferrari C, Girelli VA, Gusella L, Masi S, Mognol A. Pattern temporale del limite
- 648 altitudinale dei boschi di faggio nell'Appennino settentrionale. Un'analisi di dati fotogrammetrici.
- 649 Forest@. 2007 Mar 21;4(1):79-87.
- 650 Pezzi G, Maresi G, Conedera M, Ferrari C. Woody species composition of chestnut stands in the
- Northern Apennines: the result of 200 years of changes in land use. Landscape ecology. 2011 Dec
 1;26(10):1463-76.
- 653 Piccioli L. Selvicoltura. U.E.E.T., Torino. 1923.
- 654 Piermattei A., Lingua E., Urbinati C., Garbarino M. Pinus nigra anthropogenic treelines in the central
- Apennines show common pattern of tree recruitment. European Journal of Forest Research. 2016; in
- 656 press. Doi: 10.1007/s10342-016-0999-y.
- 657 Pignatti G La vegetazione forestale di fronte ad alcuni scenari di cambiamento climatico in Italia.
 658 Forest@. 2011;8:1-12
- 659 Piotti A, Borghetti M, Schettino A, Vendramin GG, Studio della variabilità genetica adattativa in specie
- 660 forestali: un approccio genomico per la definizione di strategie di conservazione per il pino loricato nel
- 661 Parco Nazionale del Pollino. L'Italia Forestale e Montana. 2014; 69(2):115-24.

- Piovesan G, Biondi F, Di Filippo AD, Alessandrini A, Maugeri M. Drought-driven growth reduction in
 old beech (*Fagus sylvatica* L.) forests of the central Apennines, Italy. Global Change Biology. 2008 Jun
 1;14(6):1265-81.
- Plini P, Tondi G. La distribuzione appenninica della Betulla bianca. Natura e Montagna. 1989;36(34):21-8.
- Puddu A, Luisi N, Capretti P, Santini A. Environmental factors related to damage by *Heterobasidion abietinum* in *Abies alba* forests in Southern Italy. Forest Ecology and Management. 2003 Jul
 17;180(1):37-44.
- 670 Ravazzi C. Late Quaternary history of spruce in southern Europe. Review of Palaeobotany and
- 671 Palynology. 2002 Jul 15;120(1):131-77.
- Ray N, Adams J. A GIS-based vegetation map of the world at the last glacial maximum (25,000-15,000
 BP). Internet Archaeology. 2001;11. URL:
- 674 <u>http://www.ncdc.noaa.gov/paleo/pubs/ray2001/ray_adams_2001.pdf</u>. Last accessed 12th October 2016.
- 675 Regione Toscana. D.P.G.R. 8 agosto 2003, n. 48/R. Regolamento Fore- stale della Toscana. Bollettino
- 676 Ufficiale della Regione Toscana. 2003;37:32–78
- 677 Romano D. I rimboschimenti nella politica forestale italiana. Quaderni di Monti e Boschi n. 3.
- 678 Edagricole, Bologna. 1987.
- 679 Rosenbaum G, Lister GS. Neogene and Quaternary rollback evolution of the Tyrrhenian Sea, the
- Apennines, and the Sicilian Maghrebides. Tectonics. 2004 Feb 1;23(1).
- 681 Rossignol-Strick M, Planchais N. Climate patterns revealed by pollen and oxygen isotope records of a
- Tyrrhenian sea core. Nature. 1989 Nov;342:413-6.

- 683 Sadori L, Allevato E, Bellini C, Bertacchi A, Boetto G, Di Pasquale G, Giachi G, Giardini M, Masi A,
- 684 Pepe C, Russo Ermolli E, Mariotti Lippi M. Archaeobotany in Italian ancient Roman harbours. Review
- of Palaeobotany and Palynology. 2015; 218:217–230.
- 686 Sadori L, Giraudi C, Masi A, Magny M, Ortu E, Zanchetta G, Izdebski A. Climate, environment and
- 687 society in Southern Italy during the last 2000 years. A review of the environmental, historical and
- archaeological evidence. Quaternary Science Reviews. 2016 Mar 15;136:173-88.
- 689 Schulz T, Krumm F, Bücking W, Frank G, Kraus D, Lier M, Lovrić M, van der Maaten-Theunissen M,
- 690 Paillet Y, Parviainen J, Vacchiano G. Comparison of integrative nature conservation in forest policy in
- 691 Europe: a qualitative pilot study of institutional determinants. Biodiversity and Conservation. 2014 Dec

692 1;23(14):3425-50.

- Seidl R, Schelhaas MJ, Rammer W, Verkerk PJ. Increasing forest disturbances in Europe and their
 impact on carbon storage. Nature Climate Change. 2014;4:806–810.
- 695 Tellini Florenzano G. Birds as indicators of recent environmental changes in the Apennines (Foreste
- 696 Casentinesi National Park, central Italy). Italian Journal of Zoology. 2004 Jan 1;71(4):317-24.
- 697 Temperli C, Bugmann H, Elkin C. Adaptive management for competing forest goods and services under
- 698 climate change. Ecological Applications. 2013;22:2065–2077
- 699 Tinner W, Colombaroli D, Heiri O, Henne PD, Steinacher M, Untenecker J, Vescovi E, Allen JR,
- 700 Carraro G, Conedera M, Joos F. The past ecology of *Abies alba* provides new perspectives on future
- responses of silver fir forests to global warming. Ecological Monographs. 2013 Nov;83(4):419-39.
- 702 Tinner W, Conedera M. Indagini paleobotaniche sulla storia della vegetazione e degli incendi forestali
- durante l'Olocene al Lago di Origlio (Ticino Meridionale). Bollettino della Società ticinese di Scienze
- 704 naturali. 1995;83(1/2):91-106.

- 705 Tonon G, Panzacchi P, Grassi G, Gianfranco M, Cantoni L, Bagnaresi U. Spatial dynamics of late
- successional species under Pinus nigra stands in the northern Apennines (Italy). Annals of Forest
- 707 Science. 2005 Nov 1;62(7):669-79.
- 708 Travaglini D, Paffetti D, Bianchi L, Bottacci A, Bottalico F, Giovannini G, Maltoni A, Nocentini S,
- 709 Vettori C, Calamini G. Characterization, structure and genetic dating of an old-growth beech-fir forest in
- the northern Apennines (Italy). Plant Biosystems. 2012 Mar 1;146(1):175-88.
- Triglia A, Iadanza C. Rapporto di sintesi sul dissesto idrogeologico in Italia 2014. Istituto Superiore per
 la Protezione e la Ricerca Ambientale, Dipartimento Difesa del Suolo, Roma. 2014.
- 713 Tzedakis PC, Bennett KD. Interglacial vegetation succession: a view from southern Europe. Quaternary
- 714 Science Reviews. 1995 Dec 31;14(10):967-82.
- 715 Uzunov D, Conti F, Lakusic D, Gangale C. Dati preliminari sulla fitogeografia, ecologia e
- conservazione delle specie Appenino-Balcaniche. Informatore Botanico Italiano. 2005;37(1):386-7.
- 717 Vacchiano G, Berretti R, Mondino EB, Meloni F, Motta R. Assessing the effect of disturbances on the
- functionality of direct protection forests. Mountain Research and Development. 2016 Apr 15;36(1):41-
- 719 55.
- 720 Vacchiano G, Garbarino M, Mondino EB, Motta R. Evidences of drought stress as a predisposing factor
- to Scots pine decline in Valle d'Aosta (Italy). European Journal of Forest Research. 2012a Jul

722 1;131(4):989-1000.

- Vacchiano G, Magnani F, Collalti A. Modeling Italian forests: state of the art and future challenges.
 iForest. 2012b;5(3):113.
- 725 Vacchiano G, Stanchi S, Marinari G, Ascoli D, Zanini E, Motta R. Fire severity, residuals and soil
- regardle regeneration of Scots pine in the Southern Alps. Science of The Total Environment. 2014

727 Feb 15;472:778-88.

- Valese E, Conedera M, Held AC, Ascoli D. Fire, humans and landscape in the European Alpine region
 during the Holocene. Anthropocene. 2014 Jun 30;6:63-74.
- 730 Valsecchi V, Finsinger W, Tinner W, Ammann B. Testing the influence of climate, human impact and
- fire on the Holocene population expansion of Fagus sylvatica in the southern Prealps (Italy). The
- 732 Holocene. 2008 Jun 1;18(4):603-14.
- Valva VL. Aspetti corologici della flora di interesse fitogeografico nell'Apennino Meridionale. Plant
 Biosystem. 1992 Jan 1;126(2):131-44.
- van Gils H, Batsukh O, Rossiter D, Munthali W, Liberatoscioli E. Forecasting the pattern and pace of
- 736 Fagus forest expansion in Majella National Park, Italy. Applied Vegetation Science. 2008 Dec
- 737 1;11(4):539-46.
- van Gils H, Odoi JO, Andrisano T. From monospecific to mixed forest after fire?: An early forecast for
 the montane belt of Majella, Italy. Forest Ecology and Management. 2010 Jan 25;259(3):433-9.
- 740 Venanzoni R, Pedrotti F, Manzi A. A relict of vegetational landscape related to seasonal migratory
- 741 grazing in the south of Italy: the 'Bosco dell'Incoronata' (Foggia) southern Italy. Landscape and Urban
- 742 Planning. 1993 Jul 31;24(1):55-62.
- 743 Vettraino AM, Morel O, Perlerou C, Robin C, Diamandis S, Vannini A. Occurrence and distribution of
- *Phytophthora* species in European chestnut stands, and their association with Ink Disease and crown
 decline. European Journal of Plant Pathology. 2005 Feb 1;111(2):169-80.
- 746 Vignali G, Barbarotti S, Piovani P, Maresi G, Salvadori C. Analisi dell'infestazione di *Ips typographus*
- nella foresta dell'Alta Val Parma e strategie per la ricomposizione del bosco. Forest@. 2015 Jun
 22;12(1):16-24.
- 749 Vizzarri M, Tognetti R, Marchetti M. Forest ecosystem services: issues and challenge for biodiversity
- conservation and management in Italy. Forests. 2015;6(6):1810-38.

- 751 Wang L, d'Odorico P, Evans JP, Eldridge DJ, McCabe MF, Caylor KK, King EG. Dryland
- ecohydrology and climate change: critical issues and technical advances. Hydrology and Earth System

753 Sciences. 2012 Aug 9;16(8):2585-603.

- 754 Watson CS. The vegetational history of the northern Apennines, Italy: information from three new
- requences and a review of regional vegetational change. Journal of Biogeography. 1996 Nov
- 756 1;23(6):805-41.
- 757 Westphal C, Tremer N, von Oheimb G, Hansen J, von Gadow K, Härdtle W. Is the reverse J-shaped
- 758 diameter distribution universally applicable in European virgin beech forests?. Forest Ecology and
- 759 Management. 2006 Mar 1;223(1):75-83.
- 760 Zanotti Cavazzoni. Boschi e ferrovie. L'Alpe, Bologna. 1907.
- 761 Ziaco E, Biondi F, Di Filippo A, Piovesan G. Biogeoclimatic influences on tree growth releases
- identified by the boundary line method in beech (*Fagus sylvatica* L.) populations of southern Europe.
- Forest ecology and management. 2012 Dec 15;286:28-37.
- 764

765 Figure captions

- Fig. 1 Mountain systems of the Italian peninsula, bounded by the 600 m a.s.l. elevation line
- **Fig. 2** Simplified pollen diagram (% total pollen) from Lagdei, northern Apennines (modified from
- 768 Ravazzi 2002; attribution of periods by Bertoldi 1981).
- **Fig. 3** Forest cover change in the 20th century in the Apennine from selected studies.
- Fig. 4 Disturbance chronologies produced for Valle Cervara using the boundary line method for high
- mountain beech populations in the Apennines. Moderate and major growth releases are those falling
- within 20–49.9% and 50–100% of the boundary line, respectively. Disturbance event dates (only when
- >=20 sampled trees) grouped into 5-year intervals (modified from Ziaco et al. 2012)
- 774

775 Appendices

- 776 Additional supporting information in the online version of this article (see "Supplementary
- 777 Material") contains the following: Appendix 1 List of references reviewed on Apennine forests.

778

Appendix 1 Click here to download Supplementary Material for online publication only: appendix1.docx







