

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Application of municipal biowaste derived products in Hibiscus cultivation: Effect on leaf gaseous exchange activity, and plant biomass accumulation and quality

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1619351> since 2022-01-24T20:14:38Z

Published version:

DOI:10.1016/j.scienta.2016.03.033

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Application of municipal biowaste derived products in *Hibiscus* cultivation: effect**
2 **on leaf gaseous exchange activity, and plant biomass accumulation and quality**

3

4 Daniele Massa^{1*}, Domenico Prisa¹, Enzo Montoneri², Daniele Battaglini¹, Marco Ginepro³, Michele
5 Negre⁴, Gianluca Burchi¹.

6

7 ¹ Council for Agricultural Research and Economics (CREA), Landscaping plants and nursery research unit,
8 Via dei Fiori 8, 51012 Pescia (PT), Italy

9 ²Biowaste Processing, Via XXIV Maggio 25, 37126 Verona, Italy

10 ³Dipartimento di Chimica, Università di Torino, Via P. Giuria 7, 10125 Torino, Italy

11 ⁴Dipartimento di Scienze Agrarie Forestali e Alimentari, Università di Torino, Largo P. Braccini 2, 10095
12 Grugliasco (TO), Italy

13

14 *Corrisponding author:

15 Daniele Massa

16 Tel +39 0572 451033

17 Fax +39 0572 453309

18 daniele.massa@entecra.it

19 **Abstract**

20 Two urban biowaste materials, fermented under anaerobic and aerobic conditions, their
21 soluble and insoluble alkaline hydrolysates, and a commercial biostimulant product, were compared
22 for their capacity to boost *Hibiscus* plants crop production and quality. Plants were grown in 4-litre
23 pots, containing peat and pumice as substrate, under optimal nutrient. A randomized-block
24 experimental design was adopted. Two types (main factors of variability) of treatments were applied,
25 i.e. by blending the above products with the substrate at transplant, and by fertigation using the soluble
26 hydrolysates only. Plant biomass characteristics and ecophysiological parameters were measured.
27 Mean effect of factors and their interactions were assessed by two-way ANOVA. Principal
28 component analysis was performed by using the different dependent variables to summarize the most
29 relevant differences among treatments. Compared with the control (untreated plants), the applied
30 treatments enhanced most of the investigated parameters. The most valuable effects were observed
31 for total biomass accumulation (+25 %), plant height (+10 %), leaf chlorophyll SPAD index (+15 %)
32 and net photosynthesis (+24 %). The hydrolysates performed better than the pristine materials. The
33 former ones were comparable to the commercial biostimulant. The results confirmed the hypothesis
34 that biowaste derived products induce biostimulant activity on *Hibiscus*; their application can
35 improve cultivation sustainability.

36 **Keywords**

37 Biowaste, biostimulant, photosynthesis, substrate culture, *Hibiscus palustris*

38

39 Abbreviations

| Acronym | Description | |
|------------------|--|--------------------------------------|
| D | Digestate obtained from organic humid fraction of urban waste | |
| DAT ₁ | Days after T ₁ | day |
| DW | Plant dry weight | g m ⁻² |
| E | Evapotranspiration rate | mmol m ⁻² s ⁻¹ |
| -F | Fertigation treatments | |
| FERT | Treatments applied during the cultivation through fertigation | |
| GC | Green compost obtained from urban gardening wastes | |
| G _s | Stomatal conductance | mmol m ⁻² s ⁻¹ |
| ID | Digestate insoluble alkaline hydrolysate | |
| IGC | Compost insoluble alkaline hydrolysate | |
| LAI | Leaf area index | m ² m ⁻² |
| NAR | Net assimilation rate | g m ⁻² d ⁻¹ |
| NO | Untreated | |
| NOVA | Commercial biostimulant | |
| OM | Organic matter | kg m ³ |
| P _n | Net photosynthetic rate | μmol m ⁻² s ⁻¹ |
| RGR | Relative growth rate | mg g ⁻¹ d ⁻¹ |
| SD | Digestate soluble alkaline hydrolysate | |
| SGC | Compost soluble alkaline hydrolysate | |
| SUB | Treatments applied to the substrate as a powder blended with the substrate | |
| T ₁ | Beginning of the experimentation (30 days after transplant) | day |

40

41

42 **Introduction**

43 Soluble biobased substances isolated from the alkaline hydrolysate of fermented urban biowastes
44 have been proven to perform as efficient ecofriendly chemical auxiliaries in diversified fields of the
45 chemical industry (Montoneri et al., 2011), agriculture (Sortino et al., 2014), and animal husbandry
46 (Montoneri et al., 2013). The products are mixtures of molecules with molecular weight from 5 to
47 several hundreds kDa, comprising aliphatic and aromatic C atoms bonded to a variety of acid and
48 basic functional groups. This is the likely reason of their multipurpose performance (Montoneri et al.,
49 2011). Particularly relevant for agriculture are the results published by Sortino et al. (2014) and Gomis
50 et al. (2014). Sortino et al. (2014) have shown that the above soluble biobased substances added to
51 soil increased tomato and red pepper plant photosynthetic activity, growth and productivity. The same
52 substances enhanced the photochemical degradation of organic pollutants in industrial effluent
53 (Gomis et al., 2014). These findings suggested that they might promote either C fixation or
54 mineralization, according to the different operational environments. In both cases, it was suggested
55 that, by their capacity to complex Fe ions and keep them in solution at slightly acidic or alkaline
56 conditions, the above soluble biobased substances may contribute to enhance photo-Fenton processes.
57 On this basis, the development of a self-sustainable ecosystem based on cycling renewable organic C
58 between wastes and added value products appears a fascinating reachable goal, certainly worthwhile
59 to pursuit. Generally, the environmental benefits of these substances lie in the fact that these products
60 are potentially alternative to commercial synthetic chemicals. Thus, by their use, one may expect a
61 reduction of fossil source depletion and of CO₂ emissions.

62 The realization of the above perspectives does not depend only on the product properties and
63 performance, but also on sustainability of the production process. Rosso et al. (2015) have reported
64 the hydrolysis of several materials recovered from different streams of a typical municipal waste
65 treatment plant located in North West Italy. These materials were the digestate of the plant biogas
66 reactor performing the anaerobic digestion of the organic humid fraction, and several composts

67 obtained from mixtures of digestate, private and public gardening residues, and/or sewage sludge.
68 The hydrolysis of these materials was carried out in water at pH 13 and 60 °C. In all cases, the soluble
69 hydrolysate was obtained together with equal or higher amounts of insoluble hydrolysate. The
70 published data showed that both the soluble and insoluble hydrolysates contain organic and mineral
71 matter, but the organic/mineral ratio is higher in the former. The soluble hydrolysates contain the
72 lowest relative amount of total mineral matter, compared with the insoluble hydrolysates and the
73 pristine sourcing materials. Although this process, performed at low temperature and with complete
74 solvent and reagents recycling, appears ecofriendly, a critical point remains the production of the
75 insoluble hydrolysate. In the absence of demonstration of a promising use of this product, its disposal
76 is an economic burden for the soluble hydrolysate production process. Yet, by its content of mineral
77 elements, also the insoluble hydrolysate product might be useful for agriculture.

78 The possibility of boosting yield and quality of crops by using biostimulants has been largely
79 investigated in the last years (e.g. Bulgari et al., 2015; Calvo et al., 2014; du Jardin, 2015). Many
80 relationships between the application of biostimulants and crop performance have been assessed in
81 several experimental trials. However, the mechanisms activated at plant level by these products are
82 largely unknown and difficult to identify. Biostimulants have been found to enhance plant growth
83 and/or production and quality (Calvo et al., 2014), nutrient uptake (Saa et al., 2015), photosynthetic
84 activity (Castro et al., 2012), plant response to biotic and abiotic stresses (Petrozza et al., 2014), and
85 several metabolic processes (Bulgari et al., 2015; Calvo et al., 2014). Most literature on biostimulants
86 reports their use by foliar application (e.g. Saa et al., 2015). However, the possibility of supplying
87 these substances directly to the root zone is valuable due to improved efficiency of distribution and
88 dosage. This is especially true if fertigation is adopted for cover fertilization; obviously, this practice
89 is restricted only to soluble compounds.

90 Much research activity on the use of biostimulants in agriculture has been carried out under soil
91 conditions for extensive crops (Calvo et al., 2014; du Jardin, 2015), while scarce literature is available
92 for substrate cultures (e.g. Bulgari et al., 2015). For the latter, root zone conditions are significantly

93 different from soil in terms of bio-organic process (e.g. microbial activity), climate (e.g. temperature,
94 humidity) and chemo-physical characteristics. In intensive agriculture, such as container-grown plant
95 systems, the required agrochemicals, water, and energy inputs per unit area are significantly higher,
96 compared with those needed for extensive crop systems. The former systems apply to ornamental
97 plants, for which the high aesthetic quality is a critical marketability parameter. In this scenario, using
98 natural products to boost plants high yield and quality could significantly improve the use efficiency
99 of non-renewable resources (e.g. fresh water and chemical fertilizers), thus increasing environmental
100 and economic sustainability. This poses worthwhile research scope on such products for
101 understanding better their action on plant through the root zone and optimizing their direct
102 administration to the substrate.

103 The present work was undertaken to address several questions posed in previous experiments on
104 the performance of the above soluble biowaste hydrolysates as promoter of plant photosynthetic
105 activity, growth and productivity, and on the sustainability of their production process. To this
106 purpose, *Hibiscus* was chosen as test plant. The soluble hydrolysates were obtained, together with the
107 corresponding insoluble hydrolysate, by alkaline hydrolysis of two different fermented municipal
108 biowaste materials. The soluble and insoluble hydrolysates, and their sourcing materials, were
109 distributed at transplant as powder blended with the substrate, and/or solubilised into the irrigation
110 water and supplied through fertigation. A commercial product, claimed by the vendor to have
111 biostimulant properties, was also tested by substrate application. This product, under the trade of
112 NOVA@@GR, is produced and distributed by Biolchim Spa, Medicina (BO), Italy. The applied
113 products were evaluated for their effects on several plant performance indicators related to plant
114 biomass accumulation and quality, and to leaf gas exchange activity. Main goals of the experiment
115 were: i) evaluation of the different urban biowaste sourced materials and/or their mixtures; ii)
116 assessment of the products application methodology; iii) evaluation of the above urban biowaste
117 sourced materials with respect to a commercial biostimulant.

118 **Materials and Methods**

119 *Location, plant material and growing conditions*

120 The experiment was carried out in the spring-summer of 2014, under typical Mediterranean
121 climate conditions, in a open-air experimental nursery located in Pescia (Tuscany, Italy; latitude 43.54
122 °N, longitude 10.42 °E) at the Landscaping Plants and Nursery Research Unit of the Italian Council
123 for Agricultural Research and Economics.

124 *Hibiscus* seedlings (*Hibiscus moscheutos* L. subsp. *Hibiscus palustris*) were transplanted on 10th
125 of April into four-L black pots (Ø 18 cm) containing a base substrate (peat 50% and pumice 50%,
126 V/V) adjusted to pH 6.

127 Plants were left 30 days under shade net (40% of shading) for acclimatisation to external (outdoor)
128 conditions. Thirty days after transplant (T₁), pots were moved to the experimental site and arranged
129 in a randomized-block experimental design, with three replicates per treatment (eight plants per
130 replicate). Pots were positioned at 0.40 x 0.60 m each other, obtaining a crop density of 4.2 plants m⁻².
131 Twenty days after T₁ (DAT₁), all plants were trimmed above the fourth true leaf to stimulate the
132 emission of lateral shoots as recommended by the standard cultivation technique in commercial
133 nursery production of *Hibiscus palustris*. The experiment was concluded at 110 DAT₁ with the
134 destructive analysis of plant material performed roughly one month before the beginning of plant
135 senescence (basing on local climate conditions).

136 Irrigation water was supplied through drip irrigation system (2 emitters per pot with a total flow
137 rate of 7.5 L h⁻¹, on average), using a timer for triggering irrigation four times per day (on average,
138 overall the cultivation period). Irrigation scheduling was adjusted weekly according to climate
139 conditions and leaching fraction (the ratio between water drainage and water supply), which was
140 calculated by measuring the volume of water drained out from a limited (three per block) number of
141 pots. Irrigation was regulated to keep leaching fraction on a target value of roughly 10-15%. Irrigation
142 water pH and electrical conductivity ranged 6.2-6.6 and 0.42-0.60 dS m⁻¹, respectively.

143 Prior to adding the biobased products under investigation, standard basic fertilization was applied
144 to the cultivation substrate, based on plant nutrient uptake, and taking into account irrigation water
145 characteristics and possible fertilizer losses due to leaching. However, the fertilization plan was
146 disposed with the aim of preventing any nutritional stress. Macro and micro-nutrients were in part
147 supplied at the transplant date, through controlled release fertilizers blended with the substrate (3 kg
148 m⁻³ of Osmocote Pro® 3-4 months + 3 kg m⁻³ of Osmocote Pro® 5-6 month). An additional
149 fertigation, between the exponential growth and incipient flowering phases, (from 40 to 50 DAT₁)
150 was performed with water soluble salts for a total supply of 0.21 N, 0.28 P₂O₅ and 0.13 K₂O kg m⁻³
151 of substrate.

152 Climate conditions were monitored continuously (5-minute basis) during the experiment by
153 recording radiation, relative humidity and temperature of the air through the on-site meteorological
154 station (Dacagon Device, Pullman, WA 99163 USA). Minimum, mean and maximum daily average
155 photosynthetic photon flux density values were 109.2, 568.3 and 750.5 μmol m⁻² s⁻¹, respectively.
156 Mean daily global radiation averaged 21.7 MJ m⁻² d⁻¹. Minimum, mean and maximum daily average
157 of air temperature values were 11.6, 20.5 and 22.3 °C, respectively. Mean daily relative humidity of
158 air averaged 64.5 %.

159 ***Biobased materials under investigation***

160 The soluble and insoluble hydrolysate products were produced and supplied by Studio Chiono
161 ed Associati in Rivarolo Canavese, Torino, Italy. They were obtained as previously reported by
162 Sortino et al. (2014). The pristine materials were the digestate recovered from the anaerobic
163 fermentation of the organic humid fraction of municipal solid waste from separate source collection,
164 and the green compost obtained by 180 days aerobic fermentation of private gardening and public
165 park trimmings. The digestate and the compost materials were processed in a pilot production facility
166 comprising a 500 litre capacity reactor. The solid material was treated under stirring with alkaline
167 water at 4 ml g⁻¹ liquid/solid and 0.02 w/w KOH/solid ratios at 60 °C for 4 h. The reaction mix was

168 allowed settling to separate the supernatant liquid containing the soluble hydrolysate fraction from
169 the solid insoluble hydrolysate residue. The latter residue was washed once with fresh water at 4 ml
170 g⁻¹ added water/insoluble hydrolysate ratio. The total collected liquid phase was centrifuged to
171 separate fine solid particles, and run through a 5 kDa cut off ultrafiltration polysulphone membrane.
172 The membrane retentate containing the soluble hydrolysate, and the solid insoluble hydrolysate were
173 dried in ventilated oven at 60 °C. The final dried products were analyzed according to previously
174 reported methods (Montoneri et al., 2011) and yielded the analytical values reported in Table 1 for
175 the sourcing digestate (D) and green compost (GC) materials, and for the respective hydrolysates, i.e.
176 the soluble (SD) and insoluble (ID) digestate hydrolysates, and the soluble (SGC) and insoluble (IGC)
177 compost hydrolysates.

178 ***Treatments***

179 Treatments were arranged based on the following criteria and/or objectives: i) to supply amounts
180 of treatments' products per unit area or plant, which were negligible, compared with the applied doses
181 of most biowaste sourced materials (such as compost), which are reported in literature (Dorais et al.,
182 2007); ii) to supply amounts of macronutrients contributed by the treatments' products, which were
183 negligible, compared with the normally applied amounts of chemical fertilizers; iii) to account for
184 effects of treatments' products as a function of their formulation and distribution mode; iv) to study
185 possible interactions between products obtained by different sourcing materials.

186 Two different types (factors of variability) of treatments were applied (Table 2): i) the substrate
187 treatments (SUB) comprising 11 treatments and the control, and the fertigation (FERT) treatments
188 comprising 2 treatments and the control. In the first case, the substrate at seedlings transplant was
189 blended with the different materials, which are listed in Table 2. These are the commercial NOVA
190 biostimulant, the green compost (GC), the insoluble compost hydrolysate (IGC), the soluble compost
191 hydrolysate (SGC), the compost mixed with its soluble hydrolysate (GC/S), the insoluble compost
192 hydrolysate mixed with the compost soluble hydrolysate (IGC/S), the digestate (D), the insoluble

193 digestate hydrolysate (ID), the soluble digestate hydrolysate (SD), the digestate mixed with the
194 soluble digestate hydrolysate (D/S), the insoluble digestate hydrolysate mixed with the digestate
195 soluble hydrolysate (ID/S). In the fertigation experiments, only the soluble compost (SGC-F) and
196 digestate (SD-F) hydrolysates versus the untreated control were tested. In this case, the total dose of
197 each soluble hydrolysate, which is reported in Table 2, was delivered through the irrigation system in
198 four weekly irrigations, starting at 50 DAT₁. The NOVA product was not included in the fertigation
199 experiments, as no claim for this application mode was reported in the vendor's product description
200 (Biolchim, 2015). The day before treatments, irrigation was interrupted, which allowed a loss of water
201 up to the same quantity of easily available water (roughly 400 ml pot⁻¹) in the pot volume. This
202 quantity was then recovered by supplying the same amount of water with the dispensed soluble
203 product solution. This expedient allowed preventing reduction of the applied dose per plant per week,
204 due to possible products' leaching.

205 Table 2 reports also dry matter, and organic matter (OM) and carbon (C) applied doses. These
206 were calculated basing on previous studies carried out with similar products in tomato and green
207 pepper cultivation trials (Sortino et al., 2013 and 2014). Based on the chemical composition of the
208 applied products (Table 1), dry matter doses were arranged in order to apply the same amount of OM
209 with each product. The only exception was the commercial NOVA product, which was inevitably
210 applied at the dose suggested by the producer. The products' mixtures for treatment 6, 7, 11 and 12
211 (Table 2) were made at 4/1 OM w/w ratio of the respective ingredients.

212 The above experimental plan, containing the two SUB and FERT factors of variability (Table 2),
213 allowed analysing 36 SUBx FERT combinations. In this fashion, three types of effects could be
214 investigated: i) effects due to the nature of the applied product; ii) effects related to the product
215 application mode; iii) effects from the possible interaction of the product nature and application mode.
216 In addition to what reported in Table 2, six more treatments were also carried out. In this case, the
217 control substrate and soluble hydrolysates in fertigation treatments were replicated but with a total
218 supply of nutrients equal to roughly 1/3 the dose reported in Table 2. These treatments were excluded

219 from statistical analyses, except for the linear regression analysis of photosynthetic rate and leaf tissue
220 characteristics. To this purpose, the additional treatments were necessary to have a wider range of
221 values, especially below optimal growing conditions (i.e. limited nutrient availability).

222 ***Plant biomass analyses***

223 Plant growth was monitored measuring leaf area index (LAI), by portable ceptometer
224 (AccuPAR LP-80, Decagon Devices Inc, Pullman WA 99163 USA), and plant height, twice per
225 month. Two destructive plant analyses, at 50 (just before the beginning of fertigation treatments) and
226 110 DAT₁ (at the end of the experiment) were performed to quantify the effect of treatments on
227 physiological and biometric parameters, biomass accumulation and partitioning, and tissue mineral
228 content of the plants. Before destructive analysis, SPAD index was measured through a portable
229 SPAD-502 (Konica Minolta Optics, 2970 Ishikawa-machi, Hachioji, Tokyo, Japan) on six healthy
230 leaves per plant, from the bottom to the top, and then averaging a total of 18 measurements per
231 replicate. Plant height, mean diameter (of the canopy projected to the soil) and number of flowers and
232 stems were measured. Plant height (cm pt⁻¹) and mean diameter (cm pt⁻¹) were used to calculate plant
233 ellipsoid volume (cm³ pt⁻¹) and the plant shape index (as the plant height to diameter ratio). Afterward,
234 the plant material was separated into flowers, stems and leaves, and weighted to assess biomass fresh
235 weight. A portion (roughly 150 g) of leaves was used for measuring unit leaf area through a leaf area
236 meter (WinDIAS Image Analysis System, Delta-T Devices, U.K.). Unit leaf area was used to
237 calculate leaf area index (LAI, m² m⁻²). Plant fresh material was dried in a forced-air oven (80°C for
238 72 h) and weighed for assessing plant dry matter partitioning.

239 Leaf tissues were analyzed for the organic nitrogen content (N) by Kjeldhal method. Leaf dry
240 matter (0.5 g) was weighted, put into pyrex tubes with a selenium catalyser (potassium sulphate 4.63
241 g, copper sulphate 0.28 g and selenium 0.09 g) and digested with 12 ml of phosphosulfuric acid at
242 370 °C for 40' using the VELP-K20 apparatus by VELP Scientifica, Usmate MB, Italy. At the end
243 of the digestion, the samples were distilled using the VELP-UDK127 apparatus (VELP Scientifica,

244 Usmate MB, Italy) after adding 40 ml of NaOH (40 % w/V). The distillate was collected in a conical
245 flask containing boric acid (4 % w/V) and bromocresol green methyl red colour indicator. Finally,
246 the content of N was determined by titration with 0.1 N HCl.

247 ***Leaf gas exchange measurements***

248 Leaf gas exchange measurements were performed in two different periods, during the
249 experiment, at 45 DAT₁, just before the first destructive plant analysis, or 105 DAT₁, at the end of
250 the experiment. A portable gas analyzer (Portable Photosynthesis System Ciras-2, PPSystems,
251 Amesbury, MA 01913 USA) was used to perform onsite measurements during the morning (between
252 9.00 and 12.00 am). During measurements, to maintain comparable analytical conditions, the
253 chamber was set at a constant value of light saturating photosynthesis (PPFD = 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$,
254 primarily determined through photosynthesis light-response curves), CO₂ (400 g m^{-3}), vapour
255 pressure deficit (VPD = 1.0 \pm 0.2 kPa) and temperature (27.5 \pm 1 °C). The temperature value was the
256 average of climate data recorded with a datalogger, in the same daily period of measurements, over
257 the three days prior to measurement start up. Two mature and healthy leaves (second and fourth leaf,
258 completely unfolded above the apex of the main stem) were chosen for gas exchange analysis. For
259 these measurements, two plants per replicate were used. A total of twelve measurements per treatment
260 combination at each stage was carried out. This procedure provided acquisition of values for current
261 net photosynthetic (carbon assimilation) rate (Pn, $\mu\text{mol m}^{-2} \text{s}^{-1}$), transpiration rate (E, $\text{mmol m}^{-2} \text{s}^{-1}$)
262 and stomatal conductance (Gs, $\text{mmol m}^{-2} \text{s}^{-1}$). The net photosynthetic rate values were finally used
263 for the computation of water use efficiency (Pn E⁻¹, $\mu\text{molCO}_2 \text{mmol}^{-1}\text{H}_2\text{O}$).

264 ***Statistics***

265 Collected data were analysed by ANOVA two-ways variance analysis, in order to assess
266 significant ($P \leq 0.05$, 0.01 and 0.001) differences among treatments. Mean values were then separated
267 by Duncan's multiple range test ($P = 0.05$). Data analysis included also multiple-variable analysis
268 (correlation analysis), variables' relationships by linear regressions and principal component analysis

269 (PCA). Statistics and graphics were supported by the programs Statgraphics Centurion XV (Stat
270 Point, Inc., Herndon, VA, USA) and Prism 5 (GraphPad Software, Inc., La Jolla, California USA).

271 **Results**

272 ***Chemical composition of biowaste and commercial products.***

273 Table 1 reports the chemical composition of the municipal biowaste and the commercial
274 products tested for their effects in the cultivation of *Hibiscus*. All contain organic and mineral matter.
275 NOVA is clearly distinguished for the highest content of OM, the relatively lowest organic C and
276 highest N contents. Indeed, the NOVA OM, C/N and OM/(C+N) values are 73.8 %, 4.5 and 2.6,
277 respectively, while the values for the biowaste products range from 50.7 to 66.4 % OM, from 6 to
278 22.1 C/N and from 1.4 to 1.8 OM/(C+N). These features reflect the different nature of the products'
279 sources. In essence, the municipal biowaste products are derived from mixtures of unsorted kitchen
280 wastes and gardening residues, pre-treated by anaerobic and/or aerobic fermentation followed by
281 chemical hydrolysis. The commercial NOVA product (Biolchim, 2015) is described by the vendor as
282 a proprietary mix of plant extract rich in fulvic acids, humic acids, amino acids and glycine betaine
283 as biostimulants. Also, the biowaste products present differences, one from the other. Compared with
284 the pristine digestate (D) and compost (GC) materials, and with the insoluble digestate (ID) and
285 compost (IGC) hydrolysates, the soluble digestate (SD) and compost (SGC) hydrolysates have the
286 highest content of organic matter, C and N. Compared with the soluble digestate hydrolysate (SD),
287 the soluble compost hydrolysate (SGC) has higher content of all mineral elements, except for P and
288 N, which are higher in the soluble digestate hydrolysate (SD). It is also worthwhile to observe that,
289 compared with all other products in Table 1, the insoluble compost hydrolysate (IGC) has the highest
290 content of Si and Fe, followed by the pristine compost (GC) in the order of decreasing Si and Fe.

291 ***Biomass accumulation and biometric parameters***

292 Data collected during the first destructive analysis, carried out in the middle of the experimental
293 period, before the initiation of fertigation treatments (50 DAT₁), did not show any significant effect
294 on biomass accumulation and other biometric parameters by the substrate treatments (data not
295 shown). On the contrary, the data (Table 3) collected by destructive analysis, performed at the end of
296 the experiment (110 DAT₁), demonstrated significant effects on dry biomass accumulation and leaf
297 area index (LAI). Table 3 shows that all substrate treatments, except treatment 7 by the mix of
298 insoluble and soluble compost hydrolysates, increased significantly the leaf dry weight (DW),
299 compared with the control no treatment 1. Significant increases in leaf DW were also observed by
300 the fertigation treatments 14 (SGC-F) and 15 (SD-F), compared with the control plants. Similar
301 results were found for LAI (Table 3). A significant positive correlation between LAI and leaf DW (R
302 = 0.90 and $P < 0.001$, data not shown) was found. In particular, in the fertigation treatments 14
303 (SGC-F) and 15 (SD-F), the soluble compost and soluble digestate hydrolysates, respectively, caused
304 30.2 % higher LAI, compared with the control plants in the no treatment 13 (NO-F).

305 The stem and flower DW, and the total shoot DW (Table 3) showed higher selectivity in
306 establishing the ranking order of the treatments effects. Shoot DW correlated better with stem and
307 flower DW ($R = 0.99$, $P < 0.001$) than with the leaf DW ($R = 0.92$, $P < 0.001$). In the substrate
308 treatments, the highest increase (25-26 %) was caused by the soluble digestate hydrolysate (SD)
309 treatment 10 only, compared with the control no treatment 1. For the stem and flower DW, the other
310 treatments did not appear to cause significantly different effects, compared with the control no
311 treatment 1. For the total shoot DW, only the commercial NOVA, and the insoluble (ID) and soluble
312 (SD) digestate hydrolysates caused significant increase, compared with the control no treatment 1. In
313 the fertigation treatments 14 (SGC-F) and 15 (SD-F) by the soluble compost and soluble digestate
314 hydrolysates caused significant 24.2 % stem and flower DW, and 22.4 % total shoot DW increase,
315 compared with the control no treatment 13 (NO-F).

316 Unlike the significant differences found in DW accumulation, the different treatments did not
317 affect at all neither biomass partitioning nor the percentage of DW in plant tissues (data not shown).
318 The latter was 17.2 % for flowers, 27.3 % for leaves, and 29.8 % for stems, calculated as the average
319 of all treatment combinations.

320 Neither the substrate nor the fertigation treatments influenced significantly the main biometric
321 parameters, which were measured during the first destructive analysis (50 DAT₁). In the substrate
322 treatments, the only statistically significant proven effect was the plant volume increase, which was
323 caused by the commercial NOVA product in treatment 2 at the end of the cultivation cycle (Table 3).
324 On the other hand, the soluble compost and soluble digestate hydrolysates, which were supplied
325 through fertigation treatments 14 (SGC-F) and 15 (SD-F), increased significantly plant height and
326 volume (Table 3), compared with the control no treatment 13 (NO-F).

327 Table 3 shows also that the plant shape index was reduced by the fertigation treatments 14
328 (SGC-F) and 15 (SD-F), performed with the soluble compost and soluble digestate hydrolysates,
329 compared with the control no treatment 13 (NO-F). Indeed, the control plants showed larger and
330 shorter vegetative habitus, compared with the plants treated by fertigation.

331 ***Leaf analysis and gas exchange activity***

332 The experimental data collected for leaf chlorophyll and N content, and for leaf gas exchange
333 activity are reported in Table 4 and 5, respectively. Chlorophyll (represented in this work by the
334 SPAD index) and N content in leaf samples, did not show significant differences among all treatments
335 in the first destructive plant analysis at 50 DAT₁. However, significant differences were observed
336 later during the last period of analyses (Table 4). In the substrate treatments, the insoluble compost
337 hydrolysate (IGC) treatment 4, the mix of the insoluble soluble hydrolysate (ID/S), and the soluble
338 compost hydrolysate (SGC) treatments 12 and 5, respectively, caused the highest SPAD value
339 compared with the control no treatment 1. For leaf N, only the insoluble compost hydrolysate (IGC)
340 treatment 4, the mix of the insoluble and soluble digestate hydrolysates (ID/S), and the commercial

341 NOVA product treatments 12 and 2, respectively, were significantly higher than the control no
342 treatment 1. In the fertigation treatments, the soluble compost hydrolysate treatment 14 (SGC-F) gave
343 the highest SPAD value. This was 4.4 % and 10.8 % higher than the respective values, which were
344 recorded for the soluble digestate hydrolysate treatment 15 (SD-F) and the control no treatment 13.
345 For leaf N, no fertigation treatment gave significantly different values from the value that was
346 recorded for the control no treatment 13 (NO-F). The data in Table 4 also show that an interaction
347 was found between the main experimental factors for chlorophyll SPAD index; such an interaction
348 was due to the significantly higher response of the soluble compost (treatment SGC-F) and the soluble
349 digestate (treatment SD-F) hydrolysates, when combined with the insoluble digestate hydrolysate
350 (ID). A heterogeneous response was observed for the other combinations (data not shown).

351 Leaf N concentration was highly correlated with SPAD ($R = 0.85$, $P < 0.001$) while a poor,
352 although significant, correlation was found with leaf DW ($R = 0.46$, $P = 0.002$) and total shoot DW
353 ($R = 0.59$, $P < 0.001$) as can be also deduced by Fig. 1. The relationship between SPAD and N was
354 significantly ($P < 0.001$) represented by a linear equation ($\text{SPAD} = 13.09N + 0.72$) with a
355 determination coefficient explaining 73 % of the experimental variability (Fig. 1).

356 Gas exchange activity measured at 45 DAT₁ did not show any significant difference among
357 treatments for the measured photosynthetic rate (Pn), stomatal conductance (Gs), and
358 evapotranspiration (E). However, the measurements at 105 DAT₁ (Table 5) showed significant
359 treatment effects. Substrate treatments caused significant increases of the photosynthetic activity,
360 compared with the control no treatment 1. Photosynthetic rate of treated plant averaged $20.4 \mu\text{mol m}^{-2}$
361 s^{-1} , compared to $17.4 \mu\text{mol m}^{-2} \text{s}^{-1}$ for untreated plants. The compost (GC) treatment 3 gave the
362 highest 24.3 % Pn increase, compared with the control no treatment 1. In the fertigation treatments,
363 the soluble compost and soluble digestate treatments 14 (SGC-F) and 15 (SD-F) gave significantly
364 higher average 22.7 % Pn increase, compared with the control no treatment 13 (NO-F). A
365 heterogeneous plant response to the different treatment combinations was highlighted by a significant
366 interaction of the main experimental factors. In this case, the general tendency of soluble compost

367 and soluble digestate fertigation treatments 14 (SGC-F) and 15 (SD-F) to increase Pn was
368 significantly more pronounced for the following combinations: treatments 1x14, 5x14, 8x14, 11x14
369 and 9x15 (see Table 2 for combinations) with respect to untreated plants (data not shown). These
370 results were generally denoting that: i) soluble compost had a major effect on Pn with respect to
371 soluble digestate; ii) combinations of products (between substrate and fertigation treatments) derived
372 from the same organic matrixes had major effects respect to their mixtures. The collected data showed
373 a strong correlation between Pn and SPAD ($R = 0.87$, $P < 0.001$) as confirmed by data analysis
374 reported in Fig. 1, and a poorer although significant correlation with total shoot DW accumulated at
375 the end of the cultivation cycle ($R = 0.65$, $P < 0.001$).

376 Stomatal conductance (Table 5) was significantly increased, by 45 % on average, only by the
377 insoluble compost hydrolysate (IGC) and the GC treatments 4 and 3, respectively, compared with the
378 control no treatment 1. In the fertigation treatments, the soluble compost hydrolysate treatment 14
379 (SGC-F) increased Gs by 25.7 % and 33.5 %, compared with the soluble digestate hydrolysate
380 treatment 15 (SD-F) and to the control no treatment 13 (NO-F), respectively. A roughly similar plant
381 response to the different treatments was observed for the evapotranspiration rate (E) reported in Table
382 7. In the fertigation treatments, the soluble compost hydrolysate treatment 14 (SGC-F) increased the
383 evapotranspiration rate by 28.2 and 44.4 %, compared with the soluble digestate hydrolysate
384 treatment 15 (SD-F) and the control no treatment 13 (NO-F), respectively. The evapotranspiration
385 rate and the stomatal conductance were found significantly correlated ($R = 0.61$, $P < 0.001$; data not
386 shown).

387 Finally, in the substrate treatments, the crop water use efficiency values (Table 5) showed no
388 significant difference among most of treatments. The only exception was treatment 12 by the mix of
389 the insoluble and soluble digestate hydrolysates (ID/S), which caused a 27.1 % significantly higher
390 water use efficiency value, compared with the control no treatment 1. In the fertigation treatments,
391 the soluble digestate hydrolysate (SD) treatment 15 (SD-F) caused 16.9 % significantly higher water
392 use efficiency value compared with the control no treatment 13 (NO-F). On the contrary, the soluble

393 compost hydrolysate treatment 14 (SGC-F) caused 20.1 % significant decrease of water use
394 efficiency, compared with the control no treatment 13 (NO-F).

395 ***Treatments' ranking order and principal component analysis***

396 Summarizing the data reported in Table 3, 4 and 5, it may be observed that, for substrate
397 treatments, the biowaste sourced products rank significantly first in most cases. Exceptions are the
398 plant volume, for which the commercial NOVA biostimulant ranks first, and the plant height and
399 shape index, which showed no treatment effects. The soluble (SD) and/or insoluble (ID) digestate
400 hydrolysates rank significantly first for their effects on four indicators connected to plant biomass
401 accumulation and biometric parameters (Table 3). These are the leaf, stem and flowers, and total
402 shoot DW, and LAI. The insoluble compost hydrolysate (IGC) and/or the compost (GC) rank
403 significantly first for the effects connected to the Pn, leaf N and SPAD index (Table 4 and 5). The
404 ranking order of the fertigation treatments summarizes the results described in the above subsections.
405 In essence, the soluble digestate and compost hydrolysates treatments 14 (SGC-F) and 15 (SD-F),
406 compared with the control no treatment 13 (NO-F), showed significant effects on all indicators. The
407 only exception was the leaf N content, for which no significant difference occurred among treatments
408 14 and 15, and the no treatment 13 (Table 4). Treatment 15 was better than treatment 14, for the effect
409 on plant height and volume. On the contrary, for the effects on Gs, E and SPAD index, the soluble
410 compost hydrolysate in treatment 14 (SGC-F) performed better than the soluble digestate hydrolysate
411 in treatment 15 (SD-F).

412 The above-summarised results were used as a base for the interpretation of relationships among
413 treatments, between treatments and nature of the pristine materials, and between treatments and
414 investigated variables. To this end, principal component analysis (PCA) was performed by using the
415 investigated parameters to summarise main differences among treatments and sourcing materials
416 under investigation (Fig. 1). As reported in Fig. 1, the first principal component 1 (PC1) separates
417 treated and untreated plants. Such a separation is more marked for fertigation treatments that are

418 grouped in three main areas related to: i) no treatment 13 (100 % of its combinations); ii) soluble
419 green compost hydrolysate treatment 14 (92 % of its combinations); iii) soluble digestate hydrolysate
420 treatment 15 (83 % of its combinations). The latter two treatments were then separated by the second
421 principal component 2 (PC2). According to the loadings (Fig. 1), treatment combinations with soluble
422 green compost hydrolysate (treatment 14) were more correlated with ecophysiological (Pn, E and Gs)
423 parameters and tissue characteristics (leaf N content and SPAD) than combinations treatments
424 obtained by soluble digestate treatment 15. However, by PC1 it was clear that both treatments were,
425 in general, more correlated with biomass accumulation, plant height and volume, SPAD, leaf N
426 content, and photosynthesis than untreated plants (no treatment 13). Within the three main clusters,
427 treatment combinations with NOVA and hydrolysate products were more correlated with the above
428 parameters than untreated plants (control treatment 1), which performed at the lowest level. More in
429 details, hydrolysates products (treatment 4, 5, 9 and 10) were, on average, more correlated with the
430 above parameters than pristine materials (treatment 3 and 8) and/or mixed treatments (treatment 6, 7,
431 11 and 12) that instead gave heterogeneous responses.

432 **Discussion**

433 To avoid any nutritional plant stress, the crop received an optimal amount of nutrients by standard
434 chemical fertilization (see Materials and Methods, growing conditions section). The applied products
435 under investigation (Table 1) contained a quantity of potential plant nutrients (for instance roughly
436 2-6 % N on dry matter basis) that varied, as a function of product chemical nature. As well known,
437 any increase in nutrient supply, above an optimal threshold, does not produce any significant effect
438 on plant yield and dry biomass accumulation (Massa et al., 2009; Reid 2002; Silberbush et al., 2005).
439 Therefore, the amounts of potential additional nutrients, which were contributed by the added
440 experimental biowaste and commercial NOVA products (Table 1 and 2), were deemed negligible
441 (2-5 %) compared with the amounts of nutrients, which were available to the crop by the standard
442 chemical fertilization. This consideration appears highly relevant for the following discussion.

443 The data reported in Table 3-7 show that the plant response to the applied treatments was
444 generally positive, showing higher accumulation of fresh (data not shown) and dry biomass and
445 photosynthetic activity, compared with untreated plants. Considering the relatively low amount
446 (Table 2) of products blended with the substrate (in the substrate treatments), and/or delivered through
447 the fertigation system (in the fertigation treatments), the increase in most of the measured plant
448 parameters (Tables 3-5) was remarkable. In fact, the total quantity of products used was in the order
449 of 5-15 % of the minimum dose for common organic fertilizers and/or generic amendments, which
450 are normally applied in agriculture (Dorais, 2007).

451 In recent on-field red pepper and tomato cultivation trials, Sortino et al. (2013 and 2014) found
452 that the application of 700 kg ha⁻¹ of urban biowastes' hydrolysates did not alter significantly the soil
453 chemical composition. Yet, plant photosynthetic activity, growth, and productivity were enhanced
454 significantly by the added soluble hydrolysates. The authors concluded that the tested products
455 enhanced the plant photosynthetic activity and that, in turn, the increase of photosynthetic activity
456 was the main factor increasing plant growth and productivity. More recently (Fascella et al., 2015),
457 this hypothesis was confirmed by applying the same soluble hydrolysates for the cultivation of
458 *Euphorbia* in pots. The data collected on *Hibiscus* seem to support the role of the tested products as
459 promoters of the plant photosynthetic activity (Table 5). In relation to these previous works, and to
460 the present work, there are two key issues deserving further discussion. These are (i) the relationship
461 between leaf chemical features, gas exchange activity, and biomass accumulation, and (ii) the
462 relationship of the observed effects with the applied products chemical nature and/or composition.

463 Several authors (Bulgari et al., 2015; Calvo et al., 2014; du Jardin, 2015; Ertani et al., 2013a)
464 have associated the increase in dry biomass of agricultural crops, treated with products derived from
465 organic matrixes, to the presence of organic molecules that stimulate plants. More generally,
466 biostimulant substances augment biomass accumulation and yield, improve nutrient uptake and
467 enhance metabolic functions. Also, many biostimulant organic substances, such as algae and/or
468 seaweed extract, soil and water humic and fulvic acid, have been found to improve photosynthetic

469 capacity in several different plant species, which are grown either under optimal conditions (Castro
470 et al., 2012; Jannin et al., 2012) or in presence of abiotic stress (Anjum et al., 2011; Ertani et al.,
471 2013b). Humic-like substances, obtained from different organic wastes, fall in this category, either
472 for their chemical nature or for their biostimulant activity (Eyheraguibel et al., 2008; Morard et al.,
473 2011). The biowaste derived products tested in this work belong to the category of “complex organic
474 material” (du Jardin, 2015) and bear structural similarities with natural humic substances (Montoneri
475 et al., 2011). They are rich in molecules, which are typically found in products claimed to have
476 biostimulant activity (e.g. Biolchim, 2015).

477 For the indicators connected to *Hibiscus* biomass accumulation in the substrate treatments,
478 the products’ ranking in order of decreasing effect changes, depends on the plant parameter and organ
479 (Table 3). By comparison, in the fertigation treatments, the soluble products (SGC-F and SD-F)
480 improved most of the above indicators, compared with the control (NO-F). However, it should be
481 highlighted that fertigation treatments practically received a quantity of products higher than substrate
482 treatments (Table 2), which could raise the hypothesis of an additive and/or dose response effect of
483 the tested products as observed for example by Sortino et al. (2013) with soluble hydrolysates. As
484 matter of fact, plant response to fertigation treatments was more pronounced and defined than for
485 substrate treatments (Tables 3-5 and Fig. 1).

486 Attempts to correlate the observed effect ranking order with the products’ chemical
487 characteristics (Table 1) did not allow assessing definite clear product-properties relationships. It is
488 true that the investigated products differ for the concentration of macro- and micro-nutrients.
489 However, when the products are supplied to the plants at the doses reported in Table 2, the mineral
490 composition differences are likely to be levelled out by the higher relatively amount of nutrients
491 supplied by the conventional chemical fertilizers. It was worth to determine the amount of beneficial
492 elements such as Se and Si among the oligoelements in the tested products. Selenium can have a
493 growth-promoting effect for many plant species (Pilon-Smits et al., 2009), for example by increasing
494 N reductase activity (Nowak et al., 2004). It naturally occurs as trace element in most soils, at typical

495 levels below 1 mg kg⁻¹ (Pilon-Smits et al., 2009). Selenium in the investigated products (Table 1) has
496 been found slightly above this level, only in the soluble digestate hydrolysate (SD) and the
497 commercial product (NOVA). However, neither any significant variation of Se nor correlation with
498 N content was found in leaf tissues (data not shown). These findings did not help to explain the
499 products' performance ranking order and were in agreement with previous works (Hawrylak-Nowak
500 et al., 2015; Rios et al., 2013).

501 Leaf N correlated (Fig. 1) well with SPAD, but less well with Pn, and biomass accumulation
502 parameters. Weak correlations between Pn and leaf N were observed by other authors (e.g. Kenzo et
503 al., 2015). In the present experiment, it is likely that the optimal nutrient availability in the root
504 allowed plants taking up the needed quantity of nutrients to support the higher growth rate induced
505 in treated plants, without affecting leaf N content (Table 4). The high correlation between Pn and
506 SPAD, as observed in the present work (Fig. 1), is usual. Very often, the increase in Pn, in
507 biostimulated plants, is coupled to higher chlorophyll and nutrient content in plant tissues (Bulgari et
508 al., 2014; Calvo et al., 2014). To this purpose, most of literature is based on the study of extensive
509 crops, cultivated on soil and treated through foliar application (e.g. Anjum et al. 2011; Jannin et al.,
510 2013). By comparison, very little is known on potted ornamental plants cultivated in substrate and
511 treated directly in the root zone.

512 The positive influence of tested products on chlorophyll content and Pn (Table 4 and 5) is in
513 agreement with previous works conducted on plant biostimulation obtained with various substances
514 and products (e.g. Amanda et al., 2009; Anjum et al. 2012; Ertani et al., 2013b; Fascella et al., 2015;
515 Jannin et al., 2013). Furthermore, the relationship between SPAD index and leaf colour (e.g.
516 Papasavvas et al., 2008; Shibayama et al., 2012) helps to assess a relevant commercial parameter for
517 ornamental plants such as leaf greenness. To this purpose, Loh et al. (2002) proposed SPAD meter as
518 effective tool for the evaluation of plant quality in ornamental crops.

519 The data collected on tissue characteristics and leaf gas exchanges (Table 4 and 5) support the
520 intriguing role of the biowaste derived products as photosensitizers (Bianco Prevot et al., 2011, and

521 Gomis et al., 2014), promoters of photosynthetic activity, in agreement with Sortino et al. (2013 and
522 2014) and Fascella et al. (2015). This hypothesis is consistently with similar findings for humic acids
523 (Bulgari et al., 2015; Calvo et al., 2014), bearing structural similarities with the above biowaste
524 product (Montoneri et al., 2011).

525 The compost (GC) and the insoluble compost hydrolysate (IGC) rank first for the effects on
526 Pn in the substrate treatments (Table 5). These data were consistent with PCA shown in Fig. 1. Table
527 1 shows that these products contain the highest amounts of Fe and Si, compared with all other tested
528 products. The contribution of the compost (GC) and insoluble compost hydrolysate (IGC) to the total
529 amount of Fe and Si was roughly 64 % and 177 %, respectively, higher than the average value
530 contributed by the other applied products listed in Table 1. Both Si (Houben et al., 2013) and Fe
531 (Miller et al., 1995) are known to have important role in photosynthesis. Gomis et al. (2014) have
532 proposed that Fe in the soluble compost hydrolysate is responsible of the photosensitizing properties
533 of this material, which were demonstrated for the remediation of chemical industrial waste waters
534 containing organic pollutants. Other works (du Jardin, 2015; Li et al., 2015; Pilon-Smits et al 2009)
535 report that Si is a beneficial element for plants, inasmuch as it plays a key role against several biotic
536 and abiotic stress, and promotes plant growth and development, chlorophyll concentration and
537 photosynthetic activity. The data collected in the present work seem consistent with the above
538 literature findings. However, considering all treatments, a definite correlation of photosynthetic
539 activity and/or SPAD index vs. the amounts of applied Si and Fe amounts could not be established.
540 Very likely, the effects of the above mineral elements do not rely to the applied amounts, but depend
541 also on the nature and solubility of the organic matter to which they are bonded.

542 The confirmation that the biowaste-derived products, which were tested in the present work,
543 promote plant photosynthetic activity is certainly relevant. Nevertheless, under the adopted
544 experimental conditions, many other factors could contribute to the observed plant growth and
545 development. The weak, although significant, correlation found between Pn and the total shoot DW
546 ($R = 0.65$, $P < 0.001$; see also Fig. 1) leads to suppose that not all applied products in the present work

547 were able to induce a real improved efficiency in the conversion of carbon assimilates in plant
548 biomass. As matter of fact, only the soluble (SD) and insoluble (ID) digestate hydrolysates, and the
549 commercial NOVA biostimulant, in the substrate treatments, and the soluble compost (SGC) and
550 soluble digestate (SD), in fertigation treatments, caused a significant higher accumulation in total
551 shoot DW (Table 3). This could also imply likely further positive effects by the above mentioned
552 treatments on different aspects of plant primary and secondary metabolism, other than photosynthesis.
553 For example, other complex organic materials have been found improving nutrient uptake (e.g. Ertani
554 et al., 2013b; Morard et al., 2011) and other metabolic processes (du Jardin, 2015; Ertani et al.,
555 2013a). On the other hand, a high carbon intake does not necessarily results in high long-term carbon
556 storage. Indeed, not all assimilated CO₂ is converted in structural biomass due to different respiration
557 efficiency during dark hours and carbon allocation to the secondary metabolism, as for example
558 related to volatile organic compounds (Herms and Mattson 1992). However, the use of products
559 stimulating photosynthetic activity appears of great interest to ornamental cultivation for boosting
560 high yield and quality, and for improving the efficiency of the production process. Photosynthesis is
561 a process very sensitive to biotic and abiotic stresses (Ashraf and Harris, 2013). The application of
562 the tested products could be a practice for overcoming and/or containing moderate stressful
563 conditions, which affect photosynthesis (e.g. Flexas et al., 2004; Lieth and Pasian 1991; Yamori et
564 al., 2005), with positive effects on production sustainability and market competition (e.g. shorter
565 cultivation cycle).

566 In the present work, together with the highest enhancement of Pn, the compost-derived
567 products caused the highest E and Gs values compared to untreated plants, thus decreasing crop water
568 use efficiency (Tables 5). Digestate-derived products resulted more efficient in enhancing the crop
569 water use efficiency. Generally speaking, the supply of water can not be a limiting factor in
570 container-grown plants. This is especially the case of ornamentals where the aesthetic value of the
571 plant is the fundamental market parameter. However, water use efficiency must be highly considered
572 in agriculture for its environmental implications.

573 The collected data allow drawing some conclusions on some important issues, which have
574 both economic and environmental relevance. These are related not only to agriculture, but also to
575 urban biowastes processes and product development. One main issue is the value of the biowaste
576 hydrolysates, compared with their sourcing materials. The data reported in Tables 3-5 and Fig. 1
577 indicate that the soluble and/or insoluble hydrolysates appear more efficient than the sourcing
578 digestate or compost materials. Undoubtedly, the most remarkable and defined effects were obtained
579 with soluble hydrolysates applied by fertigation (Fig. 1); this effect was also due to a likely higher
580 bioavailability of biostimulant substances. As matter of fact, PCA (Fig. 1) highlighted the presence
581 of three main clusters, related to fertigation treatments; treated plants were more correlated with
582 biomass accumulation, plant height and volume, SPAD, leaf N content, and photosynthesis than
583 control treatment 13. Within these macro-clusters, hydrolysate products were more correlated with
584 the above parameters than untreated plants and/or plants treated only with the pristine materials. The
585 evidence that both the soluble and insoluble hydrolysates are valuable products, more than their
586 sourcing biowaste materials, offers positive prospects for their production process.

587 A further issue addressed by the present work is how general the effects of the
588 biowaste-derived products are for different plant species (ornamentals), considering previous
589 experiments reported on other plant species with the same or similar fermented biowastes and
590 hydrolysates. Fascella et al. (2015) compared the same soluble digestate hydrolysates (SD) with the
591 soluble hydrolysate obtained from a composted mix of the above digestate, and urban gardening
592 wastes and sewage sludge. They showed that, in the case of Euphorbia cultivation, the soluble
593 hydrolysate obtained from the compost was more efficient than the soluble digestate hydrolysate
594 (SD). Sortino et al. (2013 and 2014) compared the same above soluble digestate (SD) and soluble
595 compost (SGC) hydrolysates for greenhouse tomato cultivation. They found that the soluble compost
596 hydrolysate (SGC) as well or better than the soluble digestate hydrolysate (SD). Rovero et al. (2015)
597 compared the composted mix of the above digestate, and urban gardening wastes and sewage sludge,
598 and its soluble and insoluble hydrolysates, in maize cultivation trials performed in open field. No

599 added benefits were shown from the use of the soluble and insoluble compost hydrolysates, compared
600 with the source compost. The results of the above reported studies, and the results of the present work,
601 demonstrate that product performance depends much on the cultivated plant and on the boundary
602 (environmental) conditions. However, different products may be obtained from a variety of different
603 municipal biowastes sourced from different locations and pre-treated by anaerobic and aerobic
604 fermentation (Rosso et al., 2015). This offers the prospect to obtain a wide variety of products, which
605 can be used *ad hoc* for different cultivations. It seems therefore that the results and conclusions of the
606 present work offer worthwhile scope for further research aiming to develop biowaste-sourced
607 products tailored for specific plants' studies and cultivation. To this purpose, it should be underlined
608 that many commercial biostimulant products offer scarce and often insufficient and or vague
609 information on product dosage and distribution. One of the reasons for selecting the NOVA product,
610 as reference commercial product in the present work, was the detailed information reported by the
611 producer about the dosage for substrate cultivations. The comparative data reported in this work, for
612 the biowaste-sourced products and for the commercial NOVA biostimulant, prove that the
613 biowaste-derived products have great potential in intensive agriculture.

614 **Conclusions**

615 The results obtained in the present work assess the effect of biowaste-sourced products on
616 container-grown *Hibiscus*. All treatments have been found to enhance most of the investigated plant
617 parameters, at different degree and depending on their nature, compared to untreated plants. The most
618 valuable effects were observed on biomass accumulation, relative growth rate, net assimilation rate,
619 SPAD index and gas exchange activity. The comparison of the above biowaste-sourced products with
620 the commercial NOVA biostimulant demonstrates that the biowaste-sourced products have a
621 commercially exploitable potential. The comparison of the results obtained in the present work, with
622 those previously reported on the performance of similar products in the cultivation of other
623 ornamental and food plants, shows that a wide variety of biowaste-sourced products is potentially

624 obtainable, which may be targeted to the cultivation of different plant species. This work offers scope
625 for further worthwhile investigation on other plant species, using other different biowastes, in order
626 to assess the full potential of municipal biowaste as source of added value products for use in
627 agriculture. This would also involve a better understanding of the biomolecular mechanisms ruling
628 the effects of these products.

629 **Acknowledgements**

630 This work was carried out partly with funds from the Italian Ministry of Agriculture as part of
631 the “Agrienergia” project. The authors are grateful to Dr. A. Di Nardo, Dr. L. Incrocci and Dr. B.
632 Nesi for their manuscript revision and comments.

633 **References**

- 634 Amanda, A., Ferrante, A., Valagussa, M., Piaggese, A., 2009. Effect of biostimulants on quality of
635 baby leaf lettuce grown under plastic tunnel. *Acta. Hortic.* 807, 407-412.
- 636 Anjum, S.A., Wang, L., Farooq, M., Xue, L., Ali, S., 2011. Fulvic acid application improves the
637 maize performance under well-watered and drought conditions. *J. Agron. Crop Sci.* 197, 409-417.
638 <http://dx.doi.org/10.1111/j.1439-037X.2011.00483.x>.
- 639 Ashraf, M., Harris, P.J.C., 2013. Photosynthesis under stressful environments: An overview.
640 *Photosynthetica* 51, 163-190. <http://dx.doi.org/10.1007/s11099-013-0021-6>.
- 641 Bianco Prevot, A., Avetta, P., Fabbri, D., Laurenti, E., Marchis, T., Perrone, D.G., Montoneri, E.,
642 Boffa, V., 2011. Waste-derived bioorganic substances for light-induced generation of reactive
643 oxygenated species. *ChemSusChem* 4, 85-90. <http://dx.doi.org/10.1002/cssc.201000237>.
- 644 Biolchim, 2015. NOVA@@GR Granular biostimulant based on plant extracts,
645 http://www.biolchim.it/index.php?lang=en&page=products/biostimulants/novagr_granular_bios
646 [timulant_based_on_plant_extracts.html](http://www.biolchim.it/index.php?lang=en&page=products/biostimulants/novagr_granular_bios) (last access 1st September 2015).
- 647 Bulgari, R., Cocetta, G., Trivellini, A., Vernieri, P., Ferrante, A., 2015. Biostimulants and crop
648 responses: a review. *Biol. Agric. Hortic.* 31, 1-17.
649 <http://dx.doi.org/10.1080/01448765.2014.964649>.
- 650 Calvo, P., Nelson, L., Kloepper, J.W., 2014. Agricultural uses of plant biostimulants. *Plant Soil* 383,
651 3-41.

652 Castro, J., Vera, J., González, A., Moenne, A., 2012. Oligo-carrageenans stimulate growth by
653 enhancing photosynthesis, basal metabolism, and cell cycle in tobacco plants (var. Burley). *J Plant*
654 *Growth. Regul.* 31, 173-185. <http://dx.doi.org/10.1007/s00344-011-9229-5>.

655 Dorais, M., 2007. Organic production of vegetables: State of the art and challenges. *Can. J. Plant Sci.*
656 87, 1055-1066.

657 du Jardin, P., 2015. Plant biostimulants: Definition, concept, main categories and regulation. *Sci.*
658 *Hortic.* 196, 3-14. <http://dx.doi.org/http://dx.doi.org/10.1016/j.scienta.2015.09.021>.

659 Ertani, A., Nardi, S., Altissimo, A., 2013a. Review: Long-term research activity on the biostimulant
660 properties of natural origin compounds. *Acta Hortic.* 1009, 181-188.

661 Ertani, A., Schiavon, M., Muscolo, A., Nardi, S., 2013b. Alfalfa plant-derived biostimulant stimulate
662 short-term growth of salt stressed *Zea mays* L. plants. *Plant Soil* 364, 145-158.
663 <http://dx.doi.org/10.1007/s11104-012-1335-z>.

664 Eyheraguibel, B., Silvestre, J., Morard, P., 2008. Effects of humic substances derived from organic
665 waste enhancement on the growth and mineral nutrition of maize. *Bioresour. Technol.* 99, 4206-
666 4212. <http://dx.doi.org/10.1016/j.biortech.2007.08.082>.

667 Fascella, G., Montoneri, E., Ginepro, M., Francavilla, M., 2015. Effect of urban biowaste derived
668 soluble substances on growth, photosynthesis and ornamental value of *Euphorbia x lomi*. *Sci.*
669 *Hortic.* 197, 90-98. <http://dx.doi.org/10.1016/j.scienta.2015.10.042>.

670 Flexas, J., Bota, J., Loreto, F., Cornic, G., Sharkey, T.D., 2004. Diffusive and metabolic limitations
671 to photosynthesis under drought and salinity in C3 plants. *Plant Biol.* 6, 269-279.
672 <http://dx.doi.org/10.1055/s-2004-820867>.

673 Gardner, F.P., Pearce, R.B., Mitchell, R.L., 1985. *Physiology of crop plants*, 1st ed. Iowa State
674 University Press, Ames.

675 Gomis, J., Bianco Prevot, A., Montoneri, E., González, M.C., Amat, A.M., Martínez, D.O., Arques,
676 A., Carlos, L., 2014. Waste sourced bio-based substances for solar-driven wastewater

677 remediation: Photodegradation of emerging pollutants. Chem. Eng. J. 235, 236-243.
678 <http://dx.doi.org/10.1016/j.cej.2013.09.009>.

679 Hawrylak-Nowak, B., Matraszek, R., Pogorzelec, M., 2015. The dual effects of two inorganic
680 selenium forms on the growth, selected physiological parameters and macronutrients
681 accumulation in cucumber plants. Acta Physiol. Plant. 37. [http://dx.doi.org/10.1007/s11738-015-](http://dx.doi.org/10.1007/s11738-015-1788-9)
682 1788-9.

683 Herms, D.A., Mattson, W.J., 1992. The Dilemma of Plants: To Grow or Defend. Q. Rev. Biol. 67,
684 283-335. <http://dx.doi.org/10.2307/2830650>.

685 Houben, D., Sonnet, P., Cornelis, J.T., 2014. Biochar from Miscanthus: A potential silicon fertilizer.
686 Plant Soil 374, 871-882. <http://dx.doi.org/10.1007/s11104-013-1885-8>.

687 Jannin, L., Arkoun, M., Ourry, A., Laine, P., Goux, D., Garnica, M., Fuentes, M., Francisco, S.S.,
688 Baigorri, R., Cruz, F., Houdusse, F., Garcia-Mina, J.M., Yvin, J.C., Etienne, P., 2012. Microarray
689 analysis of humic acid effects on *Brassica napus* growth: involvement of N, C and S metabolisms.
690 Plant Soil 359, 297-319.

691 Kenzo, T., Inoue, Y., Yoshimura, M., Yamashita, M., Tanaka-Oda, A., Ichie, T., 2015. Height-related
692 changes in leaf photosynthetic traits in diverse Bornean tropical rain forest trees. Oecologia 177,
693 191-202. <http://dx.doi.org/10.1007/s00442-014-3126-0>.

694 Li, P., Song, A., Li, Z., Fan, F., Liang, Y., 2015. Silicon ameliorates manganese toxicity by regulating
695 both physiological processes and expression of genes associated with photosynthesis in rice
696 (*Oryza sativa* L.). Plant Soil. <http://dx.doi.org/10.1007/s11104-015-2626-y>.

697 Lieth, J.H., Pasian, C.C., 1991. A simulation model for the growth and development of flowering rose
698 shoots. Sci. Hortic. 46, 109-128.

699 Loh, F.C.W., Grabosky, J.C., Bassuk, N.L., 2002. Using the SPAD 502 meter to assess chlorophyll
700 and nitrogen content of benjamin fig and cottonwood leaves. Horttechnology 12, 682-686.

701 Massa, D., Mattson, N.S., Lieth, H.J., 2009. Effects of saline root environment (NaCl) on nitrate and
702 potassium uptake kinetics for rose plants: a Michaelis-Menten modelling approach. *Plant Soil*
703 318, 101-115. <http://dx.doi.org/10.1007/s11104-008-9821-z>.

704 Miller, G.W., Huang, I.J., Welkie, G.W., Pusnik, J.C., 1995. Function of iron in plants with special
705 emphasis on chloroplast and photosynthetic activity, In: Abadia, J. (Ed.), *Iron Nutrition in Soils*
706 *and Plants*. Kluwer Academic Publishers, Dordrecht, pp. 19-28.

707 Montoneri, C., Montoneri, E., Tomasso, L., Piva, A. 2013. Compost derived substances decrease feed
708 protein N mineralization in swine cecal fermentation. *J Agric Sci* 13, 31-44.
709 <http://dx.doi.org/10.5539/jas.v5n3p>

710 Montoneri, E., Mainero, D., Boffa, V., Perrone, D.G., Montoneri, C., 2011. Biochemenergy: A
711 project to turn an urban wastes treatment plant into biorefinery for the production of energy,
712 chemicals and consumer's products with friendly environmental impact. *Int. J. Global Environ.*
713 *Issues* 11, 170-196. <http://dx.doi.org/10.1504/ijgenvi.2011.043528>.

714 Morard, P., Eyheraguibel, B., Morard, M., Silvestre, J., 2011. Direct effects of Humic-Like substance
715 on growth, water, and mineral nutrition of various species. *Journal Plant Nut.* 34, 46-59.
716 <http://dx.doi.org/10.1080/01904167.2011.531358>.

717 Nowak, J., Kaklewski, K., Ligocki, M., 2004. Influence of selenium on oxidoreductive enzymes
718 activity in soil and in plants. *Soil Biol. Biochem.* 36, 1553-1558.
719 <http://dx.doi.org/10.1016/j.soilbio.2004.07.002>.

720 Papasavvas, A., Triantafyllidis, V., Zervoudakis, G., Kapotis, G., Samaras, Y., Salahas, G., 2008.
721 Correlation of SPAD-502 meter readings with physiological parameters and leaf nitrate content
722 in *Beta vulgaris*. *J. Environ. Prot. Ecol.* 9, 351–356.

723 Pilon-Smits, E.A., Quinn, C.F., Tapken, W., Malagoli, M., Schiavon, M., 2009. Physiological
724 functions of beneficial elements. *Curr. Opin. Plant Biol.* 12, 267-274.
725 <http://dx.doi.org/10.1016/j.pbi.2009.04.009>.

726 Reid, J.B., 2002. Yield response to nutrient supply across a wide range of conditions. 1. Model
727 derivation. *Field Crop. Res.* 77, 161-171.

728 Rios, J.J., Blasco, B., Leyva, R., Sanchez-Rodriguez, E., Rubio-Wilhelmi, M.M., Romero, L., Ruiz,
729 J.M., 2013. Nutritional balance changes in lettuce plant grown under different doses and forms of
730 selenium. *Journal of Plant Nutrition* 36, 1344-1354.
731 <http://dx.doi.org/10.1080/01904167.2013.790427>.

732 Rosso, D., Fan, J., Montoneri, E., Negre, M., Clark, J., Mainero, D., 2015. Conventional and
733 microwave assisted hydrolysis of urban biowastes to added value lignin-like products. *Green*
734 *Chemistry* 17, 3424-3435. <http://dx.doi.org/10.1039/c5gc00357a>.

735 Rovero, A., Vitali, M., Rosso, D., Montoneri, E., Chitarra, W., Tabasso, S., Ginepro, M., Lovisolo,
736 C., 2015. Sustainable maize production by urban biowaste products. *Int. J. Agron. Agric. Res.* 6,
737 75-91.

738 Saa, S., Olivos-Del Rio, A., Castro, S., Brown, P.H., 2015. Foliar application of microbial and plant
739 based biostimulants increases growth and potassium uptake in almond (*Prunus dulcis* [Mill.] D.
740 A. Webb). *Front. Plant Sci.* 6, 87. <http://dx.doi.org/10.3389/fpls.2015.00087>.

741 Shibayama, M., Sakamoto, T., Takada, E., Inoue, A., Morita, K., Yamaguchi, T., Takahashi, W.,
742 Kimura, A., 2012. Estimating rice leaf greenness (SPAD) using fixed-point continuous
743 observations of visible red and near infrared narrow-band digital images. *Plant Prod. Sci.* 15, 293-
744 309. <http://dx.doi.org/10.1626/pps.15.293>.

745 Silberbush, M., Ben-Asher, J., Ephrath, J.E., 2005. A model for nutrient and water flow and their
746 uptake by plants grown in a soilless culture. *Plant Soil* 271, 309-319.

747 Sortino, O., Dipasquale, M., Montoneri, E., Tomasso, L., Avetta, P., Bianco Prevot, A., 2013. 90%
748 yield increase of red pepper with unexpectedly low doses of compost soluble substances. *Agron.*
749 *Sustainable Dev.* 33, 433-441.

750 Sortino, O., Montoneri, E., Patanè, C., Rosato, R., Tabasso, S., Ginepro, M., 2014. Benefits for
751 agriculture and the environment from urban waste. *Sci. Tot. Environ.* 487, 443-451.
752 <http://dx.doi.org/10.1016/j.scitotenv.2014.04.027>.

753 Yamori, W., Noguchi, K., Terashima, I., 2005. Temperature acclimation of photosynthesis in spinach
754 leaves: analyses of photosynthetic components and temperature dependencies of photosynthetic
755 partial reactions. *Plant Cell Environ.* 28, 536-547.

756

757

758 **Figure captions**

759 **Fig. 1** Variable loadings (Fig. 1a) and data scores (Fig. 1b) obtained by principal component analysis.
760 Acronyms (Fig 1a) represent leaf stomatal conductance (Gs), transpiration (E) and photosynthesis
761 (Pn), water use efficiency (Pn/E), leaf chlorophyll (SPAD) and nitrogen (N) content, plant volume
762 (V), height (H) and shape index (SI), and leaf (LeD), stem (StD) and total shoot (stems, leaves and
763 flowers) dry weight (TD). Points and numbers (Fig. 1b) represent treatment combinations (see Table
764 2): i) green numbers show combinations with control treatment 1; ii) red numbers show combinations
765 with NOVA treatment 2; iii) blue numbers show combinations with non-mixed hydrolysate products
766 (treatment 4, 5 9, and 10). Dashed lines group all combinations with fertigation treatments: i) dashed
767 circle for control treatment 13; ii) top dashed rectangle for soluble green compost hydrolysate
768 treatment 14; iii) bottom dashed rectangle for soluble digestate hydrolysate treatment 15.

769