AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Identification of exposure to toxic metals by means of segmental hair analysis: a case report of alleged chromium intoxication

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1623020> since 2017-01-25T16:03:53Z

Published version:

DOI:10.1007/s11419-016-0340-y

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Identification of exposure to toxic metals by means of segmental hair analysis. A case report of alleged chromium intoxication

Federica D'Urso^{1,2}, Alberto Salomone^{1*}, F. Seganti¹, Marco Vincenti^{1,2}

¹*Centro Regionale Antidoping “A. Bertinaria”, Regione Gonzole 10/1, 10043 Orbassano, Turin, Italy*

²*Dipartimento di Chimica, Università degli Studi di Torino, via P. Giuria 7, 10125 Turin, Italy*

***Corresponding author:** Alberto Salomone, PhD

Centro Regionale Antidoping “A. Bertinaria”,

Regione Gonzole, 10/1 – 10043 Orbassano (Turin) – Italy

Tel: +3901190224232, Fax: +3901190224242, Mobile: +393489330145

E-mail address: alberto.salomone@antidoping.piemonte.it

Abstract

Hair mineral analysis has become an interesting diagnostic tool in biomonitoring of exposure to toxic elements, in the assessment of health and nutritional status. The most inconvenience of this matrix is the lack of sufficient information to define normal ranges of metal levels in a general healthy population. In this study, segmental hair analysis was used to depict a chronological scheme of exposure to arsenic, cobalt, cadmium, chromium, copper, manganese, nickel and lead in a 16-year-old girl showing signs of potential intoxication. The quantitative results obtained from consecutive segments of hair proved the exposure to chromium. In particular, segment A (0-6 cm), approximately reflecting the last 6 months of exposure, resulted in the chromium level at 5.60 µg/g. The technique of segmental analysis allowed us to establish “intra-individual” physiological variation ranges for each heavy metal hair concentration. As a consequence, these “confidence” intervals could be used as

individualized references to highlight the occurrence of atypical metal levels in any specific hair segment, possibly identifying a period of anomalous exposure and/or intoxication.

Keywords. Metals, Intoxication, Hair, Segmental analysis, Atomic Absorption Spectrophotometry

1 Introduction

2
3 Hair analysis currently represents a reliable and well-established means of clinical
4 and forensic investigation [1]. The memory property of hair due to sequential
5 accumulation of chemicals in its inner structure, together with the opportunity of
6 conducting retrospective analysis, accounted for its success in several application
7 contexts such as the confirmation of drug-facilitated crimes, the assessment of drug
8 consumption history in addiction treatments, the environmental and occupational
9 exposure to pollutants [1, 2-4]. Also workplace drug testing, driving re-licensing,
10 withdrawal control, **postmortem** toxicology, **prenatal** exposure to drugs, and doping
11 control extensively apply hair analysis for screening and confirmation purposes [5-
12 13]. **Upon investigating** extended time-windows, as opposed to biological fluids, hair
13 analysis **has been recognized as** the most powerful tool in the assessment of chronic
14 consumption or exposure to various chemicals. The relatively constant head hair
15 growth, with an estimated average rate of about 1.0 cm/month, allows to trace the
16 chronological exposure profile by segmental hair analysis [14, 15]. As a matter of
17 fact, segmental hair analysis has been repeatedly used to ascertain occasional abuse
18 of drugs, alcohol and doping agents [1], verify the compliance of enforced abstinence
19 [8], and outline a chronological sequence of drug exposure [16]. Lastly, hair analysis
20 has been used to estimate the nutritional status of individuals and to assess poisoning
21 and environmental intoxication of exposed subjects [17-19] from a variety of organic
22 and inorganic substances, including heavy metals. From another point of view,
23 human hair can be considered as a secondary excretion vehicle of toxic substances
24 from the body. For example, the concentrations of heavy metals in hair are up to 10-
25 times higher than in blood and urine [20-21]. Heavy metals, such as chromium, lead,
26 mercury, cadmium, and arsenic – whenever biologically available - are extremely
27 toxic to most living organisms even at very low concentrations. The presence of
28 heavy metals in human hair usually reflects their bioavailability and therefore is
29 generally found at ultra-trace **levels**. Significant excess of these elements in human

hair with respect to the expected population's average may reflect the degree of body exposure to these poisons, either from environmental pollution, workplace or food chain [17,18,22]. Hair mineral analysis has also become an interesting diagnostic tool in biomonitoring the exposure to toxic elements and the health and nutritional status assessment. Likewise organic substances and drugs, trace metal analysis on hair material presents several advantages over biological fluids, because it may provide an historical overview on the individual exposure to these elements, recognize acute vs. chronic intoxication, and monitor the nutritional status of the investigated subject over extended periods of time [23]. On the other hand, hair analysis also presents some limitations, including the lack of well-defined and generally accepted reference concentration ranges [23]. This uncertainty arises from the large differences existing in the elements' level as a function of sex, age, residence area, ethnicity, hair color, dietary habits, and individual physiological variability [24]. The present study was addressed to the evaluation of arsenic (As), cobalt (Co), cadmium (Cd), chromium (Cr), copper (Cu), manganese (Mn), nickel (Ni), and lead (Pb) concentrations in the human scalp hair collected from a subject showing signs of potential intoxication, allegedly arising from exposure to toxic metals. Segmental hair analysis was performed to obtain information about the history of the patient's exposure, approximately in the preceding 3 years. Taking into account that (i) considerable interest exists in the toxicological perspectives opened by hair analysis toward the confirmation of heavy metal poisoning, and (ii) insufficient data are available in the literature about heavy metals acceptable especially for physiological hair concentrations, we decided to validate the whole analytical method and to establish an independent hair reference range for these metals in scalp hair, based on specimen collected from laboratory personnel volunteers ($n=10$). Furthermore, the technique of segmental analysis allowed us to establish "intra-individual" physiological variation ranges for each heavy metal hair concentration. As a consequence, these "confidence" intervals could be used as individualized references to highlight the

occurrence of atypical metal levels in any specific hair segment, possibly identifying a period of anomalous exposure and/or intoxication.

Materials and methods

Case history

A 16-year-old girl came to our laboratory to verify the possible past exposure to toxic metals by means of hair analysis. The girl had spent the previous 9 months abroad, where she lived, spent most of her time, and took her meals mostly inside a college. During this period, she presented severe symptoms, including metabolic disorders, skin irritation, nose bleeding, bronchitis and dysentery. Furthermore, recent blood tests evidenced sub-standard levels of glycemia (63 mg/dL) and increased potassium level at 5.94 mEq/L (range 3.5-5.5). The patient's parents suspected that the disorders presented by their daughter were to be attributed to a possible exposure to toxic substances arising either from the environment or the food, not only to the change in eating habits and climatic conditions.

Sample preparation

Two locks (diameter approximately 0.5 cm) of patient's hair (color: blond) were sampled in its entire length (about 40 cm) from the vertex region of the head [25], using stainless steel scissors and then stored at room temperature in a plastic box. The hair was segmented as follow: starting from the skull extremity, we considered six segments of 6 cm each and a final segment of 4 cm. The analyzed weight for each segment was in the range 400-500 mg. The hair aliquots were washed to remove potential external contaminations using the method proposed by Ohmori [26], consisting of three consecutive steps with 3 ml each of acetone, water and again

acetone. After decontamination, the samples were dried at room temperature overnight. The procedure adopted for hair digestion was based on the method commonly adopted in several literature studies [27-29]. Briefly, the dried hair aliquots were digested with a mixture of 65 % nitric acid (6 mL) and 67-72% perchloric acid (1 mL) at 70-80°C, until the hair was completely dissolved and the solution became clear (about 25 min). Lastly, each sample solution was diluted to 50 mL with demineralized Milli-Q water.

Standard solutions for calibration were prepared from a 1000 mg/L ICP Multielement solution (Merck, Milan, Italy), ranging from 0.10 to 10 µg/g. (0.50 to 50 µg/g for Cu).

Instrumentation

All analyses were performed using a Perkin-Elmer Analyst 800 atomic absorption spectrophotometer (Perkin Elmer, Norwalk, USA) equipped with an AS-800 autosampler and THGA graphite tubes with end caps (Perkin-Elmer). The instrumental parameters are described in Table 1.

Validation of analytical methods

The characteristic validation parameters for the analytical methods were determined from the analysis of blank water and standard solutions at different concentrations for each metal. These parameters, following the recommendations of ISO/IEC 17025:2005 international standard and others guidelines [25,30], included the limit of detection (LOD), limit of quantification (LOQ), linear range, precision (as CV %) and accuracy (as bias %). The linearity interval was evaluated by checking the linear regression coefficient (r^2) of the calibration curve. The linearity was considered acceptable when $r^2 > 0.995$. The linear calibration model was checked by analyzing (two replicates) blank water spiked with the working solution at five concentrations

levels in the range of 0.10-10 µg/g (0.50 to 50 µg/g for Cu). The LODs and the LOQs were extrapolated by Hubaux and Vos approach [31]. For all elements, intraday precision (expressed as percent variation coefficient, CV%) and accuracy (expressed as bias %) were evaluated by spiking blank solution at low and high concentration levels at 0.10 µg/g (0.50 µg/g for Cu only) and 10 µg/g (50 µg/g for Cu only). Intraday precision was satisfactory when CV% values were below 15%. Satisfactory accuracy was achieved when the experimentally determined average concentration lied within ±15% from the expected value.

Results and discussion

Validation parameters

All the validation results are reported in Table 2. Each element showed a coefficient of determination higher than 0.995 indicating good fit and linearity for the calibration curves (0.10-10 µg/g for all elements, except for Cu, 0.50-50 µg/g). LOD values ranged from 0.012 µg/g for Pb to 0.35 µg/g for Cu, while LOQ values lied between 0.021 for Mn µg/g and 0.70 for Cu µg/g. Intraday precision and accuracy were satisfactory for most but not all analytes. In particular, at the lowest calibration level, accuracy (as bias %) for As, Cd, Cu, and Mn exceeded the accepted values, while at high concentration level (10 µg/g), only Cd exceeded the accepted values. At last, intraday precision (as CV %) exceeding the accepted interval of ±15% was observed for Co, Cr, Mn, and Pb at low concentrations, while modest deviation from 15% was observed at high concentration for As.

Analysis of real hair samples

Physiological reference values and ranges for all heavy metals in hair were determined on ten healthy volunteers, four females and six males. For women's hair, segmental hair analysis was used whenever possible, but no concentration difference was found among the segments arising from the same subject. Table 3 reports the median and reference ranges obtained thereby. Table 3 also reports the expected concentration ranges of trace metals in the hair of healthy subjects, as were determined within a systematic ICP-MS study reported in the literature ($n=45$) [32]. These values can be compared with those obtained within the present study from young laboratory personnel hair ($n=10$, six of which were browns, two blondes and two blacks) using electro-thermal atomization AAS. While the lower limits of the reference concentration ranges appear to be similar in the two studies, the same did not apply to the upper limits of Co, Cr, and Mn, that are 4-8 times higher in the present study than in the one previously published [32], despite the lower number of subjects involved in the present study. For these metals, also the median values are appreciably higher. The results obtained on the hair sample from our case in question are summarized in Table 4 and visually represented in Fig. 1. The visual examination of the data obtained from the seven segments reveals that the level of Cr in segment A (0-6 cm), approximately reflecting the last 6 months of exposure, is much higher than in the preceding segments. In fact, segments B-G have an average Cr concentration of $0.74 \pm 0.39 \mu\text{g/g}$; a Student t -test made with the level found in segment A ($5.6 \mu\text{g/g}$) produces a residual probability $\alpha < 10^{-20}$. Analogous t -tests calculated with respect to the reference Cr levels of both the present and Goullé et al. [32] studies (see Table 3) show a highly significant difference for segment A ($\alpha < 10^{-12}$). Although no literature data are available on the Cr scalp hair concentration in patients suffering from subchronic Cr intoxication, the exceedingly high Cr concentration found in hair segment A - corresponding to the period that the patient spent abroad - is consistent with the patient's symptoms in that period and the

hypothesis of her intoxication by Cr exposure, which can produce the severe health effects actually observed (metabolic disorders, severe skin irritation, and dysentery). Indeed chromium is an essential nutrient in our diet that helps insulin to maintain normal glucose level [33]. **Because** Cr^(III) is poorly absorbed by any route, the toxicity of chromium is mainly attributable to the Cr^(VI) form. It can be absorbed by the lung and gastrointestinal tract, and even to a certain extent by intact skin. **It is known** that **Cr toxicity is commonly** associated with stomach upsets, ulcer, **and kidney and liver damages** [17]. Inhaling high levels of Cr can cause irritation to the lining of the nose, nose ulcers and breathing problems. Long term exposure can cause **damages to the liver and kidney, and can cause** circulatory and nerve disorders, as well as severe skin irritation [34]. For the remaining metals (As, Cd, Co, Cu, Ni, Pb), the concentrations found in the patient's hair segments are consistently within the expected reference ranges (see **Tables 3 and 4**). Only the Mn level is somehow higher than the published [32] upper reference limit, but this **result is not** considered significant because the Mn hair concentration was found approximately constant throughout the seven segments and lower than the upper reference limit observed in the present study (Table 3).

Conclusions

The human exposure to toxic metals is generally monitored by determining their concentrations in conventional body fluids such as blood and urine. Just recently, other non-conventional matrices, in particular scalp hair, are gaining importance in the investigation of possible excessive exposure to toxic metals. Unlike conventional matrices, human scalp hair, and particularly segmental analysis **on long hair**, provide historical and chronological information on trace **element concentrations in the body**, that portrays a unique profiling of exposure. **Compared to the screening for drugs in the keratin matrix, the detection of (heavy) metals is less influenced by the redistribution along the hair shaft and less affected by washing out effects. Therefore, a segmental hair analysis is much more effective in order to obtain an over months**

chronological information about poisoning/exposure with/to heavy metals. In reporting an interesting real case of alleged intoxication by Cr, the present study demonstrates that segmental hair analysis allows to compare the heavy metal exposure during a specific period of time with that during other time intervals, possibly corresponding to different external conditions (i.e., different environmental, occupational, or domiciliary exposure, or even deliberate poisoning). This unique intra-individual comparison is referred to the subject under study, and proves to be more specific than any comparison made with a generic reference population. The patient under examination presented severe symptoms possibly associated to heavy metal intoxication, and indeed the concentration of Cr in her hair segment grown during the period when she changed her domicile and living habits was found to be exceedingly high, as compared with those in the hair segments grown in preceding time periods. Moreover, the most severe symptoms that she accused (i.e. hypoglycemia, diarrhea and erythema), were consistent with the hypothesis of sub-chronic intoxication by Cr. The present study is likely to contribute to the scientific knowledge about the relationship existing between exposure to toxic metals and their expected concentration in the scalp hair grown in the corresponding period.

Conflict of Interest. The authors declare that they have no conflict of interest.

Ethical approval. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study

227

228

229

230 **Figure caption**

231 **Figure 1** Graphical representation of metal concentrations in each segment. The
232 segmentation (cm) is reported on x-axis. The y-scale for Cu was reduced by one order
233 of magnitude

234

235

References

1. Kintz P, Salomone A, Vincenti M (2015) Hair analysis in clinical and forensic toxicology. Academic Press/ Elsevier, London
2. Appenzeller BMR, Tsatsakis AM (2012) Hair analysis for biomonitoring of environmental and occupational exposure to organic pollutants: state of the art, critical review and future needs. *Toxicol Lett* 210:119-140
3. Vincenti M, Salomone A, Gerace E, Pirro V (2013) Application of mass spectrometry to hair analysis for forensic toxicological investigation. *Mass Spectrom Rev* 32:312-332
4. Vincenti M, Salomone A, Gerace E, Pirro V (2013) Role of LC-MS/MS in hair testing for the determination of common drugs of abuse and other psychoactive drugs. *Bioanalysis* 5:1919-1938
5. Pragst F, Balikova MA (2006) State of the art in hair analysis for detection of drug and alcohol abuse. *Clin Chim Acta* 370:17-49
6. Cone EJ (2001) Legal, workplace, and treatment drug testing with alternative biological matrices on a global scale. *Forensic Sci Int* 121:7-15
7. Caplan YH, Goldberger BA (2001) Alternative specimens for workplace drug testing. *J Anal Toxicol* 25:396-399
8. Kintz P, Villain M, Cirimele V (2006) Hair analysis for drug detection. *Ther Drug Monit* 28 442-446
9. Kintz P (2007) Bioanalytical procedures for detection of chemical agents in hair in the case of drug-facilitated crimes. *Anal Bioanal Chem* 388:1467-1474
10. Lozano J, Garcia-Algar O, Vall O, de la Torre R, Scaravelli G, Pichini S (2007) Biological matrices for the evaluation of in utero exposure to drugs of abuse. *Ther Drug Monit* 29:711-734
11. Barroso M, Dias M, Vieira DN, Lopez-Rivadulla M, Queiroz JA (2012) Simultaneous quantitation of morphine, 6-acetylmorphine, codeine, 6-acetylcodeine

and tramadol in hair using mixed-mode solid-phase extraction and gas chromatography–mass spectrometry. *Anal Bioanal Chem* 396:3059–3069

12. Gerace E, Salomone A, Pellegrino S, Vincenti M (2012) Evidence of Haldol (haloperidol) long-term intoxication. *Forensic Sci Int* 215:121-123

13. Salomone A, Gerace E, Di Corcia D, Martra G, Petrarulo M, Vincenti M (2012) Hair analysis of drugs involved in drug-facilitated sexual assault and detection of zolpidem in a suspected case. *Int J Legal Med* 126:451-459

14. Beumer JH, Bosman IJ, Maes RA (2001) Hair as a biological specimen for therapeutic drug monitoring. *Int. J Clin Pract* 55:353-357

15. Kintz P, Alvarez JC, Deveaux M, Dumestre V, Gaillard Y, Gaulier JM (2014) Conflicting hair testing results can have an impact in courts: interpretation of single exposure to zolpidem. *J Anal Toxicol* 38:304-305

16. Pirro V, Fusari I, Di Corcia D, Gerace E, De Vivo E, Salomone A, Vincenti M (2014) Hair analysis for long-term monitoring of buprenorphine intake in opiates withdrawal. *Ther Drug Monit* 36:796-807

17. Onuwa OP, Ishaq SE, Rufus S (2012) Analysis of heavy metals in human hair using atomic absorption spectrometry (AAS). *Am J Anal Chem* 3:770-773

18. Randall JA, Gibson RS (1989) Hair chromium as an index of chromium exposure of tannery workers. *Br J Ind Med* 46:171-175

19. Momčilović B, Prejac J, Višnjević V, Brundić S, Skalny AA, Mimica N (2016) High hair selenium mother to fetus transfer after the Brazil nuts consumption. *J Trace Elem Med Biol* 33:110–113

20. Bader M, Dietz MC, Ihrig A, Triebig G (1999) Biomonitoring of manganese in blood, urine and axillary hair following low-dose exposure during the manufacture of dry cell batteries. *Int Arch Occup Environ Health* 72:521-527.

21. Omedo P, Pla A, Hernández AF, Lòpez-Guarnida O, Rodrigo L, Gil F (2010) Validation of a method to quantify chromium, cadmium, manganese, nickel and lead in human whole blood, urine, saliva and hair samples by electrothermal atomic absorption spectrometry, *Anal Chim Acta*. 659:60-67.

22. Queipo Abad S, Rodríguez-González P, García Alonso JI (2016) Evidence of the direct adsorption of mercury in human hair during occupational exposure to mercury vapour, *J Trace Elem Med Biol*, 36:16–21
23. Ozmen H, Akarsu S, Polat F, Cukurovali A (2013) The levels of calcium and magnesium, and of selected trace elements, in whole blood and scalp hair of children with growth retardation. *Iran J Pediatr* 23:125-130
24. Szykowska MI, Pawlaczyk A, Wojciechowska E, Sypniewski S, Paryjczak T (2009) Human hair as a biomarker in assessing exposure to toxic metals. *Polish J Environ Stud* 18:1151-1161
25. Salomone A, Tsanaclis L, Agius R, Kintz P, Baumgartner MR (2016) European guidelines for workplace drug and alcohol testing in hair. *Drug Test Anal*, doi 10.1002/dta.1999
26. Ohmori S (1984) Hair as a monitor for environmental pollution. Variations in element concentrations for different lead-processing workers and different washing methods of their hair. *J Radioanal Nucl Chem* 84:451-459
27. Srogi K (2006) Determination of trace elements in the human hair. *Wiadom Chem* 3:279-300
28. Teresa M, Vasconcelos SD, Tavares HM (1997) Trace element concentrations in blood and hair of young apprentices of a technical-professional school. *Sci Total Environ* 205:189-199
29. Chojnacka K, Górecka H, Górecki H (2006) The effect of age, sex, smoking habit and hair color on the composition of hair. *Environ Toxicol Pharmacol* 22:52-57
30. Society of Forensic Toxicologists/American Academy of Forensic Sciences. (2006) Forensic toxicology laboratory guidelines. Mesa/Colorado Springs
31. Hubaux A, Vos G (1970) Decision and detection limits for calibration curves. *Anal Chem* 42:849–855
32. Goullé JP, Mahieu L, Castermant J, Neveu N, Bonneau L, Lainé G, Bouige D, Lacroix C (2005) Metal and metalloid multi-elementary ICP-MS validation in whole blood, plasma, urine and hair: reference values. *Forensic Sci Int* 153:39-44

- 323 33. Schwartz K, Mertz W (1959) Chromium(III) and the glucose tolerance factor.
324 Arch Biochem Biophys 85:292-295
- 325 34. Senofonte O, Violante N, Caroli S (2000) Assessment of reference values for
326 elements in human hair of urban schoolboys. J Trace Elem Med Biol 14:6-13
327

Table 1 Experimental parameters used in electro-thermal atomization for atomic absorption spectrophotometry (AAS) analyses.

Parameters	As	Cd	Co	Cr	Cu	Mn	Ni	Pb
Wavelength (nm)	193.7	228.8	240.7	357.9	324.8	279.5	232.0	283.3
Slit-width (nm)	0.7	0.7	0.2	0.7	0.7	0.2	0.2	0.7
Pretreatment temperature (°C)	1200	700	1400	1500	700	900	1500	1900
Atomization temperature (°C)	2000	1900	2450	2300	2250	2400	2300	2450
Sample volume (μL)	20	20	20	20	20	20	20	20

As: arsenic, *Cd*: cadmium, *Co*: cobalt, *Cr*: chromium, *Cu*: copper, *Mn*: manganese, *Ni*: nickel, *Pb*: lead

Table 2 Limits of detection (LODs), limits of quantitation (LOQs), coefficient of determination (r^2), and data of accuracy and precision for determination of As, Cd, Co, Cr, Cu, Mn, Ni, and Pb by electro-thermal atomization AAS

Parameter	As	Cd	Co	Cr	Cu	Mn	Ni	Pb
LOD ($\mu\text{g/g}$)	0.10	0.050	0.10	0.12	0.35	0.021	0.050	0.012
LOQ ($\mu\text{g/g}$)	0.20	0.10	0.20	0.25	0.70	0.042	0.10	0.024
r^2	^a 0.9988	0.9962	0.9988	0.9984	0.9972	0.9999	0.9997	0.9999
Accuracy	^b +37.3	+21.4	+7.4	+15.7	+18.8	+18.2	+1.0	+13.3
(bias %)	^c -0.04	-39.9	+0.07	-2.8	+5.5	-1.9	+3.2	-7.9
Precision	^b 8.9	2.0	21.2	31.1	13.4	22.3	11.0	26.3
(CV%)	^c 16.5	5.4	0.8	2.2	0.6	1.3	4.4	4.2

Limits of Detection and Quantification, Squared Correlation Coefficient, Accuracy and Precision data are reported.

^aThe linearity range tested was 0.10-10 $\mu\text{g/g}$ (0.50 -50 $\mu\text{g/g}$ for Cu)

^bLow concentration 0.10 $\mu\text{g/g}$ (0.50 $\mu\text{g/g}$ for Cu)

^cHigh concentration 10 $\mu\text{g/g}$ (50 $\mu\text{g/g}$ for Cu)

Table 3 Comparison of heavy metal levels in hair obtained from the healthy subjects in the present study with those reported by Goullé et al. [32]

Element	Present study (<i>n</i> =10)		Goullé, J.P. et al. [32] (<i>n</i> =45)	
	Reference range ($\mu\text{g/g}$ or ppm)	Median ($\mu\text{g/g}$ or ppm)	Reference range ($\mu\text{g/g}$ or ppm)	Median ($\mu\text{g/g}$ or ppm)
As	< 0.10	-	0.03-0.08	0.05
Cd	< 0.050	-	0.04-0.17	0.011
Co	0.10-1.1	0.19	0.004-0.14	0.023
Cr	0.12-2.4	0.34	0.11-0.52	0.20
Mn	0.022-3.8	0.35	0.016-0.57	0.067
Ni	0.05-1.9	0.66	0.08-0.90	0.23
Cu	6.5-42	11	9.0-61	20
Pb	0.013-1.4	0.42	0.13-4.6	0.41

Table 4 Analytical results for the case under study. Concentration of trace elements in the scalp hair of the patient are reported as $\mu\text{g/g}$ or ppm.

Segment (cm)		As	Cd	Co	Cr	Ni	Pb	Cu	Mn
A	0-6	< 0.10	< 0.10	< 0.10	5.6	0.29	0.38	16	1.1
B	6-12	< 0.10	< 0.10	< 0.10	0.26	0.15	0.32	17	1.1
C	12-18	< 0.10	0.26	< 0.10	1.1	0.26	0.31	17	1.1
D	18-24	< 0.10	0.15	0.29	0.55	0.37	0.56	19	1.8
E	24-30	< 0.10	0.25	0.22	0.39	0.94	0.77	23	1.7
F	30-36	< 0.10	0.25	< 0.10	1.1	0.57	1.6	27	1.9
G	36-40	< 0.10	0.52	< 0.10	1.1	0.52	1.8	25	2.0

Figure 1

[Click here to download Figure 1 BN.tif](#)