

UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Novel Synthetic, Host-defense Peptide Protects Against Organ Injury/Dysfunction in a Rat Model of Severe Hemorrhagic Shock

(Article begins on next page)

This is the author's final version of the contribution published as:

Yamada, Noriaki; Martin, Lukas B.; Zechendorf, Elisabeth; Purvis, Gareth S. D.; Chiazza, Fausto; Varrone, Barbara; Collino, Massimo; Shepherd, Joanna; Heinbockel, Lena; Gutsmann, Thomas; Correa, Wilmar; Brandenburg, Klaus; Marx, Gernot; Schuerholz, Tobias; Brohi, Karim; Thiemermann, Christoph. Novel Synthetic, Host-defense Peptide Protects Against Organ Injury/Dysfunction in a Rat Model of Severe Hemorrhagic Shock. ANNALS OF SURGERY. None pp: 1-9. DOI: 10.1097/SLA.0000000000002186

The publisher's version is available at: http://Insights.ovid.com/crossref?an=00000658-900000000-96193

When citing, please refer to the published version.

Link to this full text: http://hdl.handle.net/2318/1629885

This full text was downloaded from iris - AperTO: https://iris.unito.it/

A novel synthetic, host-defence peptide protects against organ injury/dysfunction in a rat-model of severe hemorrhagic shock

Noriaki Yamada, MD PhD^{1#}, Lukas Martin, MD PhD^{1,2#*}, Elisabeth Zechendorf, MSc², Gareth

Samuel Desmond Purvis, BSc¹, Fausto Chiazza BSc³, Barbara Varrone MSc³, Massimo Collino,

PhD³, Joanna Shepherd, MD⁴, Lena Heinbockel, PhD⁵, Thomas Gutsmann, PhD⁵,

Wilmar-Alexander Correa-Vargas, PhD⁵, Klaus Brandenburg, PhD⁵, Gernot Marx, MD²,

Tobias Schuerholz, MD⁶, Karim Brohi, MD⁴, Christoph Thiemermann, MD PhD^{1*}

1 The William Harvey Research Institute, Barts & The London School of Medicine & Dentistry, Queen Mary University of London, London, United Kingdom;

²Department of Intensive Care and Intermediate Care, University Hospital RWTH Aachen,

Aachen, Germany;

³Departmentof Drug Science & Technology, University of Turin, Turin, Italy;

4 Barts Centre for Trauma Sciences, Blizard Institute, Queen Mary University of London, London, United Kingdom;

5 Division of Biophysics, Forschungszentrum Borstel, Borstel, Germany

6 Department of Anaesthesia and Intensive Care, University of Rostock, Rostock, Germany

both authors worked equally

***Addresses for correspondence and requests for reprints**

Dr Lukas Benjamin Martin MD Prof Christoph Thiemermann MD PhD FBPhS FRCP FMedSci Queen Mary University of London Barts and The London School of Medicine and Dentistry, William Harvey Research Institute Centre for Translational Medicine and Therapeutics, Charterhouse Square London, EC1M 6BQ, UK. Phone: +44 (0) 20 78822107 E-mail: [l.martin@qmul.ac.uk;](mailto:l.martin@qmul.ac.uk) c.thiemermann@qmul.ac.uk

Running head: Pep19-4LF in hemorrhagic shock

MINI-ABSTRACT (50 words)

This study evaluated the role of host-defence/antimicrobial peptides in trauma-associated hemorrhagic shock (HS). Trauma-associated HS resulted in the release of the host defence peptide LL-37. The synthetic host defence peptide Pep19-4LF attenuated the HS-associated organ injury/dysfunction in the rat by activating pro-survival pathways and by inhibiting local and systemic inflammation.

ABSTRACT

Objective: To evaluate (i) levels of the host-defence/antimicrobial peptide LL-37 in patients with trauma and hemorrhagic shock (HS) and (ii) the effects of a synthetic host-defence peptide on multiple organ failure (MOF) associated with HS in rats.

Background: There are no specific interventions, which reduce MOF in HS. Injury and infection triggers the release of host-defence/antimicrobial peptides, the function of which in HS is unknown.

Methods: LL-37 was measured in plasma from 48 trauma/HS patients admitted to an urban major trauma center. Rats were submitted to HS followed by resuscitation and treated during resuscitation with the synthetic host-defence/antimicrobial peptide Pep19-4LF (66 or 333 μg/kg x h) or vehicle.

Results:

Plasma levels of LL-37 were elevated in patients with trauma/HS. In anesthetized rats subjected to HS and resuscitation, Pep19-4LF attenuated the HS-induced kidney dysfunction, liver injury and lung inflammation. Pep19-4LF enhanced (kidney/liver) the phosphorylation of (i) protein kinase B (Akt) and (ii) endothelial nitric oxide synthase (eNOS) and, hence, activated the Akt/eNOS survival pathway. Pep19-4LF also attenuated the HS-induced (i) translocation of p65 from cytosol to nucleus, (ii) phosphorylation of IKK on Ser^{176/180} and (iii) phosphorylation of IκBα on Ser32/36 resulting in inhibition of NF-κB and formation of NF-κB-dependent pro-inflammatory cytokines. Pep19-4LF prevented the release of TNF α caused by heparan sulfate in human mononuclear cells by binding to this DAMP.

Conclusions: Trauma-associated HS results in release of the host-defence/antimicrobial peptide LL-37. The synthetic host-defence/antimicrobial peptide Pep19-4LF attenuates the organ injury/dysfunction associated with HS by activating pro-survival and anti-inflammatory pathways.

Keywords

Antimicrobial peptides, hemorrhagic shock, multiple organ failure, NF-κB pathway, LL-37

INTRODUCTION

Severe injuries account for 9 % of the deaths worldwide. ¹ Although guidelines for the early management of hemorrhagic shock (HS; including resuscitation and organ support strategies) have decreased the rates of immediate (on scene/within 60min) and early (emergency department and operating room/within 1-4h) deaths, ² post-injury multiple organ failure (MOF) is still associated with significant morbidity and mortality². Therapeutic agents that reduce the incidence and severity of MOF following HS could, therefore, have a major global impact on patient outcomes and resource utilization. The MOF after HS is associated with excessive systemic inflammation, secondary to the release of damage-associated molecular patterns (DAMPs) from extensive tissue damage and ischemia reperfusion injury.³ To date, there are no specific pharmacological interventions used clinically to prevent MOF following/associated with HS.

Host-defence/antimicrobial peptides are known for over 100 years and form part of the innate immune system of insects, plants, and vertebrates by defending the host against invading microorganisms. 4,5,6 Although these peptides differ in sequence and structure, they are predominantly short (10 – 50 amino acids) amphipathic molecules. ⁶ The most extensively studied host-defence/antimicrobial peptide in humans is the cathelicidin-derived peptide LL-37.⁶ LL-37 exhibits strong bactericidal properties, but at the same time neutralizes pathogenic factors released during injury/infection including lipopolysaccharide (LPS) or lipoprotein (LP).⁷ In addition to the interaction with PAMPs (pathogen associated molecular patterns), LL-37 modulates the inflammatory response induced by DAMPs and, hence, modulates many physiological host functions including inflammation, angiogenesis and wound healing.^{6,8} Thus, hostdefence/antimicrobial peptides are attractive candidates for the development of novel therapeutic interventions in infectious and inflammatory diseases. ⁶ The systemic application of LL-37 as a potential drug in man, however, is limited by its toxicity.^{9,10} The challenge is to develop synthetic

4

host-defence/antimicrobial peptides (mimetics) that have little or no adverse effects. Peptide 19- 4LF (Pep19-4LF) is one of several new synthetic host-defence/antimicrobial peptides, which belongs to the class of synthetic anti-lipopolysaccharide peptides (SALP = synthetic anti-LPS peptides). 11,12 However, in addition to binding LPS, these peptides exhibit potent anti-inflammatory effects in experimental sepsis by interacting with a variety of PAMPs and DAMPs.^{13,12,14,15}

The role of host-defence/antimicrobial peptides in HS is unknown. Therefore, the aims of the present study were to (i) investigate the plasma levels of LL-37 in patients with trauma with or without HS and (ii) to explore the effects of Pep19-4LF on the organ injury/dysfunction associated with HS. We report here for the first time, that (i) the plasma levels of LL-37 are elevated in patients with trauma/HS (when compared with trauma without HS) and that (ii) Pep19-4LF attenuates the HS-associated organ injury/dysfunction. Mechanistically, Pep19-4LF has pro-survival and antiinflammatory properties, as it activates the Akt/eNOS cell survival pathways and attenuates the activation of the nuclear factor kappa B (NF-κB) pathway in the rat *in vivo*. Moreover, Pep19-4LF exhibits its anti-inflammatory activity, at least in part, by directly interacting/binding to the DAMP heparan sulfate in human mononuclear cells (MNCs) *in vitro*. These data suggest that Pep19-4LF may prevent the MOF in patients caused by trauma-associated HS, which, in turn, may improve outcome in these patients.

METHODS

Study population and human outcome measurements

Details relating to the study population and human outcome measurements are provided in the s*upplemental.*

Use of human subjects-ethic statement

All patients or their legal representative gave written informed consent. Before inclusion of the first individual, the local National Health Service Research Ethics Committee (REC: 07/Q0603/29) approved this study, which was performed in accordance with the Declaration of Helsinki in its latest form. The use of plasma from healthy volunteers was approved by the ethics committee of the University Hospital Aachen (EC Nr. 206_09, 5 January 2010).

Use of experimental animals-ethic statement

The experimental protocols used in this study have been approved by the Animal Welfare Ethics Review Board (AWERB) of Queen Mary University of London and the study was performed under license issued by home office (Procedure Project License; PPL: 70/7348). Animal care was in accordance with the Home Office guidance on Operation of Animals (Scientific Procedures Act 1986) published by Her Majesty's Stationery Office and the Guide for the Care and Use of Laboratory Animals of the National Research Council.

Hemorrhagic shock and quantification of organ injury and dysfunction

This study was carried out on 46 male Wistar rats (Charles River Ltd, Margate, UK) weighing 230- 350 g receiving a standard diet and water *ad libitum*. Hemorrhagic shock and quantification of organ injury and dysfunction were performed as described previously in this journal (*supplemental Fig. 1)*. 16

Experimental design

Rats were randomly allocated to the following groups: Sham + vehicle ($n = 11$), sham + Pep4LF $(n = 6)$, HS + vehicle $(n = 12)$, HS + Pep4LF-LD $(n = 4)$, HS + Pep4LF-HD $(n = 8)$. Rats were administered vehicle (saline 1.5 ml/kg/h) or Pep19-4LF (low dose (LD) = 66 μg/kg x h; high dose (HD) = 333 μg/kg x h) continuously for 4 hours after resuscitation using infusion pump for rodents

(PHD2000, 70-2000; Harvard apparatus Massachusetts, U.S). The doses of Pep19-4LF used in this study were based on efficacy seen in previous *in vitro* and *in vivo* studies. 13,12,14,15

Immunoblot analysis

Western blot was performed as previously described.¹⁷

Cytokine analysis

Concentrations of serum cytokines were determined using a commercially available cytometric bead array (BD Biosciences, San Jose, CA or BioLegend, San Diego, CA) following the manufacturer/product specific protocol.

Immunohistochemistry

Lung samples were obtained at the end of the experiment and fixed in formalin for 48 h and immunohistochemistry was performed as described previously. 17

Determination of Myeloperoxidase Activity

Myeloperoxidase (MPO) activity was determined as an indicator of neutrophil accumulation into the lungs and performed as described previously. 16

Human mononuclear cells study

Mononuclear cells (MNC) were isolated from heparinized blood samples obtained from healthy donors as described previously. 11

Isothermal titration calorimetry

Microcalorimetric experiments of peptide binding to heparan sulfate were performed using isothermal titration calorimeter as described before. 15

Statistics

Unless otherwise stated, the data is expressed as median and standard error or described in box and whisker format showing medians, interquartile ranges and full ranges of *n* observations, where *n* represents the number of animals/experiments studied. Statistical analysis was carried out using Prism 6 for Mac OS X (GrapPad, San Diego, CA, USA). The distribution of the data was assessed using D'Agostino's K-squared test or Kolmogorov–Smirnov test. Unless otherwise stated, normal distributed data were assessed by 1 or 2-way analysis of variance followed by Bonferroni post hoc test. Unless otherwise stated, not normally distributed data were analyzed with a non-parametric test (Kruskal-Wallis followed by Dunn test). A p < 0.05 was considered to be significant.

RESULTS

Plasma concentrations of the host defence AMP cathelicidin LL-37

Figure 1 shows plasma concentrations of LL-37 in healthy volunteers and trauma patients recruited from an urban major trauma center. The median age of the healthy volunteers were 47 (32-53) years with 80% male. Further demographic and clinical parameters of trauma patients are described in table 1. Admission blood samples of trauma patients were obtained within 2 h of injury (Table 1). When compared to healthy volunteers, trauma patients ($n = 24$) and trauma hemorrhage patients (n = 23) (defined as patients who received greater than or equal to two units of packed red blood cells on admission) showed significantly higher plasma levels of LL-37. When compared to trauma patients, the plasma levels of LL-37 were significantly higher in trauma hemorrhage patients (Fig. 1).

Pep19-4LF attenuates the decline in blood pressure during resuscitation after HS

When compared to sham-operated rats, HS-rats treated with vehicle showed a significant decline in mean arterial pressure (MAP) after resuscitation (Fig.2). Intravenous administration of highdose Pep19-4LF (333 μg/kg x h, in 0.9% saline) significantly attenuated the decline in MAP observed during the resuscitation period in HS-rats, while the low dose of Pep19-4LF (66 μg/kg x h, in 0.9% saline) had no significant effect. In contrast, the high dose of Pep19-4LF had no effect of MAP in sham-operated rats (Fig. 2).

Pep19-4LF attenuates the organ injury and dysfunction caused by HS

When compared to sham-operated rats, rats subjected to HS treated with vehicle exhibited a renal dysfunction, as indicated by significant increases in serum urea (Fig. 3A) and creatinine (Fig. 3B), and a significant decline in creatinine clearance (Fig. 3C). HS-rats also exhibited significant increases in alanine aminotransferase, aspartate aminotransferase, amylase and lipase, indicating the development of liver and pancreatic injury, respectively (Fig. 3E-G). In addition, HSrats exhibited a significant increase in serum lactate, indicating global ischemia (Fig. 3D). When compared to vehicle-treated rats, intravenous administration of high-dose Pep19-4LF significantly attenuated the organ injury and dysfunction as well as the rise in lactate caused by HS (Fig. 3A-H). Although the lower dose of Pep19-4LF reduced pancreatic injury, it had no effect on any of the other parameters measured (Fig. 3A-H).

Pep19-4LF attenuates lung inflammation caused by HS

Having shown that Pep-4LF-treatment attenuates kidney dysfunction and liver injury, we next investigated the effects of Pep-4LF on lung inflammation measured as recruitment of macrophages (CD68-positive cells) and neutrophil activation (MPO activity) into the lung. When compared to sham-operated rats, we found a significant increase in CD68-positive cells and MPOactivity in lungs of HS-rats treated with vehicle (Fig. 4A-C). Treatment of HS-rats with Pep19-4LF significantly attenuated the recruitment of macrophages and neutrophil activation caused by HS (Fig. 4A-C).

Pep19-4LF attenuates the increase in interleukin (IL)-6 and monocyte chemotactic protein-1 (MCP-1) caused by HS

Having shown that Pep-4LF attenuates the activation of NF-κB in kidney and liver, we next investigated the effects of Pep-4LF on the formation of pro- and anti-inflammatory cytokines caused by HS. When compared to sham-operated rats, HS-rats treated with vehicle developed a significant increase in serum IL-6 and MCP-1. In HS-rats, we also observed increases in serum IL-10 and C-X-C motif ligand 1 (CXCL1), but these effects were not significant (Fig. 5A,B).

Treatment of HS-rats with Pep19-4LF abolished the increases in IL-6, MCP-1, IL-10, and CXCL1 caused by HS (Fig. 5A-D).

Pep19-4LF attenuates the activation of NF-κB (liver and kidney) caused by HS

Having shown that Pep19-4LF significantly attenuates kidney dysfunction and liver injury caused by HS, we next explored the potential mechanism(s) underlying the observed beneficial effects of high dose of Pep19-4LF. When compared to sham-operated rats, HS-rats treated with vehicle exhibited a significant increase in the nuclear translocation of the p65 subunit of NF-κB in kidney and liver (Fig. 6A,E) as well as a significantly increased degree of phosphophorylation of IκB kinase α and β (IKK α/β) on Ser^{176/180} and of I_{KB} \langle on Ser^{32/36} in both kidney and liver (Fig. 6B,C,F,G). The intravenous administration of high-dose Pep19-4LF attenuated the nuclear translocation of the NF-_KB subunit p65, the phosphorylation of IKK α/β on Ser^{176/180}, and of I_{KB α} on Ser^{32/36} in both liver and kidney (Fig. 6B,C,F,G).

Pep19-4LF increases activation of Akt and eNOS in kidney and liver after HS

As activation of the Akt-survival pathway is known to reduce HS-induced organ dysfunction^{16,17} we next investigated whether the high dose of Pep19-4LF activates Akt in kidney and liver of HS-rats (Fig. 6D,I). Moreover, we investigated the phosphorylation of eNOS in kidney and liver (Fig. 6E,J). When compared to sham-operated rats, HS-rats treated with vehicle showed a significant reduction in the phosphorylation of Akt on Ser^{473} and eNOS on Ser^{113} in both kidney and liver (Fig. 6D,E,I,J). In contrast, treatment of HS-rats with Pep19-4LF attenuated the decline in Akt phosphorylation on Ser⁴⁷³ and of eNOS phosphorylation on Ser¹¹³ in kidney and liver, when compared to HS rats treated with vehicle (Fig. 6D,E,I,J).

Pep19-4LF inhibits the heparan sulfate-induced TNFα **secretion in human peripheral blood mononuclear cells**

As discussed above, traumatic injury and trauma-associated HS result in release a variety of endogenous TLR ligands, including heparan sulfate ¹⁸. We report here that heparan sulfate stimulates the release of TNF α from human MNCs, and that this effect is reduced/abolished in a concentration-dependent manner by Pep19-4LF (Fig. 7A).

Pep19-4LF exhibits a strong binding to heparan sulfate

To gain a better understanding of the mechanism(s) by which Pep19-4LF reduces the formation of TNF α in human MNC challenged with heparan sulfate, we investigated the potential binding of Pep19-4LF to heparan sulfate by isothermal titration calorimetry. There was a strong exothermic reaction between the two reactants, running into a saturation of binding at higher mass ratios (Fig. 7B). This high affinity binding was characteristic for a chemical complex reaction, and may explain that the binding epitopes of heparan sulfate to TLR4 are hidden by the peptide.

Pep19-4LF does not show cytotoxic activity

Finally, possible cytotoxic effects of Pep19-4LF were studied in the hemolysis assay with RBC as sensitive target cells for cytotoxicity. Pep19-4LF caused only a very small degree of hemolysis in concentrations of up to 100 μg/ml (Fig. 7C).

DISCUSSION

The main findings of this study are that (i) plasma levels of the host-defence/antimicrobial peptide LL-37 are elevated in patients with trauma-associated HS, and that (ii) the synthetic hostdefence/antimicrobial peptide Pep19-4LF attenuates the renal dysfunction and liver injury caused by HS in the rat. Pep19-4LF also reduced both the local (lung) and the systemic (rise in plasma IL-6 and MCP1) inflammation caused by HS. Pep19-4LF activated the Akt/eNOS cell survival pathway and attenuated the activation of NF-κB in kidney/liver of HS-rats. Moreover, Pep4LF attenuated the release of TNF α caused by the DAMP heparan sulfate in human peripheral blood MNCs.

Trauma-hemorrhage is associated with the release of DAMPs (from the host) and PAMPs (from bacteria after bacterial translocation from the gut), which drive inflammation and contribute to tissue/organ-damage. It is unclear whether trauma-hemorrhage also triggers the release of hostdefence/antimicrobial peptides, but the serum of trauma patients does have an enhanced antimicrobial capacity, which limits the growths and ultimately kills Gram-negative and Grampositive bacteria. ¹⁹ We report here for the first time that trauma leads (within 2 h) to a significant increase in the plasma levels of the host-defence/antimicrobial peptide LL-37. Most notably, the highest levels of LL-37 were found in patients with trauma complicated by severe hemorrhage (Fig.1). LL-37 is primarily released by T cells and natural killer cells²⁰. Notably, natural killer cells are elevated and activated within 2 h of injury in trauma patients who later develop MOF, suggesting that these cells contribute to the rapid rise in plasma LL-37. 21

Having found that trauma-HS increases plasma LL-37, we next performed a reverse translational approach and investigated whether pharmacological intervention with a synthetic hostdefence/antimicrobial peptides attenuates the MOF associated with HS in rats. As the therapeutic

13

use of LL37 in man is limited by its systemic toxicity in therapeutic doses,^{9,10} we synthesized Pep19-4LF, which does not cause any significant adverse effects (hemolysis) in the doses used (Fig. 7C).

The administration of Pep19-4LF significantly attenuated the fall in blood pressure (Fig. 2) as well as the rise in serum lactate caused by HS (Fig. 3D). Thus, Pep19-4LF reduces the delayed vascular decompensation and organ/tissue ischemia probably due to increased microvascular perfusion secondary to increased perfusion pressure. Moreover, Pep19-4LF significantly attenuated the liver injury, renal dysfunction, pancreatic injury and lung inflammation caused by HS (Fig. 3 A, B, E, F, G, H). As neutrophils and macrophages play an important role in HSassociated lung inflammation, we evaluated the degree of macrophage infiltration (measured as number of CD68-positive cells) and the degree of neutrophil activation (measured as MPOactivity) in the lung. HS resulted in a significant increase in the number of macrophages in the lung as well as a significant increase in MPO-activity, both of which was attenuated by the treatment of HS-rats with Pep19-4LF (Fig. 4). Neutrophils and macrophages release (via degranulation) proinflammatory cytokines, such as IL-6 or MCP-1 which importantly contribute to acute lung injury/inflammation.²² The pro-inflammatory cytokines IL-6 and MCP-1 are important mediators of alterations associated with organ dysfunction and even lethality following HS and resuscitation.23,24,25,26 For instance, a monoclonal antibody against IL-6 reduces the organ dysfunction and inflammation caused by HS. ⁴² Indeed, we report here that Pep19-4LF also attenuates the rise in serum IL-6 and MCP-1 caused by HS in the rat (Fig. 5).

What, then, are the mechanisms by which Pep19-4LF attenuates HS-associated organ injury/dysfunction? There is good evidence that PAMPs and DAMPs released during trauma-HS interact with Toll-like receptors (i.e. TLR2 and TLR4) resulting in activation of NF-κB.^{27,28,29} Indeed,

14

we observed a significant increase in (a) the nuclear translocation of NF-κB subunit p65 (Fig. 6 A, E) and (b) the degree of phosphorylation of IKK α/β on Ser^{176/180} (Fig. 6B,F) and of I_{KB α} on Ser^{32/36} (Fig. 6C,G) in liver/kidneys of rats with HS. This activation of NF-κB in key target organs was attenuated in HS-rats treated with Pep19-4LF during resuscitation. I κ B α masks the nuclear localization signals of NF-κB proteins and sequesters NF-κB as an inactive complex in the cytoplasm, thereby inhibiting NF-κB.^{30,31} Signal-induced proteolytic degradation of IκBα, which has been phosphorylated by I_KB kinases (IKK α/β) liberates NF-_KB to translocate to the nucleus.³¹ Subsequently, NF-κB activates the transcription of a number of genes involved in producing proinflammatory cytokines and chemokines known to result in the transcription of a multitude of proinflammatory cytokines, chemokines and proteins that are widely implicated in the pathophysiology of MOF.^{17,16} Thus, the organ protective effects of Pep19-4LF in HS are associated with a significant reduction in the activation of the NF-κB pathway, which in turn accounts for the reduced formation of IL-6 and MCP-1 (see above).

We also investigated the effects of HS with or without Pep19-4LF on the degree of activation of the Akt-survival pathway (Fig. 6). When compared to sham rats, HS rats treated with vehicle showed a significantly decreased phosphorylation of Akt on Ser⁴⁷³ (indicating reduction in activity of this kinase) in kidney and liver, which makes these organs less resistant to organ injury (see below). In contrast, Pep19-4LF attenuated the decline in Akt-activation caused by HS (Fig. 6D, H). Akt is a member of the phosphoinositide-3-kinase (PI3K) signal transduction enzyme family. When activated (phosphorylated on Ser⁴⁷³) by its upstream regulator PI3K, Akt controls inflammatory response, chemotaxis and apoptosis.³² Most notably, activation of the Akt-survival pathway reduces organ injury in many conditions associated with ischemia/inflammation including ventilation-induced lung injury, sepsis-induced organ dysfunction, myocardial infarction, and HS-

induced organ dysfunction.^{33,34,35,36,16}. Moreover, activation of Akt results in phosphorylation and activation of eNOS at Ser¹¹⁷⁷ that enhances the formation of small amounts of NO, which is pivotal for the preservation of microvascular perfusion and, hence, reducing organ injury.^{37,34,38} We report here that the degree of eNOS phosphorylation on Ser¹¹⁷⁷ is significantly higher in HS-rats treated with Pep19-4LF indicating activation of eNOS and enhanced formation of NO in the microcirculation at least of kidney and liver. We propose that the enhanced formation of NO by eNOS in HS-rats treated with Pep19-4LF contributes to improved microcirculatory perfusion resulting in better tissue oxygenation and lower lactate levels (Fig 3D). Thus, the organ protective effects of Pep19-4LF in HS are associated with a significant activation of the Akt/eNOS survival pathway.

Traumatic injury and trauma-associated HS result in the release of a variety of endogenous TLR ligands, including heparan sulfates.^{27,28,29,39} Moreover, the degradation of the glycocalyx (and subsequent liberation of heparan sulfates) induces remote organ injury after trauma/hemorrhagic shock,^{40,41,42,43} suggesting heparan sulfate as a potential therapeutic target for Pep19-4LF. Indeed, using isothermal titration calorimetry, we found a strong Coulomb interaction between Pep19-4LF and heparan sulfate, as indicated by strong exothermic reaction running into a saturation characteristic (Fig. 7B). These results are in line with our recent findings, reporting a strong Coulomb interaction between the related peptide Pep2.5 and heparan sulfate. ¹⁵ To investigate whether Pep19-4LF inhibits the release of TNF α caused by heparan sulfate in human cells, we exposed human MNCs to heparan sulfate in the presence or absence of Pep19-4LF. Moist notably, Pep19-4LF attenuated the release of TNF α caused by heparin sulfate in these cells in a dose-related fashion (Fig. 7A) indicating that Pep19-4LF is able to prevent the activation of human cells by the DAMP heparin sulfate. Thus, these findings indicate, that the organ protective and anti-inflammatory effects of Pep19-4LF in HS are, at least partly, associated with its interaction with relevant DAMPs, such as heparan sulfate.

In conclusion, we report here for the first time that trauma and trauma-haemorrhage result in a significant release of the host-defence/antimicrobial peptide LL-37. As the systemic administration of higher doses of LL-37 leads to adverse effects, we have synthetized a small hostdefence/antimicrobial peptide, Pep19-4L. Like LL-37, Pep19-4LF neutralizes the effects of LPS and lipoproteins.¹² LL-37 also interacts with and attenuates the effects of several DAMPs.^{6,8} We report here that Pep19-4LF abolishes the release of TNF α caused by heparan sulfate in human mononuclear cells. In addition, Pep19-4LF attenuates the organ injury/dysfunction caused by severe hemorrhage and resuscitation in the anesthetized rat. This protective effect of Pep19-4LF was associated with activation of the Akt/eNOS-survival pathway (kidney and liver), which increases the resistance of these organs to injury. In addition, Pep19-4LF also attenuates the activation of NF-κB in these organs, resulting in the reduced formation of the pro-inflammatory cytokines IL-6 and MCP-1. Thus, we propose that Pep19-4LF may be useful to reduce the organ injury/dysfunction and inflammation caused by severe hemorrhage and resuscitations in patients with trauma.

REFERENCES

- 1. WHO. Violence and injuries: the facts 2014. *Geneva, Switzerland World Health Organization* 2015.
- 2. Sauaia A, Moore EE, Johnson JL, et al. Temporal trends of postinjury multiple-organ failure: still resource intensive, morbid, and lethal. *J Trauma Acute Care Surg* 2014; 76(3):582-92, discussion 592-3.
- 3. Lord JM, Midwinter MJ, Chen Y-F, et al. The systemic immune response to trauma: an overview of pathophysiology and treatment. *The Lancet* 2014; 384(9952):1455-1465.
- 4. Lehrer RI, Ganz T. Antimicrobial peptides in mammalian and insect host defence. *Curr Opin Immunol* 1999; 11(1):23.
- 5. Guani-Guerra E, Santos-Mendoza T, Lugo-Reyes SlO, et al. Antimicrobial peptides: general overview and clinical implications in human health and disease. *Clin Immunol* 2010; 135(1):1-11.
- 6. Martin L, Meegern Av, Domming S, et al. Antimicrobial Peptides in Human Sepsis. *Front Immunol* 2015; 6:404.
- 7. Schuerholz T, Brandenburg K, Marx G. Antimicrobial peptides and their potential application in inflammation and sepsis. *Crit Care* 2012; 16(2):207.
- 8. Hu Z, Murakami T, Suzuki K, et al. Antimicrobial cathelicidin peptide LL-37 inhibits the LPS/ATP-induced pyroptosis of macrophages by dual mechanism. *PLoS One* 2014; 9(1):e85765.
- 9. Hancock REW, Sahl H-G. Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. *Nat Biotechnol* 2006; 24(12):1551-7.
- 10. Zhang L, Falla T. Antimicrobial peptides: therapeutic potential. *Expert Opin Pharmacother* 2006; 7(6):653-663.

18

- 11. Gutsmann T, Razquin-Olazarán I, Kowalski I, et al. New antiseptic peptides to protect against endotoxin-mediated shock. *Antimicrob Agents Chemother* 2010; 54(9):3817-24.
- 12. Tejada GMd, Heinbockel L, Ferrer-Espada R, et al. Lipoproteins/peptides are sepsisinducing toxins from bacteria that can be neutralized by synthetic anti-endotoxin peptides. *Sci Rep* 2015; 5:14292.
- 13. Schuerholz T, Doemming S, Horne M, et al. The anti-inflammatory effect of the synthetic antimicrobial peptide 19-2.5 in a murine sepsis model: a prospective randomized study. *Crit Care* 2013; 17(1):R3.
- 14. Martin L, Schmitz S, De Santis R, et al. Peptide 19-2.5 inhibits heparan sulfate-triggered inflammation in murine cardiomyocytes stimulated with human sepsis serum. *PLoS One* 2015; 10(5):e0127584.
- 15. Martin L, De Santis R, Koczera P, et al. The Synthetic Antimicrobial Peptide 19-2.5 Interacts with Heparanase and Heparan Sulfate in Murine and Human Sepsis. *PLoS One* 2015; 10(11):e0143583.
- 16. Sordi R, Nandra KK, Chiazza F, et al. Artesunate Protects Against the Organ Injury and Dysfunction Induced by Severe Hemorrhage and Resuscitation. *Ann Surg* 2016; (Epub ahead of print)
- 17. Sordi R, Chiazza F, Johnson FL, et al. Inhibition of IkappaB Kinase Attenuates the Organ Injury and Dysfunction Associated with Hemorrhagic Shock. *Mol Med* 2015; 21:563-75.
- 18. Rahbar E, Cardenas JC, Baimukanova G, et al. Endothelial glycocalyx shedding and vascular permeability in severely injured trauma patients. *J Transl Med* 2015; 13:117.
- 19. Lippross S, Klueter T, Steubesand N, et al. Multiple trauma induces serum production of host defence peptides. *Injury* 2012; 43(2):137-42.
- 20. Agerberth B, Charo J, Werr J, et al. *Blood* 2000; 96(9):3086-3093.
- 21. Manson J, Cole E, De'Ath H, et al. Early changes within the lymphocyte population are associated with the development of multiple organ dysfunction syndrome in trauma patients. *Crit Care* 2016; 20(1):176.
- 22. Dewar D, Moore FAM, Moore EE, et al. Postinjury multiple organ failure. *Injury* 2009; 40:912-918.
- 23. Jarrer D, Chaudry IH, Wang P. Organ dysfunction following hemorrhage and sepsis: mechanisms and therapeutic approaches (review). *Int J Mol Med* 1999; 4:575-583.
- 24. Sordi R, Chiazza F, Patel NS, et al. 'Preconditioning' with low dose lipopolysaccharide aggravates the organ injury / dysfunction caused by hemorrhagic shock in rats. *PLoS One* 2015; 10(4):e0122096.
- 25. Bahrami S, Yao YM, Leichtfried G et al. Significance of TNF in hemorrhage-related hemodynamic alterations, organ injury, and mortality in rats. *Am J Physiol* 1997; 272:H2119.
- 26. Zhang Y, Zhang J, Korff S, et al. Delayed neutralization of interleukin 6 reduces organ injury, selectively suppresses inflammatory mediator, and partially normalizes immune dysfunction following trauma and hemorrhagic shock. *Shock* 2014; 42(3):218-27.
- 27. Tang D, Kang R, Coyne CB, et al. PAMPs and DAMPs: signal 0s that spur autophagy and immunity. *Immunol Rev* 2012; 249(1):158.
- 28. Pradeu T, Cooper E. The danger theory: 20 years later. *Front Immunol* 2012; 3:287.
- 29. Martin L, Koczera P, Zechendorf E, et al. The Endothelial Glycocalyx: New Diagnostic and Therapeutic Approaches in Sepsis. *Biomed Res Int* 2016; 2016:3758278.
- 30. Jacobs MD, Harrison SC. Structure of an IkappaBalpha/NF-kappaB complex. *Cell* 1998; 95:749-758.
- 31. Senftleben U, Karin M. The IKK/NF-kappaB pathway. *Crit Care Med* 2002; 30(Suppl1):S18-S26.
- 20
- 32. Cantley L. The phosphoinositide 3-kinase pathway. *Science* 2002; 296:1655-1657.
- 33. Shu YS, Tao W, Miao QB, et al. Improvement of ventilation-induced lung injury in a rodent model by inhibition of inhibitory kB kinase. *J Trauma Acute Care Surg* 2014; 76:1417– 1424.
- 34. Khan AI, Coldewey SM, Patel NSA, et al. Erythropoietin attenuates cardiac dysfunction in experimental sepsis in mice via activation of the beta-common receptor. *Dis Model Mech* 2013; 6(4):1021-30.
- 35. Coldewey SM, Rogazzo M, Collino M, et al. Inhibition of IkappaB kinase reduces the multiple organ dysfunction caused by sepsis in the mouse. *Dis Model Mech* 2013; 6(4):1031-42.
- 36. Cai Z, Semenza GL. Phosphatidylinositol-3-kinase signaling is required for erythropoietinmediated acute protection against myocardial ischemia/reperfusion injury. *Circulation* 2004; 109(17):2050-3.
- 37. Cabrales P, Tsai AG, Intaglietta M. Exogenous nitric oxide induces protection during hemorrhagic shock. *Resuscitation* 2009; 80(6):707-12.
- 38. Nandra KK, Collino M, Rogazzo M, et al. Pharmacological preconditioning with erythropoietin attenuates the organ injury and dysfunction induced in a rat model of hemorrhagic shock. *Dis Model Mech* 2013; 6(3):701-9.
- 39. Horst K, Hildebrand F, Pfeifer R, et al. Impact of haemorrhagic shock intensity on the dynamic of alarmins release in porcine poly-trauma animal model. *Eur J Trauma Emerg Surg* 2016; 42:67-75.
- 40. Wu H, Ma J, Wang P, et al. HMGB1 contributes to kidney ischemia reperfusion injury. *J Am Soc Nephrol* 2010; 21(11):1878-90.
- 41. Sodhi CP, Jia H, Yamaguchi Y, et al. Intestinal Epithelial TLR-4 Activation Is Required for the Development of Acute Lung Injury after Trauma/Hemorrhagic Shock via the Release of HMGB1 from the Gut. *J Immunol* 2015; 194(10):4931-9.
- 42. Torres Filho IP, Torres LN, Salgado C, et al. Plasma syndecan-1 and heparan sulfate correlate with microvascular glycocalyx degradation in hemorrhaged rats after different resuscitation fluids. *Am J Physiol Heart Circ Physiol* 2016; 310(11):H1468-1478.
- 43. Levy RM, Mollen KP, Prince JM, et al. Systemic inflammation and remote organ injury following trauma require HMGB1. *Am J Physiol Regul Integr Comp Physiol* 2007; 293(4):R1538-44.

ACKNOWLEDGEMENTS

NY was supported by Gifu University Hospital Advanced Critical Care Center. LM has received grants by the Deutsche Forschungsgemeinschaft (DFG, MA 7082/1-1). This work was supported, in part, by the University of Turin (ex-60% 2015 A and B) and by the William Harvey Research Foundation and forms part of the research themes contributing to the translational research portfolio of Barts and the London Cardiovascular Biomedical Research Unit that is supported and funded by the National Institute for Health Research. This work also contributes to the Organ Protection research theme of the Barts Centre for Trauma Sciences supported by the Barts and The London Charity (Award 753/1722).

AUTHOR CONTRIBUTIONS

Conception and design: NY, LM, and CT; Animal experiments: NY; Cell culture experiments: LH, TG, WC, and KB; Human sample analysis: NY, LM, EZ, GM, TS; Animal sample analyses: NY, LM, GP, FC, BV, MC, and TS; Clinical study and patient data analyses: LM, JS, KB, GM, and TS; Peptide development and synthesis: KB, LH, TG, WC. Statistical analyses: NY, LM, and CT; Drafting the manuscript for important intellectual content: NY, LM, EZ, GP, FC, BV, MC, JS, LH, TG, WC, KB, GM, TS, KB, and CT; All authors reviewed and approved the manuscript.

ADDITIONAL INFORMATION

Supplementary information accompanies this paper.

Competing financial interests: KB has a patent for the structure of the synthetic antimicrobial peptide 19-4LF (Brandenburg Antiinfektiva, Borstel, Germany): Patent-No: PCT/EP2009/002565. KB chief scientific officer and TS is chief medical officer of Brandenburg Antiinfektiva GmbH. All the other authors declare no conflicts of interest.

LEGENDS TO FIGURES

FIGURE 1. LL-37 plasma levels in human healthy volunteers and trauma patients

Plasma concentrations of the cathelicidin LL-37 were assessed in control healthy volunteers ($n =$ 10) and in trauma patients ($n = 24$) as well as in trauma hemorrhagic patients ($n = 23$). Data are expressed as box and whisker min to max for n number of observations. + = mean value. * P < 0.05 vs. healthy; [§]P < 0.05 vs. trauma (Kruskall-Wallis test with Dunn´s multiple comparisons test).

FIGURE 2. Pep19-4LF attenuates the decline in MAP during resuscitation after HS.

HS-rats received continuous administration of low-dose (LD; 66 μg/kg x h) or high-dose (HD; 333 μg/kg x h) of Pep-4LF or saline (vehicle) throughout 4 h after resuscitation. Sham animal were used as control and received saline or high-dose of Pep19-4LF. The MAP was recorded during the whole experiment. The following groups were studied: sham + vehicle ($n = 11$); sham + Pep4LF-HD (n = 6); HS + vehicle (n = 12), HS + Pep4LF-LD (n = 4); HS + Pep19-4LF-HD (n = 8). Data are expressed as mean ± SEM for *n* number of observations. Statistical analysis was performed using 2-way ANOVA followed by Bonferroni post hoc test. **P* < 0.05 vs HS + vehicle. ANOVA indicated analysis of variance; SEM, standard error of the mean.

FIGURE 3. Pep19-4LF attenuates the organ injury and dysfunction caused by HS.

(A) Serum urea, (B) serum creatinine, (C) creatinine clearance (CCr), (D) serum lactate, (E) serum alanine transaminase (ALT), (F) serum aspartate transaminase (AST), (G) serum amylase, and (H) serum lipase of HS or sham-operated rats. All parameters were assessed 4 h subsequent to HS. HS-rats received continuous administration of low-dose (LD; 66 μg/kg x h) or high-dose (HD; 333 μg/kg x h) of Pep-4LF or saline (vehicle) throughout 4 hours after resuscitation. Sham animal were used as control and received saline or high-dose of Pep19-4LF. Data are shown as box and whiskers, showing medians, interquartile range, and full range. The following groups were studied: sham + vehicle (n = 11); sham + Pep4LF-HD (n = 6); HS + vehicle (n = 12), HS + Pep4LF-LD (n $= 4$); HS + Pep19-4LF-HD (n = 8). Statistical analysis was performed using 1-way ANOVA followed by Bonferroni post hoc test. §*P* < 0.05 vs sham + vehicle and **P* < 0.05 vs HS + vehicle. ANOVA indicated analysis of variance.

FIGURE 4. Pep19-4LF attenuates lung inflammation caused by HS.

Representative images for CD68 as a macrophage marker (200 fold), (B) quantitative analysis of the number of CD68-positive cells/mm², (C) MPO activity in lungs of HS or sham-operated rats. The scale bar represents 100µm. All parameters were assessed 4 h subsequent to HS. HS-rats received continuous administration of vehicle (saline) or Pep19-4LF of (333 μg/kg x h) throughout 4 hours after resuscitation. Sham animal were used as control and received saline. Data are shown as box and whiskers, showing medians, interquartile range, and full range. The following groups were studied: sham + vehicle $(n = 4)$; HS + vehicle $(n = 6)$, HS + Pep19-4LF HD $(n = 6)$. Statistical analysis was performed using 1-way ANOVA followed by Bonferroni post hoc test. §*P* < 0.05 vs sham + vehicle and **P* < 0.05 vs HS + vehicle. ANOVA indicated analysis of variance.

FIGURE 5. Pep19-4LF attenuates the increase in IL-6 and MCP-1 caused by HS.

The serum concentrations of (A) interleukin (IL)-6, (B) monocyte chemotactic protein-1 (MCP-1), (C) IL-10, and (D) C-X-C motif ligand 1 (CXCL1) were determined using a cytometric bead array in sham and HS rats treated with vehicle or Pep19-4LF (333 μg/kg x h) throughout 4 h after resuscitation. All parameters were assessed 4 h subsequent to HS. Data are presented as box and whiskers format, showing medians, interquartile range, and full range. (E) Heatmap of measured cytokines. The following groups were studied: sham + vehicle (n = 11); HS + vehicle (n $=$ 12), HS + Pep19-4LF-HD (n = 8). Statistical analysis was performed using 1-way ANOVA

followed by Bonferroni post hoc test. §*P* < 0.05 vs sham + vehicle and **P* < 0.05 vs HS + vehicle. ANOVA indicated analysis of variance.

FIGURE 6. Pep19-4LF attenuates the activation of NF-κB and increases the activation of Akt and eNOS in kidney and liver after HS.

The nuclear translocation of p65 subunit of NF-κB (kidney (A), liver (F)), and the phosphorylation of Ser^{176/180} on IKKα/β (kidney (B), liver (G)), Ser ^{32/36} on IκBα (kidney (C), liver (H)), Ser⁴⁷³ on Akt (kidney (D), liver (I)), and Ser113 on eNOS (kidney (E), liver (J) of sham and HS rats treated with vehicle or high-dose of Pep19-4LF (333 μg/kg x h) upon resuscitation were determined by Western blotting. Protein expression was measured as relative OD. Data are shown as box and whiskers, showing medians, interquartile range, and full range. The following groups were studied: sham + vehicle (n =4); HS + vehicle (n = 4), HS + Pep19-4LF-HD (n = 4). Statistical analysis was performed using 1-way ANOVA followed by Bonferroni post hoc test. §*P* < 0.05 vs sham + vehicle and **P* < 0.05 vs HS + vehicle. ANOVA indicated analysis of variance; OD, optical density.

FIGURE 7. Pep19-4LF interacts with heparan sulfate

Inhibitory effect of Pep10-4LF on heparan sulfate-induced TNF α release in human peripheral blood mononuclear cells from healthy donors. Pep19-4LF was added at the indicated weight ratios of the concentrations of heparan sulfate to Pep19-4LF. (B) Enthalpy of the Pep19-4LF-heparan sulfate binding. Isothermal calorimetric titration of a $1 - 4$ mM Pep19-4LF solution into a 200 $\mu q/ml$ heparan sulfate dispersion. The enthalpy changes at each injection were measured and the area underneath each injection peak was integrated and plotted against the weight ratio of the concentrations of Pep19-4LF to heparan sulfate. A downward peak corresponds to an exothermic reaction, and an upward peak corresponds to an endothermic reaction. (C) Red blood cells, obtained from citrated human blood were suspended at a concentration equivalent to 5% of the

normal hematocrit. Pep19-4LF was added at different concentrations and the supernatants were analyzed for haemoglobin. Results are expressed as the percentage released with respect to sonicated controls (100% release) or controls processed without peptide (0% release).

Table 1. Admission Characteristics & Outcomes

PRBC: Packed Red Blood Cells; ISS: Injury Severity Score; AIS: Abbreviated Injury Severity Score; LOS: Length of Stay. Median (interquartile range) reported unless specified. *Trauma hemorrhage vs. No hemorrhage Mann-Whitney U-test unless ‡ Chi-Squared test.

SUPPLEMENTARY INFORMATION

A novel synthetic antimicrobial peptide protects against organ injury/dysfunction in a rat-model of severe hemorrhagic shock

Noriaki Yamada, Lukas Martin, Elisabeth Zechendorf, Gareth Samuel Desmond Purvis, Fausto Chiazza, Barbara Varrone, Massimo Collino, Joanna Shepherd, Lena Heinbockel, Thomas Gutsmann, Wilmar-Alexander Correa-Vargas, Klaus Brandenburg, Gernot Marx, Tobias Schuerholz, Karim Brohi, Christoph Thiemermann

METHODS

Study population and human outcome measurements

Study setting and participants

Trauma patients who presented to an urban major trauma centre were recruited to an ongoing, prospective observational cohort study called the 'Activation of Coagulation and Inflammation in Trauma Study II' (ACIT-II). This study was originally established in 2008 to investigate the biological mechanisms underlying acute traumatic coagulopathy and the inflammatory response to trauma. Adult trauma patients who require trauma team activation on admission were eligible for inclusion. Exclusion criteria included age under 16 years, transfer from another hospital, arrival-time greater than 120 min from injury, pre-hospital administration of  greater than 2000 ml crystalloid, greater than 5 % burns, severe liver disease, known bleeding abnormality (including anticoagulant medication), refused consent and vulnerable patients.

Patient selection

Trauma patients for this study ($n = 47$) were identified from the available ACIT-II cohort based on their Injury Severity Score (ISS) and blood product requirements during resuscitation. Patients were included if they had an ISS score greater than or equal to 16. Trauma Haemorrhage patients were defined as patients who received greater than or equal to two units of packed red blood cells (PRBC) on admission. Control patients (No Haemorrhage) received no PRBCs during the first 24 hours of their admission. We subsequently excluded patients with a head Abbreviated Injury Severity (AIS) score greater than three in order to ensure the groups were not skewed on the basis of severe traumatic brain injury. Furthermore, we included healthy volunteers $(n = 10)$ as a control group.

Sample and data collection

Admission data were collected on patient demographics, mechanism of injury, blood product use and baseline physiology. Arterial blood gas analysis for base deficit (BD) and lactate was performed during the trauma team resuscitation as part of normal care processes. Admission bloods were drawn within 2 hours of injury. Whole blood was collected in 4.5 ml citrated vacutainer tubes and centrifuged at 3400 rpm for 10 minutes. The plasma supernatant was centrifuged again at the same settings, and the double-spun plasma was subsequently stored in aliquots at -80 °C. Patient outcomes including 28-day mortality, ventilator days, critical care and hospital length of stay were recorded.

LL37-ELISA

Plasma-levels of the cathelicidin-derived human AMP LL-37 in patients were quantified using a commercially available ELISA kit (Cambridge Bioscience Ltd, Cambridge, UK) by following the manufacturer/product specific protocol.

Immunoblot analysis

Semi-quantitative immunoblot analyses of nuclear translocation of p65 and the phosphorylation of I_KBα, IKKα/β, Akt, and eNOS were carried out in tissue samples as described before. ¹ Briefly, lung, kidney and liver samples were homogenized in buffer and centrifuged at 4000 rpm for 5 min at 4℃To obtain the cytosolic fraction, supernatants were centrifuged at 14000 rpm at 4℃for 40 min. The pelleted nuclei were re-suspended in extraction buffer and centrifuged at 14,000 rpm for 20 min at 4℃. Protein content was determined on both nuclear and cytosolic extracts using bicinchoninic acid (BCA) protein assay (Thermo Fisher Scientific Inc, Rockford, IL). Proteins were separated by 8% sodium dodecyl sulphatepolyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinyldenediflouoride (PVDF) membrane, which was incubated with a primary antibody [mouse anti-total Akt (1:1000); mouse anti-IκBα pSer^{32/36} (1:1000); rabbit anti-NF-κB p65 (1:1000); rabbit anti-eNOS pSer¹¹³ (1:1000)]. Membranes were incubated with a secondary antibody conjugated with horseradish peroxidase (1:2000) for 30 min at room temperature and developed with ECL detection system. The immunoreactive bands were visualized by autoradiography and the densitometric analysis was performed using Gel Pro Analyzer 4.5, 2000 software (Media Cybernetics, Silver Spring, MD, USA). The membranes were stripped and incubated with β-actin monoclonal antibody (1:5000) and subsequently with an anti-mouse antibody (1:10000) to assess gel-loading homogeneity. Densitometric analysis of the related bands is expressed as relative optical density, and normalized using the related sham-operated band.

Peptide-synthesis

The synthesis and purification of Pep19-4LF was performed at the Research Center Borstel, Germany as described previously.² The amino acid sequence of this 19 $\acute{ }$ mer is GKKYRRFRWKFKGKLFLFG. Pep19-4LF was amidated at the C-terminal end and had a purity of > 95% as measured by HPLC and MALDI-TOF mass spectrometry.²

Immunohistochemistry

Lung samples were obtained at the end of the experiment and fixed in formalin for 48 h and immunohistochemistry was performed as described previously ¹. Briefly, lung tissue was embedding in paraffin and processed to obtain 4-μm sections. After deparaffinization, the slides were then incubated with rabbit anti-CD68 antibody ED1 (1:400; catalog no. MCA341R; AbD Serotec) for 1 h at 37°C and afterwards incubated for 30 min with labelled polymer-HRP antibody. Counterstaining was performed with Harris hematoxylin. Images were acquired using a NanoZoomer Digital Pathology Scanner (Hamamatsu Photonics K.K. Japan) and analyzed using the NDP Viewer software. The numbers of CD68 positive cells were counted in 10 randomly selected fields (200×) in a double-blinded manner by three independent investigators.

Hemorrhagic shock and quantification of organ injury and dysfunction

Hemorrhagic shock and quantification of organ injury and dysfunction were performed as described previously ³ . Rats were anesthetized by sodium thiopentone (120 mg/kg *i.p.* for induction, followed by 10 mg/kg *i.v.* for maintenance). We performed cannulation of the trachea, femoral artery (for measuring blood pressure), and carotid arteries (for blood withdrawal), jugular vein (for drug administration), and bladder (for collecting urine). We withdrew blood (up to 1 mL/min) via the cannula inserted in the carotid artery in order to achieve a fall in mean arterial pressure (MAP) to 27 to 32 mmHg. Thereafter, MAP was maintained at this level for a period of 90 min either by further withdrawal of blood during the compensation phase or administration shed blood during the decompensation phase. At 90 min after initiation of hemorrhage (or when 25% of the shed blood had to be reinjected to sustain MAP at 27 to 32 mmHg), animals were resuscitated with the remaining shed blood (mixed with 100 IU/mL heparinized saline) (over a period of 5 min) plus a volume of Ringer's lactate identical to the volume of blood spent during decompensation. During the last 3 h of resuscitation, urine was obtained for the estimation of creatinine clearance. Then, blood samples were collected via the carotid artery for measurement of lactate (Accutrend Plus Meter, Roche Diagnostics, West Sussex, UK) and organ injury/dysfunction parameters. Under deep anesthesia, the heart was removed to terminate the experiment. Blood samples were centrifuged to separate serum, which was used for the determination of urea, creatinine, aspartate aminotransferase (AST), alanine aminotransferase (ALT), lipase, amylase and creatine kinase (CK) by an external contract research facility (IDEXX Laboratories Ltd, West Yorkshire, UK) in a blinded fashion. In addition, lung, kidney and liver samples were taken and stored at -80 °C for further analysis. Sham-operated rats were used as control and underwent identical surgical procedures, but without hemorrhage or resuscitation. Blood pressures during the experiment were measured by powerlab®, and recorded and analyzed by Labchart® (AD instruments Ltd, Dunedin, New Zealand).

Human mononuclear cells study

Mononuclear cells (MNC) were isolated from heparinized blood samples obtained from healthy donors as described previously. ⁴ The cells were re-suspended in medium (RPMI 1640), and their number was equilibrated at 5 x 106 cells/ml. For stimulation, 200 μl MNC (1x 106 cells) was transferred into each well of a 96-well culture plate. Heparan sulfate and the mixtures of heparan sulfate and peptide were pre-incubated for 30 min at 37°C and added to the cultures at 20 μ per well. The cells were then incubated for 4 h at 37°C with 5% CO2. Supernatants were collected after centrifugation of the culture plates for 10 min at 400 x g and stored at - 20°C until immunological determination of tumor necrosis factor alpha (TNF α) was carried out with a sandwich enzyme-linked immunosorbent assay (ELISA) using a monoclonal antibody against $TNF\alpha$ (clone 6b; Intex AG, Switzerland) and described previously in detail. 4

Isothermal titration calorimetry (ITC)

Microcalorimetric experiments of peptide binding to heparan sulfate were performed on a MSC isothermal titration calorimeter (MicroCal Inc., Northampton, MA) at 37 °C as described before.⁵ Briefly, after thorough degassing of the samples, Pep19-4LF (1 to 4 mM in 20 mM HEPES, pH 7.0) was titrated to a heparan sulfate suspension (200 μg/ml in 20 mM HEPES, pH 7.0). The enthalpy change during each injection was measured by the instrument, and the area underneath each injection peak was integrated (Origin; MicroCal) and plotted against the weight ratio of the concentrations of peptide to heparan sulfate. Titration of the pure peptide into HEPES buffer resulted in a negligible endothermic reaction due to dilution. All experiments were carried out in duplicate.

Hemolysis assay

Red blood cells (RBC) were obtained from citrated human blood by centrifugation (1,500 x g; 10 min), washed three times with isotonic 20 mM phosphate- NaCl buffer (pH 7.4), and suspended in the same buffer at a concentration equivalent to 5% of the normal hematocrit. Forty-microliter aliquots of this RBC suspension were added to 0.96 ml of Pep19-4LF dilutions prepared in the same isotonic phosphate solution, incubated at 37°C for 30 min, and centrifuged (1,500 x g, 10 min). The supernatants were analyzed spectrophotometrically (with absorbance at 543 nm) for hemoglobin, and results were expressed as the percentage released with respect to sonicated controls (100% release) or controls processed without peptide (0% release).

Materials

Unless otherwise stated, all compounds were from Sigma-Aldrich Company Ltd (Poole, Dorset, UK). Ringer's Lactate was from Baxter Healthcare Ltd (Deerfield, IL); Thiopental sodium from Archimedes Pharma Limited (Reading, UK).

Supplemental Figure 1. Entire study course of hemorrhagic shock in rats.

REFERENCES

- 1. Sordi R, Chiazza F, Johnson FL, et al. Inhibition of IkappaB Kinase Attenuates the Organ Injury and Dysfunction Associated with Hemorrhagic Shock. *Mol Med* 2015; 21:563-75.
- 2. Tejada GMd, Heinbockel L, Ferrer-Espada R, et al. Lipoproteins/peptides are sepsis- inducing toxins from bacteria that can be neutralized by synthetic anti-endotoxin peptides. *Sci Rep* 2015; 5:14292.
- 3. Sordi R, Nandra KK, Chiazza F, et al. Artesunate Protects Against the Organ Injury and Dysfunction Induced by Severe Hemorrhage and Resuscitation. *Ann Surg* 2016 [Epub ahead of print].
- 4. Gutsmann T, Razquin-Olazarán I, Kowalski I, et al. New antiseptic peptides to protect against endotoxin-mediated shock. *Antimicrob Agents Chemother* 2010; 54(9):3817-24.
- 5. Martin L, De Santis R, Koczera P, et al. The Synthetic Antimicrobial Peptide 19-2.5 Interacts with Heparanase and Heparan Sulfate in Murine and Human Sepsis. *PLoS One* 2015; 10(11):e0143583.

veh

0

T₂₀

Sham HS

Pep^{kxxxx}D

B $\qquad \qquad$ **C**

Sham HS

Pep⁴-LF-XID

veh

veh

0

C

