

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

On Inverse Association between Helicobacter pylori Gastritis and Microscopic Colitis: The European Data

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1632796> since 2019-07-30T16:18:31Z

Published version:

DOI:10.1097/MIB.0000000000000704

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

On Inverse Association Between Helicobacter pylori Gastritis and Microscopic Colitis: The European data.

Davide Giuseppe Ribaldone, MD¹, Daniele Simondi, MD², Marco Astegiano, MD¹, and Rinaldo Pellicano, MD¹

¹ Department of General and Specialistic Medicine, Gastroenterologia-U, Città della Salute e della Scienza di Torino, C.so Bramante 88, 10126 Turin, Italy

² Gastroenterology, Santa Croce e Carle of Cuneo Hospital, Via Coppino 26, 12100 Cuneo, Italy

Correspondence: Davide Giuseppe Ribaldone, Department of General and Specialistic Medicine, Gastroenterologia-U, Città della Salute e della Scienza di Torino, C.so Bramante 88, 10126 Turin, Italy; tel (0039)0116335208, fax (0039)0116336752, davrib_1998@yahoo.com

Conflicts of Interest and Source of Funding: none declared.

Keywords: microscopic colitis; Helicobacter pylori; inflammatory bowel diseases; infections

To The Editors,

in a recent paper, Sonnenberg and Genta¹ have reported an inverse association (OR = 0.6, 95% CI = 0.52-0.7) between *Helicobacter pylori* (*H. pylori*) infection and the occurrence of microscopic colitis (MC), in a United States' study.

Since these data could be influenced by the epidemiology of *H. pylori* infection, we performed a single-center study in Northern Italy, an area with a higher prevalence of *H. pylori* infection (only 10.2% of prevalence in the United States' paper¹).

The inclusion criteria were: a diagnosis of lymphocytic or collagenous colitis performed by standard international criteria;² *H. pylori* infection assessed by a direct method.³

The exclusion criteria were: a diagnosis of celiac disease;⁴ a suspect of inflammatory bowel disease;⁵ a diagnosis of *H. pylori* infection preceding that of MC but lack of data about the eradication treatment before the diagnosis of MC.

We included 50 patients, 36 women (72%) and 14 men (28%), median age 51 years, affected by MC and in whom *H. pylori* status was assessed by histology (n=47) or by 13C Urea Breath Test (UBT) (n=3). Of this cohort, 40 patients were affected by lymphocytic colitis and 10 by collagenous colitis. *H. pylori* resulted positive in 18 patients (36%), of whom 13 affected by lymphocytic colitis and 5 affected by collagenous colitis; the difference was not statistically significant (Chi-squared test, P = 0.51). These data were compared with a control population of 404 subjects suffering from constipation, who underwent UBT for study purposes. This cohort included 224 women (55.4%) and 180 men (44.6%) (statistically significant difference with cases, P = 0.038), with median age of 55 years (not statistically significant different, Mann-Whitney test, P = 0.4).

H. pylori infection was reported in 166 controls (41.1%), without difference with cases (Chi-squared test, P = 0.59).

Of the 18 patients with *H. pylori* positive status in the MC populations, in 8 (44%) the *H. pylori* infection was eradicated before the diagnosis of MC. Thus, at the diagnosis of MC

only 10 patients (20%) were *H. pylori* positive. Considering patients with history of *H. pylori* positivity but with eradication before the diagnosis of MC as patients with negative *H. pylori* status, there was a statistically significant difference between cases and controls (Chi-squared test, $P = 0.006$).

In conclusion, to be *H. pylori*-negative *ab initio* or after antibiotic treatment seems to be a risk factor for the onset of MC.

REFERENCES

1. Sonnenberg A, Genta RM. Inverse Association Between Helicobacter pylori Gastritis and Microscopic Colitis. *Inflamm Bowel Dis*. 2015. [Epub ahead of print]
2. Fenoglio-Preiser CM, Noffsinger AE. *Gastrointestinal Pathology: An Atlas and Text*. 3rd ed. Philadelphia, PA: Wolters Kluwer; 2007:843–847.
3. Working Party of the European Helicobacter pylori Study Group. Guidelines for clinical trials in Helicobacter pylori infection. *Gut*. 1997;41:S1-S23.
4. Simondi D, Ribaldone DG, Bonagura GA, et al. Helicobacter pylori in celiac disease and in duodenal intraepithelial lymphocytosis: Active protagonist or innocent bystander? *Clin Res Hepatol Gastroenterol*. 2015. pii: S2210-7401. [Epub ahead of print]
5. Sonnenberg A, Melton SD, Genta RM. Frequent occurrence of gastritis and duodenitis in patients with inflammatory bowel disease. *Inflamm Bowel Dis*. 2011;17:39-44.