

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Arthroscopic Bankart Repair Versus Open Bristow-Latarjet for Shoulder Instability: A Matched-Pair Multicenter Study Focused on Return to Sport

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1633171> since 2017-05-12T09:58:47Z

Published version:

DOI:10.1177/0363546516658037

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Title: return to sport after surgery for recurrent anterior shoulder instability. Arthroscopic**
2 **stabilization using anchors versus open Bristow-Latarjet: a pair-matched multicenter study.**

3
4 **Abstract**

5 **Background:** Recurrent shoulder instability is a common shoulder pathology especially in athletes. Among
6 the available techniques, the arthroscopic Bankart and open Bristow-Latarjet procedures have been most
7 commonly used to treat recurrent instability.

8 **Hypothesis/Purpose:** The aim of this multicenter study was to compare in a case-control matched manner
9 the two techniques with particular emphasis on the return to sport after surgery.

10 **Study Design:** retrospective study.

11 **Methods:** A study was conducted in two hospitals in patients with post traumatic recurrent anterior
12 dislocations with a minimum follow-up of two years. Patients with glenoid bone loss were excluded. In one
13 hospital, patients were treated with arthroscopic Bankart repair using anchors, while patients in the other had
14 the Bristow-Latarjet procedure. The patients included in the study were matched according to age at surgery,
15 type and level of sport (DOSIS score), and number of dislocations. The primary outcomes were: rate of
16 recurrent instability, return to sport (SPORTS score), Oxford shoulder score for instability (OSSI),
17 Subjective Shoulder Value (SSV), WOSI score and range of motion (ROM). The Student t-test and the
18 Fisher test were used to compare parametric and not parametric results. A multiple regression analysis was
19 also performed to search for factors affecting the ability to return to sport.

20 **Results:** The patients that underwent arthroscopic Bankart repair obtained better results in terms of return to
21 sport (SPORTS score: 8 vs. 6, $p = 0.02$) and ROM in the throwing position (86° vs. 79° , $p = 0.01$). The
22 subjective perception of the shoulder was also better in the Bankart group (SSV= 86% vs. 75%, $p = 0.02$).
23 No differences were detectable using the OSSI and WOSI score. The rate of recurrent instability was not
24 statistically different in the two groups (Bankart 11% vs. Bristow-Latarjet 0%, $p = 0.25$). The multiple

25 regression analysis showed that the independent variables statistically related to the return to a sport were:
26 preoperative DOSIS score, type of surgery and recurrent dislocations after surgery.

27 **Conclusion:** In a cohort of patients representative of our population, the arthroscopic stabilization using
28 anchors, provided better results compared to the open Bristow-Latarjet procedure.

29 **Key Terms:** recurrent shoulder instability, arthroscopic Bankart repair, open Bristow-Latarjet procedure,
30 DOSIS score, return to sport.

31 **What is known about this subject:** Several techniques have been proposed for the treatment of anterior
32 shoulder instability and the two most commonly used techniques are the Bristow-Latarjet technique and the
33 Bankart arthroscopic stabilization. There are very few works comparing these two techniques: they are
34 typically retrospective studies focused on a single technique rather than being based on direct comparison
35 between the two techniques. In only one recent study the transfer of the coracoid was compared with the
36 arthroscopic Bankart repair with slight better outcomes after coracoid transfer.

37 **What this study adds to existing knowledge:** While the literature commonly focuses on outcomes such as
38 pain, range of motion and rate of recurrent instability, we put particular emphasis on the return to sport after
39 surgery. In order to optimize the pair-matched process, a modified Tegner score for upper extremity, called
40 the Degree of Shoulder Involvement in Sport (DOSIS), was developed. The arthroscopic stabilization seems
41 to provide better return to sport, better external rotation in the throwing position, and better subjective
42 perception of the affected shoulder compared to the open Bristow-Latarjet procedure.

43

44 **Introduction**

45 Anterior shoulder instability can be a disabling condition especially when it affects active patients. If
46 conservative treatment fails and an active patient is not able to return to their sport, surgical treatment is
47 usually indicated^{6, 7, 18, 21}.

48 Several techniques have been proposed for the treatment of anterior shoulder instability. Among the
49 available techniques, the two most commonly used are the transposition of the coracoid (Bristow-Latarjet
50 technique)¹¹ and the arthroscopic stabilization by tensioning of the capsule and repair of the labral lesion⁸.

51 Evidence-based data exists supporting the efficacy of both of these procedures^{2, 7, 9, 11, 15, 18, 23}, and the
52 surgeon's personal preference plays a significant role in the choice of one of these techniques over the other.
53 Supporters of the transposition of the coracoid justify their choice based mainly on a lower recurrence rate
54 and a better return to the patients' pre-injury sport activity levels especially if the patient participated in
55 collision sports^{2, 7, 11, 15, 18}. On the other hand, the arthroscopic stabilization strategy restores the anatomy of
56 the shoulder, preserves the range of motion and, with modern instruments, is as effective as the transfer of
57 the coracoid with respect to its recurrence rate^{9, 23}. However, most of these opinions have been deduced from
58 retrospective studies focused on a single technique rather than being based on direct comparison between the
59 two techniques.

60 Very few studies have compared the open Bristow-Latarjet procedure to the arthroscopic stabilization
61 procedure^{15, 24}. Hovelius, *et al*, reported in their study including 185 shoulders that the Bristow-Latarjet
62 procedure had a lower recurrence rate and better subjective-based outcomes compared to the open Bankart
63 repair¹⁵. However, Petrera, *et al*, reported that modern arthroscopic stabilizations have led to better outcomes
64 compared to open Bankart repair¹⁹. In only one recent study² the transfer of the coracoid was compared with
65 the arthroscopic Bankart repair with slight better outcomes after coracoid transfer.

66 The relative paucity of studies directly comparing these two procedures can be attributed to several factors.
67 Any single surgeon usually has more experience with the one technique that they tend to use for the majority
68 of their cases. This can result in an unintended technical bias toward their favored technique.

69 Another significant challenge in conducting comparative studies is the extreme variability of patients
70 undergoing surgery for shoulder instability. It is clear that a high-level rugby player cannot be compared with
71 a sedentary patient in this context. This confounding factor could be controlled through a ranking of sports
72 activity into groups with similar involvement of the upper extremity. The Tegner activity scale²² is a score
73 designed with this specific aim, however, it has been developed for ligamentous injuries of the knee.
74 Unfortunately a similar score has not been developed for the shoulder.

75 Despite these difficulties, a comparison of the two techniques remains important for its potential benefit in
76 the decision making process for the treatment of recurrent anterior shoulder instability.

77 The aim of our study was to compare, in a pair-matched manner, the return to sport after open Bristow-
78 Latarjet versus arthroscopic shoulder stabilization. In order to optimize the pair-matched process, a modified
79 Tegner score for upper extremity was developed. This score was called the Degree of Shoulder Involvement
80 in Sport (DOSIS) score and was tested and then applied in the same cohort of patients.

81

82 **Material and Methods**

83 This study was organized in two phases. The first was a basic research phase to develop and test the DOSIS
84 score. This part was essential for the second clinical phase that was a retrospective multicenter pair-matched
85 study, designed to compare the arthroscopic shoulder stabilization procedure using suture anchors to the
86 open Bristow-Latarjet procedure.

87

88 **First phase: design of the Degree Of Shoulder Involvement In Sport score (DOSIS score)**

89 The DOSIS score was developed by the SPORT committee of the SIGASCOT (Società Italiana di
90 Ginocchio, Artroscopia, Sport, Cartilagine e Tecnologie Ortopediche) to score the sport activity based on
91 three parameters: 1) *the type of sport* classified as: a) no/minimal demand, b) moderate demand, c) high
92 demand; 2) *the frequency* the sport was played: a) occasionally, b) at least twice a week; 3) *the level* at which
93 the sport was played: a) recreational, b) low level of competition, c) high level of competition (Table I,
94 Figure 1). Using these parameters the DOSIS score is calculated by the researcher using an allocation table
95 (Appendix 1).

96 During the construction of the DOSIS score it was clear that the parameter “*type of sport*” was the parameter
97 more predisposed to disagreement. To reduce potential disagreement the ranking of the sports followed a 2-
98 step process.

99 In the first step a questionnaire including a list of 20 common sports and specific positions/roles (e.g. goal
100 keeper in soccer) was sent by email to 7 international experts on the shoulder and elbow (from USA, Italy,
101 Germany, Switzerland, Belgium, Spain, and Turkey). This questionnaire included three questions for each
102 sport that assessed three factors: a) whether or not the sport requires full or almost full range of motion
103 (ROM), b) whether or not the sport is an overhead sport, c) the amount of stress and contact forces to the
104 upper extremity while playing that sport.

105 Two of the experts suggested that some sports required a significantly different involvement between the
106 dominant and the non-dominant arm. In a second step, based on their suggestions, a new questionnaire that

107 separated the dominant and non-dominant arms for some specific sports or positions/roles were made. In
108 addition, a question regarding the degree of expertise in the sport of the respondent was added.

109 This questionnaire was sent by email to the members of the SIGASCOT. This society includes members with
110 different backgrounds but with a comparable expertise in sport related trauma. Using the answers to the
111 questionnaire, the sports were initially classified into either 3 categories (no/minimal demand, moderate
112 demand, high demand, Table II) or 5 categories (no/minimal demand, low demand, moderate demand, high
113 demand, extremely high demand). The classification of a sport into a category was done by measuring the
114 amount of points given for each characteristic (see Appendix 2). In summary, the category of “high demand
115 sport” was defined as a sport that required full range of motion, that included an overhead gesture, and that
116 involved a high stress or contact force on the upper extremity (e.g., water polo, baseball pitcher). The
117 decision to use a 3-group or 5-group classification was made after measuring the inter-observer reliability.

118 Using the classification of sports table (Table II), the frequency the sport was played and the level at which
119 the sport was played, the DOSIS score was then calculated using the allocation table.

120 In a subsequent phase the intraobserver reliability of the DOSIS score was tested in a subgroup of 41 patients
121 during their follow up examination after Bristow-Latarjet procedure. These patients belonged to the cohort
122 used for the clinical study. Three weeks before the scheduled appointment an envelope was sent to them
123 containing the DOSIS score questionnaire (Table I). The patients were instructed to answer the questions for
124 the DOSIS score, the same day or the day after they had received the envelope. Two weeks before the
125 planned visit they were re-contacted by phone in order to be sure that they had received the questionnaire
126 and completed it. During the subsequent appointment in the outpatient clinic, the patients were then
127 interviewed and asked to complete again the DOSIS score questionnaire. The answers to the DOSIS score
128 questionnaire were used to calculate the DOSIS score and to measure reliability.

129 **Statistical analysis**

130 The interobserver reliability of the classification of the sports into three categories and into five categories
131 was assessed by calculating the Fleiss' kappa¹² - a statistical measure for assessing the reliability of
132 agreement between a fixed number of raters. A limit of agreement of 0.7 for the experts was predetermined

133 as the limit to accept the classification. If both the classifications had reached the level of 0.7 of agreement,
134 the 5-group classification would have been used.

135 The intraobserver reliability of the DOSIS score completed at home was compared to the scores completed
136 in the clinic. The data were analyzed using Bland and Altman analysis³.

137

138 **Second phase (pair-matched retrospective study)**

139 The second phase was a retrospective multicenter study conducted in two university hospitals. The two
140 hospitals are located in the same region and, during the period of the study, each hospital had one shoulder
141 unit staffed by two expert surgeons and one or two surgeons in training. The principal investigator (initials)
142 was a member of the shoulder unit that used the Bristow-Latarjet procedure in the majority of the cases.
143 However, he participated in the data analysis only and not the data collection. The follow up was performed
144 by two researchers not involved in the care of the patients.

145 This study was made possible by the fact that the two shoulder units have had similar surgical indications for
146 recurrent anterior shoulder instability but different approaches with respect to the techniques used (Figure 2).

147 The inclusion criteria were: 1) post traumatic recurrent anterior dislocation with a minimum of two episodes
148 of documented dislocations, 2) minimum follow-up of two years, 3) patients older than 18 years old.

149 Cases for which both shoulder units would use the same treatment were excluded. These included: a) patients
150 treated with a coracoid transfer for revision surgery or in cases of 20% or greater glenoid surface bone loss
151 and b) patients treated with arthroscopic shoulder stabilization for recurrent anterior shoulder instability in
152 patients younger than 25 year old not involved in collision sport with less than 5 episodes of dislocations.
153 Patients with other concomitant pathologies (cuff tear, SLAP lesion) were also excluded.

154 A review of the data of the patients that underwent surgery from April 2004 to December 2010 was
155 performed. The selected patients were then contacted by phone and asked to participate in this study. If they
156 agreed an appointment was scheduled.

157 During the follow up examination the patients were evaluated using the following scores: the SPORTS score
158 questionnaire^{4, 5}, the Oxford Instability Shoulder Score (OISS)¹⁰, the Subjective Shoulder Value (SSV)¹³, the
159 Western Ontario Shoulders Instability Index (WOSI) score¹⁷ and the DOSIS score.

160 The SPORTS score was the primary outcome of this study. This is a score designed to address the ability to
161 resume the pre-injury sport level. The available scores range from 0 points (unable to resume the same sport)
162 to 10 points (able perform/resume the same sport at the same level of effort and performance as before onset
163 of impairment without pain). Patients who did not resume their sport for reasons other than the shoulder were
164 asked to answer considering the best level reached before leaving their sport.

165 The patients were also asked to report any episode of recurrent shoulder dislocation or subluxation.
166 Moreover the active ROM in external rotation with elbow at the side (ER1) and at 90° of abduction (ER2)
167 was measured with a handheld goniometer.

168

169 **Surgical techniques and post-operative rehabilitation**

170 The arthroscopic stabilization including Bankart repair and re-tensioning of the anterior capsule was
171 performed in lateral decubitus using the classic three portals⁸. Different anchors were used over the course of
172 the study period. The type, number of anchors, length of surgery and perioperative complications were
173 recorded. Postoperatively the patients were placed in a standard sling for 4 weeks. From 5 to 8 weeks
174 patients were allowed to perform passive and active motion up to 0° of external rotation and 90° of
175 abduction. Starting from 9 weeks post-op, full ROM was allowed. Non-collision sports were allowed starting
176 from 3 to 5 months post-op. Collision sports were permitted after 6 months.

177 The Bristow-Latarjet procedure was performed similarly to the technique described by Walch G.¹¹ Unlike
178 the original technique, a single 4.5mm malleolar screw without washer was used in all the cases to fix the
179 coracoid to the glenoid. A drain was always kept in place for approximately 24 hours. The length of surgery
180 and perioperative complications were recorded for comparison with the arthroscopic stabilization. After
181 surgery, the patients were placed in a standard sling for 3 weeks. After 3 weeks the sling was removed and

182 the patients were allowed progressive full range of motion. Non-collision sports were allowed starting from 2
183 months post-op. Collision sports were permitted after 6 months postoperatively.

184 **Statistical analysis**

185 Patients that underwent arthroscopic repair and Bristow-Latarjet procedure were pair-matched according to
186 age (≤ 22 and >22 year old), the number of episodes of anterior shoulder dislocations before surgery (≤ 10 , $>$
187 10 episodes), and the level of sport activity before the onset of shoulder instability (DOSIS ± 0).

188 The differences between continuous and non-continuous variables were tested using independent samples
189 Student t-Test and the Fisher test for the unmatched cohort of patients. The differences between pair-
190 matched patients were tested using paired Student t-Test and the McNemar test for continuous and non-
191 continuous variables respectively.

192 A multiple regression analysis was performed to measure correlation between return to sport and recurrent
193 instability after surgery and the following independent variables: age at surgery, involvement of the
194 dominant arm, number of dislocations before surgery, DOSIS score, and recurrent dislocation. The number
195 of anchors was not analyzed since in all but a few cases three anchors were consistently used. P-values of
196 ≤ 0.05 were considered to be significant for all analyses.

197

198 **Results**

199 A total of 131 surgeries were performed during the study period (62 arthroscopic stabilizations and 60
200 Bristow-Latarjet procedures). Of these 131 patients, 36 were met one or more of the exclusion criteria and
201 were not included in the study: 10 patients had a multidirectional instability, 9 a common indication for
202 arthroscopic procedure, 3 had a concomitant SLAP lesion and 3 had surgery after a primary dislocation. An
203 additional 11 cases of coracoid transfer were excluded because they were performed in patients with glenoid
204 bone loss of greater than 20%, or because they were performed as a revision surgeries. This left 95 patients
205 for inclusion. Of these, 90 were traceable and 85 consented to participate in the study and had a follow-up
206 examination. The 5 patients that declined to participate reported no recurrent shoulder instability. Their data,
207 however, were not included in this study. The demographic data and clinical results of the unmatched cohort
208 of patients are reported in Tables III and IV.

209

210 **DOSIS score**

211 One hundred and twenty five members of SIGASCOT replied to the online questionnaire. However, 15
212 questionnaires were incomplete and were excluded from the analysis. The remaining 110 questionnaires
213 were filled by the SIGASCOT members including 57 whom considered themselves to be experts in shoulder
214 pathology. The classification of the sports into five categories had a moderate inter-rater reliability among
215 the shoulder pathology experts ($k = 0.51$) and non-experts ($k = 0.45$). The classification of the sports into
216 three subgroups increased the reliability of the classification for both experts ($k = 0.71$) and non-expert raters
217 ($k = 0.63$). Therefore, the classification of the sports into three subgroups was adopted in the definitive
218 version of the DOSIS score and use for the analysis of intra-observer reliability and for the clinical part of
219 this study.

220 The DOSIS score had excellent intra-observer reliability. The systematic error between the first and the
221 second assessment of the DOSIS score was 0 points with a 95% upper limit of agreement of 0.88 points. Of
222 the 41 patients, 5 changed their answers when they completed the DOSIS score for the second time. Of these,

223 1 patient changed the frequency he practiced the sport, 1 patient changed the type of sport, 1 patient changed
224 from “no sports” to “occasional jogging” and 2 patients changed their level of competition.

225

226 **Open Bristow-Latarjet vs. Arthroscopic Stabilization**

227 In the un-matched cohort of patients (average follow-up of $\approx 5 \frac{1}{2}$ years) the recurrence rate was higher in the
228 patients that underwent arthroscopic stabilization (4 dislocations in the arthroscopic stabilization group (9%)
229 vs. 0 dislocations in the Bristow-Latarjet group) but the difference was not statistically significantly ($p =$
230 0.12, Table IV). Of the 4 patients that had recurrent dislocation 2 had revision surgeries: 1 with a coracoid
231 transfer and 1 with an arthroscopic stabilization. None of the patients reported subluxation after surgery.

232 Despite the higher recurrence rate seen with the arthroscopic stabilization patients, this group had a better
233 return to sport (SPORT score 8 versus 6, $p = 0.09$) and a better subjective opinion of their operated shoulder
234 (SSV = 84% vs. 74%, $p = 0.004$). No differences were found in the active external rotation.

235 No acute complications were reported in the arthroscopic group while one patient in the Bristow-Latarjet
236 group experienced a postoperative hematoma that resolved spontaneously. An average of 3 bio-absorbable
237 anchors were used in the patients that underwent an arthroscopic stabilization (range 2 to 4). All the anchors
238 were loaded with high-strength sutures (Orthocord®, Fiberwire®, Tigerwire®, Tigertail®). Three types of
239 bio-absorbable sutures were used: LUPINE™ (DePuy-Synthes Raynham, Massachusetts, USA), Bio-
240 SutureTak™ (Arhtrex, Naples, Florida, USA), Bio-FASTak® (Arhtrex, Naples, Florida, USA).

241

242 **Pair-matched analysis**

243 Fifty-six patients were successfully pair-matched. The demographic data and results are reported in Table III
244 and IV. Age, number of preoperative anterior dislocations and DOSIS scores were similar in the two groups
245 confirming the efficacy of the matching process. The average length of surgery was 96 minutes (range 35 to
246 210 minutes) in the arthroscopic stabilization group and 71 minutes (range 50-120 minutes) in the Bristow-
247 Latarjet group ($p = 0.001$).

248 The pair-matched analysis confirmed the better clinical results seen with arthroscopic stabilization with
249 respect to the return to sport and SSV. The external rotation in abduction was also better after arthroscopic
250 treatment (Table IV). Three patients in the arthroscopic stabilization group (11%) experienced a recurrent
251 dislocation. Three anchors were used in the patients that had recurrent dislocations. The patients with
252 recurrent instability had significantly more preoperative dislocations (average dislocation 44 vs. 11, $p =$
253 0.02). Among the patients that underwent arthroscopic stabilization, the patients with recurrent dislocations,
254 had significantly worse SPORTS scores (5 vs. 9, $p= 0.06$), SSVs (67 vs. 88, $p= 0.003$) and WOSI scores
255 (64% vs. 86%, $p= 0.03$).

256 Two patients in each pair-matched group (7%) had a SPORTS score of 10 points even though they did not
257 play the same sport at the same level at the time of the follow-up. They were interviewed to investigate the
258 reason of these inconsistent results. It was determined that they quit their sport or changed level (from
259 competitive to recreational) for reasons not associate with their shoulder. Five patients in the Bristow-
260 Latarjet group (18%) and three patients in the arthroscopic stabilization group (11%) were not able to return
261 to sport (SPORTS score 0) ($p = 0.7$).

262 The multiple regression analysis showed that the independent variables statistically related to the return to a
263 sport were: preoperative DOSIS score, type of surgery and recurrent dislocations after surgery. Although
264 external rotation at 90° was not statistically related to the return to sport, its p-value approached significance
265 ($p < 0.08$) (Table V). Figure 3 illustrates the relationship between the degree of shoulder involvement in
266 sport (DOSIS) and the ability to the return to sport after surgery. It seems that the higher the shoulder
267 involvement was in the sport before surgery the lower the ability was to resume it (Figure 3B). The type of
268 surgery was the only independent variable related to recurrent instability after surgery (Bristow-Latarjet,
269 correlation coefficient = -0,13, $t = -2.016$, $p = 0.047$).

270

271 **Discussion**

272 This pair-matched retrospective study has been designed to compare two of the most prevalent techniques for
273 the treatment of recurrent posttraumatic anterior shoulder instability. The principal outcome that we used was
274 the return to sport. The return to sport is usually the first goal of patients seeking treatment for shoulder
275 instability. Warth *et al* in a study of 313 shoulders treated by an arthroscopic procedure, found that the most
276 important expectation was to continue participation in sporting activities²⁵. Interestingly, the patients that
277 sought treatment for shoulder instability valued their continued participation in sports more than stopping
278 their shoulder from dislocating.

279 With respect to the SPORTS score, the patients that underwent arthroscopic stabilization showed a better
280 return to sport compared to those with the open Bristow-Latarjet procedure even though both the techniques
281 provided an high rate of return to sport. More than 80% of the patients returned to their sport for both repair
282 techniques. However, the level at which they returned to sport was significantly in favor of arthroscopic
283 stabilization. Interestingly, the difference in SPORTS score was higher in the middle range of the DOSIS
284 scores and tended to disappear for sports with a high degree of shoulder involvement.

285 Patients that played sports with high upper extremity involvement (e.g. swimming, rugby, martial arts) at
286 competitive level (DOSIS score 9 or 10) have a lower level of return to sport with both repair techniques
287 (Figure 3). The ability to resume a sport after arthroscopic stabilization seemed, in our series, to be more
288 dependent on the type of sport rather than the use of the Bristow-Latarjet technique. A study by Kim, *et al*.
289 showed a similar correlation between Bankart repair, level of activity/demand and return to previous activity
290 level¹⁶.

291 In contrast to our findings, the return to sport after coracoid transfer was better in studies focused on
292 competitive rugby and soccer players^{7, 18} in comparison to our results. The reasons for this could be a
293 technical issue or the fact that our population was more heterogeneous, including different sports practiced at
294 a recreational level. It is conceivable that recreational athletes had invested less time and effort into their
295 post-operative rehabilitation and had also less motivation to resume their sport compared to professional
296 athletes.

297 Return to sports after surgery for shoulder instability depends on several factors, including perceived
298 shoulder stability while the sport is being played. Other factors, however, such as the recovery of ROM,
299 proprioception, and the type of sport, are equally important. Some studies have already demonstrated that
300 after open and arthroscopic repair the joint position sense improved significantly, to the level of a normal,
301 healthy shoulder^{20,27}.

302 Regarding ROM, we found a significant loss of external rotation in the throwing position in patients after
303 coracoid transfer. This finding could contribute to the difference in SPORTS score between the surgical
304 techniques in our study. The multivariate regression analysis seemed to support this hypothesis. A recent
305 systematic review paper has describes similar results regarding external rotation after the Bristow-Latarjet
306 procedure¹⁴.

307 The different levels of return to sport between arthroscopic stabilization and open Bristow-Latarjet procedure
308 is not corroborated by previously published studies in the literature. For this reason our outcome has to be
309 considered with great caution. There is only one study available in the literature that compared arthroscopic
310 Bankart repair and coracoid transfer with similar return to sport results². One possible reason for this
311 discrepancy between our return to sport findings and those previously reported in other studies is that the
312 other studies did not use a dedicated score to measure the return to sport.

313 In our study a new score (DOSIS score) has been used to reach this outcome while other well-known scores
314 such as the Oxford score for instability and the WOSI score did not show significant differences. However,
315 in support of the better return to sport for the arthroscopic technique, we also found a better subjective
316 perception of the operated shoulder compared to a normal shoulder. The SSV in fact was 86% in the
317 arthroscopic group and 75% in the Bristow-Latarjet group ($p = 0.02$). A combination of the limitation in
318 external rotation, different return to sport outcome, as well as cosmetic reasons could have contributed to the
319 difference in SSVs.

320 Similar limitations in the use of specific scores for shoulder instability have been already reported in the
321 recent literature. Netto, *et al.*¹, comparing arthroscopic and open techniques for the treatment of Bankart
322 lesions found no differences when using established scores for instability (ROWE and UCLA scores).

323 Remarkably, they found differences in scores when using the Disabilities of the Arm, Shoulder and Hand
324 (DASH) questionnaire -- a score not specifically designed for shoulder instability. A recent study by Stein, *et*
325 *al.*²¹, has been focused on shoulder sport-specific impairments after arthroscopic Bankart repair. They
326 concluded sport-specific impairments are not detectable by established clinical score systems. In this
327 scenario, therefore, it seems reasonable to search for new tools for a more complete assessment of outcomes
328 after surgery for shoulder instability.

329 The DOSIS score was developed with the specific aim to match the patients according to the level of sport
330 before shoulder impairment. Without this tool, in an unmatched population, the differences between
331 arthroscopic stabilization and coracoid transfer were not significant and similar to the outcomes reported by
332 Bessier, *et al.*².

333 In this study, both repair techniques were able to restore stability in the majority of the patients. The rate of
334 recurrent instability after surgery was 9% in the arthroscopic group and 0% in the Bristow-Latarjet group.
335 Similar rates were reported in the literature^{1, 7, 16}. However, the rate of 0% in the Bristow-Latarjet group
336 deserves some further discussion. The patients with glenoid loss higher than 20% were excluded from this
337 study as well as patients with prior surgery. Moreover the average age of patients was 33 years old and
338 adolescent athletes were absent. This could easily explain this low rate of recurrent instability and the low
339 rate of complications²⁶.

340 Even if the different in rate of recurrent instability between arthroscopic stabilization and Bristow-Latarjet
341 procedure was not statistically significant, there was a tendency toward a better rate of recurrent instability in
342 the Bristow-Latarjet group. The failure to achieve statistical significance is likely due to the small numbers
343 of patients included with the consequent under-powering of the study. This trend was confirmed by the
344 regression analysis that showed a correlation between technique and rate of recurrent instability after
345 surgery.

346 Overall we observed a mismatch between subjective clinical outcomes that was in favor of the arthroscopic
347 procedure and rate of recurrent instability that favored the Bristow-Latarjet procedure. Although a definite
348 explanation for this is not available, it is possible that having only 3 patients with recurrent instability,

349 resulted in a sample size that was too small to negatively affect the overall outcomes after the arthroscopic
350 stabilization.

351 This study has some limitations. It is a retrospective study based on new score. This score, however, was
352 designed carefully and tested in the same cohort of patients where it showed good psychometric features. Not
353 all the psychometric features of the DOSIS score were analyzed and other studies are needed to further
354 investigate this.

355 Another limitation is the minimum follow up of 2 years. More dislocations could be observed with longer
356 follow-up. This study, however, was focused on the return to sport that generally does not require a lengthy
357 follow-up to be assessed. Moreover, Griesser, *et al.*, has reported that more than 70% of dislocations occur
358 within one year of the Bristow-Latarjet procedure¹⁴. Another limitation is that we did not assess the presence
359 and extent of Hill-Sachs lesions. This data was not reported in a reliable way in the documentation of most of
360 the patients treated before 2007.

361 The strength of this study is that it is a matched-pair study where the patients were carefully matched
362 according to age, degree of shoulder involvement in sport and number of prior shoulder dislocations.

363

364 **Conclusions**

365 In a cohort of patients representative of our population, without significant glenoid erosion, the arthroscopic
366 stabilization using anchors, provided better return to sport, better external rotation in the throwing position,
367 and better subjective perception of the affected shoulder compared to the open Bristow-Latarjet procedure.
368 The difference in return to sport disappeared when the patients practiced sports with high involvement of the
369 shoulder and at competitive level. If the patient is greatly concerned about recurrent dislocation the open
370 Bristow-Latarjet procedure is preferred.

371

372 **Table legends**

373 **Table I:** DOSIS score

374 *: For seasonal sports considered the frequency during the season

375 The DOSIS score can be completed either with respect to the sport performed during the month before the
376 questionnaire is filled out, or with respect to the sport performed during the period before the onset of the
377 shoulder disease. In this study we used the DOSIS to investigate the status before the onset of the shoulder
378 instability.

379

380 **Table II:** classification of sports

381 The list of sports included in each group has to be considered an open list. Other sports not listed here can be
382 added using the instructions in the appendix.

383

384 **Table III:** demographic data

385 Demographic data of the un-matched and matched patients.

386

387 **Table IV:** outcomes

388 Bold values are statistically significant ($p < 0.05$). ER1: external rotation with the elbow at the side. ER2:
389 external rotation in the throwing position (90° of abduction).

390

391 **Table V:** Multiple regression analyses for return to sport

392 Bold values are statistically significant ($p < 0.05$).

393

394 **Figure legends**

395 **Figure 1:** design of the DOSIS score

396 The DOSIS score was developed by the SPORT committee of the SIGASCOT to score the sport activity
397 based on three parameters: 1) *the type of sport* (in blue) classified as: a) no/minimal demand, b) moderate
398 demand, c) high demand; 2) *the frequency* the sport was played (in orange) : a) occasionally, b) at least twice
399 a week; 3) *the level* at which the sport was played (in yellow): a) recreational, b) low level of competition, c)
400 high level of competition.

401 The classification of sports in groups with similar involvement of the upper extremity was done taking into
402 account whether or not the sport requires full or almost full range of motion (ROM), b) whether or not the
403 sport is an overhead sport, c) the amount of stress and contact forces to the upper extremity while playing
404 that sport. To address this issue an online questionnaire was sent to the members of the SIGASCOT.

405

406 **Figure 2:** inclusion and exclusion criteria

407

408 **Figure 3A&B:** relationship between DOSIS score and SPORT score.

409 The difference in SPORTS scores tends to disappear for higher DOSIS scores.

410

411 **Appendix legends**

412 **Appendix 1:** Allocation table

413 Using this table the researcher scores patient according to type of sport, frequency in which the sport is
414 played, and level of the sport. For example an occasional tennis player (high demand sport-Table II), with
415 involvement of the dominant arm, is assigned a DOSIS score of 6 (in gray).

416

417 **Appendix 2:** Classification of sports based on the online questionnaire

418 According to the three characteristics of each sport (ROM, overhead gesture, amount of stress and contact
419 forces) the sports were ranked into three groups. This is an open list. New sports can be added and ranked in
420 one of three groups. The ranking will be defined by the total points given for each characteristic. For
421 example if a researcher wants to add windsurfing he/she has to define a)the ROM needed to perform
422 windsurfing (not full = 0 points), b) if windsurfing can be considered an overhead sport (no = 0 points),
423 amount of stress and contact forces (high = 2 points). The total points for windsurfing would therefore be 2
424 points = moderate demand. The researcher has to decide if the involvement of the shoulder is significantly
425 different between dominant and non-dominant arms. For windsurfing the involvement is approximately the
426 same, hence the distinction between dominant and non-dominant arms is not necessary.

427

428 **References**

- 429 1. Archetti Netto N, Tamaoki MJ, Lenza M, et al. Treatment of Bankart lesions in traumatic anterior
430 instability of the shoulder: a randomized controlled trial comparing arthroscopy and open techniques.
431 Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy
432 Association of North America and the International Arthroscopy Association. 2012;28(7):900-908.
- 433 2. Bessiere C, Trojani C, Pelegri C, Carles M, Boileau P. Coracoid bone block versus arthroscopic
434 Bankart repair: a comparative paired study with 5-year follow-up. Orthopaedics & traumatology,
435 surgery & research : OTSR. 2013;99(2):123-130.
- 436 3. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical
437 measurement. Lancet. 1986;1(8476):307-310.
- 438 4. Blonna D, Castoldi F, Delicio D, et al. Validity and reliability of the SPORTS score. Knee surgery,
439 sports traumatology, arthroscopy : official journal of the ESSKA. 2012;20(2):356-360.
- 440 5. Blonna D, Lee GC, O'Driscoll SW. Arthroscopic restoration of terminal elbow extension in high-
441 level athletes. The American journal of sports medicine. 2010;38(12):2509-2515.
- 442 6. Castagna A, Delle Rose G, Borroni M, et al. Arthroscopic stabilization of the shoulder in adolescent
443 athletes participating in overhead or contact sports. Arthroscopy : the journal of arthroscopic &
444 related surgery : official publication of the Arthroscopy Association of North America and the
445 International Arthroscopy Association. 2012;28(3):309-315.
- 446 7. Cerciello S, Edwards TB, Walch G. Chronic anterior glenohumeral instability in soccer players:
447 results for a series of 28 shoulders treated with the Latarjet procedure. Journal of orthopaedics and
448 traumatology : official journal of the Italian Society of Orthopaedics and Traumatology.
449 2012;13(4):197-202.
- 450 8. Cole BJ, Romeo AA. Arthroscopic shoulder stabilization with suture anchors: Technique,
451 technology, and pitfalls. Clinical orthopaedics and related research. 2001(390):17-30.
- 452 9. Cole BJ, Warner JJ. Arthroscopic versus open Bankart repair for traumatic anterior shoulder
453 instability. Clinics in sports medicine. 2000;19(1):19-48.

- 454 10. Dawson J, Fitzpatrick R, Carr A. The assessment of shoulder instability. The development and
455 validation of a questionnaire. *The Journal of bone and joint surgery. British volume.* 1999;81(3):420-
456 426.
- 457 11. Edwards TB, Walch G. The Latarjet procedure for recurrent anterior shoulder instability: Rationale
458 and technique. *Operative Techniques in Sports Medicine.* 2002;10(1):25-32.
- 459 12. Fleiss J. Measuring nominal scale agreement among many raters. *Psychological Bulletin.*
460 1997;75(6):378-382.
- 461 13. Gilbert MK, Gerber C. Comparison of the subjective shoulder value and the Constant score. *Journal*
462 *of shoulder and elbow surgery / American Shoulder and Elbow Surgeons ... [et al.].* 2007;16(6):717-
463 721.
- 464 14. Griesser MJ, Harris JD, McCoy BW, et al. Complications and re-operations after Bristow-Latarjet
465 shoulder stabilization: a systematic review. *Journal of shoulder and elbow surgery / American*
466 *Shoulder and Elbow Surgeons ... [et al.].* 2013;22(2):286-292.
- 467 15. Hovelius L, Vikerfors O, Olofsson A, Svensson O, Rahme H. Bristow-Latarjet and Bankart: a
468 comparative study of shoulder stabilization in 185 shoulders during a seventeen-year follow-up.
469 *Journal of shoulder and elbow surgery / American Shoulder and Elbow Surgeons ... [et al.].*
470 2011;20(7):1095-1101.
- 471 16. Kim SH, Ha KI. Bankart repair in traumatic anterior shoulder instability: open versus arthroscopic
472 technique. *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the*
473 *Arthroscopy Association of North America and the International Arthroscopy Association.*
474 2002;18(7):755-763.
- 475 17. Kirkley A, Griffin S, McLintock H, Ng L. The development and evaluation of a disease-specific
476 quality of life measurement tool for shoulder instability. *The Western Ontario Shoulder Instability*
477 *Index (WOSI). The American journal of sports medicine.* 1998;26(6):764-772.
- 478 18. Neyton L, Young A, Dawidziak B, et al. Surgical treatment of anterior instability in rugby union
479 players: clinical and radiographic results of the Latarjet-Patte procedure with minimum 5-year
480 follow-up. *Journal of shoulder and elbow surgery / American Shoulder and Elbow Surgeons ... [et*
481 *al.].* 2012;21(12):1721-1727.

- 482 19. Petrera M, Patella V, Patella S, Theodoropoulos J. A meta-analysis of open versus arthroscopic
483 Bankart repair using suture anchors. *Knee surgery, sports traumatology, arthroscopy : official journal*
484 *of the ESSKA*. 2010;18(12):1742-1747.
- 485 20. Potzl W, Thorwesten L, Gotze C, Garmann S, Steinbeck J. Proprioception of the shoulder joint after
486 surgical repair for Instability: a long-term follow-up study. *The American journal of sports medicine*.
487 2004;32(2):425-430.
- 488 21. Stein T, Linke RD, Buckup J, et al. Shoulder sport-specific impairments after arthroscopic Bankart
489 repair: a prospective longitudinal assessment. *The American journal of sports medicine*.
490 2011;39(11):2404-2414.
- 491 22. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clinical*
492 *orthopaedics and related research*. 1985(198):43-49.
- 493 23. Tjountakaris FP, Bradley JP. The rationale for an arthroscopic approach to shoulder stabilization.
494 *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy*
495 *Association of North America and the International Arthroscopy Association*. 2011;27(10):1422-
496 1433.
- 497 24. Vander Maren C, Geulette B, Lewalle J, et al. [Coracoid process abutment according to Latarjet
498 versus the Bankart operation. A comparative study of the results in 50 cases]. *Acta orthopaedica*
499 *Belgica*. 1993;59(2):147-155.
- 500 25. Warth RJ, Briggs KK, Dornan GJ, Horan MP, Millett PJ. Patient expectations before arthroscopic
501 shoulder surgery: correlation with patients' reasons for seeking treatment. *Journal of shoulder and*
502 *elbow surgery / American Shoulder and Elbow Surgeons ... [et al.]*. 2013.
- 503 26. Weber SC. The Latarjet procedure: the patient population makes all the difference: commentary on
504 an article by Anup A. Shah, MD, et al.: "Short-term complications of the Latarjet procedure". *The*
505 *Journal of bone and joint surgery. American volume*. 2012;94(6):e37.
- 506 27. Zuckerman JD, Gallagher MA, Cuomo F, Rokito A. The effect of instability and subsequent anterior
507 shoulder repair on proprioceptive ability. *Journal of shoulder and elbow surgery / American*
508 *Shoulder and Elbow Surgeons ... [et al.]*. 2003;12(2):105-109.

Table I

1) What sports did you play before the onset of your shoulder problem? List the sports below and indicate which was the most important/predominant for you:		
List of sports	Most important/predominant	
1)	yes	no
2)	yes	no
3)	yes	no
...	yes	no
2) How frequently did you participate in sports?*		
Occasionally		
≥ 2 times a week, most of the weeks of the year		
3) What level of sport did you play?		
Recreational		
Low level of competition (regional, local)		
High level of competition (national or international or professional)		
4) Which was your dominant arm during your sports activities?		

Table II

No-minimal involvement of the upper extremity	Moderate involvement of the upper extremity	High involvement of the upper extremity
Jogging	Soccer	Swimming
Street cycling	Bowling (dominant arm)	Martial art
Horseback riding	Nordic ski	Gymnastic
Bowling (non-dominant arm)	Rowing	Volleyball (dominant arm)
Mountain bike	Motocross	Tennis/squash (dominant arm)
Alpine ski	Golf	Baseball pitcher
Nordic Walking	Track and field (running and jumping)	Baseball (dominant arm)
Hiking	Track and field (throwing) (non-dominant arm)	American football quarterback
	Kayak	American football
	Dance	Rugby
	Basketball	Water polo
	Volleyball (non-dominant arm)	Track and field (throwing) (dominant arm)
	Tennis/squash (non-dominant arm)	Soccer, Goal keeper
	Baseball (non-dominant arm)	Rock Climbing

Table III

		N	Age	Sex (Male)	DOSIS	N of shoulder dislocations before surgery	Surgery on dominant shoulder	Collision Sport	Follow up (years)
Unmatched patients	Bankart	44	34 (20 to 53)	39 (89%)	6 (2 to10)	14 (2 to >50)	25 (57%)	20 (45%)	5.4 (2.9 to 9.8)
	Bristow-Latarjet	41	33 (19 to 63)	36 (88%)	5 (2 to10)	20 (2 to >50)	23 (56%)	22 (54%)	5.1 (2 to 8.7)
p			0.72	1	0.37	0.26	1	0.5	0.5
Matched population	Bankart	28	31.5 (20 to 53)	24 (86%)	6 (2 to10)	15 (2 to >50)	17 (61%)	17 (61%)	5.3 (2.9 to 9)
	Bristow-Latarjet	28	31.5 (19 to 45)	24 (86%)	6 (2 to10)	19 (2 to >50)	18 (64%)	15 (54%)	5.3(2 to8.7)
p			0.96	1	1	0.44	1	0.79	0.89

Table IV

		Oxford	SPORTS Score	WOSI					SSV	ER1 (°)	ER2 (°)	Recurrent dislocation	Revision surgery
				A	B	C	D	%					
Un-matched patients	Bankart N= 44	42	8	118 (0-480)	58 (0-290)	62 (0-240)	53 (0-270)	84% (44-100%)	84	57° (10°-75°)	84° (60°-105°)	4 (9%)	2 (5%)
	Bristow- Latarjet N= 41	40	6	166 (0-540)	73 (0-290)	80 (0-359)	49 (0-300)	82% (37-99%)	74	54 (30°-75°)	80° (30°-100°)	0 (0%)	0
	p	0.24	0.09	0.13	0.35	0.29	0.75	0.77	0.004	0.35	0.24	0.12	0.1
Matched patients	Bankart N= 28	41	8	107 (0-480)	51 (0-290)	51 (0-210)	41 (0-190)	84 (46-100%)	86	57° (10°-75°)	86° (70°-105°)	3 (11%)	2
	Bristow- Latarjet N= 28	40	6	169 (0-540)	83 (0-290)	85 (0-285)	50 (0-260)	82 (37-99%)	75	56° (30°-70°)	79° (50°-100°)	0 (0%)	0
	p	0.36	0.02	0.13	0.101	0.07	0.63	0.588	0.02	0.76	0.01	0.25	0.5

Table V

Independent variables for return to SPORT	Coefficient	Std. Error	t	P
Bristow-Latarjet	-2.74	0.82	-3.36	0.0012
DOSIS score	-0.44	0.15	-2.95	0.0042
Recurrent dislocations	-3.75	1.37	-2.73	0.008
ER2	0.06	0.03	1.80	0.08
Age	-0.07	0.05	-1.51	0.13
ER1	-0.04	0.04	-1.0	0.28
Dominant Arm	-0.74	0.81	-0.91	0.36
Number of preoperative dislocations	0.005	0.02	0.24	0.8

Excluded

Stabilization surgery, minimum follow-up of two years, >18 y, correct phone contact (**131 patients**)

9 arthroscopic Bankart repair patients:

- younger than 25 year old
- not involved in collision sport
- less than 5 episodes of dislocations

11 Bristow-Latarjet procedure patients:

- As a revision surgery
- $\geq 20\%$ glenoid surface bone loss

3 Patients with concomitant SLAP lesion, cuff tear

10 Patients with multidirectional instability

3 patients with surgery after a primary dislocation

95 patients


10 lost to follow up



85 patients (44 Bankart, 41 Bristow-Latarjet)

Matching process

Design and testing of the DOSIS score (Figure 2)

56 patients (28 Bankart, 28 Bristow-Latarjet)





Appendix 2

SPORTS	ROM Full/almost full		Overhead		Stress/Contact				Total	Classifications
	Yes (1)	No (0)	Yes (1)	No (0)	No-minimal (0)	Moderate (1)	High (2)	Extremely high (3)		
Jogging		*		*	*					No-Minimal Demand
Road cycling		*		*	*					
Horseback riding		*		*	*					
Bowling (non-dominant arm)		*		*	*					
Mountain bike		*		*		*				
Alpine ski		*		*		*				
Nordic Walking		*		*		*				
Hiking		*		*		*				
Soccer		*		*			*			Moderate Demand
Bowling (dominant arm)		*		*			*			
Nordic ski	*			*		*				
Rowing		*		*			*			
Motocross		*		*			*			
Golf	*			*		*				
Track and field (running and jumping)	*			*		*				
Track and field (throwing), (non-dominat arm)	*			*		*				
Kayak	*			*			*			
Dance	*			*			*			
Basketball	*			*			*			
Volleyball (non-dominant arm)	*		*			*				
Tennis/squash (non-dominant arm)	*		*			*				
Baseball (non-dominant arm)	*		*			*				
Swimming	*		*				*			
Rugby	*			*				*		
American football (non-dominant arm)	*			*				*		
Martial art	*			*				*		
Gymnastic	*			*				*		
Volleyball (dominant arm)	*		*				*			
Tennis/squash (dominant arm)	*		*				*			
Water polo (non-dominant arm)	*		*				*			
Baseball pitcher (non-dominant arm)	*		*				*			
American football quarterback (non-dominant arm)	*		*				*			
Soccer, goal keeper	*			*				*		
Rock Climbing	*			*				*		
Baseball (dominant arm)	*		*					*		
Water polo(dominant arm)	*		*					*		
Baseball pitcher (dominant arm)	*		*					*		
American football quarterback (dominant arm)	*		*					*		
Track and field (throwing), (dominant arm)	*		*					*		