

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Transmission of Grapevine virus A and Grapevine leafroll-associated viruses 1 and 3 by *Planococcus ficus* and *Planococcus citri* fed on mixed-infected plants

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1636400> since 2017-05-28T20:53:02Z

Published version:

DOI:10.1111/aab.12279

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Transmission of Grapevine virus A and Grapevine leafroll-associated viruses 1 and 3 by *Planococcus ficus* and *Pl. citri* fed on mixed-infected plants

| | |
|-------------------------------|---|
| Journal: | <i>Annals of Applied Biology</i> |
| Manuscript ID: | AAB-2015-0237 |
| Manuscript Type: | Research paper |
| Date Submitted by the Author: | 14-Aug-2015 |
| Complete List of Authors: | Bertin, Sabrina; Università degli Studi di Torino, DISAFA - Entomologia Pacifico, Davide; CNR, Istituto per la Protezione Sostenibile delle Piante Cavalieri, Vincenzo; Università degli Studi di Torino, DISAFA - Entomologia Marzachi, Cristina; CNR, Istituto per la Protezione Sostenibile delle Piante Bosco, Domenico; Università degli Studi di Torino, DISAFA - Entomologia |
| Key Words: | Entomology, Epidemiology, Plant-Virus-Vector Interactions, Virology |
| | |

1
2
3 **Transmission of *Grapevine virus A* and *Grapevine leafroll-associated***
4
5
6 ***viruses 1* and *3* by *Planococcus ficus* and *Pl. citri* fed on mixed-infected**
7
8
9 **plants**
10

11
12
13
14 **Sabrina Bertin¹, Davide Pacifico², Vincenzo Cavalieri¹, Cristina Marzachi², and**
15
16

17 **Domenico Bosco¹**
18

19
20 ¹ DISAFA – Entomologia, Università di Torino, Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy
21

22 ² CNR - IPSP, Strada delle Cacce 73, 10135 (TO), Italy
23
24

25 Corresponding author:

26 D. Bosco

27 DISAFA – Entomologia

28 Università di Torino

29 Largo Paolo Braccini, 2

30 10095 Grugliasco (TO) - Italy

31 Phone: 39-0116708529

32 Fax: 39-0116708535

33 E-mail: domenico.bosco@unito.it
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

The *Grapevine virus A* (GVA) and *Grapevine leafroll-associated viruses 1 and 3* (GLRaV-1 and GLRaV-3) are associated with grapevine diseases that induce severe reductions in yield and berry quality. These three viruses are known to coexist in both grapevine and insect vectors, but their co-transmission has been poorly characterized so far. This study investigates the acquisition and transmission of GLRaV-1, GLRaV-3 and GVA by *Planococcus ficus* and *Pl. citri* (Hemiptera: Pseudococcidae) following feeding on multiple-infected plants. The retention and load of each virus in insects were analysed, and a method for the quantification of GVA and GLRaVs in the vectors was set up for the first time. After feeding onto GVA, GLRaV-1 and GLRaV-3 mixed-infected grapevines, nymphs of *Pl. ficus* and *Pl. citri* showed similar virus acquisition rates and retained low quantities of viruses until the third post-acquisition day. Despite the similar acquisition patterns, the two vectors differed in transmission efficiency: *Pl. ficus* showed a higher efficiency in transmitting GVA and GLRaV-3, whereas *Pl. citri* transmitted more efficient in GLRaV-1. When focusing on the virus co-transmission, it appears that GVA could be transmitted to grapevine without GLRaV-1 and/or GLRaV-3 and that the GLRaVs transmission could take place in absence of GVA. This comparative study involving different viruses and vector species improves the current knowledge of the semi-persistent transmission of these three viruses and contribute to the understanding of the grapevine virus epidemiology.

Keywords

Mealybug, *Vitis vinifera*, Leafroll, Rugose wood.

Introduction

Grapevine (*Vitis vinifera* L.) is one of the most important fruit crop worldwide and represents a highly valuable commodity. The global grapevine production is threaten by many pests and pathogens, including viruses. Among the virus-induced diseases, the Grapevine Leafroll (GLD) and Rugose Wood (RW) are the most widespread and are responsible for severe reductions in yield and quality (Naidu *et al.*, 2014).

GLD and RW diseases are caused by positive single-stranded RNA viruses belonging to two distinct taxonomical groups and are frequently found in mixed infections. GLD is associated with a complex of virus species in the family *Closteroviridae* that are collectively referred to as *Grapevine leafroll-associated viruses* (GLRaV-1, -2, -3, -4, -7; Martelli *et al.*, 2012). RW includes different syndromes associated with virus species belonging to the family *Betaflexiviridae*, genus *Vitivirus* (*Grapevine virus A, B, D, E and F*; GVA, GVB, GVD, GVE and GVF) and *Foveavirus* (*Grapevine rupestris stem pitting associated virus*; GRSaV) (Martelli, 2014b). Both GLRaVs and RW-associated viruses are phloem-restricted and cause distinctive symptoms in grapevine, such as colour alteration and downward rolling of the leaves (GLD), stem grooving (RW), as well as delay in fruit ripening and alteration of final products (Martelli, 2014a; Martelli, 2014b).

Plant-to-plant transmission of GLRaVs and RW-associated viruses is mediated by phloem-sucking insects, mealybugs (Hemiptera: *Pseudococcidae*) and soft scales (Hemiptera: *Coccidae*). Compared to soft scales, mealybugs are likely to play a major role in virus spread because of their higher mobility and they are responsible for severe damages even at low infestation levels (Golino *et al.*, 2008; Cabaleiro *et al.*, 2008; Cabaleiro and Segura, 2006). Although all life stages of mealybugs are capable of transmission, first-instar nymphs are known to be the most efficient vectors (Tsai *et al.*, 2008; Petersen and Charles, 1997; Le Maguet *et al.*, 2012a).

The closteroviruses GLRaV-1 and GLRaV-3, known to be closely related within the genus

1
2
3 *Ampelovirus* (Maliogka *et al.*, 2008), are regarded as critical pathogens of grapevine because of
4
5 severity of the associated symptoms and their wide distribution. The transmission of these two
6
7 GLRaVs often occurs together with GVA, the vitivirus that has been associated with the Kolber
8
9 stem grooving syndrome (a disease of the RW complex) and recently with the Shiraz disease. The
10
11 distribution of the three viruses overlap in many regions across all the continents (Sforza *et al.*,
12
13 2003; Bertin *et al.*, 2010; Le Maguet *et al.*, 2012b) and their co-infections in grapevine are
14
15 frequently reported (Pacífico *et al.*, 2011; Voncina *et al.*, 2011; Fuchs *et al.*, 2009).
16
17
18 GLRaV-1, GLRaV-3 and GVA also share insect vectors, among both mealybug and soft scale species
19
20 (Fuchs *et al.*, 2009; Bertin *et al.*, 2010). *Planococcus ficus* Signoret and *Planococcus citri* Risso are
21
22 involved in GLRaVs and GVA transmission and are regarded as a serious threat for viticulture
23
24 (Cabaleiro and Segura, 1997a; Rosciglione and Castellano, 1985; Tsai *et al.*, 2010; Scotto *et al.*,
25
26 2009; Engelbrecht and Kasdorf, 1990; Golino *et al.*, 2002). These two mealybugs often coexist in
27
28 several areas: Africa, Northern, Central and Southern America, Oriental Asia, Middle East and
29
30 Mediterranean countries (Ben-Dov, 2001). The two species show similar ecological and biological
31
32 features, and are known to easily hybridize in experimental conditions (Kol-Maimon *et al.*, 2014).
33
34 Their morphology-based discrimination is awkward and it has been recently unravelled by the
35
36 development of DNA markers (Cavaliere *et al.*, 2008; Daane *et al.*, 2011; Malausa *et al.*, 2011;
37
38 Saccaggi *et al.*, 2008). The current data on virus acquisition, retention and inoculation indicate that
39
40 both *Pl. ficus* and *Pl. citri* transmit the viruses in a semi-persistent manner (Cabaleiro and Segura,
41
42 1997b; Tsai *et al.*, 2008), although evidence of GLRaV-3 presence in the salivary glands of *Pl. citri*
43
44 has been reported (Cid *et al.*, 2007).
45
46
47 Despite the occurrence of GLRaVs and RW-associated viruses mixed infections in vineyard, the co-
48
49 transmission of these viruses has been poorly characterized. Indeed most of the studies focused on
50
51 transmission of the sole GLRaV-3, that has emerged as the key virus of grapevine throughout the
52
53
54
55
56
57
58
59
60

1
2
3 world (Maree *et al.*, 2013). Moreover, the surveys of other grape viruses often did not go beyond
4
5 the identification of new vector species and rarely investigated the relative efficiency of a vector in
6
7 transmitting different viruses as well as the possible interaction among viruses during transmission
8
9 (Tsai *et al.*, 2010; Le Maguet *et al.*, 2012a). Therefore, comparative studies involving different
10
11 GLRaVs / RW-associated viruses and different vector species at a time would improve the current
12
13 knowledge of transmission process and contribute to understand the epidemiology of diseases.
14
15 In the present study, the acquisition and transmission rate of GLRaV-1, GLRaV-3 and GVA by two
16
17 different mealybugs, *Pl. ficus* and *Pl. citri*, were investigated under controlled conditions.
18
19 Moreover, the virus retention was investigated and the virus load was measured by quantitative
20
21 real-time RT-PCR (qRT-PCR) in the mixed-infected source grapevine as well as in the vectors. qRT-
22
23 PCR was already applied to estimate the viral load in the host plants (Pacifico *et al.*, 2011; Osman
24
25 *et al.*, 2012; Tsai *et al.*, 2012) as well as in insect vectors (Mason *et al.*, 2008; Debreczeni *et al.*,
26
27 2011), but this is the first application for the study of virus load in mealybug vectors.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Material and Methods

Insect and plant material

Specimens of *Pl. ficus* and *Pl. citri* were collected on vines in Liguria and Sicily regions (Italy) respectively, and were used to establish colonies in climatic chambers. The colonies were maintained on sprouted potatoes in the dark at 20–30°C. The two species were identified by means of morphological observations as well as amplification and sequence analysis of the *Planococcus* mitochondrial *cytochrome c oxidase subunit 1* gene (Saccaggi *et al.*, 2008).

Infected plant material was from an experimental vineyard planted with *Vitis vinifera* cv. Nebbiolo and situated in the Langhe, a traditional wine-producing area of Piemonte (North-western Italy).

The sanitary status of the vineyard was regularly monitored for the presence of ampelovirus, vitivirus, maculovirus and nepovirus infection by serological (DAS-ELISA) and molecular analysis (PCR) (Gambino and Gribaudo, 2006; Gribaudo *et al.*, 2009). Based on these data, several grapevines carrying the GLRaV-1, -3 and GVA mixed-infection were identified. One of these plants served as “mother plant” and provided the source cuttings for the laboratory experiments. In detail, dormant shoots collected from this plant were rooted and maintained in a greenhouse, until they were about 50 cm tall and then used as source plants. Each sprouting cutting was further checked for GLRaV-1, GLRaV-3 and GVA infection before being used for the virus acquisition.

Healthy grapevines obtained through micropropagation of clonal lines of *V. vinifera* of cv Barbera were used as test plants in the transmission assays.

Transmission experiments

Three repetitions of transmission trials were performed for *Pl. ficus* as well as for *Pl. citri*. In each experiment, a leaf-strewn branch from one of the virus-source cuttings generated from the mother plant was laid on the mealybug rearing. The mealybugs were allowed to move onto the branch for

1
2
3 four hours; then the branch was removed and maintained in a flask of water for 24 hours. After
4
5 this 24h acquisition access period (AAP), the mealybugs were gently shaken off the source branch
6
7 onto a dark paper. The first instar nymphs were isolated, and partly kept for RNA extraction and
8
9 virus detection and partly transferred to the test plants for virus transmission. The transmission
10
11 was carried out by caging five first-instar nymphs on the upper leaf surface of each test plant for
12
13 48h. Only the first-instar nymphs were used as they are known to be the most efficient vectors of
14
15 grapevine viruses. Since the position of the leaves where the mealybugs fed during AAP (i.e. basal,
16
17 intermediate or apical leaves) does not affect the transmission rate (Tsai *et al.*, 2011), the nymphs
18
19 used for transmission were randomly chosen among the whole batch recovered on the dark paper.
20
21 After the virus inoculation access period (IAP) of 48h, the first instars were removed from the test
22
23 plants and these were drench-treated with a systemic insecticide (Actara, Syngenta Crop
24
25 Protection). The test grapevines were maintained in a greenhouse and sprayed regularly with
26
27 insecticide and fungicide for four-five months, until the RNA extraction and GLRaV-1, -3 and GVA
28
29 detection. All plants were periodically pruned to avoid overgrowth.

30
31 Thirty, 18 and 10 test plants were inoculated by *Pl. ficus* in three transmission experiments and 27,
32
33 14 and 10 grapevines respectively lived through the growing period until the GVA, GLRaV-1 and
34
35 GLRaV-3 detection. In the three experiments, a total of 71 first instar nymphs (8, 38 and 25 per
36
37 trial) were assayed after 24h-AAP to estimate the virus acquisition rate. Three transmission trials
38
39 were also performed for *Pl. citri*: 26, 19 and 10 test plants per experiment were inoculated and 25,
40
41 17 and 7 grapevines respectively **were surviving** for virus diagnosis. A total of 74 first instar nymphs
42
43 (30, 29 and 15 per experiment) **were analysed at the same time** for GVA, GLRaV-1 and GLRaV-3
44
45 presence. Specimens from healthy colonies of *Pl. ficus* and *Pl. citri* were used as negative controls.
46
47
48 To confirm the virus-free sanitary status of the test plants and to ensure that virus spread had not
49
50 occurred within the greenhouse during experimental periods, *in vitro*-derived grapevines from the
51
52
53
54
55
56
57
58
59
60

1
2
3 same batch of test plants were not exposed to mealybug vectors and served as negative controls.
4
5 Differences in virus acquisition and virus transmission frequencies within and between the two
6
7 mealybug species were tested by chi-square (χ^2) analysis (SigmaPlot 11.0). Fisher Exact Test was
8
9 used instead of χ^2 when the frequencies of one or more observations were less than five.
10
11

12 13 14 15 **Retention experiments**

16
17 The retention and the load of GLRaV-1, -3 and GVA were investigated in *Pl. citri* and *Pl. ficus*. Each
18
19 mealybug species was caged to feed onto a virus-infected vine branch as described above. After an
20
21 AAP of 24 hours, the first instar nymphs were partly collected for virus diagnosis (T_0) and partly
22
23 transferred onto potatoes for post-acquisition feeding. The potato is known to be a non-host of
24
25 grapevine viruses. Further nymphs were taken from potato sprouts after 24, 48, 72 and 96 hours
26
27 from the end of the AAP (T_{24} , T_{48} , T_{72} , T_{96}), and tested for virus presence. Samples that resulted
28
29 positive to at least one of the three viruses at different post-acquisition times were further
30
31 analysed by qRT-PCR to estimate the viral load.
32
33
34
35
36
37

38 39 **RNA purification from insects and plants**

40
41 The RNA for virus detection was purified from the *Pl. citri* and *Pl. ficus* nymphs collected after 24h-
42
43 AAP in the transmission experiments and collected at T_0 , T_{24} , T_{48} , T_{72} , T_{96} in the retention
44
45 experiments. Total RNA was extracted from single nymphs using the TRIzol® Reagent (Invitrogen)
46
47 and following the manufacturer's instructions. Samples were treated with 2 units of RNase-Free
48
49 DNase I (Applied Biosystems) in the supplied buffer to avoid residual DNA contamination. After
50
51 DNA digestion, DNase was inactivated by phenol/chloroform extraction. RNA was finally
52
53 resuspended in 20 μ l of RNase-free water containing diethylpyrocarbonate (DEPC) 0.1 %.
54
55
56
57
58
59
60

1
2
3 Concentration and purity of extracts were evaluated using the ND-1000 Spectrophotometer
4
5 (Nanodrop Technologies). The RNA was then diluted to $10 \text{ ng } \mu\text{l}^{-1}$ and stored at $-80 \text{ }^\circ\text{C}$.
6

7
8 The plant RNA was extracted from sprouting cuttings before each experiment, in order to select
9
10 the source plants carrying GLRaV-1, GLRaV-3 and GVA, and from the test grapevines at the end of
11
12 the transmission trials. The RNA was purified from a total of 0.1 g of midribs from both basal and
13
14 apical leaves. The extraction was performed using the Concert™ Plant RNA Isolation Reagent
15
16 (Invitrogen) and following the manufacturer's instructions. RNA was resuspended in 30 μl of DEPC
17
18 0.1 % RNase-free water, diluted to $10 \text{ ng } \mu\text{l}^{-1}$ and stored at $-80 \text{ }^\circ\text{C}$.
19
20
21
22
23

24 **Virus detection**

25
26 The virus detection from both insect and plant RNA extracts was carried out by SYBR® Green real-
27
28 time RT-PCR assays in the Chromo4 Real Time Thermal Cycler (Bio-Rad) supported by the
29
30 OpticonMonitor 3.1.32 software (Bio-Rad). The GVA, GLRaV-1 and GLRaV-3 were detected with the
31
32 following primer pairs, designed on the viral coat protein sequences: GVA-C7273 (5'-
33
34 CATCGTCTGAGGTTTCTACTA-3') / GVA-H7038 (5'-AGGTCCACGTTTGCTAAG-3') (MacKenzie *et al.*,
35
36 1997); GLRaV-1fw (5'-CGTTTGAAAATCCTATGCGTCAG-3') / GLRaV-1rev (5'-
37
38 GCAACTTCTCGTTCCGGCTTC-3') and GLRaV-3fw (5'-TTCGAGAAAGATCCAGACAAGTTC-3') / GLRaV-
39
40 3rev (5'-ATAACCTTCTTACACAGCTCCATC-3') (Gribaudo *et al.*, 2009). The real-time RT-PCR was
41
42 performed with the Iscript One-Step RT-PCR kit (Bio-Rad), using a final primer concentration of 300
43
44 nM. Ten nanograms of insect or plant total RNA were used as template. For all the primer pairs,
45
46 the thermo-cycling conditions consisted of an initial cycle at 50°C for 10 min, followed by 5 min at
47
48 95°C and 40 cycles at 95°C for 10 s and 60°C for 60 s. Melting curves were produced at the end of
49
50 the PCR to assess the reaction specificity: the PCR products were heated to 95°C for 1 min, cooled
51
52 at 65°C for 1 min and then slowly heated back to 95°C at a rate of 0.5°C per cycle.
53
54
55
56
57
58
59
60

1
2
3 RNA of mealybugs or vines carrying GVA, GLRaV-1 and GLRaV-3 were used as positive controls.
4
5 RNA of mealybugs from healthy colonies and from healthy *in vitro*-derived grapevines served as
6
7 negative controls.
8
9

10 11 12 **Virus quantification** 13

14 *Set up of a quantitative real-time RT-PCR assay.* Viral load was measured in relation to the
15
16 transcript copy number of a mealybug reference gene; this, rather than absolute quantitation, was
17
18 chosen to avoid the influence of different yields during RNA extractions (Pacifico *et al.*, 2011).
19

20 To identify the most stable insect gene upon mealybug development, the *actin β* (*ACTB*),
21
22 *glyceraldehyde 3-phosphate dehydrogenase* (*GAPDH*) and *18S ribosomal RNA* genes were selected
23
24 among those suggested in the literature for gene expression studies in virus-vector insects
25
26 (Maroniche *et al.*, 2011; Rotenberg *et al.*, 2009). The *18S rRNA* was amplified using the primers
27
28 MqFw and MqRv, available from Marzachí and Bosco (2005) (Table 1). The sequences of the other
29
30 two genes were not available for insects in the family *Pseudococcidae*. Therefore, the degenerated
31
32 primers ActinDegF1 (5'-YGAYTGGARAARATCTGGC-3') / ActinDegR2 (5'-CGTCGTAYTCYTYTTSGAG-
33
34 3') and GAPDegF1 (5'-GGTATCAATGGWTTTGGC-3') / GAPCoccR2 (5'-CAWAYTCRTTGTCTGACC-3')
35
36 were designed on the homologous genes of the aphid *Acyrtosiphon pisum*. The primers were
37
38 tested in conventional PCR using 2 µl of *Pl. ficus* and *Pl. citri* cDNA as templates, under the
39
40 following conditions: denaturation for 4 min at 94°C, followed by 35 cycles of 30 s at 94°C, 30 s at
41
42 48°C and 1 min 30 s at 72°C; and a post-dwell period of 5 min at 72°C. PCR products were analysed
43
44 by electrophoresis through a 1.5 % agarose gel in 1 × Tris-borate-EDTA (TBE) buffer along with a 1
45
46 kb plus DNA size marker (Gibco BRL), and visualised on a UV transilluminator. PCR products were
47
48 isolated from the gel using the PureLink™ (Invitrogen), ligated into the pGEM-T easy vector
49
50 (Promega) and transformed in *E. coli* DH5α. Plasmids were purified with the Fast Plasmid Mini kit
51
52
53
54
55
56
57
58
59
60

1
2
3 (Eppendorf AG) and sequenced (BioFab Research) with the universal primers M13F/R. Raw
4
5 sequence data were manually edited with the DNAMAN program vers. 4.02 (Lynnon BioSoft) after
6
7 at least twice the sequencing coverage for each nucleotide position. *ACTB* and *GAPDH* sequences
8
9 of both *Pl. citri* and *Pl. ficus* were trimmed and aligned with the DNAMAN 4.02 program (Lynnon
10
11 BioSoft). Finally, the specific primer pairs PfActFw2/PfActBRev1 and PfGapFw1/PfGapRev were
12
13 designed (Primer Express tool; Applied Biosystems) (Table 1).
14
15

16
17
18 The primer specificity for the three target genes was tested in SYBR[®] Green Real-Time PCR assays.
19
20 Two microliters of *Pl. citri* and *Pl. ficus* cDNA were added to a 2× iQ[™] SYBR[®] Green Supermix (Bio-
21
22 Rad) supplemented with 500 nM each primers and DEPC 0.1 % water to a final volume of 25 µl.
23
24 Reaction conditions were as follows: 95°C for 5 min followed by 40 cycles of 30 s at 95°C and 90 s
25
26 at 60°C (62°C for *GAPDH*). Melting curves were produced at the end of each specific real-time
27
28 amplification.
29
30

31
32 To identify the most stable insect gene upon mealybug development, total RNA was singly
33
34 extracted from three newly-hatched nymphs, three third instar nymphs, and three female adults of
35
36 both *Pl. citri* and *Pl. ficus*, following the RNA extraction procedures detailed above. RNA extracts
37
38 were then reverse transcribed into cDNA using High Capacity cDNA Reverse Transcription Kit
39
40 (Applied Biosystems) with random hexamers, according to the manufacturer's protocol.
41
42

43
44 For the calculation of reaction efficiencies (E), the cDNAs from nymphs and adults of the two
45
46 species were pooled and diluted to be used as real-time PCR standards. Tenfold serial dilutions of
47
48 standard cDNA (1:1, 1:10, 1:100 and 1:1000) in DEPC 0.1% RNase-free water were run in triplicate
49
50 for each PCR system (*ACTB*, *GAPDH* and 18S rRNA). Standard curves were constructed by linear
51
52 regression analysis of the threshold cycle (Ct) value of each of the standard dilution replicates
53
54 versus the Log of arbitrary concentration values attributed to each cDNA dilution (1000, 100, 10, 1,
55
56
57
58
59
60

1
2
3 respectively). SYBR[®] Green Real-Time PCR conditions were the same described above. Data
4
5 acquisition and analysis were handled by the OpticonMonitor 3.1.32 software (Bio-Rad).
6

7
8 To determine the best reference gene, the cDNA of each sample was run in duplicate under the
9
10 same reaction conditions of the standards. The average Ct value of each sample was converted
11
12 using the Δ Ct method and the calculated PCR efficiency, then analysed with geNorm
13
14 (Vandesompele *et al.*, 2002), Bestkeeper (Pfaffl *et al.*, 2004) and Normfinder (Andersen *et al.*,
15
16 2004) software, according to the authors' instructions.
17

18
19 *Virus quantification.* Within the retention experiments, a subset of *Pl. citri* and *Pl. ficus* nymphs
20
21 carrying at least one virus among GVA, GLRaV-1 and GLRaV-3 was selected for virus quantification.
22

23
24 The virus load was also assessed in the mother plant providing the source cuttings used for virus
25
26 acquisition. The RNA-dependent RNA polymerase (*RdRp*) gene was chosen as target for virus RNA
27
28 amplification. The *RdRp* gene of GVA, GLRaV-1 and GLRaV-3 was amplified by one-tube TaqMan[®]
29
30 qRT-PCR, using primers and probes specific for each virus and described in Pacifico *et al.* (2011).
31
32

33
34 The insect *GAPDH* resulted to be the most stable reference gene (see Results) and was chosen as
35
36 target for insect cDNA amplification. The *GAPDH* sequence was amplified by one-tube TaqMan[®]
37
38 qRT-PCR, using the primer pair PfGapFw1/PfGapRev and the corresponding probe PfGapProbe
39
40 (Table 1). Ten nanograms of total RNA extract from each insect were run in triplicate together with
41
42 at least three dilutions of insect *GAPDH* and virus standard RNAs. Reaction conditions were the
43
44 same reported by Pacifico *et al.* (2011). The standard RNAs for the absolute quantification of the
45
46 viral genome copies were prepared as detailed in Pacifico *et al.* (2011). The insect standard RNAs
47
48 were obtained by *in vitro* transcription of *SpeI* (Promega) linearized pGemGAPDHf with the
49
50 MAXIscript[®] *in vitro* Transcription Kit (Applied Biosystems), following the manufacturer's
51
52 instructions. To eliminate plasmid DNA, RNA was treated with two units of RNase-free DNase I
53
54 (Applied Biosystems) in the supplied buffer. Following phenol/chloroform extraction, RNA was
55
56
57
58
59
60

1
2
3 dissolved in 30 µl of DEPC 0.1 % RNase-free water, and analysed using the ND-1000
4
5 Spectrophotometer. The number of RNA copies per microlitre was calculated according to Pacifico
6
7 *et al.* (2011). RNA was diluted, distributed in aliquots and stored at –80°C. Tenfold serial dilutions
8
9 of this RNA in DEPC 0.1% RNase-free water were run in triplicate under the following conditions:
10
11 52°C for 30 m, 5 min at 95°C, 45 cycles of 15 s at 95°C, 30 s at 52°C and 30 s at 60°C.

12
13
14 The viral load in mealybugs was expressed as viral Genome Units (GU) per insect *GAPDH*: the
15
16 absolute quantity of the viral GU and *GAPDH* transcripts of each insect sample were derived from
17
18 the raw qRT-PCR data, and the concentration of each virus was finally related to the *GAPDH*
19
20 transcript copies, according to Pacifico *et al.* (2011). The GVA, GLRaV-1 and GLRaV-3 load was also
21
22 measured in the mother plant by relating the viral GU with plant *GAPDH* transcript copy. The
23
24 experimental protocol was according to Pacifico *et al.* (2011).
25
26
27
28
29
30

31 Results

32 Virus transmission

33
34
35 The results of GVA, GLRaV-1 and GLRaV-3 diagnosis in *Pl. ficus* and *Pl. citri* as well as in test plants
36
37 are summarized in Table 2. The data of the three transmission experiments performed for each
38
39 vector were cumulated as the results of the three repeats did not differ significantly. No viruses
40
41 were detected in healthy mealybugs and *in vitro*-derived grapevines used as negative controls.
42
43 Concerning the experiments carried out with *Pl. ficus*, at least one of the viruses was detected in
44
45 50 of the 71 (70%) first-instar nymphs assayed after 24h-AAP as well as in 27 of the 51 (53%)
46
47 inoculated grapevines. Out of the positive *Pl. ficus* nymphs, the GVA rate (74%) was higher than
48
49 GLRaVs rates ($\chi^2 = 9.90$, $df = 2$, $P < 0.01$). GLRaV-1 and GLRaV-3 showed similar acquisition
50
51 frequencies (46% and 48%, respectively) but greatly differed in transmission frequencies ($\chi^2 =$
52
53 17.26, $df = 1$, $P < 0.001$), occurring in the test plants at the lowest (30%) and the highest (89%) rate
54
55
56
57
58
59
60

1
2
3 respectively. GLRaV-3 transmission efficiency was higher than acquisition efficiency, but the
4
5 acquisition rate was assessed on single nymphs whereas the transmission experiments were
6
7 performed with groups of five insects per plant. The maximum-likelihood estimator P_s , calculated
8
9 according to Swallow (1985), allows to infer the proportion of infected insects within each five-
10
11 insect batch used for transmission. An estimated proportion of 0.36 *Pl. ficus* nymphs transmitted
12
13 GLRaV-3. This value approaches the proportion of single nymphs that acquired this virus after 24h-
14
15 AAP (0.48) and provides indications of the high GLRaV-3 transmission potential of *Pl. ficus*. GVA,
16
17 GLRaV-1 and GLRaV-3 were all acquired and transmitted preferably as mixed rather than single
18
19 viruses by *Pl. ficus* ($P < 0.05$ for the comparisons between single and mixed infections for all the
20
21 three viruses at both acquisition and transmission steps). Within the mixed infections, the
22
23 association of GVA + GLRaV-1 was found in the nymphs after 24h-AAP but not in the test plants
24
25 exposed to the viruliferous mealybugs (Table 2).
26
27
28
29
30

31
32 Grapevine viruses were detected in 46 of the 74 (62%) tested *Pl. citri* first-instar nymphs, both in
33
34 single and mixed infections. At least one virus was found in 34 of the 49 (69%) test plants
35
36 inoculated by *Pl. citri*. The acquisition rate of GLRaV-1 and GLRaV-3 was the same (67% of the
37
38 virus-positive nymphs; Table 2) and did not significantly differ from the GVA rate (83%; $\chi^2 = 2.09$, df
39
40 = 1, $P = 0.149$). On the other hand, the frequency of GVA, GLRaV-1 and GLRaV-3 transmission (9%,
41
42 70% and 47% of the virus-positive test plants, respectively) significantly varied ($\chi^2 = 27.10$, $df = 2$, P
43
44 < 0.001); this was mainly due to the low transmission of GVA, which was inoculated only to three
45
46 grapevines, together with GLRaV-3 or with GLRaV-1 + GLRaV-3. The relative rates of GLRaV-1 and
47
48 GLRaV-3 single and mixed infections differed in acquisition and transmission. Indeed, the two
49
50 ampeloviruses were mainly acquired by *Pl. citri* in mixed infections (acquisition rate of single
51
52 GLRaV-1 = 11%, acquisition rate of GLRaV-1 mixed infections = 56%, GLRaV-1 single vs mixed
53
54 acquisitions: $\chi^2 = 19.46$, $df = 1$, $P < 0.001$; acquisition rate of single GLRaV-3 = 2%, acquisition rate
55
56
57
58
59
60

1
2
3 of GLRaV-3 mixed infections = 64%, GLRaV-3 single vs mixed acquisitions: $\chi^2 = 38.14$, $df = 1$, $P <$
4
5 0.001). However, the GLRaV-1 transmission occurred more frequently in single, 53%, than in mixed
6
7 infections, 18%, (GLRaV-1 single vs mixed transmissions: $\chi^2 = 7.79$, $df = 1$, $P < 0.01$) and the
8
9 transmission rates of GLRaV-3 single and mixed infections were the same (24%, Table 2). Within
10
11 the mixed infections, the association GVA + GLRaV-1 + GLRaV-3 was acquired and transmitted by
12
13 *Pl. citri* at different rates: the GVA + GLRaV-1 + GLRaV-3 frequency was the highest in nymphs
14
15 (43%; $P < 0.05$ for all the comparisons: GVA + GLRaV-1 + GLRaV-3 vs GVA + GLRaV-1, GVA + GLRaV-
16
17 1 + GLRaV-3 vs GVA + GLRaV-3 and GVA + GLRaV-1 + GLRaV-3 vs GLRaV-1 + GLRaV-3), but did not
18
19 vary significantly in test plants (3%) compared to the other mixed infections ($P > 0,05$ for all the
20
21 comparisons: GVA + GLRaV-1 + GLRaV-3 vs GVA + GLRaV-1, GVA + GLRaV-1 + GLRaV-3 vs GVA +
22
23 GLRaV-3 and GVA + GLRaV-1 + GLRaV-3 vs GLRaV-1 + GLRaV-3).

24
25
26
27
28
29 The comparison between *Pl. ficus* and *Pl. citri* experimental data showed that the acquisition
30
31 efficiency of the two species did not significantly vary for either GVA, or GLRaV-1, or GLRaV-3 ($P >$
32
33 0.05 for all the comparisons), though *Pl. citri* acquired the three viruses together with a higher
34
35 efficiency ($\chi^2 = 12.26$, $df = 1$, $P < 0.001$). On the other hand, the two mealybugs differently
36
37 transmitted the viruses to the test plants. Indeed, the overall GVA and GLRaV-3 transmission
38
39 frequencies were higher for *Pl. ficus* than *Pl. citri* ($\chi^2 = 17.63$, $df = 1$, $P < 0.001$ for GVA; $\chi^2 = 9.88$, df
40
41 $= 1$, $P < 0.01$ for GLRaV-3), whereas *Pl. citri* was significantly more efficient in transmitting GLRaV-1
42
43 ($\chi^2 = 8.55$, $df = 1$, $P < 0.01$), mainly as single infection ($\chi^2 = 14.79$, $df = 1$, $P < 0.001$). Looking at the
44
45 transmission of virus mixed infections, it is worthy to note that *Pl. ficus* transmitted GVA + GLRaV-
46
47 3 at a higher frequency than *Pl. citri* ($\chi^2 = 8.92$, $df = 1$, $P < 0.01$) and that both the vectors carried
48
49 GVA + GLRaV-1 but they did not transmit this virus association to test plants.
50
51
52
53
54
55
56
57
58
59
60

Virus retention by *Pl. citri* and *Pl. ficus*

Both *Pl. citri* and *Pl. ficus* were daily assayed for virus retention up to four days after a 24h-AAP.

A total of 218 individuals of *P. ficus* were analysed at T₀ (n=46), T₂₄ (n=45), T₄₈ (n=43), T₇₂ (n=44) and T₉₆ (n=40) by SYBR[®] Green real-time RT-PCR assays. The 72%, 46% and 11% of the samples tested at T₀ were positive to GVA, GLRaV-1 and GLRaV-3, respectively. The infection rate of GVA, GLRaV-1 and GLRaV-3 decreased from T₀ to T₇₂ and none of the viruses was detected at T₉₆ (Fig. 1a).

One hundred individuals of *Pl. citri* were analysed, twenty at each sampling point. In this species, the trends of GVA and GLRaV3 retention were similar to those observed in *Pl. ficus*. A slightly different retention trend of GLRaV-1 was observed (with a peak of virus presence at T₇₂), likely due to the small sample size (Fig. 1b).

Virus quantification

The absolute quantity of GLRaV-1, GLRaV-3 and GVA genomes in infected *Pl. ficus* and *Pl. citri* as well as in the source plant were measured by qRT-PCR. The *RdRp* gene of each virus was quantified and normalized on the expression level of insect/plant mRNA.

To choose the best insect reference gene, the expression stability of the *ACTB*, *GAPDH* and *18S rRNA* genes was verified. Based on the SYBR[®] Green Real-Time PCR efficiency calculated for each amplification system, Bestkeeper, GeNorm and Normfinder softwares ranked the *GAPDH* as the most stable gene (Table 1). Therefore the viral load in the mealybugs was expressed as viral GU per insect *GAPDH* transcript copy.

For the absolute quantification of viral GU and insect *GAPDH* transcripts, specific standard curves were obtained by running 10-fold serial dilutions of insect and virus standards RNAs. The *GAPDH* standard curve covered a range between 10⁹ and 10³ RNA copies, showing a correlation coefficient (R²) = 0.996 and a reaction efficiency of 75%. For the GLRaV-1, GLRaV-3 and GVA absolute

1
2
3 quantification, the standard curves covered a range between 10^8 and 10^2 copies of transcript RNA,
4
5 with the same R^2 and reaction efficiencies reported in Pacifico *et al.* (2011). The calculated *GAPDH*
6
7 mRNA mean absolute quantities were 3.19×10^5 (SE = 1.06×10^5) in *Pl. ficus* and 1.67×10^5 (SE =
8
9 3.10×10^4) in *Pl. citri*, respectively.

10
11
12 GVA positive *Pl. ficus* (19 individuals) and *Pl. citri* (18 individuals) were analysed to quantify virus
13
14 load. Virus load was above the quantification threshold in five *Pl. ficus* and four *Pl. citri* at T_0 , and in
15
16 two *Pl. ficus* at T_{48} , only (Table 3). GLRaV-1 positive *Pl. ficus* (23 individuals) and *Pl. citri* (7
17
18 individuals) were analysed to quantify virus load. As occurred for GVA, GLRaV-1 load was below
19
20 the quantification threshold in most tested mealybugs (Table 3). GLRaV-3 load was below the
21
22 quantification threshold in all the six *Pl. ficus* and six *Pl. citri* infected individuals. No amplification
23
24 was obtained from water control devoid of template RNA or RT-PCR mix devoid of MuLV reverse
25
26 transcriptase. Due to the low number of samples above the quantification threshold, a statistical
27
28 comparison of virus accumulation between *Pl. ficus* and *Pl. citri* was not conducted.
29
30
31
32
33
34 Viral loads measured in the grapevine used as virus source were 0.52, 2.40 and 2.54 for GVA,
35
36 GLRaV-1 and GLRaV-3 GU per 100 plant *GAPDH* transcript, respectively.

37 38 39 40 41 Discussion

42
43 This study investigates the co-transmission of GLRaVs and RW-associated viruses to grapevine by
44
45 the mealybugs *Pl. ficus* and *Pl. citri*. After feeding onto GVA, GLRaV-1 and GLRaV-3 mixed-infected
46
47 plants, the vectors were tested for their transmission efficiencies. The two species shared high
48
49 rates of virus acquisition, being 70% and 62% of the tested first-instar nymphs positive to at least
50
51 one virus for *Pl. ficus* and *Pl. citri* respectively. Looking at the overall virus transmission rates, *Pl.*
52
53 *citri* was able to inoculate a higher number of test plants compared to *Pl. ficus* (70% vs 53%).
54
55
56
57
58 Anyway, both the species proved to be efficient vectors under our experimental conditions.
59
60

1
2
3 To provide a more accurate description of transmission patterns, the overall transmission data
4
5 were detailed for each single- and mixed-virus association and the retention and load of each virus
6
7 in insects were analysed as well. To measure the virus load, a method for the quantification of GVA
8
9 and GLRaVs in the vectors was set up for the first time. In this assay the absolute quantity of viral
10
11 genomes was related to the copy numbers of a suitable insect reference gene to reduce the
12
13 experimental bias due to RNA extraction and PCR set up. To this aim, the expression stability of
14
15 three candidate reference genes proposed for gene expression studies of planthopper and thrip
16
17 vectors was evaluated and the *GAPDH* gene was selected as the best insect reference gene in our
18
19 experimental conditions. The virus RNA quantification was performed by using the *RdRp* gene,
20
21 already used as amplification target in grapevine (Pacífico *et al.*, 2011; Tsai *et al.*, 2012; Velasco *et*
22
23 *al.*, 2014). This target avoids the overestimation of viral loads as *RdRp* copy numbers are directly
24
25 related to the number of viral genomes.
26
27
28
29

30
31 For the transmission experiments, insects were fed on source cuttings derived from a mother plant
32
33 with a known amount of the three viruses. The virus load in plant may vary as a consequence of
34
35 specific interaction virus-host and may influence the acquisition and transmission by the same
36
37 vector (Martin and Elena, 2009). For example, different studies reported that crinivirus
38
39 transmission efficiency by whitefly vectors to herbaceous plants was directly influenced by the
40
41 virus concentration in the acquisition source (Ng *et al.*, 2004; Wintermantel *et al.*, 2008). In this
42
43 study, the qRT-PCR assay performed on the mother plant showed that the quantities of the three
44
45 viruses were similar: the load of GLRaV-1 and GLRaV-3 was almost the same (2.40 - 2.54 GU / 100
46
47 *GAPDH*) and GVA was slightly less concentrated (0.52 GU / 100 *GAPDH*). These can be reasonably
48
49 considered as similar loads, especially when compared with the GVA, GLRaV-1 and GLRaV-3
50
51 amounts measured in grapevines by Giribaldi *et al.* (2011) and Pacífico *et al.* (2011).
52
53
54

55
56 Besides having the same load in plant, GLRaV-1 and GLRaV-3 also showed similar acquisition
57
58
59
60

1
2
3 patterns. Indeed the acquisition frequencies of the two viruses were very close in the experiments
4
5 carried out with *Pl. ficus* and *Pl. citri* and their uptake preferably occurred together with GVA.
6
7 However, their transmission patterns diverged in the two species. The GLRaV-3 transmission by *Pl.*
8
9 *ficus* occurred at very high frequencies. The percentage of GLRaV-3 infected plants (89%) as well as
10
11 the proportion of single infecting nymphs ($P_s = 0.36$) were higher than the estimates reported in
12
13 previous studies dealing with GLRaV-3 transmission by first- and second-instar nymphs of *Pl. ficus*
14
15 (Douglas and Krüger, 2008; Tsai *et al.*, 2008; Tsai *et al.*, 2010; Tsai *et al.*, 2011; Mahfoudhi *et al.*,
16
17 2009). Almeida *et al.* (2013) matched up all these transmission data and provided an overall P_s
18
19 interval covering the information from several studies: these P_s values just ranged from 0.04 to 0.2
20
21 and therefore did not include our value. The high transmission efficiency observed for *Pl. ficus*
22
23 nymphs is not associated with a high virus load in the insect, as GLRaV-3 load never reached the
24
25 quantification threshold and was detected only up to 48h after AAP. These data suggest that *Pl.*
26
27 *ficus* may have a high capability to transmit GLRaV-3, in spite of the accumulation of a very low
28
29 virus load.
30
31
32
33
34

35
36 *Planococcus citri* is known as GLRaV-3 vector in vineyards, and high percentages of *Pl. citri*
37
38 individuals carrying the virus can be trapped in the field (Cid *et al.*, 2010). However, only Cabaleiro
39
40 and Segura (1997b) performed laboratory experiments to estimate the GLRaV-3 transmission
41
42 efficiency by this mealybug. The authors reported low GLRaV-3 transmission rates even against
43
44 high acquisition rates (10% of test plants resulted positive to GLRaV-3 following exposure to groups
45
46 of *Pl. citri* nymphs that were expected to have 80% of GLRaV-3 viruliferous individuals) and showed
47
48 that nymphs quickly lost their infectivity (1h after leaving the infected grapevine). In our study, the
49
50 GLRaV-3 transmission efficiency was higher and virus retention lasted longer. Therefore, this
51
52 species proved to be an efficient vector, consistently with the fast GLRaV-3 spread that was
53
54 recorded even in case of low field-population densities of *Pl. citri*. As occurred for *Pl. ficus*, *Pl. citri*
55
56
57
58
59
60

1
2
3 nymphs hosted a GLRaV-3 load below the quantification threshold. This confirms that the GLRaV-3
4
5 transmission to plants could efficiently occur even if the vector nymphs carry a very low virus load.
6
7 Anyway, the comparison between the transmission rates performed by the two mealybug species
8
9 showed that *Pl. ficus* is a more efficient GLRaV-3 vector than *Pl. citri*, as previously suggested by
10
11 Tsai *et al.* (2008).
12
13

14
15 A large genetic variability was described for GLRaV-3 and different virus variants were found within
16
17 single plants (Sharma *et al.*, 2011). It is known that these variants can follow different patterns of
18
19 vector transmission and plant infection and that the disease spread and severity can be affected by
20
21 the virus genotype (Almeida *et al.*, 2013; Blaisdell *et al.*, 2015). Therefore, future studies should be
22
23 addressed to re-assess the GLRaV-3 transmission efficiency of *Pl. ficus* and *Pl. citri* in the presence
24
25 of different virus isolates.
26
27

28
29 GLRaV-1 is less widespread than GLRaV-3, and the virus transmission mechanisms received little
30
31 attention. So far, only few soft scale and mealybug species are recognized as competent vectors,
32
33 including *Pl. ficus* (Sforza *et al.*, 2003; Tsai *et al.*, 2010). In this study, *Pl. ficus* nymphs transmitted
34
35 GLRaV-1 but the rates were significantly lower than those observed for GLRaV-3. The low
36
37 transmission performance occurred although the nymphs retained relatively high quantities of
38
39 virus throughout the 48h-IAP.
40
41

42
43 *Planococcus citri* was initially excluded from the list of GLRaV-1 vectors (Golino *et al.*, 2002), but
44
45 later (Scotto *et al.*, 2009) proved the ability of *Pl. citri* nymphs and adults to acquire and transmit
46
47 GLRaV-1. Our study confirms these results and indicates that both acquisition and transmission
48
49 occurred with a high efficiency. In detail, GLRaV-1 was acquired mainly together with GLRaV-3 and
50
51 GVA but was then inoculated to grapevine preferably as single infection, and the number of test
52
53 plants infected by the sole GLRaV-1 was significantly higher than the number of plants carrying the
54
55 other two viruses, either alone or in associations. *Planococcus citri* was a more efficient vector of
56
57
58
59
60

1
2
3 GLRaV-1 than *Pl. ficus* (70% vs 30% inoculated plants), thus suggesting that it could contribute to
4
5 the GLRaV-1 spread in vineyard more efficiently than the most common vine mealybug, *Pl. ficus*.
6
7 GVA infections are widespread in all major wine-producing regions and the virus is often detected
8
9 in grapevine together with GLRaVs. Indeed, GVA and GLRaVs share several vectors among
10
11 mealybugs and soft scales, including *Pl. ficus* and *Pl. citri* (Rosciglione and Castellano, 1985). This
12
13 study shows that both insects acquired GVA very efficiently, and the GVA rate in *Pl. ficus* nymphs
14
15 was even significantly higher than GLRaV-1 and -3 rates, although the three viruses had similar
16
17 loads in the source plant. The two mealybugs also showed common patterns of GVA retention,
18
19 since the virus was detectable until 72h after AAP in both species and was accumulated at similar
20
21 load after acquisition. *Planococcus ficus* kept high level of efficiency also in transmitting GVA,
22
23 especially when the vitivirus was inoculated to grapevine together with GLRaV-3 and GLRaV-1 +
24
25 GLRaV-3. On the contrary, *Pl. citri* poorly transmitted GVA: no single infections were recorded in
26
27 test plants and the mixed infections were limited to few plants. The lack of transmission was even
28
29 more evident looking at the trend of the GVA + GLRaV-1 + GLRaV-3 association: the rate of *Pl. citri*
30
31 nymphs carrying all the three viruses prevailed against the other virus combinations, but this
32
33 predominance could not be observed in test plants after inoculation. This leads to suppose that
34
35 the nymphs carrying GVA in mixed infections may lose the ability to inoculate GVA to grapevine but
36
37 maintain the ability to transmit GLRaVs.
38
39
40
41
42
43

44
45 The coexistence of GVA and GLRaVs in source plants offers the chance to investigate possible
46
47 interactions among the viruses during transmission. To date, both competition or facilitation
48
49 among GLRaVs and vitiviruses were hypothesized without coming to any conclusive evidences
50
51 (Almeida *et al.*, 2013). Some authors suggested that GVA may require the presence of GLRaVs in
52
53 the source plant to be transmitted by mealybugs and soft scales and establish infection in a
54
55 susceptible plant (Hommay *et al.*, 2008; Engelbrecht and Kasdorf, 1990), whereas other studies
56
57
58
59
60

1
2
3 indicated that GLRaVs would benefit from GVA for transmissibility (Fortusini *et al.*, 1997; Zorloni *et*
4 *al.*, 2006; Tsai *et al.*, 2010). Our study confirms the finding of Blaisdell *et al.* (2012), who reported
5 that *Pl. ficus* can transmit GVA from infected to susceptible grapevines without simultaneous
6 transmission of GLRaVs. However, this was not confirmed by the transmission experiments with *Pl.*
7 *citri*. At the same time, we did not find evidences that GVA would mediate the GLRaVs
8 transmission: GLRaV-1 and GLRaV-3 can be acquired and transmitted without GVA by *Pl. ficus* and
9 *Pl. citri*. Namely, GLRaV-1 did not benefit from the presence of GVA, since the association GVA +
10 GLRaV-1 was acquired but never transmitted by both vectors. Therefore, a clear picture about virus
11 interactions is still lacking and further experiments are needed to better characterize the biology of
12 GVA and GLRaVs associations in mealybugs as well as in grapevine. For example, transmission
13 experiments from GVA singly infected plants would indicate whether the known patterns are due
14 to mechanisms of interaction or are merely circumstantial.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31 This study also provides information about the mode of GLRaV-1, GLRaV-3 and GVA transmission
32 by mealybugs. The vitiviruses GVA and GVB are known to be semi-persistently transmitted by their
33 vectors (Adams *et al.*, 2004) as well as all closteroviruses studied so far (Ng and Falk, 2006).

34
35
36
37
38 However, less information are available for the GLRaVs and concern GLRaV-3 only: it was reported
39 that *Pl. ficus* need less than 24h to acquire GLRaV-3 and that both *Pl. ficus* and *Pl. citri* quickly lose
40 the ability to transmit the virus (GLRaV-3 retention time of *Pl. ficus* = 3 days; GLRaV-3 retention
41 time of *Pl. citri* = 24h; Tsai *et al.*, 2008; Cabaleiro and Segura, 1997b). These data are consistent
42 with the characteristics of the semi-persistent transmission (Ng and Falk, 2006), even though the
43 presence of GLRaV-3 particles observed in the *Pl. citri* salivary glands raised the question that the
44 virus might circulate in the vector body (Cid *et al.*, 2007). Our study confirms that a 24h feeding
45 period is enough for efficient acquisition of GLRaV-1 and GLRaV-3 by *Pl. ficus* and *Pl. citri*, and
46 shows that the retention of the two viruses gradually decreased over time, and at the fourth post-
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 acquisition day viruses cannot be detected anymore in both vectors. This supports the hypothesis
4
5 that the GLRaVs transmission by mealybugs is semi-persistent rather than circulative. Moreover,
6
7 circulative viruses are known to have a relatively high load in their vectors because they colonize
8
9 the haemocoel and several organs, whereas our data from qRT-PCR suggest that the amount of
10
11 GLRaVs and GVA is low in both *Pl. ficus* and *Pl. citri*. Indeed, GLRaV-3 was below the detection
12
13 threshold in all our tested insects, and most of the mealybugs that were positive to GVA and
14
15 GLRaV-1 in real-time RT-PCR assay (that targeted the coat protein gene) did not provide detectable
16
17 signals in the qRT-PCR assay. This latter is less sensitive because it targets the single copy *RdRp*
18
19 gene, that has no subgenomic RNAs unlike coat protein gene.
20
21
22
23

24 In conclusion, this study provides new insights into the mealybug-mediated transmission of GLRaVs
25
26 and RW- associated viruses. It is already known that this transmission lacks of vector-virus
27
28 specificity, since a same insect species can transmit different viruses and a same virus can be
29
30 transmitted by many vector species (Tsai *et al.*, 2010; Le Maguet *et al.*, 2012a). However, the
31
32 relative efficiency of mealybugs towards GLRaVs and RW-associated viruses remained largely
33
34 unknown because of the low number of comparative transmission studies. This study steps
35
36 forward the question and deals with different sides of the story at a time: the “virus side”, by
37
38 comparing transmission patterns of different co-infecting viruses, and the “vector side”, by
39
40 comparing the transmission efficiency of two different mealybug species. Looking at the vectors,
41
42 nymphs of *Pl. ficus* and *Pl. citri* that fed on the same source plant showed similar acquisition but
43
44 different transmission patterns of GVA, GLRaV-1 as well as of GLRaV-3. Indeed *Pl. ficus* was mainly
45
46 efficient in GLRaV-3 and GVA transmission, whereas *Pl. citri* was efficient in GLRaV-1 transmission
47
48 and inefficient in GVA transmission. The “virus side” would provide information about possible
49
50 antagonisms or synergisms between GLRaVs and vitiviruses. We observed that GVA could be
51
52 transmitted to grapevine without GLRaV-1 and/or GLRaV-3 by *Pl. ficus* but not by *Pl. citri* and that
53
54
55
56
57
58
59
60

1
2
3 the GLRaVs transmission could take place also in absence of GVA.
4

5 Virus-vector relationships, together with new insights into the genetic characterization of the virus
6
7 strains, may allow a better interpretation of GLD and RW epidemiology, and contribute to the
8
9 development of control strategies against these virus-associated diseases.
10
11

12 13 14 **Acknowledgements**

15
16
17 The authors thank dr Ivana Gribaudo, IPSP-CNR, for providing *in vitro* grapevine used as test plants.
18

19
20 This study was funded by the Piemonte Region under the grant “Spread of viral re-infections in the
21
22 vineyard and role of mealybug vectors”.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Adams M. J., Antoniw J. F., Bar-Joseph M., Brunt A. A., Candresse T., Foster G. D., Martelli G. P., Milne R. G., Fauquet C. M. (2004) The new plant virus family Flexiviridae and assessment of molecular criteria for species demarcation. *Archives of Virology*, **149**, 1045-1060.
- Almeida R. P. P., Daane K. M., Bell V. A., Blaisdell G. K., Cooper M. L., Herrbach E., Pietersen G. (2013) Ecology and management of grapevine leafroll disease. *Frontiers in Microbiology*, **4**.
- Andersen C. L., Jensen J. L., Orntoft T. F. (2004) Normalization of real-time quantitative reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Research*, **64**, 5245-5250.
- Ben-Dov Y. M., D.R. Gibson, G.A.P (2001) *ScaleNet, Life Histories* [Online]. [Accessed 08/13 2015].
- Bertin S., Cavalieri V., Graziano C., Bosco D. (2010) Survey of mealybug (Hemiptera: Pseudococcidae) vectors of Ampelovirus and Vitivirus in vineyards of northwestern Italy. *Phytoparasitica*, **38**, 401-409.
- Blaisdell G. K., Zhang S., Bratburd J. R., Daane K. M., Cooper M. L., Almeida R. P. P. (2015) Interactions Within Susceptible Hosts Drive Establishment of Genetically Distinct Variants of an Insect-Borne Pathogen. *Journal of Economic Entomology*, **108**, 1531-1539.
- Blaisdell K., SimingDaane, KentAlmeida, Rodrigo P P (2012) Patterns of virus transmission from hosts with mixed infections. In: 17th Congress of the International Council for the Study of Virus and Virus-like Diseases of the Grapevine (ICVG), 2012 Ed B. Ferguson Davis, California, USA. Foundation Plant Services, University of California, Davis, pp. 178-179.
- Cabaleiro C., Couceiro C., Pereira S., Cid M., Barrasa M., Segura A. (2008) Spatial analysis of epidemics of Grapevine leafroll associated virus-3. *European Journal of Plant Pathology*, **121**, 121-130.
- Cabaleiro C., Segura A. (1997a) Field transmission of grapevine leafroll associated virus 3 (GLRaV-3) by the mealybug *Planococcus citri*. *Plant Disease*, **81**, 283-287.
- Cabaleiro C., Segura A. (1997b) Some characteristics of the transmission of grapevine leafroll associated virus 3 by *Planococcus citri* Risso. *European Journal of Plant Pathology*, **103**, 373-378.
- Cabaleiro C., Segura A. (2006) Temporal analysis of grapevine leafroll associated virus 3 epidemics. *European Journal of Plant Pathology*, **114**, 441-446.
- Cavalieri V., Mazzeo G., Garzia G. T., Buonocore E., Russo A. (2008) Identification of *Planococcus ficus* and *Planococcus citri* (Hemiptera: Pseudococcidae) by PCR-RFLP of COI gene. *Zootaxa*, **1816**, 65-68.
- Cid M., Pereira S., Cabaleiro C., Faoro F., Segura A. (2007) Presence of Grapevine leafroll-associated virus 3 in primary salivary glands of the mealybug vector *Planococcus citri* suggests a circulative transmission mechanism. *European Journal of Plant Pathology*, **118**, 23-30.
- Cid M., Pereiro S., Cabaleiro C., Segura A. (2010) Citrus mealybug (Hemiptera: Pseudococcidae) movement and population dynamics in an arbor-trained vineyard. *Journal of Economic Entomology*, **103**, 619-630.
- Daane K. M., Middleton M. C., Sforza R., Cooper M. L., Walton V. M., Walsh D. B., Zaviezo T., Almeida R. P. P. (2011) Development of a multiplex PCR for identification of vineyard mealybugs. *Environmental Entomology*, **40**, 1595-1603.
- Debreczeni D. E., Ruiz-Ruiz S., Aramburu J., Lopez C., Belliure B., Galipienso L., Soler S., Rubio L. (2011) Detection, discrimination and absolute quantitation of Tomato spotted wilt virus isolates using real time RT-PCR with TaqMan (R) MGB probes. *Journal of Virological Methods*, **176**, 32-37.

- 1
2
3 Douglas N., Krüger K. (2008) Transmission efficiency of Grapevine leafroll-associated virus 3
4 (GLRaV-3) by the mealybugs *Planococcus ficus* and *Pseudococcus longispinus* (Hemiptera:
5 Pseudococcidae). *European Journal of Plant Pathology*, **122**, 207-212.
- 6 Engelbrecht D. J., Kasdorf G. G. F. (1990) Transmission of grapevine leafroll disease and associated
7 closteroviruses by the vine mealybug, *Planococcus ficus*. *Phytophylactica*, **22**, 341–346.
- 8 Fortusini A. G., Scattini G., Prati S., Cinquanta S., Belli G. (1997) Transmission of *Grapevine leafroll*
9 *virus-1* (GLRaV-1) and *Grapevine virus A* (GVA) by scale insects. In: 12th Meeting of ICVG,
10 1997 Lisbon, Portugal. pp. 121–122.
- 11 Fuchs M., Marsella-Herrick P., Loeb G. M., Martinson T. E., Hoch H. C. (2009) Diversity of
12 ampeloviruses in mealybug and soft scale vectors and in grapevine hosts from leafroll-
13 affected vineyards. *Phytopathology*, **99**, 1177-1184.
- 14 Gambino G., Gribaudo I. (2006) Simultaneous detection of nine grapevine viruses by multiplex
15 reverse transcription-polymerase chain reaction with coamplification of a plant RNA as
16 internal control. *Phytopathology*, **96**, 1223-1229.
- 17 Giribaldi M., Purrotti M., Pacifico D., Santini D., Mannini F., Caciagli P., Rolle L., Cavallarin L.,
18 Giuffrida M. G., Marzachi C. (2011) A multidisciplinary study on the effects of phloem-
19 limited viruses on the agronomical performance and berry quality of *Vitis vinifera* cv.
20 Nebbiolo. *Journal of proteomics*, **75**, 306-315.
- 21 Golino D. A., Sim S. T., Gill R., Rowhani A. (2002) Grapevine leafroll disease can be spread by
22 California mealybugs. *California Agriculture*, **56**, 196–201.
- 23 Golino D. A., Weber E., Sim S., Rowhani A. (2008) Leafroll disease is spreading rapidly in a Napa
24 Valley vineyard. *California Agriculture*, **62**, 156-160.
- 25 Gribaudo I., Gambino G., Bertin S., Bosco D., Cotroneo A., Mannini F. (2009) Monitoring the spread
26 of viruses after vineyard replanting with heat-treated clones of *Vitis Vinifera* 'Nebbiolo'.
27 *Journal of Plant Pathology*, **91**, 741-744.
- 28 Hommay G., Komar V., Lemaire O., Herrbach E. (2008) Grapevine virus A transmission by larvae of
29 *Parthenolecanium corni*. *European Journal of Plant Pathology*, **121**, 185-188.
- 30 Kol-Maimon H., Ghanim M., Franco J. C., Mendel Z. (2014) Evidence for gene flow between two
31 sympatric mealybug species (Insecta; Coccoidea; Pseudococcidae). *PLoS ONE*, **9**.
- 32 Le Maguet J., Beuve M., Herrbach E., Lemaire O. (2012a) Transmission of Six Ampeloviruses and
33 Two Vitiviruses to Grapevine by *Phenacoccus aceris*. *Phytopathology*, **102**, 717-723.
- 34 Le Maguet J., Fuchs J. J., Chadœuf J., Beuve M., Herrbach E., Lemaire O. (2012b) The role of the
35 mealybug *Phenacoccus aceris* in the spread of Grapevine leafroll-associated virus -1
36 (GLRaV-1) in two French vineyards. *European Journal of Plant Pathology*, **135**, 415-427.
- 37 MacKenzie D. J., McLean M. A., Mukerji S., Green M. (1997) Improved RNA extraction from woody
38 plants for the detection of viral pathogens by reverse transcription-polymerase chain
39 reaction. *Plant Disease*, **81**, 222-226.
- 40 Mahfoudhi N., Digiario M., Dhouibi M. H. (2009) Transmission of grapevine leafroll viruses by
41 planococcus ficus (Hemiptera: Pseudococcidae) and *Ceroplastes Rusci* (Hemiptera:
42 Coccidae). *Plant Disease*, **93**, 999-1002.
- 43 Malausa T., Fenis A., Warot S., Germain J., Ris N., Prado E., Botton M., Vanlerberghe-Masutti F.,
44 Sforza R., Cruaud C., Couloux A., Kreiter P. (2011) DNA markers to disentangle complexes of
45 cryptic taxa in mealybugs (Hemiptera: Pseudococcidae). *Journal of Applied Entomology*,
46 **135**, 142-155.
- 47 Maliogka V. I., Dovas C. I., Katis N. I. (2008) Evolutionary relationships of virus species belonging to
48 a distinct lineage within the Ampelovirus genus. *Virus Research*, **135**, 125-135.
- 49 Maree H. J., Almeida R. P. P., Bester R., Chooi K. M., Cohen D., Dolja V. V., Fuchs M. F., Golino D. A.,
50 Jooste A. E. C., Martelli G. P., Naidu R. A., Rowhani A., Saldarelli P., Burger J. T. (2013)

- 1
2
3 Grapevine leafroll-associated virus 3. *Frontiers in Microbiology*, **4**.
- 4 Maroniche G. A., Sagadin M., Mongelli V. C., Truol G. A., del Vas M. (2011) Reference gene selection
5 for gene expression studies using RT-qPCR in virus-infected planthoppers. *Virology Journal*,
6 **8**.
- 7 Martelli G. P. (2014a) Grapevine leafroll. *Journal of Plant Pathology*, **96**, S51-S70.
- 8 Martelli G. P. (2014b) Rugose wood complex. *Journal of Plant Pathology*, **96**, S73-S88.
- 9 Martelli G. P., Abou Ghanem-Sabanadzovic N., Agranovsky A. A., Al Rwahnih M., Dolja V. V., Dovas
10 C. I., Fuchs M., Gugerli P., Hu J. S., Jelkmann W., Katis N. I., Maliogka V. I., Melzer M. J.,
11 Menzel W., Minafra A., Rott M. E., Rowhani A., Sabanadzovic S., Saldarelli P. (2012)
12 Taxonomic revision of the family Closteroviridae with special reference to the Grapevine
13 Leafroll-associated members of the genus Ampelovirus and the putative species unassigned
14 to the family. *Journal of Plant Pathology*, **94**, 7-19.
- 15
16
17 Martin S., Elena S. F. (2009) Application of game theory to the interaction between plant viruses
18 during mixed infections. *Journal of General Virology*, **90**, 2815-2820.
- 19 Marzachi C., Bosco D. (2005) Relative quantification of chrysanthemum yellows (16Sr I)
20 phytoplasma in its plant and insect host using real-time polymerase chain reaction.
21 *Molecular Biotechnology*, **30**, 117-127.
- 22
23 Mason G., Caciagli P., Accotto G. P., Noris E. (2008) Real-time PCR for the quantitation of Tomato
24 yellow leaf curl Sardinia virus in tomato plants and in Bemisia tabaci. *Journal of Virological
25 Methods*, **147**, 282-289.
- 26
27 Naidu R., Rowhani A., Fuchs M., Golino D., Martelli G. P. (2014) Grapevine Leafroll: A complex viral
28 disease affecting a high-value fruit crop. *Plant Disease*, **98**, 1172-1185.
- 29 Ng J. C. K., Falk B. W. (2006) Virus-vector interactions mediating nonpersistent and semipersistent
30 transmission of plant viruses. *In: Annual Review of Phytopathology*, **44**, 183-212.
- 31 Ng J. C. K., Tian T., Falk B. W. (2004) Quantitative parameters determining whitefly (Bemisia tabaci)
32 transmission of Lettuce infectious yellows virus and an engineered defective RNA. *Journal
33 of General Virology*, **85**, 2697-2707.
- 34
35 Osman F., Olineka T., Hodzic E., Golino D., Rowhani A. (2012) Comparative procedures for sample
36 processing and quantitative PCR detection of grapevine viruses. *Journal of Virological
37 Methods*, **179**, 303-310.
- 38
39 Pacifico D., Caciagli P., Palmano S., Mannini F., Marzachi C. (2011) Quantitation of Grapevine
40 leafroll associated virus-1 and -3, Grapevine virus A, Grapevine fanleaf virus and Grapevine
41 fleck virus in field-collected Vitis vinifera L. 'Nebbiolo' by real-time reverse transcription-
42 PCR. *Journal of Virological Methods*, **172**, 1-7.
- 43
44 Petersen C. L., Charles J. G. (1997) Transmission of grapevine leafroll-associated closteroviruses by
45 Pseudococcus longispinus and P. calceolariae. *Plant Pathology*, **46**, 509-515.
- 46 Pfaffl M. W., Tichopad A., Prgomet C., Neuvians T. P. (2004) Determination of stable housekeeping
47 genes, differentially regulated target genes and sample integrity: BestKeeper - Excel-based
48 tool using pair-wise correlations. *Biotechnology Letters*, **26**, 509-515.
- 49 Rosciglione B., Castellano M. A. (1985) Further evidence that mealybugs can transmit grapevine
50 virus A (GVA) to herbaceous hosts. *Phytopathologia Mediterranea*, **24**, 186-188.
- 51 Rotenberg D., Krishna Kumar N. K., Ullman D. E., Montero-Astua M., Willis D. K., German T. L.,
52 Whitfield A. E. (2009) Variation in Tomato spotted wilt virus Titer in Franklaniella
53 occidentalis and Its Association with Frequency of Transmission. *Phytopathology*, **99**, 404-
54 410.
- 55
56 Saccaggi D. L., Krüger K., Pietersen G. (2008) A multiplex PCR assay for the simultaneous
57 identification of three mealybug species (Hemiptera: Pseudococcidae). *Bulletin of
58 Entomological Research*, **98**, 27-33.
- 59
60

- 1
2
3 Scotto C., Conti B., Panattoni A., Triolo E. (2009) Acquisition and transmission of GLRaV-1
4 (Grapevine leafroll associated ampelovirus 1) by *Planococcus citri* (Risso) (Hemiptera
5 Pseudococcidae). In: 16th meeting of ICVG, 2009 Ed E. Boudon-Padieu Dijon, France. Le
6 progrès agricole et viticole, pp. 322, 323.
- 7
8 Sforza R., Boudon-Padieu E., Greif C. (2003) New mealybug species vectoring Grapevine leafroll-
9 associated viruses-1 and -3 (GLRaV-1 and -3). *European Journal of Plant Pathology*, **109**,
10 975-981.
- 11
12 Sharma A. M., Wang J. B., Duffy S., Zhang S. M., Wong M. K., Rashed A., Cooper M. L., Daane K. M.,
13 Almeida R. P. P. (2011) Occurrence of Grapevine Leafroll-Associated Virus Complex in Napa
14 Valley. *Plos One*, **6**, 7.
- 15
16 Swallow W. H. (1985) Group testing for estimating infection rates and probabilities of disease
17 transmission. *Phytopathology*, **75**, 882-889.
- 18
19 Tsai C. W., Bosco D., Daane K. M., Almeida R. P. P. (2011) Effect of host plant tissue on the vector
20 transmission of grapevine leafroll-associated virus 3. *Journal of Economic Entomology*, **104**,
21 1480-1485.
- 22
23 Tsai C. W., Chau J., Fernandez L., Bosco D., Daane K. M., Almeida R. P. P. (2008) Transmission of
24 Grapevine leafroll-associated virus 3 by the Vine Mealybug (*Planococcus ficus*).
25 *Phytopathology*, **98**, 1093-1098.
- 26
27 Tsai C. W., Daugherty M. P., Almeida R. P. P. (2012) Seasonal dynamics and virus translocation of
28 Grapevine leafroll-associated virus 3 in grapevine cultivars. *Plant Pathology*, **61**, 977-985.
- 29
30 Tsai C. W., Rowhani A., Golino D. A., Daane K. M., Almeida R. P. P. (2010) Mealybug transmission of
31 grapevine leafroll viruses: An analysis of virus-vector specificity. *Phytopathology*, **100**, 830-
32 834.
- 33
34 Vandesompele J., De Preter K., Pattyn F., Poppe B., Van Roy N., De Paepe A., Speleman F. (2002)
35 Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of
36 multiple internal control genes. *Genome biology*, **3**, RESEARCH0034.
- 37
38 Velasco L., Bota J., Montero R., Cretazzo E. (2014) Differences of three ampeloviruses'
39 multiplication in plant may explain their incidences in vineyards. *Plant Disease*, **98**, 395-400.
- 40
41 Voncina D., Badurina D., Preiner D., Cvjetkovic B., Maletic E., Kontic J. K. (2011) Incidence of virus
42 infections in grapevines from Croatian collection plantations. *Phytopathologia
43 Mediterranea*, **50**, 316-326.
- 44
45 Wintermantel W. M., Cortez A. A., Anchieta A. G., Gulati-Sakhuja A., Hladky L. L. (2008) Co-
46 Infection by Two Criniviruses Alters Accumulation of Each Virus in a Host-Specific Manner
47 and Influences Efficiency of Virus Transmission. *Phytopathology*, **98**, 1340-1345.
- 48
49 Zorloni A., Prati S., Bianco P. A., Belli G. (2006) Transmission of grapevine virus A and grapevine
50 leafroll-associated virus 3 by *Heliococcus bohemicus*. *Journal of Plant Pathology*, **88**, 325-
51 328.
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 1** Pattern of virus retention in *Planococcus ficus* (a) and *Pl. citri* (b) after a 24h acquisition
4 access period (AAP). Proportion of nymphs positive to *Grapevine virus A* (GVA), *Grapevine leafroll-*
5 *associated virus 1* (GLRaV-1) and *Grapevine leafroll-associated virus 3* (GLRaV-3) at different post-
6 acquisition times (T_0 , T_{24} , T_{48} , T_{72} , T_{96}).
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1 *Actin β (ACTB)*, *glyceraldehyde 3-phosphate dehydrogenase (GAPDH)* and *18S ribosomal RNA (18S rRNA)* genes of *Planococcus ficus* and *Pl. citri*: amplification systems and expression stability. Sequences of the primers used for *ACTB*, *GAPDH* and *18S rRNA* amplification in SYBR[®] Green Real-Time PCR; sequence of the probe used for *GAPDH* amplification in TaqMan[®] qRT-PCR. Ranking of gene stability derived from GeNorm, Bestkeeper and Normfinder analysis. GeNorm: M values; Bestkeeper: Correlation coefficient, $p = 0.001$; Normfinder: Stability value and Standard error (SE).

| Genes | Amplification systems | | Expression stability | | |
|-----------------|-----------------------|-------------------------------|----------------------|------------|--------------------|
| | | | GeNorm | Bestkeeper | Normfinder |
| <i>ACTB</i> | PfActFw2 | 5'-TGAYTTAACCGACTACTTGA-3' | 1.287 | 0.965 | 0.203 (SE = 0.060) |
| | PfActBRev1 | 5'-TCCAAAGCGACATAGCAC-3' | | | |
| <i>GAPDH</i> | PfGapFw1 | 5'-AAGAAAGTYATCATCTCAGC-3' | 1.035 | 0.985 | 0.148 (SE = 0.069) |
| | PfGapRev | 5'-GCGTGACGGTGGTCATTA-3' | | | |
| | PfGapProbe | FAM-ATGTACGTCTGCGGTGTCAA-BHQ1 | | | |
| <i>18S rRNA</i> | MqFw* | 5'-AACGGCTACCACATCCAAGG-3' | 1.312 | 0.982 | 0.251 (SE = 0.059) |
| | MqRv* | 5'-GCCTCGGATGAGTCCCG-3' | | | |

*from Marzachi and Bosco (2005)

Table 2 Rate of *Planococcus ficus* and *Pl. citri* first-instar nymphs positive to *Grapevine virus A* (GVA), *Grapevine leafroll-associated virus 1* (GLRaV-1) and *Grapevine leafroll-associated virus 3* (GLRaV-3) in single and mixed infections after a 24h-AAP on triple-infected grapevine; rate of test grapevines positive to GVA, GLRaV-1 and GLRaV-3 in single and mixed infections after a 48h-IAP by *Pl. ficus* and *Pl. citri*. Virus detection was performed by SYBR[®] Green real-time RT-PCR.

| | <i>Pl. ficus</i> | | <i>Pl. citri</i> | |
|-------------------------|--|--|--|--|
| | Positive nymphs (n = 50 ^a) | Positive grapevines (n = 27 ^b) | Positive nymphs (n = 46 ^c) | Positive grapevines (n = 34 ^d) |
| GVA | 0.20 | 0.07 | 0.13 | - |
| GLRaV-1 | 0.10 | 0.04 | 0.11 | 0.53 |
| GLRaV-3 | 0.12 | 0.22 | 0.02 | 0.24 |
| GVA + GLRaV-1 | 0.22 | - | 0.09 | - |
| GVA + GLRaV-3 | 0.22 | 0.41 | 0.17 | 0.06 |
| GLRaV-1 + GLRaV-3 | 0.04 | 0.11 | 0.04 | 0.15 |
| GVA + GLRaV-1 + GLRaV-3 | 0.10 | 0.15 | 0.43 | 0.03 |
| Total GVA | 0.74 | 0.63 | 0.83 | 0.09 |
| Total GLRaV-1 | 0.46 | 0.30 | 0.67 | 0.70 |
| Total GLRaV-3 | 0.48 | 0.89 | 0.67 | 0.47 |

^a Total number of *Pl. ficus* nymphs positive to at least one virus, out of the 71 tested samples.

^b Total number of grapevines positive to at least one virus after inoculation by *Pl. ficus*, out of the 51 tested samples.



^c Total number of *Pl. citri* nymphs positive to at least one virus, out of the 74 tested samples.

^d Total number of grapevines positive to at least one virus after inoculation by *Pl. citri*, out of the 49 tested samples.

Table 3 Mean loads of *Grapevine virus A* (GVA) and *Grapevine leafroll-associated virus 1* (GLRaV-1) in *Planococcus ficus* and *Pl. citri* at different post-acquisition times. The mean loads are expressed as viral genome units per 100,000 insect *GAPDH* transcripts \pm the Standard Error. *Grapevine leafroll-associated virus 3* (GLRaV-3) titre was below the quantification threshold in all the 6 *Pl. ficus* and 6 *Pl. citri* individuals that tested positive in SYBR[®] Green real-time RT-PCR. *n*: number of quantified samples.

| Post-acquisition times | Total samples | GVA mean loads (viral GU/100,000 <i>GAPDH</i> \pm SE) | | GLRaV1 mean loads (viral GU/100,000 <i>GAPDH</i> \pm SE) | |
|------------------------|---------------|--|-----------------------------------|---|-------------------------------------|
| | | <i>Pl. ficus</i> | <i>Pl. citri</i> | <i>Pl. ficus</i> | <i>Pl. citri</i> |
| T ₀ | 16 | 1.07 (\pm 1.5) <i>n</i> = 5 | 2.34 (\pm 1.1) <i>n</i> = 4 | 400 (\pm 521.0) <i>n</i> = 7 | |
| T ₂₄ | 4 | | | 1,170 (\pm 893.0) <i>n</i> = 2 | 84.00 (\pm 69.0) <i>n</i> = 2 |
| T ₄₈ | 3 | 2.91 (\pm 0.69) <i>n</i> = 2 | | 8,670 <i>n</i> = 1 | |
| T ₇₂ | 1 | | | | 83.1 <i>n</i> = 1 |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



83x146mm (300 x 300 DPI)