

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Comparison of fortified, sfursat, and passito wines produced from fresh and dehydrated grapes of aromatic black cv. Moscato nero (Vitis vinifera L.)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1641550> since 2019-08-25T15:28:12Z

Published version:

DOI:10.1016/j.foodres.2016.11.012

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

Food Research International 98 (2017) 59–67;

<https://doi.org/10.1016/j.foodres.2016.11.012>

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>),
<http://www.sciencedirect.com/science/article/pii/S0963996916305518>

Comparison of *fortified*, *sfursat*, and *passito* wines produced from fresh and dehydrated grapes of aromatic black cv. Moscato nero (*Vitis vinifera* L.)

Carolina Ossola^{1#}, Simone Giacosa^{1#*}, Fabrizio Torchio², Susana Río Segade¹, Alberto Caudana¹, Enzo Cagnasso¹, Vincenzo Gerbi¹, Luca Rolle¹

¹ Università degli Studi di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari. Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy.

² Università Cattolica del Sacro Cuore, Istituto di Enologia e Ingegneria Agro-Alimentare. Via Emilia Parmense 84, 29122 Piacenza, Italy.

* Corresponding author. E-mail: simone.giacosa@unito.it

These authors contributed equally to the study.

ABSTRACT

Moscato nero d'Acqui is an Italian aromatic black winegrape variety characterized by a low content of anthocyanins (mostly tri-substituted), a satisfactory content of high molecular mass tannins, and a fair amount of terpenes. The grapes were subjected to a postharvest dehydration process under controlled thermohygrometric conditions (16-18 °C, 55-70 RH%, 0.6 m/s air speed). with the aim to produce three different special wine types (*fortified*, *sfursat*, and *passito*) from fresh, partially dehydrated (27 °Brix), and withered (36 °Brix) grapes, respectively. Chemical traits of produced grapes and wines were then evaluated through spectrophotometric, HPLC, and GC-MS methods. Increased contents of skin phenolic compounds and reduced extractable contents of seed phenolic compounds were observed as dehydration progressed. Few significant differences were found in the anthocyanin profile of grapes, although the relative abundance of coumaroylated anthocyanins was higher in dehydrated grapes. The predominant free volatile compound found in grapes was geraniol, which decreased with increasing water loss, whereas the contents of major glycosylated volatile compounds increased even above the concentration effect. The changes in the phenolic composition among wines agreed with those among grape skins. *Fortified* wines were chromatically unsatisfactory probably due to the low content of total anthocyanins, whereas *sfursat* and *passito* wines meet good chromatic characteristics as a result of the concentration effect during grape dehydration. *Fortified* and *sfursat* wines had free aroma profiles richer in 2-phenylethanol and citronellol, whereas *passito* wines were mainly composed of 2-phenylethanol and 2-phenylethyl acetate, citronellol being the predominant terpenol in all the wine types studied.

Keywords: Moscato nero d'Acqui; Phenolic compounds; Free and bound volatile compounds; Dehydrated grapes; Special red wines.

1. INTRODUCTION

Nowadays, one of the key factors of the economical development of the viticulture and wine industry in specific limited geographical areas is the exploitation of ancient, local grape varieties for the development of oenological products with special features. In recent years, the growing interest to rediscover ancient and minor varieties has promoted many studies in order to retrain these realities (Maul et al., 2015; Urcan et al., 2016). This is particularly true in Italy, a country with a rich varietal endowment due to a centuries-old wine-growing tradition.

Vitis vinifera L. cv. Moscato nero d'Acqui is an ancient and local aromatic red grape variety, which could be found sporadically in old vineyards in the provinces of Asti and Alessandria, and takes its name from the town of Acqui, located at the center of this production zone (North-West Italy). The Moscato nero d'Acqui cultivar was described for the first time in 1875 (De Maria & Leardi 1875), while in 1971 it was included in the National register of vine varieties (Mipaaf). Nowadays, the spread of this variety is limited, but new vineyards are being planted in Tortona (South-East Piedmont; Raimondi, Valota, & Schneider, 2009) and the nursery production is active (Pecile, Zavaglia, & Ciardi, 2016). The Moscato nero d'Acqui cultivar can be called simply Moscato nero. It is not a synonym of Moscato d'Amburgo (Muscat Hamburg, one of the most known table grape varieties), even though the two cultivars belong to the large family of Muscat vines that have in common the characteristic Muscat flavor. Monoterpenes are responsible for the varietal aroma of Muscat cultivars (Selli, Canbas, Cabaroglu, Erten, & Gunata, 2006). In the past, Moscato nero d'Acqui grapes were consumed as table grapes, but the oenological potential was noted (Mannini et al. 2012). From the comparison among 34 *Vitis vinifera* genotypes emerges that the studied cultivar is characterized by low content of total anthocyanins, although the accumulation of these compounds is significantly year-dependent (Ferrandino, Carra, Rolle, Schneider & Schubert, 2012). The vine is known by low productivity and good vigor. The bunches are loose and usually resistant to rot, which makes them suitable to the dehydration process used for special (sweet) wines production.

During postharvest grape dehydration, important metabolic changes occur due to water loss and, in specific conditions, to the development of *Botrytis cinerea* (as noble rot). Water stress induces an active metabolism that affects the chemical composition and physical properties of grape berries (Rolle et al., 2013, Toffali et al. 2011), while *Botrytis cinerea*, when developing as noble rot, favors the release of aroma compounds from the berry skin (Genovese, Gambuti, Piombino, & Moio, 2007) and other skin modifications (Carbajal-Ida, Maury, Salas, Siret, & Mehinagic, 2016; Rolle et al., 2012). Therefore, the dehydration process promotes the release, concentration, synthesis, and oxidation of phenolic and volatile compounds (Bellincontro, De Santis, Botondi, Villa, & Mencarelli, 2004; Costantini, Bellincontro, De Santis, Botondi, & Mencarelli, 2006; Mencarelli et al., 2010), although the varietal aroma of winegrapes is particularly influenced in botrytized berries because of the mould-induced oxidation of monoterpenes (Cámara, Herbert, Marques, & Alves, 2004). In fact, the interaction between the level of *B. cinerea* infection and the degree of grape withering is of great relevance for modulating the aroma of *passito* wines (Tosi et al., 2013). Furthermore, the increased production of glycerol enhances the wine mouthfeel (Rolle et al., 2012; Vincenzi et al., 2012).

To gain knowledge on the enological potential of the Moscato nero d'Acqui variety, the two main aims of this work were: *i)* to investigate the phenolic and aromatic composition of fresh grapes and dehydrated grapes using two different postharvest treatment times, *ii)* to evaluate the aptitude to the production of special wines, particularly *fortified* (sweet wine from fresh grapes), *sfursat* (dry wine from partially dehydrated grapes), and *passito* (sweet wine from dehydrated grapes). *Fortified* wine belongs to the category of liqueur wines that are made from grape musts (including partially fermented grape musts) and/or wine, to which distillates, spirits and alcohol of vitivinicultural origin are added alone or in a mixture (OIV, ECO 2/2007). *Sfursat* and *passito* wines are produced

by using partially dehydrated and withered grapes, respectively. The knowledge of the enological potential and special features of Moscato nero d'Acqui would allow the better exploitation of this particular red aromatic variety enhancing the distinctive character of special wines.

2. MATERIALS AND METHODS

2.1. Grape and wine samples

Red grapes of *Vitis vinifera* L. cv. Moscato nero d'Acqui were harvested at ripeness (about 200 g/L must sugars concentration) in a vineyard located in Santo Stefano Belbo (Cuneo province, Piedmont, North-West Italy) in September 2013. Three sets of small clusters were selected (3 replicates for fresh grape sample). Six batches of 125 kg of grapes (3 replicates for each of the two withered grape samples) were placed in a single layer in plastic boxes (60 cm x 40 cm x 15 cm, with bottom holes of about 1.5 cm x 5 cm to improve air flow), in quantities of about 1.5 kg grapes for each box, and dehydrated in a thermohygrometrically controlled chamber (16-18 °C, 55-70% relative humidity, 0.6 m/s air speed). The dehydration process was carried out for 23 days for the production of *sfursat* wine (until reaching 27 °Brix, "partially dehydrated" grapes sample) and for 52 days for the production of *passito* wine (until reaching 36 °Brix, "withered" grapes sample).

Before the analysis of each sample, the randomly-taken berries were manually separated from the stalk and then used as follows: three replicates of 100 berries for the determination of technological ripeness parameters, three replicates of 10 berries for phenolic compound determination, and three replicates of 150 berries for volatile compound determination.

Micro-scale vinifications were made at the experimental winery of the University of Torino. For each type of wine (*fortified*, *sfursat*, and *passito*), two replicates of about 100 kg of grapes each were processed. The vinification protocol was the same for all the six trials (2 replicates × 3 wines) but using fresh grapes for *fortified* wine, partially dehydrated grapes for *sfursat* wine, and withered grapes for *passito* wine. The grapes were destemmed and crushed, the mash was then placed into a fermenter saturated with CO₂, and 20 mg/L of sulfur dioxide were added. After about 6 hours, 20 g/hL of yeast (LSA ES181 previously activated, Esseco, Trecate, IT), and 8 g/hL of yeast nutrients (ammonium sulfate and ammonium phosphate) were added. Alcoholic fermentation was carried out at controlled temperature (22±2 °C) until reaching 5% v/v of ethanol. The pomace was pressed and the resulting must-wine continued the fermentation without the solid parts. In the case of *fortified* wine, the fermentation was stopped, when the residual sugar content in the fermenting must was about 100 g/L, with the addition of 100 mg/L of sulfur dioxide and 95% v/v food-grade ethanol up to a total alcohol content of about 13% v/v, and the resulting wine was stored at 10 °C for one month. In the case of *sfursat* wine, the fermentation progressed until less than 5 g/L of sugars were present and 50 mg/L of sulfur dioxide were added, while in *passito* wine the fermentation was stopped when the residual sugar content was about 110 g/L by adding 100 mg/L of sulfur dioxide. *Fortified*, *sfursat* and *passito* wines were stored at 0 °C for 2 weeks, filtered (Seitz K300 grade filter sheets, Pall Corporation, Port Washington, NY, USA), and finally bottled.

2.2. Chemical analysis of grapes and wines

2.2.1 Reagents and standards

Solvents of HPLC-gradient grade and all other chemicals of analytical-reagent grade were purchased from Sigma-Aldrich (Milan, Italy). The solutions were prepared in deionized water produced by a Milli-Q system (Merck Millipore, Darmstadt, DE). Chemical standards of delphinidin-3-O-glucoside chloride, cyanidin-3-O-glucoside chloride, petunidin chloride, peonidin-3-O-glucoside chloride, malvidin-3-O-glucoside chloride, and cyanidin chloride were supplied by Extrasynthèse (Genay, France), whereas those of (+)-catechin, linalool oxide (mixture of isomers), linalool, α -terpineol, citral, citronellol, nerol, geraniol, rose oxide (mixture of isomers), geranyl acetate, geranic acid, benzyl alcohol, 2-phenylethanol, and 2-phenylethyl acetate were purchased from Sigma-Aldrich.

2.2.2 Standard parameters

In the grape musts resulting from manual grape crushing and centrifugation, total soluble solids content (°Brix) was measured using an Atago °Brix temperature compensating refractometer (Atago Corporation, Tokyo, Japan). In the grape musts and in the wines obtained after five months from bottling, reducing sugars (glucose and fructose, g/L) and organic acids (tartaric acid and malic acid, g/L) were determined by high performance liquid chromatography (HPLC) (Agilent Technologies, Santa Clara, CA, USA) using a refractive index detector and a diode array detector (DAD) set to 210 nm (Giordano, Rolle, Zeppa, & Gerbi, 2009). Citric acid (g/L) in grape musts, and glycerol (g/L) and ethanol (% v/v) in wines were determined following the same HPLC methodology. pH was determined by potentiometry using an InoLab 730 pH meter (WTW, Weilheim, Germany), and titratable acidity (g/L tartaric acid) and volatile acidity (g/L acetic acid) were estimated according to the International Organization of Vine and Wine methods (OIV, 2008). Gluconic acid (g/L) in grape musts was determined using an enzymatic kit (R-Biopharm Italia, Cerro al Lambro, MI, Italy) and a UV-1800 spectrophotometer (Shimadzu Corporation, Kyoto, Japan).

2.2.3 Extraction and determination of phenolic compounds

The grape berries were weighed, and the skins and seeds were manually separated from the pulp using a laboratory spatula. The berry skins were quickly put into 25 mL of a hydroalcoholic buffer solution of pH 3.2 composed of 5 g/L of tartaric acid, 2 g/L of sodium metabisulfite, and 12% v/v of ethanol (Torchio, Cagnasso, Gerbi, & Rolle, 2010). The skins were homogenized for 1 minute at 8000 rpm with an Ultra-Turrax T25 high-speed homogenizer (IKA Labortechnik, Staufen, Germany) and subsequently centrifuged in a PK 131 centrifuge (ALC International, MI, Italy) for 15 minutes at 3000 x g and 20 °C. The supernatant was then used for skin analysis. The berry seeds were put into 10 mL of the same buffer solution used for the skins and then placed in an oven at 30 °C for 1 week (Torchio et al., 2010). Afterwards, the seeds were discarded and the solution was then used for seed analysis.

Spectrophotometric methods were used to determine total anthocyanins (mg malvidin-3-O-glucoside chloride/kg grape or L wine, as TA), absorbance at 280 nm (as A_{280} /kg grape or L wine), total flavonoids [mg (+)-catechin/kg grape or L wine, as TF], flavanols reactive to vanillin [mg (+)-catechin/kg grape or L wine, as FRV], and proanthocyanidins (mg cyanidin chloride/kg grape or L wine, as PRO) (Torchio et al., 2010). A UV-1800 spectrophotometer (Shimadzu Corporation) was used.

The anthocyanin profile of berry skins and wines was determined by HPLC-DAD following the protocol described by Rolle, Torchio, Giacosa, & Río Segade (2015). The skin extracts or the wines were previously diluted with 0.05 mol/L of sulfuric acid to have ethanol contents less than 4% v/v and submitted to reverse-phase solid-phase extraction using a 1 g Sep-Pak C18 cartridge (Waters Corporation, Milford, MA, USA). Anthocyanin compounds were recovered with methanol. The HPLC-DAD system and chromatographic conditions were those reported by Rolle et al. (2015). The chromatographic separation was performed on a LiChroCART column (250 mm × 4 mm i.d.) (Merck, Darmstadt, Germany), which was packed with LiChrospher 100 RP-18 (5 µm) particles (Alltech, Deerfield, IL, USA), using two mobile phases (A, formic acid/water, 10:90 v/v; B, formic acid/methanol/water, 10:50:40 v/v/v) at a flow rate of 1 mL/min. After identification at 520 nm, the amounts of individual anthocyanins were expressed as percentages.

2.2.4 Wine color parameters

The wine color was evaluated by the color intensity, color hue, and CIELab parameters including lightness (L^*), red/green color coordinate (a^*), and yellow/blue color coordinate (b^*) according to the methods proposed by OIV (2008). A UV-1800 spectrophotometer (Shimadzu Corporation) was used with a 2-mm path length cuvette.

2.2.5 Extraction and determination of volatile compounds

The grape berries were treated following the method reported by Rolle et al. (2015). The berries were crushed under a nitrogen atmosphere with a laboratory blender (Waring Laboratory, Torrington, USA) (1 min for fresh berries, 2 min for partially dehydrated and withered berries) and then centrifuged (7000 x g, 15 min, 4°C). For the determination of free volatile compounds, a 5 mL-aliquot of the supernatant was diluted with 5 mL of deionized water, adjusted at pH 5, and placed into a 20 mL glass headspace sampling vial containing 2 g of sodium chloride and 200 µL of a 1-heptanol solution (1.55 mg/L in 10% v/v ethanol) as internal standard (Rolle et al., 2015). In the case of wines, the same treatment was carried out replacing the supernatant by 5 mL of the wine sample previously diluted twice with a 0.2 mol/L citrate-phosphate buffer solution of pH 5.

For the determination of glycosylated volatile compounds, the method reported by Wang, Kang, Xu, & Li (2011) was slightly modified. Briefly, 10 mL of the supernatant or 10 mL of the wine were diluted with 10 mL of deionized water and then submitted to reverse-phase solid phase extraction using a 1 g Sep-Pak C18 cartridge (Waters Corporation). The free volatile compounds were released with 10 mL of dichloromethane, while the glycosylated volatile compounds were recovered with 10 mL of methanol and transferred to an evaporating flask. Subsequently, methanol was evaporated using a vacuum rotavapor (BÜCHI R-210, BÜCHI Labortechnik AG, Flawil, CH) at 35 °C and the residue obtained was re-dissolved in 10 mL of the buffer solution of pH 5. The enzymatic hydrolysis was carried out by adding 50 mg of an AR-2000 commercial preparation with β -glycosidase activity (DSM Oenology, Heerlen, NL) and 0.1 g of polyvinylpyrrolidone (PVPP) to avoid the inhibitory effect of tannins, heating at 40 °C for 24 h. Finally, the extract was placed into a 20 mL glass headspace sampling vial containing 2 g of sodium chloride and 200 µL of 1-heptanol internal standard solution.

The determination of free and glycosylated volatile compounds was separately carried out using head space solid phase microextraction coupled with gas chromatography-mass spectrometry (HS-SPME-GC-MS). A 50/30 µm DVB/CAR/PDMS (divinylbenzene-carboxen-polydimethylsiloxane) fibre from Supelco (Bellefonte, PA, USA) was exposed to the headspace of the capped vial for 20 min at 40 °C (Sánchez-Palomo, Díaz-Maroto, & Pérez-Coello, 2005) and the thermal desorption was performed at 250 °C for 5 min. An Agilent 7890C gas chromatograph (Little Falls, DE, USA) equipped with a DB-WAXETR capillary column (30 m × 0.25 mm, 0.25 µm, J&W Scientific Inc., Folsom, CA, USA) and coupled to an Agilent 5975 mass selective detector was used. The chromatographic and MS conditions were previously reported (Sánchez-Palomo et al., 2005). Identification was done according to retention indices previously reported by Urcan et al. (2017) as well as to mass spectra of pure standards and/or to the NIST database (<http://webbook.nist.gov/chemistry/>). Semi-quantitative determinations (µg/kg berries or µg/L wine) were performed using the internal standard method (Englezos et al., 2016).

2.3. Statistical analysis

Statistical analyses were carried out using the SPSS Statistics software package, version 19.0 (IBM Corporation, Armonk, NY, USA). The Tukey-b test for $p < 0.05$ was used to assess significant differences by one-way analysis of variance (ANOVA).

3. RESULTS AND DISCUSSION

3.1. Standard parameters of fresh and dehydrated grapes

The chemical parameters for Moscato nero d'Acqui fresh grapes, partially dehydrated grapes (27 °Brix), and withered grapes (36 °Brix) are shown in Table 1. During the postharvest dehydration process, the berry weight decreased due to water loss and the grape must components,

such as reducing sugars and organic acids, were concentrated with the exception of tartaric acid. In fact, the previously reported decrease of malic acid as a consequence of gluconeogenesis and increased respiration (Centioni, Tiberi, Pietromarchi, Bellincontro, & Mencarelli, 2014) was compensated in partially dehydrated berries and even overcome in withered grapes by the concentration effect. Nevertheless, the berries are metabolically reactive to water stress and so chemical modifications also occur. In addition, it is important to consider that the presence of mycelial mass (*Botrytis cinerea* and other moulds) affects the content of different grape metabolites such as organic acids and sugars (Lorenzini, Azzolini, Tosi, & Zapparoli, 2012). In botrytized grapes, glucose oxidase enzymes oxidize glucose to gluconic acid. Furthermore, the hydrogen peroxide released in the above reaction forms, via a Fenton reaction, a scavenger able to oxidize tartaric acid to glyoxylic acid (Vivas et al., 2010). Therefore, in withered grapes, the glucose/fructose ratio and tartaric acid content decreased significantly, whereas the gluconic acid content and pH value increased in relation to fresh grapes. During dehydration, the trend of titratable acidity was quite similar to that of tartaric acid content in agreement with the findings of Rolle et al. (2013), although the differences were less significant for titratable acidity.

3.2. Grapes phenolic composition

The grape phenolic composition gives important information on the potential of Moscato nero d'Acqui cultivar (Table 2). The grapes are characterized by a satisfactory concentration of tannins, whereas the anthocyanin content is somewhat low. Both skins and seeds showed a quite low FRV/PRO ratio, which means that the grape tannins of Moscato nero d'Acqui are highly polymerized, and therefore low bitter and more astringent wines could be obtained (Cheynier et al., 2006; Peleg, Gacon, Schlich, & Noble, 1999).

The evolution of phenolic compounds for the skins and seeds during the grape dehydration process was different (Table 2). For the skins, an increase of the A_{280} value, which is usually used as a fast index for total phenolic compounds, was observed with increasing water loss. Furthermore, TA, TF, FRV, and PRO were more abundant in withered grapes, followed by partially dehydrated grapes, in relation to fresh grapes as a consequence of the concentration effect (Centioni et al., 2014; Moreno et al., 2008). A production factor (PF; Ruiz, Zea, Moyano, & Medina, 2010) was calculated by dividing the average content of phenolic compounds in partially dehydrated and withered grapes between their content in fresh grapes (1.3 and 1.7, respectively). Considering a 20% of error for PF resulting from the concentration effect by water evaporation (Noguerol-Pato, González-Álvarez, González-Barreiro, Cancho-Grande, & Simal-Gándara, 2013), synthesis or degradation reactions did not occur during the postharvest dehydration process of Moscato nero d'Acqui grapes. Mencarelli et al. (2010) have demonstrated that the temperature is of great importance to avoid anthocyanin oxidation induced by the activity of polyphenol oxidase and peroxidase enzymes during grape dehydration. Temperatures between 10 and 20 °C were required for reaching higher anthocyanin contents in Aleatico grapes dehydrated from 10 to 30% weight loss (WL) when compared with fresh grapes. The dehydration temperature used in the present study (16–18 °C) reduced the risk of anthocyanin oxidation in Moscato nero d'Acqui grapes because the increase underwent in the anthocyanin content corresponds to the concentration effect by water loss. Nevertheless, other researchers found no significant change in Raboso Piave grapes dehydrated from 10 to 30% WL and in Nebbiolo grapes dehydrated at 20% WL even at 20 °C (Bonghi et al., 2012; Nicoletti et al., 2013), but an increase was observed in Nebbiolo grapes dehydrated at 10 °C (Nicoletti et al., 2013) in relation to fresh berries. Using different dehydration rates, Bonghi et al. (2012) showed that key genes involved in anthocyanin biosynthesis were unaffected or down-regulated, whereas Mencarelli et al. (2010) reported that they were up-regulated at 10 and 20 °C.

Regarding skin flavanols, contradictory results were found for different cultivars. Increased catechin content was observed during the dehydration of Aleatico grapes at 10 and 20 °C (Mencarelli et al., 2010), whereas some authors pointed out a low molecular mass flavanol decrease

in Raboso Piave and Corvina berries (Bonghi et al., 2012; Rolle et al., 2013) probably related to oxidative reactions by increased activity of polyphenol oxidase and laccase enzymes. However, Moreno et al. (2008) observed no significant effect on skin proanthocyanidin content in Pinot noir grapes. These discrepancies seem to be mainly associated with a genotype effect on the content and profile of skin phenolic compounds (Torchio et al., 2016). In the present study, the higher FRV/PRO ratio obtained at the greater dehydration degree showed that the increase of skin low polymerized flavanols (FRV) was higher than that of high molecular mass flavanols (PRO) during grape dehydration. This agreed with the decrease of the skin proanthocyanidin average degree of polymerization previously reported by other researchers (Moreno et al., 2008).

For the seeds, the A_{280} index and the TF, FRV, and PRO contents decreased as dehydration progressed. This trend differed from that reported for Pinot noir grapes (Moreno et al., 2008) where the PRO content in the seeds increased significantly during dehydration, and for Corvina (Rolle et al., 2013) where TF, FRV, and PRO contents also increased using similar thermohygro-metric conditions to the present study. The PRO content also increased progressively during the dehydration of Cesanese grapes (Centioni et al., 2014). Nevertheless, Torchio et al. (2016) reported a significantly lower PRO content in Avanà seeds from grapes dehydrated using slow and fast processes than that found in fresh grapes, although the FRV content was similar. Río Segade et al. (2016) also pointed out that the lowest extractable FRV and PRO contents of seeds from Nebbiolo grapes were found at 30 and 45% WL when the results were expressed on a seed weight basis. This decrease could be justified by the consistence of the changes underwent in the seeds during grape dehydration with those observed from extended ripening, such as intensive lignification of the medium integument and the dehydration of the outer integument, which could prevent increasingly flavanols from being extracted (Bautista-Ortín et al., 2012; Cadot, Miñana-Castelló, & Chevalier, 2006). Furthermore, the degradation of low molecular mass flavanols and the hydrolysis of larger oligomers could occur depending on genotype and dehydration conditions, these reactions overcoming the concentration effect due to water loss (Rolle et al., 2013). The FRV/PRO ratio of the seeds decreased when dehydration progressed, as previously observed in Corvina grapes (Rolle et al., 2013), but in disagreement with lowering seed proanthocyanidin average degree of polymerization reported for other cultivars (Moreno et al., 2008; Río Segade et al., 2016). It is important to take into account that the contribution of seed flavanols decreased with increasing the dehydration level, which might affect the sensorial properties of resulting wines, particularly astringency and bitterness.

The anthocyanin profile of Moscato nero d'Acqui skins from fresh and dehydrated grapes is characterized by a prevalence of tri-substituted anthocyanins (Table 3), particularly malvidin-3-O-glucoside and its derivatives accounted for about 43-47% of total anthocyanin forms. A high percentage of peonidin-3-O-glucoside derivatives (about 28% of total forms) was also found in all grape samples. On the other hand, the relative abundance of acylated anthocyanins was low (6.6-8.0%). Some significant differences were found in the percentages of coumaroylated anthocyanins among fresh, partially dehydrated, and withered grapes, which increased as dehydration progressed in agreement with a previous study performed at 10 and 20 °C up to 30% WL (Mencarelli et al., 2010). Nicoletti et al. (2013) hypothesized that anthocyanin acylation increases in response to berry thermal stress to stabilize these red pigments.

3.3. *Grapes aroma composition*

In Moscato nero d'Acqui grapes, fifteen free and sixteen glycosylated volatile compounds were identified and quantified (Table 4). The aroma composition of fresh, partially dehydrated, and withered grapes is essentially characterized by a high prevalence of geraniol in both free and glycosidically-bound forms (80.8-88.0% of total free volatile compounds and 77.8-79.4% of total glycosylated volatile compounds). Regarding free volatile compounds, the second most important contribution in fresh grapes is given by nerol representing 6.8%, followed by citral and linalool.

Terpenes are responsible for the characteristic Muscat aroma (Selli et al., 2006), linalool, geraniol, and nerol being also the major free monoterpenes in Muscat cultivars, such as Muscat Hamburg, Moscatuel, and Bimeijia (Fenoll, Martinez, Hellin, & Flores, 2012; Rolle et al., 2015; Yang, Wang, Wu, Fang, & Li, 2011). The contents found of free geranic acid, *cis*-furan-linalool oxide, and *cis*-pyran-linalool oxide increased significantly as dehydration progressed, whereas geraniol decreased. Furthermore, a PF of 3 for geranic acid and *cis*-furan-linalool oxide in withered grapes, and 3.9 and 6.9 for *cis*-pyran-linalool oxide in partially dehydrated and withered grapes, respectively, which were greater than those expected by the effect of water evaporation ($1.0 < \text{PF} < 1.6$ for partially dehydrated grapes, $1.4 < \text{PF} < 2.1$ for withered grapes), suggest that synthesis reactions occurred during dehydration. Linalool oxides are originated from the oxidation of linalool and also from the metabolic activity of *Botrytis cinerea* (Bock, Benda, & Schreier, 1986; Rapp & Marais, 1993). A study performed on non-aromatic Erbaluce white cultivar showed a significant decrease of the geraniol content in botrytized grapes (Rolle et al., 2012). Thereby the second most abundant free volatile compound in partially dehydrated grapes was nerol accounting for 4.0%, followed by *cis*-pyran-linalool oxide (3.7%), whereas the content of this free linalool oxide (7.2%) was higher than that of nerol (5.6%) in withered grapes. The total content of free volatile compounds in partially dehydrated and withered grapes (260 and 239 $\mu\text{g/kg}$, respectively) was lower than that of fresh berries (430 $\mu\text{g/kg}$) mainly due to the decrease of geraniol.

The total content of glycosidically-bound volatile compounds was higher than that of free forms and increased during the postharvest dehydration process from 2452 $\mu\text{g/kg}$ in fresh grapes to 4019 $\mu\text{g/kg}$ and 4761 $\mu\text{g/kg}$ in partially dehydrated and withered grapes, respectively. Various glycosylated volatile compounds increased progressively during dehydration, such as linalool, citral, citronellol, nerol, geraniol, and benzyl alcohol, whereas a significantly higher content of geranic acid was only found in partially dehydrated grapes in relation to fresh berries. The increase of citronellol and geranic acid contents exceeded the expected by water loss in partially dehydrated ($\text{PF} = 2.4$ and 9.1 , respectively) and withered ($\text{PF} = 4.3$ and 3.2 , respectively) grapes, as well as that of nerol only in partially dehydrated grapes ($\text{PF} = 1.7$), showing synthesis reactions or easier release of these compounds from the berry skin in relation to fresh berries. On the other hand, the content of glycosylated hotrienol decreased ($\text{PF} < 0.3$) indicating that it was degraded or transformed into other compounds during postharvest dehydration. As occurred for free volatile compounds, the second most important contribution of glycosylated volatiles in fresh grapes corresponded to nerol representing 12.2%, followed by citral and linalool. Nerol continued to be the second most abundant glycosylated volatile compound in partially dehydrated and withered grapes (12.9 and 11.7%, respectively), followed by geranic acid in partially dehydrated grapes but citronellol and citral in withered grapes.

3.4. Standard parameters of fortified, *sfursat*, and *passito* wines

The compositional differences among fresh, partially dehydrated, and withered grapes, and winemaking affected wine standard characteristics (Table 5). Residual sugar and ethanol contents, as well as the glucose/fructose ratio, were directly influenced by the vinification protocol because fermentation was almost complete only in *sfursat* wines (Table 5). Also, a significantly higher content of glycerol was found in *sfursat* and *passito* wines probably due to the development of *Botrytis cinerea* during postharvest grape dehydration (Rolle et al., 2012).

3.5. Wines phenolic composition and chromatic characteristics

Significant differences were found in the phenolic composition of the three types of Moscato nero d'Acqui wines (Table 5). The A_{280} value and TA, TF, FRV, and PRO contents were lower in *fortified* wines than in *sfursat* and *passito* wines and, in turn, *sfursat* wines showed a lower richness in phenolic compounds than *passito* wines. It is important to take into account that this trend agreed

with that observed in skins of fresh, partially dehydrated, and withered grapes as consequence of the concentration effect by water loss. Other researchers also pointed out an increased content of phenolic compounds in red naturally sweet wines in relation to red sweet fortified wines made from Garnacha tintorera grapes as consequence of water evaporation during dehydration (Figueiredo-González, Cancho-Grande, & Simal-Gándara, 2013). Furthermore, skin phenolic compounds diffuse to the pulp during postharvest dehydration because of cell wall degradation and cells disruption (Marquez, Serratos, Lopez-Toledano, & Merida, 2012), facilitating their extraction and leading to more red-colored wines. According to the FRV/PRO ratio, *fortified* wines contained a significantly higher proportion of highly polymerized proanthocyanidins, whereas *sfursat* wines were richer in monomeric and oligomeric flavanols. This disagrees with the slightly lower value of proanthocyanidin average degree of polymerization found for red sweet fortified wines than for red naturally sweet wines (Figueiredo-González, Regueiro, Cancho-Grande, & Simal-Gándara, 2014). Regarding the anthocyanin profile of Moscato nero d'Acqui wines (Table 3), if compared to that of the grapes there are some differences. Indeed, during the first days of fermentation, the diffusion in the grape-juice of di-substituted anthocyanins is higher than the diffusion of tri-substituted forms (González-Neves, Gil, & Barreiro, 2008), cyanidin being particularly oxidable and its concentration decreases rapidly. Instead the wines showed high percentages of malvidin, which is less prone to oxidation (Cheynier, Souquet, Kontek, & Moutounet, 1994). Malvidin-3-O-glucoside was the predominant anthocyanin form in all the wines obtained with percentages ranging from 70.6 to 74.7% of total anthocyanins. Among all of the three wines studied, significant differences were observed for non-acylated forms. The lower percentage of malvidin-3-O-glucoside corresponded to *passito* wines. Instead, the lower relative abundances of peonidin-3-O-glucoside and cyanidin-3-O-glucoside were found for *fortified* wines, which suggest that lower losses of di-substituted anthocyanins occurred during winemaking of dehydrated grapes in relation to fresh grapes.

As can be seen from the chromatic characteristics (Table 5), *sfursat* and *passito* wines were darker according to the significantly lower values of L^* and higher color intensity, when compared with *fortified* wines. Furthermore, the two first wines showed significantly lower values of color hue and higher values of a^* and b^* coordinates, which indicate that they exhibited more reddish hue with higher red and yellow color components. In fact, the chromatic characteristics of *fortified* wines were not satisfactory. The overall colorimetric differences among the wines made from fresh grapes (*fortified* wines) and those made from dehydrated grapes (*sfursat* and *passito* wines) were over the perceptibility threshold because of the high values obtained of ΔE^* parameter (45.0 and 49.0). This parameter was calculated from the average values of L^* , a^* , and b^* coordinates (OIV, 2008). However, the difference among *sfursat* and *passito* wines was hardly perceptible by the human eye (ΔE^* parameter = 4.6; Gonnet, 2001; Torchio, Río Segade, Gerbi, Cagnasso, & Rolle, 2011).

3.6. Wines aroma composition

The aroma profile of the three wines studied is shown in Table 6. A total of 18 free and 16 glycosylated volatile compounds were identified and quantified because geranyl acetate and 2-phenylethyl acetate were not detected in glycosylated form. Regarding free volatile compounds, 2-phenyl ethanol was the predominant compound in *fortified*, *sfursat*, and *passito* wines, representing 75.5, 81.3, and 68.0% of total free volatile compounds, respectively. This aromatic alcohol is mainly formed from 2-phenylalanine during alcoholic fermentation. Its significantly higher content in *sfursat* and *passito* wines than in *fortified* wines suggested that the formation of 2-phenyl ethanol is related to the accumulation of amino acids during grape postharvest dehydration (Noguerol-Pato et al., 2013). Nevertheless, free 2-phenyl ethanol cannot contribute actively to the aroma of the wines as a consequence of its high olfactory threshold (14'000 $\mu\text{g/L}$) as also occurred in Garnacha tintorera naturally sweet wine and sweet fortified wine (Noguerol-Pato, González-Álvarez,

González-Barreiro, Cancho-Grande, & Simal-Gándara, 2012). Furthermore, 2-phenyl ethanol prevailed also in dry red Amarone wines produced from withered grapes (Tosi et al., 2012).

Other major free volatile compounds in all the three Moscato nero d'Acqui wines were citronellol (the second most abundant compound in *fortified* and *sfursat* wines), linalool, and 2-phenylethyl acetate (the second most abundant compound in *passito* wines), the two terpenols being significantly more abundant in *sfursat* wines but the acetate was in *passito* wines. Free citronellol and linalool contents were above their odor threshold (40 and 6 µg/L, respectively) and they were active odorants providing pleasant nuances of fruit and flowers. Instead, free 2-phenylethyl acetate was a key odorant (odor threshold of 250 µg/L) only in *passito* wines, giving floral notes.

Another free volatile compound found in concentrations above its odor threshold (0.5 µg/L) was rose oxide. *Passito* wines were significantly richer in both *trans* and *cis* isomers, and in turn *sfursat* wines showed *trans*-rose oxide contents higher than *fortified* wines. This monoterpenoid might be considered a marker of characteristic Muscat aroma (Ruiz-García, Hellín, Flores, & Fenoll, 2014), but its abundance in wines is influenced by the yeast metabolism (Koslitz, Renaud, Kohler, & Wüst, 2008). Free geranyl acetate also contributed to the final aroma of *fortified* wines (odor threshold of 9 µg/L) with floral nuances, although significant differences were not found in its content among the three wines studied. Contrarily, *trans*-furan-linalool oxide was significantly more abundant in *fortified* wines and benzyl alcohol was in *passito* wines, but the two free volatile compounds did not contribute actively to the wine aroma. Tosi et al. (2012) reported a significant increase of benzyl alcohol in wines made from botrytized grapes probably due to fungal enzymatic activity. It is important to take into account that higher total contents of terpenes were obtained for the wines made from dehydrated grapes (*sfursat* and *passito* wines). The enzymatic hydrolysis of aroma precursors by β -glucosidases and chemical oxidation reactions, which are favored by mould grape infection, can modify the terpene content (Tosi et al., 2012).

The predominant glycosylated volatile compound in the three Moscato nero d'Acqui wines was geraniol (75.2-76.2%), followed by nerol (14.8-15.6%). The profile of glycosylated volatile compounds in *fortified*, *sfursat*, and *passito* wines was quite similar to that found in fresh, partially dehydrated, and withered grapes, respectively. Nevertheless, many significant differences were observed. The contents of linalool, citronellol, nerol, and geraniol increased progressively and significantly from *fortified* wines to *sfursat* and *passito* wines according to the dehydration level of the grapes used for their production and in agreement with the trend already observed in grapes (Table 4). *Fortified* wines showed citral contents significantly lower than *sfursat* and *passito* wines but higher contents of geranic acid.

4. CONCLUSIONS

On the basis of the results obtained in the present study, it can be stated that the Moscato nero d'Acqui variety has a good potential for the production of *sfursat* and *passito* wines. The positive relation between grape dehydration and wine composition (by means of phenolic and aroma traits) showed the impact of water evaporation on the quality of Moscato nero d'Acqui wine obtained. In particular, the highest concentration of phenolic compounds in dehydrated grapes, due to concentration effects, is matched by the highest concentration of these compounds in *sfursat* and *passito* wines compared to *fortified* wines. In turn, *sfursat* and *passito* wines had better chromatic characteristics than *fortified* wines, which showed unsatisfactory color characteristics. Regarding the aroma profile of wines, a concentration effect was also observed with grape dehydration, but water stress could have also promoted synthesis reactions and/or facilitated the release of glycosylated terpenes (citronellol, geranic acid, and nerol) in the grapes. Therefore, *sfursat* and *passito* wines showed higher aroma richness than *fortified* wines. For the production of a sweet red wine from Moscato nero d'Acqui fresh grapes, additional techniques may be tested as the

implementation of cold pre-fermentative maceration, which could be useful to promote the extraction of phenolic and aroma compounds.

References

- Bautista-Ortín, A. B., Rodríguez-Rodríguez, P., Gil-Muñoz, R., Jiménez-Pascual, E., Busse-Valverde, N., Martínez-Cutillas, A., et al. (2012). Influence of berry ripeness on concentration, qualitative composition and extractability of grape seed tannins. *Australian Journal of Grape and Wine Research*, 18, 123–130.
- Bellincontro, A., De Santis, D., Botondi, R., Villa, I., & Mencarelli, F. (2004). Different postharvest dehydration rates affect quality characteristics and volatile compounds of Malvasia, Trebbiano and Sangiovese grapes for wine production. *Journal of the Science of Food and Agriculture*, 84, 1791–1800.
- Bock, G., Benda, I., & Schreier, P. (1986). Biotransformation of linalool by *Botrytis cinerea*. *Journal of Food Science*, 51, 659–662.
- Bonghi, C., Rizzini, F. M., Gambuti, A., Moio, L., Chkaiban, L., & Tonutti, P. (2012). Phenol compound metabolism and gene expression in the skin of wine grape (*Vitis vinifera* L.) berries subjected to partial postharvest dehydration. *Postharvest Biology and Technology*, 67, 102–109.
- Cadot, Y., Miñana-Castelló, M. T., & Chevalier, M. (2006). Anatomical, histological, and histochemical changes in grape seeds from *Vitis vinifera* L. cv Cabernet franc during fruit development. *Journal of Agricultural and Food Chemistry*, 54, 9206–9215.
- Câmara, J. S., Herbert, P., Marques, J. C., & Alves, M. A. (2004). Varietal flavour compounds of four grape varieties producing Madeira wines. *Analytica Chimica Acta*, 513, 203–207.
- Carbajal-Ida, D., Maury, C., Salas, E., Siret, R., & Mehinagic, E. (2016). Physico-chemical properties of botrytised Chenin blanc grapes to assess the extent of noble rot. *European Food Research and Technology*, 242, 117–126.
- Centioni, L., Tiberi, D., Pietromarchi, P., Bellincontro, A., & Mencarelli, F. (2014). Effect of postharvest dehydration on content of volatile organic compounds in the epicarp of Cesanese grape berry. *American Journal of Enology and Viticulture*, 65, 333–340.
- Cheynier, V., Souquet, J.M., Kontek, A., & Moutounet, M. (1994). Anthocyanin degradation in oxidising grape musts. *Journal of the Science of Food and Agriculture*, 66, 283–288.
- Cheynier, V., Dueñas-Paton, M., Salas, E., Maury, C., Souquet, J. M., Sarni-Manchado, P., & Fulcrand, H. (2006). Structure and properties of wine pigments and tannins. *American Journal of Enology and Viticulture*, 57, 298–305.
- Costantini, V., Bellincontro, A., De Santis, D., Botondi, R., & Mencarelli, F. (2006). Metabolic changes of Malvasia grapes for wine production during postharvest drying. *Journal of Agricultural and Food Chemistry*, 54, 3334–3340.
- De Maria, P. P., & Leardi, C. (1875). *Ampelografia della provincia di Alessandria*. Negro, Torino, Italy.
- Englezos, V., Torchio, F., Cravero, F., Marengo, F., Giacosa, S., Gerbi, V., et al. (2016). Aroma profile and composition of Barbera wines obtained by mixed fermentations of *Starmerella bacillaris* (synonym *Candida zemplinina*) and *Saccharomyces cerevisiae*. *LWT-Food Science and Technology*, 73, 567–575.

Fenoll, J., Martinez, C. M., Hellin, P., & Flores, P. (2012). Changes of free and glycosidically bound monoterpenes and aromatic alcohols in Moscatuel and Ruby Seedless table grapes during development. *Journal International des Sciences de la Vigne et du Vin*, 46, 41–50.

Ferrandino, A., Carra, A., Rolle, L., Schneider, A., & Schubert, A. (2012). Profiling of hydroxycinnamoyl tartrates and acylated anthocyanins in the skin of 34 *Vitis vinifera* genotypes. *Journal of Agricultural and Food Chemistry*, 60, 4931–4945.

Figueiredo-González, M., Cancho-Grande, B., & Simal-Gándara, J. (2013). Garnacha Tintorera-based sweet wines: Chromatic properties and global phenolic composition by means of UV–Vis spectrophotometry. *Food Chemistry*, 140, 217–224.

Figueiredo-González, M., Regueiro, J., Cancho-Grande, B., & Simal-Gándara, J. (2014). Garnacha Tintorera-based sweet wines: Detailed phenolic composition by HPLC/DAD–ESI/MS analysis. *Food Chemistry*, 143, 282–292.

Genovese, A., Gambuti, A., Piombino, P., & Moio, L. (2007). Sensory properties and aroma compounds of sweet Fiano wine. *Food Chemistry*, 103, 1228–1236.

Giordano, M., Rolle, L., Zeppa, G., & Gerbi, V. (2009). Chemical and volatile composition of three Italian sweet white Passito wines. *Journal International des Sciences de la Vigne et du Vin*, 43, 159–170.

Gonnet, J. F. (2001). Colour effects of co-pigmentation of anthocyanin revisited-3. A further description using CIELAB differences and assessment of matched colours using the CMC model. *Food Chemistry*, 75, 473–485.

González-Neves, G., Gil, G., & Barreiro, L. (2008). Influence of grape variety on the extraction of anthocyanins during the fermentation on skins. *European Food Research and Technology*, 226, 1349–1355.

Koslitz, S., Renaud, L., Kohler, M., & Wüst, M. (2008). Stereoselective formation of the varietal aroma compound rose oxide during alcoholic fermentation. *Journal of Agricultural and Food Chemistry*, 56, 1371–1375.

Lorenzini, M., Azzolini, M., Tosi, E., & Zapparoli, G. (2012). Postharvest grape infection of *Botrytis cinerea* and its interactions with other moulds under withering conditions to produce noble-rotten grapes. *Journal of Applied Microbiology*, 114, 762–770.

Mannini, F., Mollo, A., Santini, D., Schneider, A., Raimondi, S., Ruffa, P., et al. (2012). *I principali vitigni aromatici del Piemonte a bacca colorata*. Regione Piemonte, Torino, Italy.

Marquez, A., Serratos, M. P., Lopez-Toledano, A., & Merida, J. (2012). Colour and phenolic compounds in sweet red wines from Merlot and Tempranillo grapes chamber-dried under controlled conditions. *Food Chemistry*, 130, 111–120.

Maul, E., Töpfer, R., Carka, F., Cornea, V., Crespan, M., Dallakyan, M., et al. (2015). Identification and characterization of grapevine Genetic Resources maintained in Eastern European collections. *Vitis*, 54, 5–12.

Mencarelli, F., Bellincontro, A., Nicoletti, I., Cirilli, M., Muleo, R., & Corradini, D. (2010). Chemical and biochemical change of healthy phenolic fractions in winegrape by means of postharvest dehydration. *Journal of Agricultural and Food Chemistry*, 58, 7557–7564.

Moreno, J. J., Cerpa-Calderón, F., Cohen, S. D., Fang, Y., Qian, M., & Kennedy, J. A. (2008). Effect of postharvest dehydration on the composition of Pinot noir grapes (*Vitis vinifera* L.) and wine. *Food Chemistry*, 109, 755–762.

Nicoletti, I., Bellincontro, A., De Rossi, A., De Sanctis, F., Tiberi, D., Pietromarchi, P., et al. (2013). Postharvest dehydration of Nebbiolo grapes grown at altitude is affected by time of defoliation. *Australian Journal of Grape and Wine Research*, 19, 358-368.

Noguerol-Pato, R., González-Álvarez, M., González-Barreiro, C., Cancho-Grande, B., & Simal-Gándara, J. (2012). Aroma profile of Garnacha Tintorera-based sweet wines by chromatographic and sensorial analyses. *Food Chemistry*, 134, 2313-2325.

Noguerol-Pato, R., González-Álvarez, M., González-Barreiro, C., Cancho-Grande, B., & Simal-Gándara, J. (2013). Evolution of the aromatic profile in *Garnacha Tintorera* grapes during raisining and comparison with that of the naturally sweet wine obtained. *Food Chemistry*, 139, 1052-1061.

OIV. (2008). *Recueil international des méthodes d'analyse des vins et des moûts*. Organisation Internationale de la Vigne et du Vin, Paris, France.

Pecile, M., Zavaglia, C., & Ciardi, A. (2016) Moscato Nero di Acqui - Scheda della varietà. In: *Registro Nazionale delle Varietà di Vite*. MIPAAF, Rome, Italy.

Peleg, H., Gacon, K., Schlich, P., & Noble, A.C. (1999). Bitterness and astringency of flavan-3-ol monomers, dimers and trimers. *Journal of the Sciences of Food and Agriculture*, 79, 1123-1128.

Raimondi, S., Valota, G., & Schneider, A. (2009). Lo studio dei vitigni autoctoni minori nella collezione ampelografica di Grinzani Cavour. *Quaderni della Regione Piemonte – Agricoltura*, 62, 20-24.

Rapp, A., & Marais, J. (1993). The shelf life of wine: Changes in aroma substances during storage and ageing of white wines. In G. Charalambous (Ed.). *Shelf life studies of food and beverages. Chemical, biological, physical and nutritional aspects* (pp. 891-921). Amsterdam: Elsevier Science Publishers.

Río Segade, S., Torchio, F., Gerbi, V., Quijada-Morín, N., García-Estévez, I., Giacosa, S., et al. (2016). Impact of postharvest dehydration process of winegrapes on mechanical and acoustic properties of the seeds and their relationship with flavanol extraction during simulated maceration. *Food Chemistry*, 199, 893–901.

Rolle, L., Giordano, M., Giacosa, S., Vincenzi, S., Río Segade, S., Torchio, F., et al. (2012). CIEL*a*b* parameters of white dehydrated grapes as quality markers according to chemical composition, volatile profile and mechanical properties. *Analytica Chimica Acta*, 732, 105–113.

Rolle, L., Giacosa, S., Río Segade, S., Ferrarini, R., Torchio, F., & Gerbi, V. (2013). Influence of different thermohygrometric conditions on changes in instrumental texture properties and phenolic composition during postharvest withering of “Corvina” winegrapes (*Vitis vinifera* L.). *Drying Technology*, 31, 549-564.

Rolle, L., Torchio, F., Giacosa, S., & Río Segade, S. (2015). Berry density and size as factors related to the physicochemical characteristics of Muscat Hamburg table grapes (*Vitis vinifera* L.). *Food Chemistry*, 173, 105-113.

Ruiz, M. J., Zea, L., Moyano, L., & Medina, M. (2010). Aroma active compounds during the drying of grapes cv. Pedro Ximenez destined to the production of sweet Sherry wine. *European Food Research and Technology*, 230, 429–435.

Ruiz-García, L., Hellín, P., Flores, P., & Fenoll, J. (2014). Prediction of Muscat aroma in table grape by analysis of rose oxide. *Food Chemistry*, 154, 151-157.

Sánchez-Palomo, E., Díaz-Maroto, M. C., & Pérez-Coello, M. S. (2005). Rapid determination of volatile compounds in grapes by HS-SPME coupled with GC-MS. *Talanta*, 66, 1152-1157.

Selli, S., Canbas, A., Cabaroglu, T., Erten, H., & Gunata, Z. (2006). Aroma components of cv. Muscat of Bornova wines and influence of skin contact treatment. *Food Chemistry*, 94, 319–326.

Toffali, K., Zamboni, A., Anesi, A., Stocchero, M., Pezzotti, M., Levi, M., et al. (2011). Novel aspects of grape berry ripening and post-harvest withering revealed by untargeted LC-ESI-MS metabolomics analysis. *Metabolomics*, 7, 424–436.

Torchio, F., Cagnasso, E., Gerbi, V., & Rolle, L. (2010). Mechanical properties, phenolic composition and extractability indices of Barbera grapes of different soluble solids contents from several growing areas. *Analytica Chimica Acta*, 660, 183–189.

Torchio, F., Río Segade, S., Gerbi, V., Cagnasso, E., & Rolle, L. (2011). Changes in chromatic characteristics and phenolic composition during winemaking and shelf-life of two types of red sweet sparkling wines. *Food Research International*, 44, 729–738.

Torchio, F., Urcan, D. E., Lin, L., Gerbi, V., Giacosa, S., Río Segade, S., et al. (2016). Influence of different withering conditions on phenolic composition of Avanà, Chatus and Nebbiolo grapes for the production of ‘Reinforced’ wines. *Food Chemistry*, 194, 247–256.

Tosi, E., Fedrizzi, B., Azzolini, M., Finato, F., Simonato, B., & Zapparoli, G. (2012). Effects of noble rot on must composition and aroma profile of Amarone wine produced by the traditional grape withering protocol. *Food Chemistry*, 130, 370–375.

Tosi, E., Azzolini, M., Lorenzini, M., Torriani, S., Fedrizzi, B., Finato, F., et al. (2013). Induction of grape botrytization during withering affects volatile composition of Recioto di Soave, a ‘passito’-style wine. *European Food Research and Technology*, 236, 853–862.

Urcan, D.E., Lung, M.-L., Giacosa, S., Torchio, F., Ferrandino, A., Vincenzi, S., et al. (2016). Phenolic substances, flavor compounds, and textural properties of three native Romanian wine grape varieties. *International Journal of Food Properties*, 19, 76–98.

Urcan, D.E., Giacosa, S., Torchio, F., Río Segade, S., Raimondi, S., Bertolino, M., et al. (2017). “Fortified” wines volatile composition: Effect of different postharvest dehydration conditions of wine grapes cv. Malvasia moscata (*Vitis vinifera* L.). *Food Chemistry*, 219, 346–356.

Vincenzi, S., Tolini, S., Cocolin, L., Rantsiou, K., Curioni, A., & Rolle, L. (2012). Proteins and enzymatic activities in Erbaluce grape berries with different response to the withering process. *Analytica Chimica Acta*, 732, 130–136.

Vivas, N., Vivas de Gaulejac, N., Vitry, C., Bourden-Nonier, M. F., Chauvet, S., Donèche, B., et al. (2010). Occurrence and specificity of glucose oxidase (E. C: 1.1.3.4) in botrytized sweet white wine. Comparison with laccase (E. C: 1.10.3.2), considered as the main responsible factor for oxidation in this type of wine. *Vitis*, 49, 113–120.

Wang, Y., Kang, W., Xu, Y., & Li, J. (2011). Effect of different indigenous yeast β -glucosidases on the liberation of bound aroma compounds. *Journal of the Institute of Brewing*, 117, 230–237.

Yang, C., Wang, Y., Wu, B., Fang, J., & Li, S. (2011). Volatile compounds evolution of three table grapes with different flavour during and after maturation. *Food Chemistry*, 128, 823–830.

Table 1. Standard physico-chemical parameters of Moscato nero d'Acqui fresh and dehydrated grapes.

Parameter	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a
Berry weight (g) ^b	3.27 ± 0.50 c	2.53 ± 1.20 b	1.88 ± 0.70 a	***
Reducing sugars (g/L)	202 ± 1 a	283 ± 1 b	392 ± 5 c	***
Glucose/Fructose ratio	0.92 ± 0.01 b	0.89 ± 0.01 ab	0.86 ± 0.01 a	*
pH	3.20 ± 0.01 a	3.40 ± 0.03 ab	3.60 ± 0.12 b	*
Titrateable acidity (g/L as tartaric acid)	5.8 ± 0.0 b	6.1 ± 0.1 b	5.5 ± 0.1 a	**
Malic acid (g/L)	1.8 ± 0.1 a	1.8 ± 0.1 a	3.0 ± 0.5 b	*
Tartaric acid (g/L)	6.1 ± 0.1 b	7.2 ± 0.1 c	5.1 ± 0.1 a	***
Citric acid (g/L)	0.21 ± 0.01 a	0.30 ± 0.04 a	0.56 ± 0.06 b	**
Gluconic acid (g/L)	0.10 ± 0.07 a	0.33 ± 0.04 ab	0.54 ± 0.11 b	*

Values are expressed as average ± standard deviation (n = 3). ^aSign: *, **, and *** indicate significance at p < 0.05, 0.01, and 0.001, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). ^bCalculated on sets of 10 berries.

Table 2. Skin and seed phenolic composition of Moscato nero d'Acqui fresh and dehydrated grapes.

Parameter	Skins				Seeds			
	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a
TA (mg malvidin-3-O-glucoside chloride/kg)	314 ± 49 a	497 ± 28 b	585 ± 93 b	***	-	-	-	-
A ₂₈₀ (1/kg)	25.4 ± 2.7 a	37.1 ± 0.4 b	47.5 ± 5.8 c	***	12.8 ± 2.5 b	8.5 ± 1.3 a	6.8 ± 1.6 a	**
TF (mg (+)-catechin/kg)	1542 ± 113 a	2210 ± 108 b	2926 ± 278 c	***	847 ± 157 b	667 ± 90 a	699 ± 59 ab	*
FRV (mg (+)-catechin/kg)	515 ± 61 a	648 ± 63 a	1012 ± 157 b	***	527 ± 113 c	341 ± 56 b	214 ± 59 a	***
PRO (mg cyanidin chloride/kg)	1628 ± 138 a	2116 ± 55 b	2636 ± 295 c	***	888 ± 157 b	618 ± 100 a	494 ± 99 a	**
FRV/PRO ratio	0.31 ± 0.01 a	0.32 ± 0.03 ab	0.38 ± 0.02 b	**	0.59 ± 0.03 b	0.55 ± 0.01 b	0.43 ± 0.04 a	*

Values are expressed as average ± standard deviation (n = 3). ^aSign: *, **, and *** indicate significance at p < 0.05, 0.01, and 0.001, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). TA = total anthocyanins, A₂₈₀ = absorbance measured at 280 nm, TF = total flavonoids, FRV = flavanols reactive to vanillin, PRO = proanthocyanidins.

Table 3. Anthocyanin profile (as percentage of total forms) of Moscato nero d'Acqui fresh and dehydrated grapes and of the wines produced from them.

Anthocyanin form	Grapes				Wines			
	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a	Fortified wines	Sfursat wines	Passito wines	Sign. ^a
Delphinidin-G	7.5 ± 1.0	9.2 ± 1.5	7.7 ± 1.0	ns	3.6 ± 0.3	3.1 ± 0.3	4.3 ± 0.3	ns
Cyanidin-G	7.5 ± 0.6	9.2 ± 0.9	8.0 ± 1.6	ns	1.9 ± 0.1 a	3.1 ± 0.1 b	3.2 ± 0.1 b	***
Petunidin-G	7.6 ± 1.0	8.6 ± 1.0	8.0 ± 0.9	ns	8.2 ± 0.1 a	8.0 ± 0.1 a	9.0 ± 0.2 b	*
Peonidin-G	26.9 ± 3.1	26.5 ± 2.5	25.4 ± 4.1	ns	7.3 ± 0.2 a	8.2 ± 1.1 ab	10.7 ± 0.2 b	*
Malvidin-G	44.0 ± 2.0	39.8 ± 2.0	42.9 ± 4.3	ns	74.7 ± 1.2 b	74.3 ± 0.3 b	70.6 ± 0.8 a	*
Delphinidin acetylG	0.1 ± 0.1 a	0.1 ± 0.1 b	0.1 ± 0.1 b	*	0.1 ± 0.1 a	0.4 ± 0.1 b	0.2 ± 0.1 ab	*
Cyanidin acetylG	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	ns	nd	nd	nd	ns
Petunidin acetylG	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	ns	nd	0.7 ± 0.4	0.2 ± 0.1	ns
Peonidin acetylG	0.3 ± 0.1	0.3 ± 0.1	0.3 ± 0.1	ns	0.2 ± 0.1	0.2 ± 0.1	0.2 ± 0.1	ns
Malvidin acetylG	0.7 ± 0.1	0.6 ± 0.1	0.7 ± 0.1	ns	0.6 ± 0.1	0.5 ± 0.1	0.4 ± 0.1	ns
Delphinidin-p-coumaroylG	0.4 ± 0.1	0.4 ± 0.1	0.4 ± 0.1	ns	0.2 ± 0.1	nd	nd	ns
Cyanidin-p-coumaroylG	0.7 ± 0.1 a	0.8 ± 0.1 b	0.9 ± 0.1 b	**	0.4 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	ns
Petunidin-p-coumaroylG	0.4 ± 0.1 a	0.4 ± 0.1 ab	0.5 ± 0.1 b	**	0.2 ± 0.1	0.3 ± 0.2	0.1 ± 0.1	ns
Peonidin-p-coumaroylG	1.2 ± 0.3 a	1.4 ± 0.2 ab	1.7 ± 0.2 b	**	0.8 ± 0.1	0.3 ± 0.3	0.3 ± 0.3	ns
Malvidin-p-coumaroylG	2.3 ± 0.3 ab	2.2 ± 0.3 a	2.8 ± 0.4 b	*	1.8 ± 0.3	0.6 ± 0.7	0.8 ± 0.4	ns
Peonidin-caffeoylG	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	ns	nd	nd	nd	ns
Malvidin-caffeoylG	0.3 ± 0.1 b	0.2 ± 0.1 a	0.2 ± 0.1 b	**	nd	nd	nd	ns

Percentage values are expressed as average ± standard deviation (n = 3 for grapes; n = 2 for wines).

^aSign: *, **, ***, and "ns" indicate significance at p < 0.05, 0.01, 0.001, and not significant, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). G = 3-O-glucoside. nd = not detectable.

Table 4. Free and glycosidically-bound aroma compounds of Moscato nero d'Acqui fresh and dehydrated grapes.

Compound (µg/kg berries)	Free volatile compounds				Glycosidically-bound volatile compounds			
	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a	Fresh grapes	Partially dehydrated grapes (~27°Brix)	Withered grapes (~36°Brix)	Sign. ^a
<i>l</i> -Rose oxide	0.5 ± 0.3	0.6 ± 0.3	0.5 ± 0.7	ns	25.3 ± 1.9	21.5 ± 5	26.9 ± 11.8	ns
<i>c</i> -Rose oxide	nd	nd	nd	ns	6.5 ± 3.3	6.3 ± 2.5	4.9 ± 1.6	ns
<i>l</i> -furan-linalool oxide	0.04 ± 0.01	0.04 ± 0.02	0.04 ± 0.02	ns	3.4 ± 2.4	4.7 ± 0.1	3.5 ± 1.9	ns
<i>c</i> -furan-linalool oxide	0.02 ± 0.02 a	0.02 ± 0.02 a	0.06 ± 0.02 b	*	0.9 ± 0.2 b	0.4 ± 0.2 a	1.1 ± 0.2 b	**
Linalool	7.8 ± 1.3	3.9 ± 2.4	3.8 ± 1.5	ns	46.1 ± 0.1 a	65.8 ± 9.8 b	84.1 ± 4.5 c	**
Hotrienol	0.12 ± 0.04	0.04 ± 0.02	0.06 ± 0.02	ns	3.3 ± 2.3 b	0.9 ± 0.3 a	0.3 ± 0.2 a	*
α-terpineol	0.02 ± 0.01	0.04 ± 0.06	0.02 ± 0.02	ns	12.1 ± 0.7	14.3 ± 1.8	13.6 ± 4.1	ns
Citral	13.6 ± 6.4	3.7 ± 0.4	5.1 ± 0.9	ns	51.7 ± 2.3 a	66.0 ± 23.1 a	102.1 ± 4.8 b	*
<i>l</i> -pyran-linalool oxide	0.1 ± 0.1	0.1 ± 0.1	0.2 ± 0.1	ns	2.1 ± 0.4	2.1 ± 0.6	2.4 ± 1.2	ns
<i>c</i> -pyran-linalool oxide	2.5 ± 2.2 a	9.7 ± 1.4 b	17.2 ± 1.5 c	**	0.7 ± 0.7	0.3 ± 0.4	0.6 ± 0.5	ns
Citronellol	7.1 ± 4.5	2.5 ± 0.4	5.3 ± 2.3	ns	25.3 ± 13.0 a	61.4 ± 17.1 a	109.2 ± 20.3 b	**
Nerol	29.2 ± 0.7	10.4 ± 3.9	13.4 ± 7.8	ns	300.2 ± 12.3 a	518.4 ± 50.8 b	557.6 ± 46.1 b	***
Geraniol	368.2 ± 9.3 c	228.4 ± 15.4 b	193.2 ± 3.6 a	***	1938.5 ± 99.8 a	3127.2 ± 370.2 b	3781.0 ± 190.4 c	***
Benzyl alcohol	0.8 ± 0.6	0.1 ± 0.1	0.2 ± 0.1	ns	16.7 ± 1.4 a	17.9 ± 1.6 a	22.9 ± 2.6 b	*
2-phenylethanol	0.04 ± 0.01	0.04 ± 0.02	0.04 ± 0.02	ns	8.4 ± 7.9	16.2 ± 3.5	17.8 ± 2.4	ns
Geranic acid	0.02 ± 0.02 a	0.02 ± 0.02 a	0.06 ± 0.02 b	*	10.5 ± 1.8 a	95.8 ± 24.0 b	33.4 ± 15.8 a	**

Values are expressed as average ± standard deviation (n = 3). ^aSign: *, **, ***, and "ns" indicate significance at p < 0.05, 0.01, 0.001, and not significant, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). nd = not detectable.

Table 5. Standard parameters, phenolic composition, and chromatic characteristics of Moscato nero d'Acqui wines.

Parameter	Fortified wines	Sfursat wines	Passito wines	Sign. ^a
Ethanol (% v/v)	13.3 ± 0.1 a	15.9 ± 0.1 c	13.7 ± 0.1 b	***
Residual sugars (g/L)	94 ± 5 b	5 ± 1 a	114 ± 2 c	***
Glucose/Fructose ratio	0.6 ± 0.1 a	4.6 ± 1.2 b	0.3 ± 0.1 a	*
pH	3.30 ± 0.01 a	3.60 ± 0.01 b	3.70 ± 0.01 c	***
Titrateable acidity (g/L as tartaric acid)	3.9 ± 0.1 a	5.8 ± 0.1 c	5.4 ± 0.1 b	***
Volatile acidity (g/L as acetic acid)	0.2 ± 0.1 a	0.3 ± 0.1 b	0.6 ± 0.1 c	***
Malic acid (g/L)	1.3 ± 0.1	1.4 ± 0.1	1.6 ± 0.1	ns
Tartaric acid (g/L)	1.6 ± 0.1	1.4 ± 0.1	1.4 ± 0.3	ns
Glycerol (g/L)	4.6 ± 0.2 a	10.0 ± 0.4 b	12.3 ± 0.1 c	***
TA (mg malvidin-3-O-glucoside chloride/L)	72 ± 2 a	101 ± 3 b	165 ± 3 c	***
A ₂₈₀ (1/L)	16.2 ± 0.3 a	30.1 ± 0.5 b	45.2 ± 0.2 c	***
TF (mg (+)-catechin/L)	540 ± 17 a	979 ± 15 b	1750 ± 57 c	***
FRV (mg (+)-catechin/L)	190 ± 21 a	485 ± 42 b	1034 ± 80 c	**
PRO (mg cyanidin chloride/L)	494 ± 58 a	665 ± 46 a	1926 ± 82 b	***
FRV/PRO ratio	0.38 ± 0.01 a	0.73 ± 0.01 c	0.54 ± 0.02 b	***
L*	93.6 ± 0.1 b	63.8 ± 3.8 a	61.1 ± 1.0 a	**
a*	6.2 ± 0.9 a	36.3 ± 2.1 b	37.8 ± 0.8 b	***
b*	6.4 ± 0.3 a	21.7 ± 1.8 b	25.1 ± 0.5 b	***
Color hue	1.42 ± 0.08 b	0.98 ± 0.07 a	1.11 ± 0.02 a	*
Color intensity (10 mm optical path)	0.28 ± 0.01 a	1.70 ± 0.17 b	1.92 ± 0.05 b	**

Values are expressed as average ± standard deviation (n = 2). ^aSign: *, **, ***, and "ns" indicate significance at p < 0.05, 0.01, 0.001, and not significant, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). TA = total anthocyanins, A₂₈₀ = absorbance measured at 280 nm, TF = total flavonoids, FRV = flavanols reactive to vanillin, PRO = proanthocyanidins. L* = lightness, a* = red/green color coordinate, b* = yellow/blue color coordinate.

Table 6. Free and glycosidically-bound aroma compounds of Moscato nero d'Acqui wines.

Compound (µg/L wine)	Free volatile compounds				Glycosidically-bound volatile compounds			
	Fortified wines	Sfursat wines	Passito wines	Sign. ^a	Fortified wines	Sfursat wines	Passito wines	Sign. ^a
<i>l</i> -Rose oxide	8.2 ± 0.5 a	48.8 ± 16.7 b	95.5 ± 4.6 c	**	9.0 ± 5.2	14.4 ± 0.8	20.7 ± 2.1	ns
<i>c</i> -Rose oxide	0.2 ± 0.1 a	0.1 ± 0.1 a	10.9 ± 0.1 b	***	0.8 ± 1.1	2.3 ± 0.7	5.2 ± 2.8	ns
<i>l</i> -furan-linalool oxide	7.2 ± 0.9 b	0.1 ± 0.1 a	0.1 ± 0.1 a	**	4.7 ± 0.6	6.8 ± 3.2	9.1 ± 0.2	ns
<i>c</i> -furan-linalool oxide	0.1 ± 0.1	0.2 ± 0.1	0.3 ± 0.1	ns	1.6 ± 0.6	1.9 ± 0.1	1.0 ± 0.7	ns
Linalool	89.5 ± 0.8 a	110.5 ± 1.6 b	89.7 ± 6.6 a	*	27.4 ± 1.8 a	47.6 ± 0.9 b	56.6 ± 1.2 c	***
Ho-trienol	4.1 ± 2.9	3.2 ± 0.3	11.4 ± 5.3	ns	0.5 ± 0.1	0.5 ± 0.6	1.1 ± 0.5	ns
α -terpineol	0.2 ± 0.1	15.6 ± 9.2	11.5 ± 5.2	ns	11.3 ± 1.0	17.8 ± 0.1	13.8 ± 7.4	ns
Citral	0.2 ± 0.1	0.2 ± 0.1	0.2 ± 0.1	ns	52.2 ± 3.6 a	67.8 ± 3.9 b	78.1 ± 2.1 b	**
<i>l</i> -pyran-linalool oxide	0.1 ± 0.1	0.3 ± 0.1	0.4 ± 0.2	ns	4.9 ± 1.7	4.7 ± 2.3	4.4 ± 2.6	ns
Geranyl acetate	15.4 ± 6.8	6.4 ± 8.1	8.4 ± 0.4	ns	nd	nd	nd	-
<i>c</i> -pyran-linalool oxide	2.1 ± 1.1	2.6 ± 0.7	1.8 ± 1.4	ns	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	ns
Citronellol	107.1 ± 0.9 a	536.9 ± 58.4 c	302.5 ± 24.2 b	**	26.2 ± 2.1 a	49.7 ± 0.8 b	64.9 ± 0.9 c	***
Nerol	2.4 ± 0.8	22.7 ± 9.4	35.6 ± 20.2	ns	281.6 ± 18.8 a	444.2 ± 5.8 b	501 ± 6.1 c	***
2-phenylethyl acetate	47.2 ± 29.8 a	177.7 ± 93.6 a	526 ± 12.3 b	**	nd	nd	nd	-
Geraniol	21.4 ± 9.9	19.9 ± 3.6	27.4 ± 16.6	ns	1434.4 ± 164.9 a	2167.6 ± 9.2 b	2567.6 ± 22.5 c	**
Benzyl alcohol	0.1 ± 0.1 a	0.7 ± 0.4 ab	1.6 ± 0.4 b	*	5.2 ± 2.3	3.0 ± 0.9	2.7 ± 0.7	ns
2-phenylethanol	947.9 ± 66.0 a	4151.2 ± 194.6 c	2423.2 ± 124.7 b	***	23.7 ± 1.9	16.8 ± 10.9	25.2 ± 1.2	ns
Geranic acid	2.3 ± 1.6	9.4 ± 9.0	16.6 ± 6.1	ns	24.7 ± 0.7 c	7.1 ± 1.8 a	17.5 ± 1.9 b	**

Values are expressed as average ± standard deviation (n = 2). ^aSign: *, **, ***, and "ns" indicate significance at p < 0.05, 0.01, 0.001, and not significant, respectively. Different Latin letters within the same row indicate significant differences (Tukey-b test; p < 0.05). nd = not detectable.

Highlights

- Suitability of Moscato nero d'Acqui grapes to dehydration and winemaking was tested
- Grape dehydration increased phenolic contents with similar anthocyanin forms ratio
- Free terpene concentration decreased mainly in the first dehydration phase
- *Sfursat* and *passito* wine had higher aroma richness and better color characteristics