

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Shallow system rejuvenation and magma discharge trends at Piton de la Fournaise volcano (La Réunion Island)

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1651536> since 2017-11-10T12:29:07Z

Published version:

DOI:10.1016/j.epsl.2017.01.024

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Shallow system rejuvenation and magma discharge trends at Piton de la Fournaise volcano (La Réunion Island)

D. Coppola^{1*}, A. Di Muro², A. Peltier², N. Villeneuve², V. Ferrazzini², M. Favalli³, P. Bachèlery⁴,
L. Gurioli⁴, A.J.L. Harris⁴, S. Moune⁴, I. Vlastélic⁴, B. Galle⁵, S. Arellano⁵, A. Aiuppa^{6,7}

1) *Università degli Studi di Torino, Dipartimento di Scienze della Terra, Via Valperga Caluso
35, 10125, Torino, Italy;*

2) *Observatoire Volcanologique du Piton de la Fournaise (OVPF), Institut de Physique du
Globe de Paris, Sorbonne Paris Cité, Univ. Paris Diderot, CNRS, F-97418, La Plaine des
Cafres, La Réunion, France ;*

3) *Istituto Nazionale di Geofisica e Vulcanologia (INGV), Via della Faggiola, 32, 56126, Pisa,
Italy;*

4) *Laboratoire Magmas et Volcans, Université' Blaise Pascal – CNRS – IRD, OPGC, 5 rue
Kessler, 63038 Clermont Ferrand, France ;*

5) *Chalmers University of Technology, Department of Earth and Space Sciences, SE-412 96
Gothenburg, Sweden;*

6) *Università degli Studi di Palermo, Dipartimento di Scienze della Terra e del Mare, Via
Archirafi, 22, 90123 Palermo, Italy;*

7) *Istituto Nazionale di Geofisica e Vulcanologia (INGV), Via Ugo La Malfa, 153 90146
Palermo, Italy;*

48 **Abstract**

49 Basaltic magma chambers are often characterized by emptying and refilling cycles that influence
50 their evolution in space and time, and the associated eruptive activity. During April 2007, the
51 largest historical eruption of Piton de la Fournaise (Île de La Réunion, France) drained the shallow
52 plumbing system ($> 240 \times 10^6 \text{ m}^3$) and resulted in collapse of the 1-km-wide summit crater.
53 Following these major events, Piton de la Fournaise entered a seven-year long period of near-
54 continuous deflation interrupted, in June 2014, by a new phase of significant inflation. By
55 integrating multiple datasets (lava discharge rates, deformation, seismicity, gas flux, gas
56 composition, and lava chemistry), we here show that the progressive migration of magma from a
57 deeper (below sea level) storage zone gradually rejuvenated and pressurized the above-sea-level
58 portion of the magmatic system consisting of a vertically-zoned network of relatively small-volume
59 magma pockets. Continuous inflation provoked four small ($< 5 \times 10^6 \text{ m}^3$) eruptions from vents
60 located close to the summit cone and culminated, during August-October 2015, with a chemically
61 zoned eruption that erupted $45 \pm 15 \times 10^6 \text{ m}^3$ of lava. This two-month-long eruption evolved
62 through (i) an initial phase of waning discharge, associated to the withdrawal of differentiated
63 magma from the shallow system, into (ii) a month-long phase of increasing lava and SO_2 fluxes at
64 the effusive vent, coupled with CO_2 enrichment of summit fumaroles, and involving emission of
65 less differentiated lavas, to end with, (iii) three short-lived (~ 2 day-long) pulses in lava and gas flux,
66 coupled with arrival of cumulative olivine at the surface and deflation.

67 The activity observed at Piton de la Fournaise in 2014 and 2015 points to a new model of shallow
68 system rejuvenation and discharge, whereby continuous magma supply causes eruptions from
69 increasingly deeper and larger magma storage zones. Downward depressurization continues until
70 unloading of the deepest, least differentiated magma triggers pulses in lava and gas flux,
71 accompanied by rapid contraction of the volcano edifice, that empties the main shallow reservoir

72 and terminates the cycle. Such an unloading process may characterize the evolution of shallow
73 magmatic systems at other persistently active effusive centers.

74

75 **1 Introduction**

76 The eruptive evolution of basaltic systems is known to be controlled by several interconnected
77 factors such as supply rate, conduit instability, variation in reservoir geometry, and regional and
78 gravitational stress fields (e.g., Walker, 1993, Wadge and Burt, 2011). The temporal evolution of
79 discharge at a basaltic system has classically been explained by tapping of a single, pressurized
80 chamber (Wadge, 1982). In this case, the frequency and volume of eruptions, the rate at which
81 magma is erupted, and the effusive trends reflect century-scale and short-term (decades-to-years)
82 eruptive cycles (Dzurisin et al., 1984; Harris et al., 2000; 2011) which, in turn, have been linked to
83 the development and disruption of the magmatic system occurring at various levels in the crust and
84 with different timescales (Wadge and Burt, 2011). The waxing-waning discharge trends associated
85 with such a model have been observed at Mt. Etna (Wadge, 1982; Harris et al., 1997; Bencke and
86 Neri, 2003), Nyamulagira (Wadge and Burt, 2011), Kīlauea (Swanson et al., 2014), Mauna Loa
87 (Lipman, 1980), Stromboli (Marsella et al., 2012, Valade et al., 2016), Villarrica and Llaima
88 (Dzierma and Wehrmann, 2010), and have also characterized recent eruptions at Piton de la
89 Fournaise (e.g., Ludden 1977; Peltier et al., 2008). Recently, the role of unloading of deeper
90 magma, through eruption of dense, degassed magma residing in the shallow system, has been
91 evoked to explain explosive “paroxysms” at effusing basaltic systems (Calvari et al., 2011a,b;
92 Valade et al., 2016). This mechanism implies a downward depressurization of the plumbing system
93 that complicate the application of the classical models of output from a magma reservoir (Wadge
94 and Burt, 2011).

95 We here demonstrate the need to evolve the standard model for emptying a basaltic magma
96 reservoir during a series of effusive eruptions and to define a new model for eruption cycles at a

97 persistently active effusive system. To do this we apply a pan-disciplinary (spanning chemical,
98 textural, geophysical and remote sensing) data set to track the evolution of the plumbing system of
99 Piton de la Fournaise (Île de La Réunion, France) during its eruptive activity in 2014-15. Five
100 eruptive events characterized these two years, and culminated in the voluminous August-October
101 2015 eruption (Peltier et al., 2016). Following October 2015 the volcano inflated again in a more
102 discontinuously way, and erupted again in May and September of 2016. The data presented here
103 provide a new view on the steps of pressurization, rejuvenation and drainage of a vertically
104 stratified basaltic plumbing system during a series of effusive eruptions.

105

106

Figure 1

107

108 2 Eruption cycles and recent activity of Piton de la Fournaise

109 At Piton de la Fournaise (2632 m above sea level) we characterize an eruptive cycle as beginning
110 with a period of shallow intrusions and low-volume summit-proximal activity (vent opening inside
111 the summit crater or at less than 5 km), during which the (re)distribution of the stress field inside
112 and around the edifice favors reorganization and rejuvenation of deeper storage zones (Peltier et al.,
113 2010). This “*replenishment*” stage is followed by a period of high output rate, this being a
114 “*discharge*” phase that culminates in a major effusive eruption that represents final emptying of the
115 shallow system (Michon et al., 2007; Peltier et al., 2009; Swanson et al., 2014) and re-setting of the
116 cycle (Di Muro et al., 2014). Large-volume eruptions thus often mark the end of eruptive cycles
117 that, at Piton de la Fournaise, have been recognized as occurring over time-scales of 2-40 years
118 (Peltier et al., 2008). Early research indicated that periods of high output rate emitted olivine basalts
119 and oceanites (olivine-rich basalts with olivine > 20 vol%), with periods of low output rate emitting
120 more evolved lavas (Ludden, 1977; Lénat and Bachèlery, 1990). However, Roult et al., (2012)
121 failed in identifying a long term cyclic behavior and showed that, while evolved basalts fed short-

lived and small-volume summit to summit-proximal eruptions, eruptions with higher output rates and volumes spanned a range of bulk rock compositions from evolved aphyric basalts ($\text{MgO} \approx 6\text{wt}\%$) to oceanites (MgO up to $8.7\text{wt}\%$ when filtered for the effect of olivine accumulation). Such an eruptive pattern may reflect the interaction between different portions of a vertically-zoned plumbing system where pulses of deep-sourced magma (expected to have $>12.5\text{wt}\%$ MgO ; Albarède et al., 1997) ascending from a reservoir at the crust-mantle interface (i.e., 12-15 km below the summit) are periodically injected into the more evolved and degassed shallow portion of the system (Albarède, 1993; Battaglia et al., 2005; Peltier et al., 2009; Prôno et al., 2009; Massin et al., 2011; Di Muro et al., 2014; Vlastélic and Pietruszka, 2016). The shallowest part of the Piton de la Fournaise plumbing system is located above sea level and consists of a network of relatively small-volume ($0.1\text{-}0.3 \text{ km}^3$) magma pockets, likely a plexus of dikes and sills, whose location and shape remains a matter of debate (Lénat and Bachèlery 1990; Peltier et al. 2009; Di Muro et al., 2014, 2015; Michon et al., 2016).

A period of intense eruptive activity occurred between 1998 and 2007, when 26 eruptions (Roult et al., 2012; Staudacher et al., 2015) erupted a total bulk volume of $\sim 480 \times 10^6 \text{ m}^3$ of lava. These eruptions were subdivided into five short-term cycles of less than 1.5 years in duration by Peltier et al. (2008, 2009). Each cycle began with an initial recharge of the shallow plumbing system and inflation, followed by summit-proximal eruptions ($< 5 \text{ km}$ from the summit) and slow spreading of the eastern flank (Peltier et al., 2008; Got et al., 2013). Through time, erupted lavas became progressively richer in magnesium. The end of each cycle was represented by a distal ($> 5 \text{ km}$ from the summit) eruption of oceanite on the eastern flank (Vlastélic et al., 2005; 2007). According to Got et al. (2013), this repeated pattern of activity is related to short-term stress cycles linked to a non-linearity in the stress accumulation in the edifice, and does not necessary reflect changes in the magma supply rate.

146 During April 2007, Piton de la Fournaise experienced its largest historical eruption ($> 200 \times$
147 10^6 m^3). The shallow part of the plumbing system drained, erupting olivine-rich cumulates (olivine
148 up to 24% vol; Michon et al. 2007; Di Muro et al., 2014) and resulting in summit collapse to create
149 a caldera with an estimated volume of $\sim 90 \times 10^6 \text{ m}^3$ (Staudacher et al., 2009, 2015). Deflation of the
150 whole edifice, very low rates of seismic activity and low temperature H_2O -rich summit fumarolic
151 emissions followed the 2007 eruption, before renewed inflation and magma recharge heralded the
152 beginning of a new eruptive cycle (Di Muro et al., 2012, 2015; Peltier et al., 2009, 2016).

153 Following 2007 caldera collapse event, the redistribution of the stress field favored arrest of
154 dykes at shallow-depths (1-2 km below the summit) and allowed magma to gather in shallow
155 storage zone (Peltier et al., 2010), as has been evoked to describe the shallow system at Kīlauea
156 (e.g. Fiske and Kinoshita, 1969). These shallow magma pockets subsequently erupted eight times
157 between 2007 and 2010 (Staudacher et al., 2015), feeding small-volume ($< 3 \times 10^6 \text{ m}^3$) eruptions
158 that produced relatively evolved magmas ($\text{MgO}: 7.1 \pm 0.6 \text{ wt\%}$, Di Muro et al., 2015; Vlastélic and
159 Pietruszka, 2016) inside, or close to ($< 1 \text{ km}$), the summit (Peltier et al., 2010). During this period,
160 the main trend of deflation, consequence of the April 2007 collapse, was only interrupted by pre-
161 eruptive inflation-deflation patterns close in time to each eruptive event (Peltier et al, 2010;
162 Staudacher et al., 2015).

163 After 41 months of quiescence and continuous deflation, a new phase of activity began in
164 June 2014. The June 2014 eruption was preceded by only 11 days of weak inflation (Fig. 1c; Peltier
165 et al., 2016) and increase of CO_2 soil-degassing on the volcano flanks (Liuzzo et al., 2015). After
166 this short-lived event and until August 2015, deep seismicity increased, inflation accelerated and the
167 composition of summit fumaroles become increasingly CO_2 -rich, notably from mid-April 2015
168 (Lengliné et al.; 2016; Peltier et al, 2016). The activity that followed involved a sequence of four
169 eruptions in 2015, of which the last was by far the longest-lasting and most voluminous (Table 1). It
170 is on this data-rich 2014-2015 cycle (Fig. 1) that we focus.

Table 1

3 Datasets and results

3.1 Time Averaged Lava Discharge Rate

During 2014 and 2015, the MIROVA hot spot detection system (Coppola et al., 2016) provided systematic MODIS-derived spectral radiance data for each eruption. Following the conversion routine calibrated for Piton de la Fournaise by Coppola et al (2009; 2010; 2013), these data were converted to Time Averaged Lava Discharge Rate (*TADR*) and erupted volume (bulk), for each of the five events that comprise the 2014-2015 cycle. An uncertainty of $\pm 35\%$ in the satellite-derived values reflects distinct eruptive conditions that may characterize the effusive activity of Piton de la Fournaise, such as different emplacement styles (i.e., channel- versus tube-fed) or underlying topography (steep versus shallow slopes). Thus, during the 2014-2015 eruptions upper, median and lower bounds were set on MIROVA-derived *TADR* to take into account this uncertainty (Figure 2). The comparison between satellite-based estimations with lava volumes derived from photogrammetry and InSAR analysis (Peltier et al., 2016) is shown in Table 1. In the case of the February 2015 eruption the MODIS-derived volumes were likely underestimated, because the satellite overpasses missed the initial phase of the eruption due to the presence of clouds. On the contrary for the other four eruptions the comparison reveals that the thermal approach provides correct estimates, within the uncertainty considered. It is worth noting that during small-volume ($< 3 \times 10^6 \text{ m}^3$), short-lived (< 2 days) eruptions the MODIS-derived values were slightly underestimated, while during the longer and most voluminous eruptions, the MODIS-derived values were overestimated. This difference may be due to the low sampling time of MODIS (~ 4 overpasses per days) that is not sufficient to track with precision short-lived effusive events (Coppola et al., 2016). However, it may also reflect the different emplacement styles and insulation

196 conditions that characterize short- and long-lived eruptions. In any case, both conditions are
197 efficiently embedded within upper and lower bounds of MIROVA-derived TADR that are set
198 during each eruption.

199 Eruption duration, erupted volume (*Vol*), peak *TADR* and mean magma output rate (*MOR*) are
200 summarized in Table 1 . The first four events had *TADR* and cumulative volume trends (Fig. 2a-d)
201 that displayed the classic rapid-waxing and slow-waning forms that characterize eruption from a
202 pressurized source, as defined by Wadge (1981). For these four short-lived events, the cumulative
203 volume trends can be described using an exponential asymptotic growth trend ($R^2 > 0.9$), with the
204 form (Fig. 2),

$$205 \quad V(t) = V_e[1 - \exp(-t/\tau)]$$

206 in which V_e and τ are the final lava volume (0.4–7 million m³) and the time-decay constant (0.8–4.2
207 d), respectively (Table 1). Conversely, the fifth (final and longest) eruption in the cycle had (after a
208 short and very intense initial peak reaching 60 m³/s) an initial phase of slightly decreasing output
209 (Fig. 2e) followed by an increase of *TADR* from less than 10 m³/s between 25 August and 13
210 October to more than 20 m³/s between 14 and 17 October, when the eruption suddenly stopped. The
211 two *TADR* spikes at the end of the eruption (Fig. 2e) record two short-lived events with peaks at 32
212 and 20 m³/s, respectively, and correspond to two reactivations of the eruption from the same vent on
213 23 and 30 October, respectively. An abrupt cessation in lava effusion followed the pulse that ended
214 the main phase of effusion and separated the two reactivation events. Each reactivation event was
215 separated by exactly seven days from the preceding activity and lasted for two days. The August-
216 October 2015 event can thus be divided into three phases (Fig. 2e – 4a) defined by the following
217 volumetric characteristics:

- 218 (i) an initial slightly decreasing phase (24 August – 10 September; volume = 10.8 ± 3.7
219 $\times 10^6$ m³), following a short lived phase of intense effusion during the first day

- 220 (ii) a month-long phase of increasing TADR (11 September–13 October; volume = 22.2
221 $\pm 7.7 \times 10^6 \text{ m}^3$), and
222 (iii) three short-lived (~ 2 day-long) pulses after 14 October (total volume = $12.2 \pm 4.2 \times$
223 10^6 m^3),
224 for a total erupted volume of $45.2 \pm 15.6 \times 10^6 \text{ m}^3$.

225

226 **Figure 2**

227

228 **3.2 Deformation**

229 The inter-eruptive deformation pattern observed between 9 June 2014 (11 days before the first
230 eruption of the 2014-2015 cycle) and 24 August 2015 involved nearly continuous edifice inflation
231 (Peltier et al., 2016), which accelerated in mid-April 2015 when deep seismicity and CO₂
232 enrichment of summit fumaroles was first detected (Fig. 1c).

233 Syn-eruptive deformation revealed that the decrease in TADR observed during the first four
234 eruptive events was coupled with a parallel deflation of the summit cone (Fig. 3). The occurrence of
235 such a trend is especially evident for those events that lasted more than one week, such as the
236 February (Fig. 3b) and May (Fig. 3c) 2015 eruptions. For these eruptions, the balance between the
237 volume of the pre-eruptive sources, deduced from inversion of the deformation data (see Fig. 4 of
238 Peltier et al., 2016) and corrected for compressibility, is – to an order of magnitude – equal to the
239 erupted volume (Table 1). Conversely, for the 68-day-long August-October 2015 eruption, the
240 estimated pre-eruptive volume source (Peltier et al., 2016) is about ten times greater than the
241 erupted volume (Table 1), also taking into account a correction of $\sim 40\%$ for vesicularity (Di Muro
242 et al., (2014). This suggests that most of the erupted volume involved a significant contribution
243 from deeper (below sea level) regions (Peltier et al., 2016).

244 The deformation pattern observed during the final and largest event (Fig. 4b) was different from
245 that of all previous events. It involved a first phase of moderate summit deflation (24 August – 5
246 September), followed by summit inflation between 6 and 25 September, with deflation then
247 persisting until the end of the first eruptive phase (Fig. 4b). The two last pulses of activity (22-24
248 October and 29-31 October) were each preceded by renewed summit inflation and accompanied by
249 summit deflation. Deformation and TADR were well correlated until 25 September when they
250 became anti-correlated (Fig. 4b). This implies that, during the first half of the eruption, the shallow
251 magmatic system underwent a draining-refilling sequence, whereas the second and terminal stages
252 of the eruption were characterized by fast emptying of the plumbing system.

253

254 **Figure 3**

255

256 ***3.3 Gas fluxes at the vents***

257 The time evolution of the SO₂ emission, plume altitude and direction were captured by the scanning
258 DOAS (Differential Optical Absorption Spectroscopy) network installed by NOVAC (Galle et al.,
259 2010) during four of the five eruptions treated in this paper (Fig. 3-4). Galle et al. (2010) estimates a
260 typical uncertainty of 26% for good measurement conditions and 54% for fair measurement
261 conditions for the NOVAC-type scanning-DOAS instruments used in this study. Bad weather
262 prevented acquisition of a reliable SO₂ time series during the February 2015 eruption. During the
263 June 2014, May 2015, and July-August 2015 eruptions, SO₂ emission rates decayed in a manner
264 consistent with trends observed in the TADR time series (Fig. 3). Total SO₂ emissions for these
265 three events were at least 85, 4800, and 218 t, respectively (Table 1).

266 On 24 August 2015, the network missed a portion of the initial, short-lived, phase of high TADR
267 that characterized the August-October 2015 event, as this occurred during the night. However, from

268 the temporal trend in SO₂ emission that followed (Fig. 4c), it is possible to identify the following
269 phases:

270 (i) a sharp and short-lived increase at the eruption beginning, followed by a lower and
271 generally steady emission rate between 24 August and 8 September and then a marked decrease
272 between 8 and 12 September. SO₂ emissions remained at very low level up to 14 September;

273 (ii) a progressive increase between 14 September and 13 October. Plume geometry and
274 changing weather conditions resulted in fluctuating SO₂ emissions before 3 October;

275 (iii) three discrete pulses between 14 October and 1 November;

276 The daily SO₂ flux was highest at the very beginning of the eruption (on 24-25 August) and during
277 the pulsating period 14 – 30 October. During most of the eruption, the SO₂ flux was less than 2 kt/d,
278 punctuated by peaks at 5.5-6.7 kt/d on 25 August, 25 September and 17 October. Daily SO₂ fluxes
279 increased from <0.6 kt/d during the first phase, to 0.8 - 1.4 kt/d during the second and third phases,
280 with the most intense phase of SO₂ outgassing beginning on 14 October, and overlapping with the
281 syn-eruptive deflation recorded by the deformation network (Fig. 4c). The temporal variation in the
282 emission rate of SO₂ during this event mimics in general that of the TADR (Fig. 4). However, some
283 discrepancies can be seen during 11-17 September and 1-6 October, when an apparent reduction in
284 SO₂ emission rates is observed though the TADR remained at high levels. The first discrepancy
285 (11-17 September) corresponds to a period of hazy weather with relatively little rain and a low
286 altitude gas plume, partly confined inside the caldera. The second discrepancy (1-6 October)
287 corresponds to a period of intense rain, which likely produced wet deposition of SO₂ and poor
288 measurements conditions. Consequently, in both periods the SO₂ emission rate was most likely
289 underestimated, explaining the discrepancy with the TADR. Overall, both the SO₂ flux and TADR
290 time-series indicate that gas and lava discharge increased during the last third of the eruption, i.e.,
291 between 14 and 30 October. The increase in TADR between 15 and 17 October was also associated

292 with an increase in plume height, and the final two TADR spikes were associated with daily SO₂
293 fluxes of 1500 and 940 t/d, respectively.

294 An estimation of the pre-eruptive magma sulfur content was obtained by scaling the total SO₂
295 output to the mass of erupted lava (Table 1) while assuming a bulk vesicularity of 40 % and a melt
296 density of 2800 kg/m³, as documented by Di Muro et al., (2014) and validated by Hibert et al.,
297 (2015). Results for both the May and August-October 2015 eruptions indicate similar low pre-
298 eruptive volatile contents, of ~600 ppm. This value is about 3/5 of that recorded by most melt
299 inclusions for magmas stored near-sea-level at Piton de la Fournaise (see Di Muro et al., 2016 for
300 review). These values suggest that erupted magmas experienced significant pre-eruptive outgassing
301 at a level shallower than sea level before being erupted.

302

303 **Figure 4**

304

305 ***3.4 Summit fumarole compositions***

306 A permanent MultiGaS station installed on the southern rim of the Dolomieu caldera (see Fig. 1a
307 for location) allowed us to track the daily evolution of gas emissions at the summit of Piton de la
308 Fournaise (Di Muro et al., 2012; Peltier et al., 2016). Analysis of long-term trends reveals that,
309 during quiescent phases, emissions from low temperature fumaroles at the summit are dominated by
310 vapor water, with very small CO₂ and H₂S amounts (Di Muro et al., 2012). However, the
311 reactivation in 2014-2015 was detected by a change in the composition of summit emissions, with
312 increasing contributions of H₂S and CO₂ and low amounts of SO₂ (average H₂S/SO₂ molar ratio =
313 16). In particular, CO₂/H₂S and CO₂/H₂O ratios increased during deep seismic events (down to 9.5
314 km below the volcano summit) below the central cone in mid-April 2015 (Peltier et al., 2016). Here
315 we use the CO₂/H₂S ratio as a reference to describe the time evolution of CO₂ enrichment in summit
316 emissions during each eruption (Figs. 3-4). This ratio is selected because of the generally low and,

317 relatively stable, H₂S concentrations and because this ratio is less affected (in respect to CO₂/H₂O)
318 by condensation effects. The molar ratio is calculated after subtraction of the modeled atmospheric
319 background. CO₂/H₂S molar ratios increased before the eruptions of February, May and July 2015
320 (Figs. 3b,c,d). The peak in CO₂/H₂S was measured 2-10 days before (February and May 2015) or 1-
321 3 days after (June 2014 and July 2015) each eruption onset and decreased during the following
322 days. In contrast, during the August-October 2015 eruption, CO₂/H₂S attained a first maximum at
323 the beginning of the eruption (between 24 and 28 August). It then decreased before increasing again
324 between 22 September and 1 October (Fig. 4c). A third phase of CO₂ enrichment was detected
325 towards the end of the eruption during 17-19 October, and again after the eruption end. The
326 strongest CO₂ enrichment was broadly synchronous with the syn-eruptive resumption of inflation
327 and the increase in SO₂ emission (Fig. 4b-c).

328

329 ***3.5 Geochemical and Petrological Evolution***

330 Lavas erupted between June 2014 and May 2015 showed a decreasing trend in MgO (6.6 to 6.1 wt
331 %), Ni (92 to 65 ppm), Cr (87 to 58 ppm), and CaO/Al₂O₃ (from 0.78 to 0.73) consistent with
332 sampling a differentiating magma reservoir with time (Fig. 5). Elevated and increasing Fe₂O₃
333 contents (from 12.6 to 12.9 wt %, compared with <12.4 wt % in 2010) indicate that Fe behaved as
334 an incompatible element. This implies that olivine was no longer at the liquidus so that, probably,
335 magma temperature had decreased below 1150 °C (Villemant et al. 2009; Di Muro et al., 2016).
336 The lavas erupted on 17 and 18 May 2015 were aphyric basalts with rare tabular plagioclase
337 microphenocrysts (up to 200 µm in length, An₇₀ to An₆₀), and were amongst the most differentiated
338 of those sampled during Piton de la Fournaise historical record, resembling products emitted in
339 March 1998 after 5.5 years of quiescence (Vlastelic et al. 2005; Boivin and Bachèlery, 2009).
340 However, a change in behavior occurred during the May 2015 eruption when, between 18 and 24
341 May, the temporal trends in MgO, Cr, Ni and CaO/Al₂O₃ reversed (Fig. 5a-b). The new trend was

initially subtle but became more evident during the subsequent eruptions of July-August 2015 and, especially, during the August-October 2015 event. MgO and CaO/Al₂O₃ increased continuously from 6.6 wt% and 0.80 at the beginning of this large final eruption to 11.03 wt% and 0.89 at its end (Fig. 5a-b). The CaO/Al₂O₃ ratio of 0.894 for the last lava erupted was amongst the highest ratios measured in historical lavas of Piton de la Fournaise (Albarède et al., 1997) and indicates unusually low pyroxene fractionation or pyroxene assimilation. The compositional trend reversal that occurred in May 2015 can be ascribed to the input of less differentiated magmas. The concomitant La/Sm ratio decrease (Fig. 5c) can also be explained by the same process, by the delivery of high-degree of melting from the mantle, or both.

Mixing between (i) differentiated magma (represented by lava erupted in May 2015), (ii) new, less differentiated magma (represented by the October 2015 lavas - corrected for olivine addition), and (iii) cumulitic olivine, has been modeled in MgO-Fe₂O₃ concentration space (Fig. 5d). The model points to a temporal decrease in the differentiated component after May 2015, with the contribution of cumulitic olivine progressively increasing during October 2015 to reach around 12 wt% by the end of the August-October 2015 eruption (Fig. 5e). We stress, however, that the new mafic input (MgO 7wt%) remains relatively evolved with respect to the mafic melts (MgO 7.4-8.7 wt%) involved in the 2007 and 2008 eruptions (Villemant et al., 2009; Di Muro et al., 2014,2016).

Petrological investigations carried out on quenched samples collected during the July-August and August-October 2015 eruptions highlight a change in phenocryst content. In the July-August and August to early-September 2015 lavas, phenocrysts (up to 1 mm in size) consisted of faceted crystals of plagioclase (An₈₅ to An₆₇) and clinopyroxenes, with rare and small olivines. While plagioclase content then decreased during the second half of September, lavas emitted in October were richer in mm-sized olivines (Fo₈₃) containing Cr-spinel as inclusions. Comparison of geochemical and TADR data (Fig. 1) indicates that the compositional trend can be linked to a transient increase in lava discharge (from 4 to 7 m³ s⁻¹) that occurred on 23 May 2015. In addition,

367 while cumulitic olivine began to appear in lavas erupted between 9 and 16 October, lava discharge
368 increased from less than $10 \text{ m}^3 \text{ s}^{-1}$ on 12 October to $22 \text{ m}^3 \text{ s}^{-1}$ on 15 October (Fig. 4). Such relations
369 are consistent with coupled geochemical and lava discharge trends observed during 2002-2006
370 olivine-rich eruptions (Vlastélic et al. 2007).

371

372

Figure 5

373

374 4 Discussion

375 The 2014-2015 datasets reveal a progressive rejuvenation of the shallow plumbing system of Piton
376 de la Fournaise during which new, less differentiated magma ($\text{MgO} \approx 7\text{wt}\%$) began to mix with
377 partly outgassed and more differentiated magma ($\text{MgO} \approx 6\text{wt}\%$) resident in the shallow system. We
378 term this the “*rejuvenation (loading)*” phase of the shallow system. Increases in summit CO_2
379 emissions suggest that a simultaneous pressure buildup at shallow levels (above or close to sea
380 level) was driven by a deeper source. Gradual pressurization of the shallow system during loading
381 thus triggered the first four eruptions, from June 2014 to July-August 2015 that drained separate,
382 small-volume ($V_e = 0.4\text{--}7 \times 10^6 \text{ m}^3$) pockets of degassed magma or, alternatively, different parts
383 (compartments) of a single cooling reservoir, which was initially filled by differentiated magma (T
384 $< 1150 \text{ }^\circ\text{C}$). All four of these eruptions were characterized by lava discharge rates that decayed
385 rapidly with time ($\tau = 0.7\text{--}4.2 \text{ days}$) and which were coupled with syn-eruptive deflation of the
386 summit (Fig. 2-3). This pattern is consistent with rapid withdrawal of magma from a pressurized
387 source (Wadge et al., 1981), without any new deep input during the eruption, with a progress
388 emptying and closure of the dike. In this case the volume of erupted lava (V_e) and time-decay
389 constant (τ) are proportional to the capacity (C) of the pressurized source ($V_e = CP_0$; where P_0 is the
390 initial overpressure) and to the hydraulic resistance of the eruptive dikes ($R = 1/\tau C$) (Aki and

Ferrazzini, 2001). According to Peltier et al., (2016), the pressure sources inferred from inter-eruptive deformation data were located at depths ranging from 1.3 to 3.7 km below the summit. Our datasets reveal that the depth of these pressure sources strongly affected the effusive development of the associated eruption. In particular we find that the time-decay constant ($\tau=RC$) of these small-volume, pressurized eruptions is linearly correlated with the depth of the pressure source (Table 1), suggesting that pressure buildup at shallower levels in the magmatic system involved magma storage zones with lower capacity and/or with reduced channel resistance. Measured SO₂ fluxes mirror the effusive trends (Fig. 3-4) and total SO₂ outputs scale proportionally to the emitted lava volumes (Table 1), suggesting that sulfur was essentially released by ascending and decompressing magma without significant pre-eruptive accumulation at depth. On the other hand, our data reveal that magmas erupted after the 2010-2014 quiescence pause were largely outgassed before eruption. This is consistent with the dominantly evolved compositions of most of the erupted products.

403

4.1 Discharge (unloading) of the shallow magmatic system

The initial phase of the August-October 2015 eruption was characterized by waning lava discharge, general decrease in gas emissions and deflation (Fig. 4), typical of pressurized eruptions. Such a trend persisted until 10-13 September and is interpreted to be associated with the withdrawal of magma stored within a shallow, stratified reservoir. We can model this initial effusive trend (blue dashed line in Fig. 4a) using the pressurized eruption model of Equation 1, and find that the reservoir involved in the August-October 2015 eruption was initially drained with a decay constant of 34 days, which is 10 to 100 times longer than those of the previous four eruptions (Table 1). This likely reflects a larger capacity of this reservoir to store magma compared with previous ones. However, a magnesium-rich magmatic source became evident in exact correspondence with a new period of inflation, increase in lava and SO₂ emissions and strong CO₂ enrichment in summit fumaroles (Fig. 4) that defined the beginning of the second phase of the eruption, a phase that all

416 other eruptions in the cycle lacked (c.f. Fig. 3). The relative contribution of less differentiated
417 magma increased rapidly in the days following this transition, as witnessed by the reversal in
418 compositional trend of the erupted lavas (Fig. 5). New magmatic injection at this time is consistent
419 with enrichment in deep fluid signatures (CO_2) of the hydrothermal intra-caldera fluids emitted by
420 the summit fumaroles (Fig. 4d). In this scenario, the peak in SO_2 and CO_2 emissions, recorded on
421 late September (Fig. 4c), likely represents the buffered response of the shallow system to this deep
422 pulse. This evolution suggests that deep magma was injected into the shallow reservoir at this point.

423 However, after 25 September 2015, lava discharge and the summit deformation became
424 anti-correlated, with increasing lava flux coupled with accelerating contraction of the summit (Fig.
425 4a-b). The gradual intensification of the eruption was coupled with involvement of a more mafic
426 melt and an increase in the presence of olivine phenocrysts in the erupted lavas, as well as an
427 increase in SO_2 emissions (Fig. 4). This process accelerated on 13 October and appears to have been
428 the first of three consecutive pulses that characterized the third, terminating phase of this “*cycle-*
429 *ending*” eruption. High lava and gas discharge rates characterized these terminal pulses, which
430 lasted ~ 2 days each. The pulses were associated with short lived edifice contraction, followed by
431 abrupt cessation of effusive activity (Fig. 4). This final phase produced a volume of 12.2 ± 4.2
432 million m^3 of lava and emitted the most magnesian and cumulative olivine-rich lavas of the entire
433 cycle. Notably, the three pulses were separated by two hiatuses lasting exactly six days each, during
434 which CO_2 enrichment of the summit fumarole emission was accompanied by reappearance of
435 summit inflation (Fig. 4). The change in the TADR-deformation relationship during the final phases
436 of this terminating eruption, and the rapid edifice contraction, support a model whereby the eruption
437 rate began to exceed the magma supply rate, thus causing partial emptying of the shallow reservoir,
438 contraction of the volcano edifice and culminating in high lava discharge during the arrival of
439 cumulate-olivine in the erupted lava. We thus suggest that the “*unloading*” of the deepest mass,
440 initially injected into the base of the shallow plumbing system at the beginning of the cycle,

441 contributed to the short, high discharge rate pulses that characterized the final days of the 2014-
442 2015 cycle.

443

444 **Figure 6**

445

446 ***4.2 Shallow system rejuvenation and discharge trends at Piton de la Fournaise***

447 The sequence of events and trends observed in the datasets collected during 2014 and 2015 allows
448 us to propose a schematic model for an unloading plumbing system at a persistently active basaltic
449 system (Fig. 6). At Piton de la Fournaise, the system is composed of a main magma reservoir
450 (located ~2.5 km below the summit), above which an interconnected network of small, cooling
451 magma pockets is proposed to reside. In this model, the whole system is vertically structured,
452 chemically stratified, and filled by evolved and partly outgassed magma. While pulses of deeper
453 and less-evolved magma supply the system at its base, an olivine-rich crystal-melt mush forms at
454 the cooling margins. Plagioclase and pyroxene become important phases in these shallow outgassed
455 “pockets” of relatively evolved and cooled magma. During a cycle of magma rejuvenation and
456 discharge the ascent and arrival of deep magma into the plexus causes continuous inflation
457 accompanied by CO₂ enrichment in summit fumaroles. At the beginning of the cycle, pressurization
458 is sufficient to trigger eruption of the shallow magma pockets filled by the most evolved and
459 degassed magma. This results in short-lived, small-volume eruptions characterized by fast waxing-
460 waning discharge trends. As deeper magma continues to arrive, inflation steadily continues and
461 increasingly large volumes of magma are pressurized and erupted. This promotes a downward
462 depressurization that proceeds until the drain of the deeper stratified and volatile-rich reservoir
463 begins. A key point is that, at Piton de la Fournaise, water and sulphur exsolution accelerate when
464 magma is at pressures lower than 1 kbar (Di Muro et al., 2015a; 2016), thus providing the trigger
465 for a buoyancy increase and greater magma ascent rate. Together with the eruptions of smaller

466 magma pockets, the withdrawal of magma from the larger reservoir “*unloads*” the deepest mass,
467 which has been gradually injected into the base of the system during the cycle. This process triggers
468 a sharp intensification of the effusive phase, coupled with high discharge-rate eruption of the
469 olivine-rich mush/melt and the rapid contraction of the volcano edifice (Di Muro et al., 2014),
470 which terminate the 2014-2015 cycle.

471 The unloading process described above displays several similarities with the mechanisms supposed
472 to drive explosive (basaltic) paroxysms (e.g. downward depressurization, Calvari et al., 2011a-b,
473 Valade et al., 2016), although differs in the time scale (days rather than minutes to hours), intensity
474 ($\sim 10^1$ m³/s rather than 10^2 - 10^3 m³/s; cf. Calvari et al., 2011a-b) and final result (pulses in lava
475 effusions rather than explosive eruptive columns). Thus, we introduce the term “*effusive (basaltic)*
476 *paroxysm*” to describe an accelerating effusing phase, accompanied by sharp contraction of the
477 volcano edifice, and related to the unloading of buoyancy-driven, deep mass which ascends rapidly
478 to feed a short, highly energetic, high mass flux event.

479

480 **5 Conclusions**

481 By coupling multiple datasets recorded at Piton de la Fournaise between 2014 and 2015 we were
482 able to track and parameterize a complete cycle of shallow-system rejuvenation and discharge.
483 During the *rejuvenation* stage of June 2014 - July 2015 we observed:

- 484 • Progressive arrival of deep magma marked by deep earthquakes, continuous inflation and
485 CO₂ enrichment in summit fumarole emissions (Lengliné et al., 2016; Peltier et al., 2016);
- 486 • Downward depressurization during sporadic drainage of the shallow magmatic system
487 associated with rupturing of increasingly deep and voluminous magma pockets filled with
488 less differentiated magma.

489 Conversely, during the *discharge* phase we observe the following features:

- 490 • Initial drainage of a larger (but still shallow) magma reservoir; followed by

- Increasing lava and gas flux coupled with inflation of the summit cone, with erupted lavas becoming increasingly less evolved; to result in an
- Impulsive pattern characterized by a sequence of three effusive pulses ($TADR \approx 30 \text{ m}^3/\text{s}$ and SO_2 fluxes $\sim 5000 \text{ t/d}$), accompanied by rapid deflation of the summit (hereby defined “*effusive paroxysms*”).

Unloading of basaltic systems can thus explain not only explosive events that punctuate eruptions at effusive basaltic centers such as Stromboli and Etna (Calvari et al., 2011a-b, Valade et al., 2016), but may be at the origin of many other “*effusive paroxysms*” reported for example at Villarrica (as during the 1972 eruption), Llaïma (Bouvet de Maisonneuve, et al., 2013) and Kliuchevskoy (Van Manen et al., 2012). The real-time quantification of Time Averaged Lava Discharge Rate (TADR), gas flux, deformation and petrology during further events can thus shed light on the dynamics characterizing the rejuvenation-discharge phases at systems prone to “*effusive paroxysms*”, and can allow a degree of forewarning of such events.

Acknowledgments

We thank S. Calvari and Don Swanson, for reviewing the manuscript and for stimulating discussions about the interpretation of the data. MIROVA is a collaborative project between the Universities of Turin and Florence (Italy). We acknowledge the LANCE-MODIS data system for providing MODIS Near Real Time products. The GNSS, DOAS (from the EU-FP6 NOVAC project) and MultiGaS data used in this paper were collected by Observatoire Volcanologique du Piton de la Fournaise/Institut de Physique du Globe de Paris (OVPF/IPGP). ADM and SM contribution was partly funded by ANR “STRAP” ([ANR-14-CE03-0004](#)) This is IPGP contribution number XXXX.

References

516

- 517 Aki, K., and Ferrazzini, V., (2000). Comparison of Mount Etna, Kilauea, and Piton de la Fournaise
518 by a quantitative modeling of their eruption histories: *J Geophys Res*, v. 106 (B3), p. 4091–
519 4102, doi: 10.1029/2000JB900316.
- 520 Albarède, F., (1993). Residence time analysis of geochemical fluctuations in volcanic series.
521 *Geochem Cosmochem Acta* v. 57, p. 615–621
- 522 Albarède, F., Luais, B., Fitton, G., Semet, M., Kaminski, E., Upton, B.G.J., Bachèlery, P.,
523 Cheminée, J.L. (1997). The geochemical regimes of Piton de la Fournaise volcano (Réunion)
524 during the last 530 000 years. *Journal of Petrology* v. 38, p.171-201
- 525 Battaglia, J, Ferrazzini, V., Staudacher, T., Aki, K., Cheminée, J.L., (2005). Pre-eruptive migration
526 of earthquakes at Piton de la Fournaise volcano (Réunion Island). *Geophys J Int* v. 161, p.
527 549–558
- 528 Behncke, B., and Neri, M., (2003). Cycles and trends in the recent eruptive behavior of Mount Etna
529 (Italy). *Canadian Journal of Earth Sciences*. v. 40 (10), pp. 1405-1411
- 530 Boivin, P., Bachèlery, P. (2009). Petrology of 1977 to 1998 eruptions of Piton de la Fournaise, La
531 Réunion Island. *Journal of Volcanology and Geothermal Research* v. 184, p. 109–125
- 532 Bouvet de Maisonneuve, C., Dungan, M.A., Bachmann, O., Burgisser, A., (2012). Insights into
533 shallow magma storage and crystallization at Volcán Llaima (Andean Southern Volcanic
534 Zone, Chile). *Journal of Volcanology and Geothermal Research* v. 211-212, p. 76-91
- 535 Calvari, S., Spampinato, L., Bonaccorso, A., Oppenheimer, C., Rivalta, E., Boschi, E. (2011a).
536 Lava effusion-A slow fuse for paroxysms at Stromboli volcano? *Earth and Planetary Science*
537 *Letters* v. 301 (1-2), p. 317-323
- 538 Calvari, S., Salerno, G.G., Spampinato, L., Gouhier, M., La Spina, A., Pecora, E., Harris, A.J.L.,
539 Labazuy, P., Biale, E., Boschi, E., (2011b). An unloading foam model to constrain Etna's 11–
540 13 January 2011 lava fountaining episode. *J Geophys Res: Solid Earth* v. 116 (11), B11207

541 Coppola, D., Piscopo, D., Staudacher, T., Cigolini, C. (2009), Lava discharge rate and effusive
542 pattern at Piton de la Fournaise from MODIS data; *Journal of Volcanology and Geothermal*
543 *Research*, vol. 184 (1-2), p. 174-192. doi:10.1016/j.jvolgeores.2008.11.031.

544 Coppola, D., Laiolo, M., Cigolini, C., Delle Donne, D., Ripepe, M., (2016). Enhanced volcanic hot-
545 spot detection using MODIS IR data: results from the MIROVA system. In: Harris, A.J.L., De
546 Groeve, T., Garel, F., Carn, S.A. (Eds.), *Detecting, Modelling and Responding to Effusive*
547 *eruptions*. Geological Society, London, Special Publications, v. 426, p. 181-205.
548 dx.doi.org/10.1144/SP426.5.

549 Di Muro, A., Aiuppa, A., Burton, M., Métrich, N., Allard, P., Fougereoux, T., Giudice, G., Guida,
550 R., (2012) Intra-eruptive gas emissions and shallow magma storage after the 2007 summit
551 caldera collapse of Piton de la Fournaise, Réunion Island. EGU Meeting, Vienna

552 Di Muro, A., Métrich, N., Vergani, D., Rosi, M., Armienti, P., Fougereoux, T., Deloule, E., Arienzo,
553 I., Civetta, L., (2014). The shallow plumbing system of Piton de la Fournaise Volcano (La
554 Réunion Island, Indian Ocean) revealed by the major 2007 caldera forming eruption. *J. Petrol.*
555 v. 55, p. 1287–1315

556 Di Muro, A. D., Staudacher, T., Ferrazzini, V., Métrich, N., Besson, P., Garofalo, C. and Villemant,
557 B. (2015). Shallow Magma Storage at Piton de la Fournaise Volcano After 2007 Summit
558 Caldera Collapse Tracked in Pele's Hairs, in *Hawaiian Volcanoes: From Source to Surface*
559 (eds R. Carey, V. Cayol, M. Poland and D. Weis), John Wiley & Sons, Inc, Hoboken, NJ. doi:
560 10.1002/9781118872079.ch9

561 Di Muro, A., Métrich, N., Allard, P., Aiuppa, A., Burton, M., Galle, B., Staudacher, T (2016)
562 Magma degassing at Piton de la Fournaise volcano. “Active Volcanoes of the World” series,
563 Springer, Bachelery, P., Lenat, J.F, Di Muro, A., Michon L., Editors.

564 Dzierma, Y., and Wehrmann, H., (2010). Eruption time series statistically examined: Probabilities of
 565 future eruptions at Villarrica and Llaima Volcanoes, Southern Volcanic Zone, Chile. *Journal*
 566 *of Volcanology and Geothermal Research*. v. 193 (1–2), p. 82–92.

567 Dzurisin, D., Koyanagi, R.Y., English, T.T., 1984. Magma supply and storage at Kilauea volcano,
 568 Hawaii, 1956–1983. *J. Volcanol. Geotherm. Res.*, v. 21, p. 177–206.

569 Fiske, R.S., Kinoshita, W.T., (1969). Inflation of Kilauea Volcano prior to its 1967–1968 eruption,
 570 *Science*, v. 165, p. 341–349

571 Galle, B., Johansson, M., Rivera, C., Zhang, Y., Kihlman, M., Kern, C., Lehmann, T., Platt, U.,
 572 Arellano, S., Hidalgo, S., 2010. Network for Observation of Volcanic and Atmospheric
 573 Change (NOVAC) – A global network for volcanic gas monitoring: Network layout and
 574 instrument description. *Journal of Geophysical Research*, v. 115

575 Got, J.L., Peltier, A., Staudacher, T., Kowalski, P., Boissier, P., (2013). Edifice strength and magma
 576 transfer modulation at Piton de la Fournaise volcano. *J Geophys Res* v. 118, p. 5040–5057.
 577 doi:10.1002/jgrb.50350

578 Harris, A.J.L., Blake, S., Rothery, D.A., Stevens, N.F. (1997). A chronology of the 1991 to 1993
 579 Etna eruption using AVHRR data: implications for real time thermal volcano monitoring. *J.*
 580 *Geophys. Res.*, v. 102, p. 7985–8003.

581 Harris, A., J.L., Murray, J.B., Aries, S.E., Davies, M.A., Flynn, L.P., Wooster, M.J., Wright, R.,
 582 Rothery, D.A., (2000). Effusion rate trends at Etna and Krafla and their implications for
 583 eruptive mechanisms. *Journal of Volcanology and Geothermal Research*, v. 102, p. 237–270

584 Harris, A., J.L., Steffke, A., Calvari, S., Spampinato, L., (2011). Thirty years of satellite-derived
 585 lava discharge rates at Etna: Implications for steady volumetric output. *Journal Geophysical*
 586 *Research: Solid Earth* 116 (8), B08204

587 Lénat, J-F, Bachèlery, P., (1990). Structure et fonctionnement de la zone centrale du Piton de la
 588 Fournaise. In: Lénat J-F (ed) Le volcanisme de la Réunion, monographie. Centre de
 589 Recherches Volcanologiques, Clermont-Ferrand, p. 257–296.

590 Lengliné, O., Duputel, Z., Ferrazzini, V., (2016). Uncovering the hidden signature of a magmatic
 591 recharge at Piton de la Fournaise volcano using small earthquakes. *Geophysical Research*
 592 *Letters*, v. 43 (9), p. 4255-4262

593 Lipman, P.W., (1980). Rates of volcanic activity along the southwest rift zone of Mauna Loa
 594 Volcano, in Garcia, M. O., and Decker, R. W. (eds.), *G. A. Macdonald Special Memorial*
 595 *Issue, Bulletin of Volcanology*, v. 43, no. 4, p. 703-725.

596 Liuzzo, M., Di Muro, A., Giudice, G., Michon, L., Ferrazzini, V., Gurrieri, S., (2015). New
 597 evidence of CO₂ soil degassing anomalies on Piton de la Fournaise volcano and the link with
 598 volcano tectonic structures. *Geochemistry, Geophysics, Geosystems*, v. 16 (12), p. 4388-4404

599 Ludden, J.N., (1977). Eruptive patterns for the volcano Piton de la Fournaise, Reunion Island.
 600 *Journal of Volcanology and Geothermal Research*, v. 2 (4), p. 385-395

601 Marsella, M., Baldi, P., Coltelli, M., Fabris, M., (2012). The morphological evolution of the Sciara
 602 del Fuoco since 1868: Reconstructing the effusive activity at Stromboli volcano. *Bulletin of*
 603 *Volcanology*, v. 74 (1), p. 231-248.

604 Massin, F., Ferrazzini, V., Bachèlery, P., Nercessian, A., Duputel, Z., Staudacher, T. (2011).
 605 Structures and evolution of the plumbing system of Piton de la Fournaise volcano inferred
 606 from clustering of 2007 eruptive cycle seismicity. *Journal of Volcanology and Geothermal*
 607 *Research*, v. 202, p. 96–106

608 Michon, L., Staudacher, T., Ferrazzini, V., Bachèlery, P., Marti, J., (2007). April 2007 collapse of
 609 Piton de la Fournaise: A new example of caldera formation. *Geophysical Research Letters*, v.
 610 34(21)

611 Michon, L., Ferrazzini, V., Di Muro, A., (2016). Magma Paths at Piton de la Fournaise Volcano. In:
 612 Bachèlery, P., Lénat, J.F., Di Muro, A., Michon, L. (Eds.). Active Volcanoes of the Southwest
 613 Indian Ocean, Active Volcanoes of the World, pp. 91-106.

614 Peltier, A., Famin, V., Bachèlery, P., Cayol, V., Fukushima, Y., Staudacher, T., (2008). Cyclic
 615 magma storages and transfers at Piton de La Fournaise volcano (La Réunion hotspot) inferred
 616 from deformation and geochemical data. *Earth and Planetary Science Letters*, v. 270 (3-4), p.
 617 180-188

618 Peltier, A., P. Bachèlery, T. Staudacher, T., (2009a). Magma transfer and storage at Piton de La
 619 Fournaise (La Réunion Island) between 1972 and 2007: a review of geophysical and
 620 geochemical data. *Journal of Volcanology and Geothermal Research*, v. 184(1-2), p. 93-108.

621 Peltier, A., Staudacher, T., Bachèlery, P., (2010). New behaviour of the Piton de La Fournaise
 622 volcano feeding system (La Réunion Island deduced from GPS data: influence of the 2007
 623 Dolomieu crater collapse, *Journal of Volcanology and Geothermal Research*, v. 192, p. 48-56.

624 Peltier, A., F. Beauducel, N. Villeneuve, V. Ferrazzini, A. Di Muro, A. Aiuppa, A. Derrien, K.
 625 Jourde, Taisne, B., (2016). Deep fluid transfer evidenced by surface deformation during the
 626 2014-2015 unrest at Piton de la Fournaise volcano. *Journal of Volcanology and Geothermal*
 627 *Research*, v. 321, p. 140-148, doi :10.1016 :j.jvolgeores.2016.04.031

628 Prôno, E., Battaglia, J., Monteiller, V., Got, J-L, Ferrazzini, V., (2009). Pwave velocity structure of
 629 Piton de la Fournaise volcano deduced from seismic data recorded between 1996 and 1999.
 630 *Journal of Volcanology and Geothermal Research*, v. 184(1-2), p. 49-62

631 Roult, G., Peltier, A., Taisne, B., Staudacher, T., Ferrazzini, V., Di Muro, A., and OVPF team,
 632 (2012). A new comprehensive classification of the Piton de la Fournaise activity spanning the
 633 1985–2010 period. Search and analysis of short-term precursors from a broad-band
 634 seismological station. *Journal of Volcanology and Geothermal Research*, v. 241–242, p. 78–
 635 104. doi:10.1016/j.jvolgeores.2012.06.012

636 Staudacher, T., Ferrazzini, V., Peltier, A., Kowalski, P., Boissier, P., Catherine, P., Lauret, F.,
 637 Massin, F., (2009). The April 2007 eruption and the Dolomieu crater collapse, two major
 638 events at Piton de la Fournaise (La Réunion Island, Indian Ocean). *Journal of Volcanology*
 639 *and Geothermal Research*, v. 184(1–2), p. 126–137. doi:10.1016/j.jvolgeores. 2008.11.005
 640 Staudacher, T., Peltier, A., Ferrazzini, V., Di Muro, A., Boissier, P., Catherine, P., Kowalski, P.,
 641 Lauret, F. and Lebreton J., (2015). Fifteen Years of Intense Eruptive Activity (1998–2013) at
 642 Piton de la Fournaise Volcano: A Review. In: Bachèlery, P., Lénat, J.F., Di Muro, A.,
 643 Michon, L. (Eds.). *Active Volcanoes of the Southwest Indian Ocean, Active Volcanoes of the*
 644 *World*, pp. 139-170.
 645 Swanson, D.A., Rose, T.R., Mucek, A.E., Garcia, M.O., Fiske, R.S., and Mastin, L.G., (2014).
 646 Cycles of explosive and effusive eruptions at Kīlauea Volcano, Hawai‘i. *Geology*, v. 42(7),
 647 p. 631-634
 648 Valade, S., Lacanna, G., Coppola, D., Laiolo, M., Pistolesi, M., Delle Donne, D., Genco, R.,
 649 Marchetti, E., Olivieri, G., Allocca, C., Cigolini, C., Nishimura, T., Poggi, P., Ripepe, M.,
 650 2016. Tracking dynamics of magma migration in open-conduit systems. *Bull. Volcanol.*, (*in*
 651 *press*)
 652 Van Manen, S.M., Blake, S., Dehn, J., (2012). Satellite thermal infrared data of Shiveluch,
 653 Kliuchevskoi and Karymsky, 1993-2008: Effusion, explosions and the potential to forecast
 654 ash plumes. *Bulletin of Volcanology*, v. 74 (6), p. 1313-1335
 655 Villemant, B., Salaün, A., Staudacher, T., (2009). Evidence for a homogeneous primary magma at
 656 Piton de la Fournaise (La Réunion): a geochemical study of matrix glass, melt inclusions and
 657 Pélée’s hairs of the 1998–2008 eruptive activity. *Journal Volcanology and Geothermal*
 658 *Research*, v. 184, p. 79–92

- 659 Vlastélic, I., Staudacher, T., (2005). Rapid change of lava composition from 1998 through 2002 at
 660 Piton de la Fournaise (Réunion Island) inferred from Pb isotopes and trace elements: evidence
 661 for variable crustal contamination. *Journal of Petrology*, v. 46, p.79–107
- 662 Vlastélic, I., Peltier, A., Staudacher, T., (2007). Short-term (1998–2006) fluctuations of Pb isotopes
 663 at Piton de la Fournaise volcano (Réunion Island): origins and constraints on the size and
 664 shape of the magma reservoir. *Chem Geol*, v. 244, p. 202–220
- 665 Vlastélic, I., and Pietruszka, AJ, (2016). A Review of the Recent Geochemical Evolution of Piton
 666 de la Fournaise Volcano (1927–2010). In: Bachèlery, P., Lénat, J.F., Di Muro, A., Michon, L.
 667 (Eds.). *Active Volcanoes of the Southwest Indian Ocean, Active Volcanoes of the World*, pp.
 668 185-201
- 669 Wadge, G., (1981). The variation of magma discharge during basaltic eruptions. *Journal of*
 670 *Volcanology and Geothermal Research*, v. 11, p. 139–168.
- 671 Wadge, G. (1982). Steady state volcanism: Evidence from eruption histories of polygenetic
 672 volcanoes. *Journal Geophysical Research*, v. 87(B5), p. 4035-4049
- 673 Wadge, G., Burt, L. (2011). Stress field control of eruption dynamics at a rift volcano:
 674 Nyamuragira, D.R. Congo. *Journal of Volcanology and Geothermal Research*, v. 207, p. 1–
 675 15. doi:10.1016/j.jvolgeores.2011.06.012
- 676 Walker, G.P., (1993). Basaltic-volcano systems. In: Prichard, H. M., Alabaster, T., Harris, N. B. W.
 677 & Neary, C. R. (eds), 1993. *Magmatic Processes and Plate Tectonics*, Geological Society
 678 Special Publication, v. 76, p. 3-38.

679

680

681

682 **Tables**

683

684 **Table 1:** Main parameters for the 2014-2015 eruption of Piton de la Fournaise (all times UTC).

685 ⁽¹⁾ Values derived from MODIS data: *Vol*: erupted volume (bulk) estimated; max*TADR*: Maximum

686 Time Averaged Lava Discharge Rate; *MOR*: Mean Output Rate.

687 ⁽²⁾ Values derived from model: exponential parameters and correlation coefficient obtained by

688 fitting MODIS data with equation $V_{out} = V_e[1 - \exp(-t/\tau)]$; V_e is the final lava volume and τ is the

689 decay time constant. See the text for more details. * Best-fit parameters relative to the August-

690 October 2015 eruption were retrieved for eruptive days 2 to 16.

691 ⁽³⁾ Erupted bulk lava volumes (Vol_{lava}) deduced from photogrammetry and InSAR analysis. Source

692 parameters (volume and depth) inferred from pre-eruptive deformation data (from Peltier et al.,

693 2016);

694 SO₂: Total mass of SO₂ measured by the DOAS-NOVAC network.

695

696

697

698

699

700

701

702

703

704 **Figure Captions**

705 **Figure 1** - (a) Shaded relief map of Piton de la Fournaise and monitoring network used in this work.
706 Red, green and blue transect represent the GNSS baselines shown in (c). (b) Time Averaged lava
707 Discharge Rate (TADR) derived from MODIS data (red line) during the five eruptions between
708 2014 and 2015. The mean output rate (MOR) of each eruption is represented by the black line. (c)
709 Deformation of the volcano edifice as recorded by GNSS baseline variations (syn-eruptive
710 deformation has been removed). (d) SO₂ fluxes (daily average) measured by the permanent DOAS-
711 NOVAC network. The gray bars outline the periods of effusive activity related to the June 20-21,
712 2014, February 4-15, May 17-30, July 31 – August 2, and August 24 – October 31, 2015 eruptions.
713

714 **Figure 2** - Time-series of MODIS-derived TADR (black circles) and cumulative volumes (red
715 circle) for the 2014–2015 eruptions of Piton de la Fournaise (uncertainties are represented by thin
716 lines). Note the waxing and waning trends that characterize the first four eruptions (a-d), typical of
717 pressurized magmatic systems. The best-fit exponential curve for cumulative volumes is calculated
718 using equation 1. The exponential coefficients for each eruption are reported in Table 1. The
719 August-October 2015 eruption (e) display a composite trend, reflecting an eruptive dynamic more
720 complex than the one for the previous eruptions (see the text for details).
721

722 **Figure 3** – Multi-parametric dataset recorded during the first four eruptions (gray fields) of the
723 2014-2015 cycle. The exponential decrease in discharge of lava (red) is accompanied by a coeval
724 decrease of SO₂ flux (green) and by deflation of the summit cone (SNEG-DSRG baseline; black),
725 consistent with the fast drainage of magma. CO₂/H₂S molar ratios (blue) increased after the June
726 2014 eruption, and before the eruption of February, May and July 2015 and decreased to
727 background values afterwards. No evident sign of magmatic refilling is observed during the course
728 of these four eruptions.
729

730 **Figure 4** – Multi-parametric dataset covering the August-October 2015 eruption. (a) Time
731 Averaged Lava Discharge Rate (red squares) and temporal evolution of MgO content in the erupted
732 lavas (yellow diamond). The blue dashed line outlines the effusive trend modeled during the initial
733 drainage of the shallow reservoir. (b) Syn-eruptive deformation of the summit as shown by distance
734 change on the SNEG-DSRG baseline (black circle). Note the correlation and anti-correlation with
735 TADR (dashed line) recorded before and after 25 September, respectively. (c) SO₂ flux in the plume
736 emitted at the vent (green triangles) and recorded by the NOVAC network. (d) CO₂/H₂S molar ratio
737 in the summit fumaroles (blue circles) recorded by the MultiGaS station. Dark gray fields mark the
738 three main eruptive phases discussed in the text.

739

740 **Figure 5** - Whole rock (a-b) major and (c) trace element composition plotted versus time. Gray
741 fields outline the five eruptive periods. Dashed lines indicate (1) the compositional trend reversal
742 (between 19 and 23 of May 2015), which is ascribed to the delivery of new, less differentiated
743 magma, and (2) the onset of cumulative olivine occurrence during the long-lived August-October
744 eruption (between 9 and 16 October, 2015). (d) Three component mixing model. Mixing proportions
745 are calculated in the MgO- Fe₂O₃ concentration space. Mixing components are (1) May 2015
746 magma (MgO= 6.0 wt.%, Fe₂O₃=12.87wt%), (2) October 2015 magma corrected for olivine
747 addition (MgO= 7.0 wt.%, Fe₂O₃=12.05 wt%), and olivine (MgO= 43.3 wt.%, Fe₂O₃=17.2 wt%).
748 (e) The time evolution of the mixing model point toward temporal increase in the less differentiated
749 component after May 2015, with the contribution of cumulative olivine appearing only during the
750 second half of the August-October 2015 eruption.

751

752 **Figure 6** –Sketch model of the plumbing system of Piton de la Fournaise. The progressive arrival of
753 deep MgO-rich magma pulses causes the gradual pressurization the shallow plumbing system.
754 Progressive tapping of increasingly deep and voluminous magma pockets filled with less

755 differentiated magma causes downward depressurization that continues until the largest stratified
756 reservoir starts to drain. "Unloading" of the deepest mass that was initially injected into the base of
757 the system contributes a final, short, high effusion rate phase in the final days of the cycle that
758 causes the rapid contraction of the volcanic edifice and favors the disaggregation of olivine
759 cumulate from the crystal/melt mush.
760