AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Metals in Feathers of African Penguins (*Spheniscus demersus*): Considerations for the Welfare and Management of Seabirds Under Human Care

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1660007> since 2022-12-21T11:26:42Z

Published version:

DOI:10.1007/s00128-018-2293-9

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Metals in feathers of African penguins (*Spheniscus demersus*): considerations for the welfare and management of seabirds under human care

S. Squadrone^{1*}, M. C. Abete¹, P. Brizio¹, D. Pessani², L. Favaro².

¹ *Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle d'Aosta, via Bologna 148, 10154 Torino, Italy.*

² *Department of Life Sciences and Systems Biology, University of Torino, via Accademia Albertina 13, 10123 Torino, Italy.*

*Corresponding author e-mail: stefania.squadrone@izsto.it

Abstract

Bird feathers have been proven to be reliable indicators of metal exposure originating from contaminated food and polluted environments. The concentrations of 15 essential and non-essential metals were investigated in African penguins (*Spheniscus demersus*) feathers from a Northwestern Italian zoological facility. These birds are exclusively fed with herring from the northeast Atlantic Ocean. Certain elements, such as Hg and Cd, reflected the bioaccumulation phenomena that occur through the marine food chain. The levels of Cr, Mn, and Ni were comparable to those registered in feathers of birds living in polluted areas. These results are important for comparative studies regarding the health, nutrition and welfare of endangered seabirds kept under human care.

Keywords: metal accumulation, biomonitoring, penguins, feathers.

Inorganic contaminants such as metals are major pollutants, which are persistent and ubiquitous in ecosystems due to their natural and anthropogenic origins (Abbasi et al. 2015). Essential trace elements include chromium (Cr), copper (Cu), cobalt (Co), iron (Fe), manganese (Mn), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn). These elements are necessary for life but when they exceed physiological concentrations in tissues and organs they can be toxic (Barton & Schmitz, 2009). Non-essential trace elements include arsenic (As), cadmium (Cd), mercury (Hg), and lead (Pb), that can be tolerated by biota at very low levels, but become harmful upon bioaccumulation (Eisler 1981; Burger et al. 2008).

In recent years, feathers have become the method of choice to evaluate trace elements contamination in birds (Jerez et al. 2011; Carravieri et al. 2013; Abbasi et al. 2015; Abdullah et al. 2015). Indeed, in feathers, metals are bound to keratin, a sulfur-containing protein (Dauwe et al. 2000; Metcheva et al. 2006); and several metals have a strong affinity to keratin (Dmowski 1999). During growth, feathers are perfused from blood vessels, and metals ingested with food were incorporated into feather keratin structures. Then, metals concentrations in feathers can indicate the physiological condition of the bird during the time of active feather growth (Burger 1993).

Penguins (Order: *Sphenisciformes*; Family: *Spheniscidae*) are seabirds at the top of many marine food chains. Accordingly, penguins bioconcentrate metals in biologically available forms at several orders of magnitude above environmental levels (Markowski et al. 2013). The genus *Spheniscus* comprises four different extant species, which inhabit temperate and equatorial areas of the Southern Hemisphere (Schreiber & Burger 2002) and share common morphological traits and behavioral ecology (Williams 1995; Favaro et al. 2016). The African or Jackass Penguin (*Spheniscus demersus*) is a non-migratory seabird endemic to South Africa and Namibia, and it is the only penguin species that breeds in the African continent. African penguin juveniles undergo their first molt in spring/summer, between the ages of 12 and 23 months (Kemper et al. 2008). Adult penguins molt once a year, with a feather-shedding phase of 12.7 ± 1.4 days (Randall et al. 1986). Accordingly, the discarded plumage allows investigation of the metals, which have been accumulated by the penguins since the previous molt.

In the wild, the African penguin feeds on pelagic schooling fish; prey size varies according to geographical location (Davis & Darby 1990). The current conservation status for the African penguin is “endangered”, according to the Red List of Threatened Species of the International Union for Conservation of Nature (BirdLife International, 2013). Wild African penguin populations

53 have dramatically decreased, due to loss of habitat, reduced fish stocks and environmental pollution
54 (Crawford et al. 2011). Consequently, in-situ conservation programs are becoming crucial.
55 Moreover, African penguins are also included in many ex-situ conservation programs and are
56 frequently kept and bred in zoos and aquaria worldwide African penguins are currently living in
57 captivity (Blay & Côté, 2001). Seabirds in zoos and aquaria are often subject to a variety of dietary
58 limitations. In particular, African penguins under human care are usually provided with food that is
59 not fully representative of natural prey resources (Heat & Randall, 1985). European zoos mostly
60 feed their birds with herring from the northeast Atlantic Ocean, i.e. wild-caught prey, which could
61 have elevated levels of contaminants (Pohl & Hennings, 2009). Furthermore, penguins kept in
62 captivity could be more directly exposed to anthropogenic contaminants than wild populations, due
63 to the location of many zoos close to or within metropolitan areas. Metals have been shown to be
64 related to variation in the plumage density (Eeva et al. 1998), reduction of genetic diversity (Eeva et
65 al. 2006), low fledging success (Evers et al. 2008), decreased bone mineralization degree (Gangoso
66 et al. 2009), altered humoral immune responsiveness (Snoeijs et al. 2004), aberrant incubation
67 behavior, lethargy and asymmetric wing area (Evers et al. 2008).

68 Accordingly, our main aims were:

- 69 i) to assess Al, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Sn, Zn, and V concentrations in the
70 feathers of a large captive colony of *Spheniscus demersus* in Italy;
71 ii) to increase the data available on penguin welfare in zoos, by evaluating their exposure to
72 potentially toxic concentrations of essential and non-essential elements accumulated through food
73 consumption.

74 We predict that captive penguins will bioaccumulate metal concentrations above those in their
75 provided diet.

76 **Materials and methods**

77 *Samples collection*

78 Feathers were collected before the beginning of the molting season in 2014 from 49 African
79 penguins housed at the Zoom Biopark, Torino (44°56' N, 7°25' E). In this zoological facility,
80 penguins were kept in an outdoor communal exhibit of 1500 m², which included a pond of 120 m²
81 (maximum depth: 3 m). Feather samples were collected as previously described (Squadrone et al.,
82 2016). In addition, the whole bodies of several herring (*Clupea harengus*) were collected in the
83 same period from the penguins' food stock. Fish were selected randomly, pooled, stored and then
84 maintained at -20 °C prior to analysis. All fish were from a northeast Atlantic Ocean (FAO fishing
85 area 27).

86 *Analytical methods*

87 Surface lipids and contaminants were removed from feathers following a protocol already described
88 (Squadrone et al., 2016), then minced with a stainless steel scissors. Mercury was quantified with a
89 Direct Mercury Analyzer (Milestone, Shelton, CT, USA) and the other elements by Inductively
90 Coupled Plasma-Mass Spectrometry (Thermo Scientific, Bremen, Germany) after being subjected
91 to microwave digestion as already described (Squadrone et al., 2016). Multi-elemental
92 determination was performed with ICP-MS after daily optimization of instrumental parameters and
93 using an external standard calibration curve; rodium and germanium were used as internal
94 standards. Analytical performances were verified by processing Certified Reference Materials
95 (Dogfish liver -DOLT-4 from the National Research Council of Canada, and Oyster Tissue-SRM
96 1566b from the National Institute of Standard and Technology), along with blank reagents in each
97 analytical session. The recoveries for reference materials ranged from 85 to 120% for DOLT-4 and
98 from 82 to 117% for SRM 1566b. The limit of quantitation (LOQ) was 0.010 mg Kg⁻¹ for all
99 elements.

100 *Statistical analysis*

101 Data were tested for normality by using the Kolmogorov-Smirnov test. As data distribution was
102 non-normal and could not be satisfactorily transformed, a non-parametric Spearman's rho was used
103 to test for correlations between metal concentrations in penguin feathers. All analyses were
104 performed in the SPSS version 20.0 for Macintosh. Alpha values were two-tailed and set at 0.05.

Results and discussion

The concentrations of metals in the feathers of the African penguins and in their food (herring) are shown in Table 1, and were in the decreasing order:

Fe>Zn>Ni>Al>Cu>Cr>Mn>Se>Hg>Sn>Pb>Cd>V>Co>As. In Figure 1, bioaccumulation of certain metals in penguin's feathers in comparison to penguins' food are shown, while correlations between metals are represented in Figure 2.

Mercury and arsenic

Mercury is a contaminant of great interest in marine ecosystems. This toxic element bioaccumulates and biomagnifies in the marine food web, essentially through dietary uptake (Frias et al. 2012). Effects of mercury on birds include behavioral and neurodevelopmental deficits, impaired reproduction, and even lethality: sensitive birds can experience adverse effects at dietary concentrations of 0.05 to 0.50 mg kg⁻¹ (Eisler, 1987). Seabirds are able to detoxify mercury and are therefore more resistant to its harmful effects (Ribeiro et al. 2009). Accordingly, the Hg levels we measured in penguin feathers were below the level related to harmful effects and comparable to levels reported by Falkowska et al. (2013a,b), in a colony of African penguins living in a Polish Zoo, which received an equivalent amount of herring from the Baltic Sea. The content of metals that we detected in this fish was similar to values in herring from the same Atlantic area reported by other authors (Polak-Juszczak 2009). As mercury is known to bioaccumulate through food chains (Lodenious & Solonen, 2013), it can be suggested that the presence of Hg in the feathers of this captive colony is due to the consumption of pelagic fish and the subsequent bioaccumulation in feathers (Figure 2).

Arsenic is assimilated by fish by ingesting particulate material suspended in water and by food ingestion (Višnjić-Jeftić et al. 2010). Arsenic concentration was found to decrease as the trophic level increases in food chains (Rahman et al. 2012). Accordingly, a higher As content in herring was detected compared to penguin feathers (Table 1). Currently, there are no previous reports concerning As contamination in captive seabirds, but the values found here are similar to those reported by Jerez and coauthors (2011) in feathers of wild Antarctic penguins.

Cadmium and lead

Cadmium is a very toxic element for biota and may cause reduction in growth rates and lethal effects at lower concentrations than other harmful elements such as mercury (Spahn & Sherry 1999). According to the literature, Cd levels in seabird feathers are usually less than 0.20 mg kg⁻¹ d.w (Burger & Gochfeld, 2000). Burger also reported that Cd in feathers may cause adverse and toxic effects when exceeding levels of 0.1 to 2 mg kg⁻¹, and this effect is species dependent. In this study, there was an average Cd level close to the limit considered to be toxic in the African penguin feathers (Table 1). The Cd levels found here deserve further investigation, considering that in herrings Cd concentration was close to the instrumental LOQ (Figure 2). However, we suggest that cadmium is subject to bioaccumulation following dietary intake during penguin's lifetime. In fact, it is well known that cadmium disturbs calcium homeostasis, due to the ability of Cd to mimic Ca during bone ossification and development. Thus, Cd concentration in feathers reflects the mobilization from internal tissues and may represent a biomarker of greater whole body exposure and bioaccumulation.

Lead is a neurotoxin that causes a decrease in growth, learning ability, and metabolism (Burger & Gochfeld, 2000). Eisler (1988) suggested that average Pb levels of 50 mg kg⁻¹ d.w. in the diet may produce adverse effects in avian predators, but levels as low as 0.10 mg kg⁻¹ d.w. have been correlated with learning deficits in sensitive vertebrates. Pb is not metabolically regulated and can accumulate in bird feathers at high concentrations; therefore, it is one of the most suitable metals for monitoring anthropogenic pollution using birds (Metcheva et al. 2011). Due to the high affinity of Pb for sulfur, lead is excreted in feathers, presumably bound to the sulfhydryl groups in keratin (Sterner, 2010). Harmful effects in birds were observed at levels of 4 mg kg⁻¹ d.w. in feathers (Eisler, 1988), although seabirds can often tolerate higher concentrations. Overall, the Pb levels measured here (Table 1) were below the level of concern, and in the penguins' food, the Pb content was negligible.

157 *Aluminum and tin*

158 Little is known about the toxicity of Al in birds, although high levels have been associated with
159 impaired breeding, reduction in clutch size, defective eggshell formation, and intrauterine bleeding
160 (Nyholm, 1981). Al concentrations above 1000 mg kg⁻¹ in food could be toxic for young birds
161 (Sparling et al. 1997). Al is likely to have a high affinity with feathers because several seabirds
162 exhibited the highest levels in this integumentary structure (Lucia et al. 2010). The mean
163 concentration found in this study was two orders of magnitude higher than in the penguins' food
164 (Table 1). This could be the result of an accumulation phenomenon following dietary intake.

165 Tin and its compounds are generally thought relatively immobile in food chains and data are not
166 still available for tin in captive birds. Values reported here are comparable to those obtained by
167 Burger and Gochfeld (2000) in seabirds from the northern Pacific Ocean.

168 *Iron and manganese*

169 Iron is an essential element for biota, but could become toxic in high doses (Thomas & McGill,
170 2008). Fe originates naturally from rock and soil, but anthropogenic activities also contribute to its
171 release in the environment (Abdullah et al. 2015). Iron was the most abundant element detected in
172 penguin feathers in this study. These results also indicate a high availability of this metal in the
173 penguins' food. Therefore, further investigations in other organs are needed to evaluate possible Fe
174 bioaccumulation in captive penguins, and its possible toxic effects at high concentrations.

175 Manganese concentrations (Table 1) were detected at one order of magnitude higher than those
176 reported in feathers of wild seabirds (Ribeiro et al. 2009), but were similar to those found in birds
177 living in highly contaminated areas (Abdullah et al. 2015). This trace element enters the food chain,
178 in fact, an elevated level was detected in penguin's food (Table 1, Figure 2), resulting in
179 bioaccumulation in feathers.

180 *Copper and zinc*

181 According to the literature, copper and zinc do not bioaccumulate through food chains, but are
182 regulated by organisms (Adriano, 2001). The copper levels detected in the feathers of the African
183 penguins studied here were similar to those detected in the feathers of wild seabirds from other parts
184 of the world, such as the Southwest Atlantic Coast of France and Antarctica (Barbieri et al. 2010,
185 Lucia et al. 2010). Moreover, the Cu level measured in herring was in the range reported by other
186 authors in fish from the same area, and was of no toxicological concern.

187 Zinc is an essential element in the formation of feathers, and birds have been reported to accumulate
188 large amounts of this element (Deng et al. 2007). Zn levels measured here were in accordance with
189 the high concentration ranges reported in various bird species around the world. It was suggested
190 that high Zn levels could be related to an adaptive process of the African penguins to mercury and
191 cadmium contamination, as an increase in Zn levels is known to reduce the toxic effect of these
192 heavy metals (Jerez et al. 2011).

193 *Chromium and nickel*

194 Chromium was detected in all samples, reflecting its role as an essential element. However,
195 neurotoxic effects in birds were already suggested and results reported here are within the upper
196 range of Cr concentrations found in bird feathers (Burger, 1993). In particular, the feathers of the
197 African Penguins examined showed Cr levels of one order of magnitude higher than those detected
198 in the feathers of wild seabirds (Burger & Gochfeld, 2009, 2010).

199 Nickel is essential for animal nutrition, but data on Ni levels in seabirds are still scarce. It has been
200 suggested that the tissues of wild birds from uncontaminated environments should contain between
201 0.10 and 5.0 mg kg⁻¹ d.w. (Outridge & Scheuhammer, 1993), but scarce information is available on
202 the toxicity of Ni in birds. However, adverse effects such as genotoxicity and immunotoxicity were
203 suggested for this metal (Das, 2008). The Ni levels measured here in penguin feathers are
204 comparable with those obtained by Abdullah and coauthors (2015) in birds living in an industrial
205 area in Pakistan. Anthropogenic sources like mining and waste incineration are known increase Ni
206 environmental levels (ATSDR, 2005). Comparison with Cr and Ni levels in captive seabirds was
207 not possible due to the scarcity of data regarding these trace elements, but we found that they were
208 particularly bio accumulated in penguin's feathers.

Selenium, cobalt and vanadium

Selenium is a metalloid that birds and other wildlife require in small amounts for biological functions (Ohlendorf & Heinz, 2009). However, at high concentrations, selenium can be very toxic and subject to homeostatic regulation. In feathers, levels of 3.8 to 26 mg kg⁻¹ (according to species) result in severe adverse effects, such as mortality of eggs; moreover, Heinz (1996) reported that concentrations of 1.8 mg kg⁻¹ could result in sublethal adverse effects in birds. Selenium levels in the feathers of the African penguins that were analyzed in this study were below the values reported to be toxic, and the herring content did not pose any risk for the penguins.

Cobalt is a relatively rare element of the earth's crust, essential to mammals in the form of cobalamin (vitamin B₁₂). Co is a naturally occurring element found in rocks, soil, water, plants, and animals, and has diverse industrial importance. Vanadium has variable concentrations in biota, due to different dietary and background levels. Co and V levels in penguin feathers were relatively low and were below the LOQ in the penguins' food.

There were a number of positive significant relationships between concentrations of metals in bird feathers, suggesting common uptake and storage pathways, or similar regulation and detoxification processes. Specifically, in African Penguin feathers, we found three different positive correlations between pairs of elements, suggesting that penguin feathers accumulate these metals during growth, due to the existence of a high blood flow. This accumulation in feathers allows the elimination of partial contents of toxic metals from the organism. In fact, we observed a positive correlation between Fe and Cr (Spearman's rho $\rho = 0.835$, N = 49, $p < 0.001$), Cu and Ni (Spearman's rho $\rho = 0.806$, N = 49, $p < 0.001$), Co and V (Spearman's rho $\rho = 0.770$, N = 49, $p < 0.001$).

Conclusions

Zoos and aquaria worldwide aim to contribute to the *ex-situ* conservation of a variety of endangered seabird species, including penguins. In order to increase the reproductive success, decrease the incidence of pathologies, and avoid genotoxic effects, it is essential to monitor and minimize the level of exposure to essential and non-essential heavy metals in seabirds maintained under human care. According to the literature, there is usually a link between metal levels in the diet of birds and levels detected in their feathers. The captive colony of African penguins studied here received a specific and homogeneous diet (herring from the northeast Atlantic Ocean) which revealed the effect of food on the degree of exposure to essential and non-essential metals. For this reason, it can be recommended that captive colonies of penguins and seabirds, in general, should be fed with a varied diet, where possible, which is representative of their natural diet, avoiding the use of only one pelagic fish species.

The authors would like to thank Zoom Torino S.p.A. (www.zoomtorino.it), and in particular Dr. Daniel Sanchez, Dr. Valentina Isaja, Dr. Laura Ozella, and Dr. Sara Piga for their help during collection of samples. Kim Maciej is acknowledged for providing holding data for the genus *Spheniscus*. Livio Favaro was supported during the writing of this manuscript by the University of Torino through a MIUR co-financed postdoctoral fellowship.

The authors also thank the editor and the anonymous reviewers for useful suggestions and comments on an earlier version of this manuscript.

Ethical statement - This research conformed to the Ethical Guidelines for the Conduct of Research on Animals by Zoos and Aquariums (WAZA, 2005), and was carried out with the approval of the Ethical Committee of the Istituto Zooprofilattico Sperimentale del Piemonte Liguria e Valle d'Aosta (11168; 14 July 2014).

Table 1. Metals concentrations (mg kg⁻¹ d.w., mean $\pm$ SD) in feathers and in food of the examined African Penguin specimens (n=49).

Element	Feathers	Herring
Al	38 ±16	0.27±0.05
As	0.15±0.06	2.0±0.04
Cd	0.33±0.2	<0.010
Co	0.16±0.1	<0.010
Cr	22±77	0.04±0.005
Cu	23 ±12	1.0±0.01
Fe	183±543	9.7±0.14
Hg	2.2±0.59	0.041±0.01
Mn	15±17	0.32±0.10
Ni	58±45	0.05±0.003
Pb	0.56±0.38	0.02±0.003
Se	2.4±0.52	0.55±0.01
Sn	1.2±7.3	0.02±0.003
V	0.28±0.22	<0.010
Zn	98±32	4.4±0.40

Figure 1. Metals bioaccumulation in penguin's feathers from food.

a) cadmium, chromium, manganese, nickel and mercury levels in penguins feathers (mean±SD, mg Kg⁻¹ log scale) b) cadmium, chromium, manganese, nickel and mercury levels in penguins food (mean±SD, mg Kg⁻¹ log scale)

Figure 2. Trace elements whose bioaccumulation was found to be correlated in penguin feathers. Dotted lines represent the 95% Confidence Interval.

References

- Abbasi, N.A., Chaudhry, V.L.B., Ali, S. & Malik, R.M. 2015. Influence of taxa, trophic level, and location on bioaccumulation of toxic metals in bird's feathers: A preliminary biomonitoring study using multiple bird species from Pakistan. *Chemosphere* 120: 527-537.
- Abdullah, M., Fasola, M., Muhammad, A., Malik, S.A., Bostan, N., Bokhari, B., Kamran, M.A., Shafqat, M.N., Alamdar, A., Khan, M., Ali N. & Eqan, S.A.M.A.S. 2015. Avian feathers as a non-destructive bio-monitoring tool of trace metals signatures: A case study from severely contaminated areas. *Chemosphere* 119: 553-561.
- Adriano, D.C. 2001. Trace elements in terrestrial environments: Biochemistry, bioavailability and risks of metals. New York: Springer.
- Appelquist, H., Asbirk, S. & Drabaek, I. 1984. Mercury monitoring: mercury stability in bird feathers. *Mar. Poll. Bull.* 15(1): 22-24.
- ATSDR (Agency for Toxic Substances and Disease Registry) 2005. *Toxicological Profile for Nickel*. U.S. Public Health Service. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- Barbieri, E., Passos, E., Filippini, A., dos Santos, I.S., Garcia, C.A.B. (2010). Assessment of trace metal concentration in feathers of seabird (*Larus dominicanus*) sampled in the Florianópolis, SC, Brazilian coast. *Environ. Monit. Assess.* 169: 631-638.
- Barton, C.C. & Schmitz, S.C. 2009. Environmental toxicology: wildlife. In: Wexler P. (Ed), *Information Resources in Toxicology* (pp. 251-254).
- BirdLife International, 2013. *Spheniscus demersus*. The IUCN Red List of Threatened Species. Version 2014.2. <www.iucnredlist.org>. Downloaded on 15 December 2014.
- Blay, N., Côté I.M. 2001. Optimal conditions for breeding of captive Humboldt penguins (*Spheniscus humboldti*): a survey of British zoo. *Zoo Biol.* 20: 545-555.
- Burger, J., Gochfeld, M., Sullivan, K., Irons, D. & McKnight, A. 2008. Arsenic, cadmium, chromium, lead, manganese, mercury, and selenium in feathers of Black-legged Kittiwake (*Rissa tridactyla*) and Black Oystercatcher (*Haematopus bachmani*) from Prince William Sound, Alaska. *Sci. Tot. Environ.* 398: 20-25.
- Burger, J. & Gochfeld, M. 2000. Metal levels in feathers of 12 species of seabirds from Midway Atoll in the northern Pacific Ocean. *Sci. Tot. Environ.* 257: 37-52.
- Burger, J. 1993. Metals in avian feathers: bioindicators of environmental pollution. *Rev. Environ. Toxicol.* 5: 203-311.
- Carravieri, A., Bustamante, P., Churlaud, C. & Cherel, Y. 2013. Penguins as bioindicators of mercury contamination in the Southern Ocean: Birds from the Kerguelen Islands as a case study. *Sci. Tot. Environ.* 454: 141-148.
- Crawford, R.J.M., Altwegg, R., Barham, B.J., Barham, P.J., Durant, J.M., Dyer, B.M., Geldenhuys, D., Makhado, A.B., Pichegru, L., Ryan, P.G., Underhill, L.G., Upfol, L., Waller, L.J. & Whittington, P.A. 2011. Collapse of South Africa's penguins in the early 21st century. *African J. Mar. Sci.* 33: 139-156.

306 Das, K.K. 2008. Nickel, its adverse health effects & oxidative stress. *Indian J. Med. Res.* 128: 412-
307 425.

308 Dauwe, T., Bervoets, L., Blust, R., Pinxten, R. & Eens, M. 2000. Can excrement and feathers of
309 nestling songbirds be used as biomonitors for heavy metals pollution? *Arch. Environ. Contam.*
310 *Toxicol.* 9: 541-546.

311 Davis, L.S. & Darby, J.T. 1990. *Penguin Biology*. London: Academic Press.

312 Deng, H., Zhang, Z., Chang, C. & Wang, Y. 2007. Trace metal concentration in Great Tit (*Parus*
313 *major*) and Greenfinch (*Carduelis sinica*) at the Western Mountains of Beijing, China. *Environ.*
314 *Poll.* 148: 620-626.

315 Dmowski, K. 1999. Birds as bioindicators of heavy metal pollution: review and examples
316 concerning European species. *Acta Ornithol.* 34: 1-25.

317 Eeva, T., Belskii, E. & Kuranov, B. 2006. Environmental pollution affects genetic diversity in wild
318 bird populations. *Mut. Res.* 608: 8-15.

319 Eeva, T., Lehikoinen, E. & Rönkä, M. 1998. Air pollution fades the plumage of the Great Tit. *Funct.*
320 *Ecol.* 12: 607-612.

321 Eisler, R. 1988. Lead hazards to fish, wildlife, and invertebrates: a synoptic review. Laurel, MD,
322 USA: U.S. Fish and Wildlife Service.

323 Eisler, R. 1987. Mercury hazards to fish, wildlife and invertebrates: a synoptic review. Laurel, MD,
324 USA: U.S. Fish and Wildlife Service.

325 Eisler, R. 1981. *Trace Metal Concentrations in Marine Organisms*. New York: Pergamon Press.

326 Evers, D.C., Savoy, L.J., DeSorbo, C.R., Yates, D.E., Hanson, W., Taylor, K.M., Siegel, L.S.,
327 Cooley Jr., J.H., Bank, M.S., Major, A., Munney, K., Mower, B.F., Vogel, H.S., Schoch, N.,
328 Pokras, M., Goodale, M.W. & Fair, J. 2008. Adverse effects from environmental mercury loads
329 on breeding common loons. *Ecotoxicol.* 17: 69-81

330 Heath, R.G.M., Randall, R.M. (1985). Growth of Jackass penguin chicks, *Spheniscus demersus*,
331 hand reared on different diets. *J Zool* 205: 91-105.

332 Heinz, G.H. 1996. Selenium in birds. In: Beyer, W.N., Heinz, G.H., Redmom-Norwood, A.W.
333 (Eds.), *Environmental contaminants in wildlife: interpreting tissues concentrations* (pp. 447-
334 458). Boca Raton, FL: Lewis Publishers INC.

335 Hockey, P., Dean, R. & Ryan, P. 2005. *Roberts Birds of Southern Africa*. Cape Town: John Voelcker
336 Bird Book Fund.

337 Falkowska, L., Szumilo, E., Hajdryh, J., Grajewska, A., Beldowska, M. & Krause, I. 2013a. Effect
338 of diet on the capacity to remove mercury from the body of a penguin (*Spheniscus demersus*) living
339 in the zoo. E3S Web of Conferences 1, 12002, doi: 10.1051/e3sconf/20130112002

340 Falkowska, L., Reindl, A.R., Szumiło, E., Kwaśniak, J., Staniszevska, M., Beldowska, M.,
341 Lewandowska, A. & Krause, I. 2013b. Mercury and chlorinated pesticides on the highest level of
342 the food web as exemplified by herring from the southern Baltic and African Penguins from the
343 zoo. *Water Air Soil Poll.* 224: 1549.

344 Favaro, L., Gamba, M., Alfieri, C., Pessani, D. & McElligott, A.G. 2016. Vocal individuality cues in
345 the African penguin (*Spheniscus demersus*): a source-filter theory. *Sci Rep* 5:17255 | DOI:
346 10.1038/srep17255.

347 Gangoso, L., Alvarez-Lloret, P.A., Rodriguez-Navarro, A.A.B., Mateo, R., Hiraldo, F. & Donazar,
348 J.A. 2009. Long-term effects of lead poisoning on bone mineralization in vultures exposed to
349 ammunition sources. *Env. Poll.* 157: 569-574.

350 Lodenious, M. & Solonen, T. 2013. The use of feathers of birds of prey as indicators of metal
351 pollution. *Ecotoxicology* 22: 1319-1334.

352 Lucia, M., André, J.M., Gontier, K., Diot, N., Veiga, J. & Davail, S. 2010. Trace element
353 concentrations (mercury, cadmium, copper, zinc, lead, aluminium, nickel, arsenic, and selenium)
354 in some aquatic birds of the Southwest Atlantic Coast of France. *Arch. Environ. Contam. Toxicol.*
355 58: 844-853.

356 Jakimska, A., Konieczka, P., Skora, K. & Namieśnik, S. 2011. Bioaccumulation of metals in tissues
 357 of marine animals. Part I: The role and impact of heavy metals on organisms. *Pol. J. Environ. St.*
 358 20:1117-1125.

359 Jerez, S., Motas, M., Palacios, M.J., Valera, F., Cuervo, J.J. & Barbosa, A. 2011. Concentration of
 360 trace elements in feathers of three Antarctic penguins: geographical and interspecific differences.
 361 *Environ. Poll.* 159: 2412-2419.

362 Kemper, J., Roux, J.P., Underhill & L.G. 2008. Effect of age and breeding status on molt phenology
 363 of adult African penguins (*Spheniscus demersus*) in Namibia. *The Auk* 125: 809-819.

364 Metcheva, R., Yurukova, L. & Teodorova, S.E. 2011. Biogenic and toxic elements in feathers, eggs,
 365 and excreta of Gentoo penguin (*Pygoscelis papua ellsworthii*) in the Antarctic. *Env. Monit.*
 366 *Asses.* 182(1-4): 571-85.

367 Metcheva, R., Yurukova, L., Teodorova, S. & Nikolova, E. 2006. The penguin feathers as
 368 bioindicator of Antarctica environmental state. *Sci. Tot. Environ.* 362: 259-265.

369 Markowski, M., Banbura, M., Kalinski, A., Markowski, J., Skwarska, J., Zielinski, P., Wawrzyniak,
 370 J. & Ban, J. 2013. Avian feathers as bioindicators of the exposure to heavy metal contamination
 371 of food. *Bull. Environ. Contam. Toxicol.* 91: 302-305.

372 Nyholm, N.E.I. 1981. Evidence of involvement of Aluminum in causation of defective formation of
 373 eggshells and of impaired breeding in wild passerine birds. *Environ. Res.* 26: 363-371.

374 Outridge, P.M. & Scheuhammer, A.M. 1993. Bioaccumulation and toxicology of nickel:
 375 implications for wild mammals and birds. *Environ. Rev.*, 1(2): 172-197.

376 Pohl, C. & Hennings, U. 2009. Trace metal concentrations and trends in Baltic surface and deep
 377 waters. Baltic Sea Environment Fact Sheet, HELCOM Baltic Sea Environment Fact Sheets.
 378 Available at: <http://www.helcom.fi/baltic-sea-trends/environment-fact-sheets/>

379 Polak-Juszczak, L. 2009. Temporal trends in bioaccumulation of trace metals in herring, sprat, and
 380 cod from the southern Baltic Sea in the 1994-2003 period. *Chemosphere* 76: 1334-1339.

381 Rahman, M.A., Hasegawa, H. & Lim, R.P. 2012. Bioaccumulation, biotransformation and trophic
 382 transfer of arsenic in the aquatic food chain. *Environ. Res.* 116: 118-135.

383 Randall, R.M., Randall, B.M., Cooper, J. & Frost, P.G.H. 1986. A new census method for penguins
 384 tested on Jackass Penguins (*Spheniscus demersus*). *Ostrich* 57: 211-215.

385 Ribeiro, A.R., Eira, C., Torres, J., Mendes, P., Miquel, J., Soares, A.M. & Vingada, J. 2009. Toxic
 386 element concentrations in the razorbill Alca torda (*Charadriiformes, Alcidae*) in Portugal. *Arch.*
 387 *Environ Contam Toxicol.* 56: 588-595.

388 Schreiber, E.A. & Burger, J. 2002. *Biology of Marine Birds*. Boca Raton, Florida, USA: CRC Press.

389 Snoeijs, T., Dauwe, T., Pinxten, R., Vandesande, F. & Eens, M. 2004. Heavy metal exposure affects
 390 the humoral immune response in a free-living small songbird, the Great Tit (*Parus major*). *Arch.*
 391 *Environ. Contam. Toxicol.* 46: 399-404.

392 Spahn, S.A. & Sherry, T.W. 1999. Cadmium and lead exposure associated with reduced growth
 393 rates, poorer fledging success of little blue heron chicks (*Egretta cerulea*). *Arch. Environ.*
 394 *Contam. Toxicol* 37: 377-384.

395 Sparling, D.W., Lowe, T.P. & Campbell, P.G.C. 1997. Ecotoxicology of aluminum to fish and
 396 wildlife. In: Yokel, R.A., Golub, M.S. (Eds), *Research Issues in Aluminum Toxicity* (pp 47-68).
 397 Washington, DC: Taylor and Frances Publishers.

398 Squadrone, S., Abete, M.C., Brizio, P., Monaco, G., Colussi, S., Biolatti, C., Modesto, P., Acutis,
 399 P.L., Pessani, D. & Favaro, L. 2016. Sex- and age-related variation in metal content of penguin
 400 feathers. *Ecotoxicology* 25(2):431-438.

401 Sterner, O. 2010. *Chemistry, health, and environment*. Weinheim: Wiley-Blackwell.

402 Thomas, V.G. & McGill, I.R. 2008. Dissolution of copper, tin, and iron from sintered tungsten-
 403 bronze spheres in a simulated avian gizzard, and an assessment of their potential toxicity to
 404 birds. *Sci.Tot. Environ.* 394: 283-289.

405 WAZA (World Association of Zoos and Aquariums) (2005). Ethical guidelines for the conduct of
 406 research on animals by zoos and aquariums. 60th Annual Conference of the World Association of
 407 Zoos and Aquariums, New York, USA. Available at:

408 <http://www.waza.org/en/site/conservation/code-of-ethics-and-animal-welfare>. Accessed 13
409 January 2014.

410 Whittington, P.A., Dyer, B.M. & Klages, N.T.W. 2000. Maximum longevity of African penguin
411 *Spheniscus demersus* based on banding records. *Mar. Ornithol.* 28: 81–82.

412 Williams, T.D. 1995. *The Penguins*. Oxford: Oxford University Press.

413 Wilson, R.P. 1985. Seasonality in diet and breeding success of the Jackass Penguin *Spheniscus*
414 *demersus*. *J. Ornithol.* 126: 53-62.

415 Višnjić-Jeftić, Ž., Jarić, I., Jovanović, L.J., Skorić, S., Smederevac-Lalić, M. & Lenhardt, M. 2010.
416 Heavy metal and trace element accumulation in muscle, liver, and gills of the Pontic shad (*Alosa*
417 *immaculata* Bennet 1835) from the Danube River (Serbia). *Microchem. J.* 95: 341-344.
418
419