

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Listeria innocua and Listeria monocytogenes strains from dairy plants behave similarly in biofilm sanitizer testing**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1662912> since 2019-02-15T15:42:38Z

*Published version:*

DOI:10.1016/j.lwt.2018.02.073

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 ***Listeria innocua* and *Listeria monocytogenes* strains from dairy plants behave similarly in**  
2 **biofilm sanitizer testing**

3 **Annalisa Costa<sup>a,b</sup>, Antonio Lourenco<sup>b,1</sup>, Tiziana Civera<sup>a</sup>, Luísa Brito<sup>b,\*</sup>**  
4

5 <sup>a</sup>Dipartimento di Scienze Veterinarie, University of Turin, Grugliasco (TO), Italy.

6 <sup>b</sup>LEAF-Linking Landscape, Environment, Agriculture and Food/DRAT-Departamento dos  
7 Recursos Naturais, Ambiente e Território, Instituto Superior de Agronomia, University of Lisbon,  
8 1349-017 Lisbon, Portugal

9 <sup>1</sup>Current address: Food Safety Department, Teagasc Food Research Centre, Moorepark, Fermoy,  
10 Cork, Ireland

11 E-mail addresses:

12 annalisa.costa@unito.it (A. Costa)

13 tiziana.civera@unito.it (T. Civera)

14 antonio.lourenco@teagasc.ie (A. Lourenco)

15  
16  
17  
18 <sup>\*</sup>Corresponding author: Laboratório de Microbiologia, DRAT, Instituto Superior de Agronomia,  
19 University of Lisbon, Tapada da Ajuda 1349-017 Lisbon, Portugal.

20 Tel.:+ 351 21 365 3435/3240; fax: +351 21 365 3383

21 E-mail address: lbrito@isa.ulisboa.pt (L. Brito)

22 **ABSTRACT**

23 Five *L. innocua* and five *L. monocytogenes*, including persistent and non-persistent isolates  
24 collected from Gorgonzola processing plants, were compared regarding their biofilm-forming  
25 ability and their biofilm susceptibility to two hydrogen peroxide (HP) based disinfectants in use at  
26 the plants. No significant difference in biofilm-forming ability by both species was observed  
27 ( $P>0.05$ ) in crystal violet staining and viable count assays. The susceptibility to HP disinfectants of  
28 the *L. monocytogenes* and *L. innocua* biofilms was determined. In order to mimic clean and soiled  
29 biofilm forming conditions, biofilms were grown, respectively, in 1/10 diluted TSB-YE and in  
30 TSB-YE. The results showed no significant differences between species or conditions ( $P>0.05$ )  
31 regardless of whether the isolates were classified as persistent or non-persistent. A hierarchical  
32 clustering based on Principal Component Analysis performed on the tested variables, indicated the  
33 presence of two major clusters. Persistent and non-persistent isolates from both species were  
34 allocated in both clusters, suggesting that they behaved in a similar way in response to the tested  
35 conditions. This study showed that biofilms of in-house *L. innocua* could monitor the effectiveness  
36 of HP-based disinfectants. Moreover, biofilms of *L. innocua* could be used as surrogates of *L.*  
37 *monocytogenes* in sanitizer-based biofilm eradication trials simulating dairy processing  
38 environments, whenever the use of the pathogen is not an option.

39

40

41

42 **KEYWORDS**

43 *Listeria innocua*; *Listeria monocytogenes*; biofilms; disinfectants; surrogate.

## 44 1. INTRODUCTION

45 *Listeria innocua* and *Listeria monocytogenes* have been isolated from the same foods (Lappi et al.,  
46 2004; Moshtaghi & Mohamadpour, 2007; Simmons et al., 2014; Vongkamjan, Fuangpaiboon,  
47 Turner, & Vuddhakul, 2016) and food processing environments (Chambel et al., 2007; Lappi et al.,  
48 2004; Nucera, Morra, & Grassi, 2011; Rørvik, Caugant, & Yndestad, 1995). Sauders et al. (2012)  
49 found that both species were associated with urban environments, in contrast with *L. seeligeri* and  
50 *L. welshimeri* that were associated with natural environments. Also phylogenetic analysis showed  
51 that *L. innocua* and *L. monocytogenes* are closely related species (den Bakker et al., 2010; Glaser et  
52 al., 2001; Schmid et al., 2005).

53 Reviews on *L. monocytogenes* sanitizer susceptibility and persistence have pointed out the difficulty  
54 to extrapolate from laboratory-based results to a food processing environment and the need to better  
55 understand the involved mechanisms (Carpentier & Cerf, 2011; Ferreira, Wiedmann, Teixeira, &  
56 Stasiewicz, 2014). Since *L. innocua* is nonpathogenic, it might be possible to employ this species  
57 directly in trials in processing plant environments to investigate its persistence and sanitizer  
58 susceptibility rendering valuable data to predict *L. monocytogenes* behavior. In fact, *L. innocua* has  
59 already been proposed as a surrogate, although not in biofilm state, in order to predict the response  
60 of *L. monocytogenes* to chemical and physical stresses (Delaquis, Stanich, & Toivonen, 2005;  
61 Fairchild & Foegeding, 1993; Friedly et al., 2008; Silva-Angulo et al., 2015).

62 *L. monocytogenes* is a concern in the production of Gorgonzola cheese, an Italian blue-veined  
63 cheese made of pasteurized cow's milk. Some *L. monocytogenes* strains can persist in Gorgonzola  
64 processing plants, suggesting niche adaptation to the dairy environment (Lomonaco et al., 2009).

65 The reasons for persistence are not known, but strong biofilm forming ability and disinfectant  
66 susceptibility do not seem to be prerequisites (Costa, Bertolotti, Brito, & Civera, 2016). The present  
67 study aimed to investigate whether *L. innocua* biofilms may be used as surrogates of the pathogenic  
68 *L. monocytogenes* biofilms. Persistent and non-persistent *L. innocua* and *L. monocytogenes* isolates

69 from Gorgonzola processing environments were selected and their biofilm-forming ability as well  
70 as biofilm susceptibility to two in use hydrogen peroxide (HP) based disinfectants were compared.

## 71 **2. MATERIALS AND METHODS**

### 72 **2.1 Bacterial isolates**

73 The strains used in this work, five *L. innocua* and five *L. monocytogenes*, were isolated from  
74 Gorgonzola cheese processing plants and the farms supplying milk, located in Piedmont and  
75 Lombardy, Italy (Table1). Isolates were classified as persistent if they were found repeatedly within  
76 the collection periods indicated in Table 1, and were at least 95% similar according to repetitive  
77 element sequence-based PCR assays (rep-PCRs) (Nucera et al., 2011; Nucera, Lomonaco, Costa,  
78 Morra, & Grassi, 2013) using ERIC and REP primers (Versalovic, Koeuth, & Lupski, 1991)

### 79 **2.2 Evaluation of biofilm-forming ability and biofilm susceptibility**

80 The evaluation of biofilm-forming ability was performed by quantification of biofilm biomass on  
81 microtiter plates (polystyrene), using both the crystal violet (CV) method (Borucki, Peppin, White,  
82 Loge, & Call, 2003) and a method of enumeration of viable cells on stainless steel coupons (SSC),  
83 as described by Costa et al. (2016). Biofilms were grown in tryptic soy broth with 0.6% yeast  
84 extract (TSB-YE, Biokar Diagnostics, Beauvais, France) at 25 °C for 24 (CV) and 48 h (SSC). The  
85 CV approach was replicated at least three times on biologically independent cultures on distinct  
86 days (biological replicates), with six repetitions under identical conditions each (technical  
87 replicates). For the cell enumeration, two biological replicates were performed, with two technical  
88 replicates each.

89 The evaluation of the biofilm susceptibility to the disinfectants was performed on biofilms grown  
90 on SCC in nutrient-limiting (1/10 diluted TSB-YE) and in nutrient-rich (TSB-YE) medium at 25 °C  
91 for 48 h, as described by Costa et al. (2016). Two commercial HP-based disinfectants, commonly  
92 employed at the dairies where the isolates were collected, were used: P3-oxonia active (ECOLAB  
93 S.r.l) containing acetic acid and peracetic acid (designated herein as P3), and Mida San 315

94 (Christeyns Food Hygiene S.r.l) containing citric acid (designated herein as MS). Disinfectants  
95 were diluted using sterile hard water, prepared according to EN 13697 (Anonymous, 2001), to  
96 achieve the concentrations recommended by the manufacturers: 0.2% (v/v) and 0.5% (v/v) for P3,  
97 and 0.5% (v/v) and 1% (v/v) for MS. Disinfectant efficiency was defined according to EN 13697  
98 (Anonymous, 2001), that states that a minimum 4-log reduction in viable cells is required. If that  
99 threshold was obtained in 2.5 min with the lowest concentration, no additional treatments were  
100 performed. When a 4-log reduction was not achieved, 5- and 7.5-min treatments with higher  
101 disinfectant concentrations, when needed, were carried out until a 4-log reduction was observed.  
102 For each isolate, a control was exposed to sterile hard water and used for the calculation of the log  
103 reduction. Each treatment was repeated under identical conditions on two distinct biofilms.

### 104 **2.3 Data analysis**

105 Agreement to a normal distribution of the data generated by the CV ( $A_{600}$ ), enumeration on SSC  
106 ( $\log \text{CFU}/\text{cm}^2$ ), and biofilm susceptibility (reduction of  $\log \text{CFU}/\text{cm}^2$ ) assays was checked using  
107 the Shapiro-Wilk test, and the homogeneity of the variance was confirmed by Levene's test.  
108 Comparisons between means were then performed via one-way ANOVA, using Scheffé test. These  
109 analyses were performed with the OriginPro 8 SR0 (Northampton, MA, USA) software.

110 Using JMP Pro 13.2.1 (SAS, Cary, NC USA) software, a principal component analysis (PCA) was  
111 performed in order to compare the isolates, based on their response to the ten following conditions  
112 of  $\log \text{CFU}/\text{cm}^2$  reduction on SSC after disinfectant (P3 or MS) treatment of biofilms grown in  
113 clean (C) and soiled (S) conditions, and of biofilm formation: 1) P3 2.5 min 0.2% C; 2) P3 2.5 min  
114 0.2% S; 3) P3 5 min 0.2% C; 4) P3 5 min 0.2% S; 5) MS 2.5 min 0.5% C; 6) MS 2.5 min 0.5% S;  
115 7) MS 5 min 0.5% C; 8) MS 5 min 0.5% S; 9) biofilm formation by CV; and 10) biofilm formation  
116 by enumeration on SSC. A hierarchical clustering using the Ward method was performed based on  
117 the component 1, which explained the majority of the variance.

118 For all tests, the confidence level for significance was 95 % ( $P < 0.05$ ).

## 119 3. RESULTS

### 120 3.1 Comparison of the biofilm-forming ability of the isolates

121 The biofilm formation of *L. innocua* isolates ranged from 0.117 and 0.170 ( $A_{600}$ ) using the CV  
122 method while the enumeration method registered values between 6.20 and 6.69 log CFU/cm<sup>2</sup> (Fig.  
123 1). When *L. innocua* isolates biofilm-formation values are presented alongside *L. monocytogenes*  
124 values (Costa et al., 2016), it is possible to observe that *L. innocua* presented a similar range for  
125 both the CV method (0.087-0.270) and enumeration method (5.65-6.74 log CFU/cm<sup>2</sup>) (Fig.1).

126 The CV assay relies on the property of the dye to bind to negatively charged surface molecules and  
127 polysaccharides in the matrix. This assay may be influenced by the amount of exocellular polymer  
128 and by cell sedimentation, which increases with planktonic growth (Lourenco, Rego, Brito, &  
129 Frank, 2012). This may explain data variability obtain with this method. However, the CV method  
130 showed no significant differences between isolates nor between species ( $P>0.05$ , Fig. 1B). The  
131 same outcome was obtained for the enumeration on SSC, except for *L. monocytogenes* isolate GR,  
132 that was significantly different from *L. monocytogenes* isolates GI and GN and from *L. innocua*  
133 isolate 2, with P-values of 0.011, 0.009 and 0.016, respectively (Fig. 1A).

### 134 3.2 Comparison of the antibiofilm activity of the disinfectants

135 The log reductions obtained after the treatment of *L. innocua* biofilms (light dots) with the  
136 disinfectants P3 and MS are shown in Fig. 2. In order to allow comparison between species, data  
137 from *L. monocytogenes* isolates (dark dots) (Costa et al., 2016) are also shown. Among the isolates  
138 that did not reach the 4-log reduction threshold, and for that reason were consequently exposed to  
139 longer treatments and/or higher concentrations of P3 and MS, *L. monocytogenes* and *L. innocua*  
140 persistent and non-persistent isolates were found. This was observed when biofilms were grown  
141 both in nutrient-limiting (1/10 diluted TSB-YE) and in nutrient-rich (TSB-YE) conditions (Fig. 2A-  
142 I, quadrants II, III and IV). Moreover, when the log reduction values obtained with both  
143 disinfectants at all the tested concentrations and contact times were compared, there were no

144 significant differences between isolates nor between species ( $P>0.05$ ). The exception was the  
145 comparison of susceptibility to P3 at the mildest exposure conditions (0.2% [v/v] for 2.5 min) in  
146 which *L. monocytogenes* presented a higher log reduction (P-value of 0.0497).

147 The disinfectant P3 showed a greater efficacy than MS at 7.5 minutes of exposure and at the lowest  
148 concentration indicated by the manufacturer (0.2% [v/v]), P3 was effective for all the isolates Fig.  
149 2A-C) and the treatment with the higher concentration (0.5% [v/v]) was not required. For the same  
150 exposure time, the lowest concentration of MS (0.5% [v/v]) was not enough to achieve a 4-log  
151 reduction by all the isolates (Fig. 2D-F) and a higher concentration (1% [v/v]) was needed (Fig. 2G-  
152 I).

### 153 **3.3 Comparison of *L. monocytogenes* and *L. innocua* by PCA**

154 Principal Component Analysis (PCA) of data from the 10 isolates regarding biofilm forming ability  
155 (CV and SSC) and disinfectant (P3 and MS) susceptibility was performed. Data were from biofilms  
156 produced in nutrient-limiting (clean) (1/10 diluted TSB-YE) and in nutrient-rich (soiled) (TSB-YE)  
157 conditions. The log reduction values of the 10 isolates exposed to the mildest disinfectant  
158 concentrations (0.2% for P3 and 0.5% for MS) for 2.5 and 5 minutes were considered (Fig. 2A, B,  
159 D and E). These were the treatments applied to the 10 isolates, since isolates that after these  
160 treatments reached the 4-log reduction threshold were not subsequently exposed.

161 By PCA, the initial 10-dimensional space (10 variables) was reduced to a plane F1F2, defined by  
162 the two first principal components. This plane accounts for about 68.5% of the variance of the  
163 original data (Fig. 3). The projection of the 10 original variables on the first two principal  
164 components is presented in Fig. 3A. Except for the treatment with P3 for 5 min at 0.2% in both  
165 clean and soiled conditions, in general, disinfectant susceptibility (log reduction) shows a positive  
166 correlation with the first component increasing along it. The second principal component is  
167 positively correlated with both CV and SSC values, i.e. biofilm production increase along this axis.



168 The projection of the different isolates in the plane F1F2 is presented in Fig. 3B. The isolates in  
169 quadrant I (2, GN, GI and 4) and in quadrant II (1) are more susceptible to P3 as all of them reached  
170 the 4-log reduction threshold after exposure for 5 min to 0.2% (v/v) (Fig. 3, and Fig. 2B). Isolates in  
171 quadrants III (99, 5, 3 and G39) and IV (GR) were less susceptible to P3. These isolates only  
172 achieved the 4-log reduction after a 7.5 min exposure to the same concentration (Fig. 3, and Fig.  
173 2C). Isolate GR can be seen isolated on quadrant IV as it presented the higher log reductions in both  
174 clean and soiled condition for the eight treatments used to perform the PCA (Fig 3, and Fig. 2A, B,  
175 D and E). Moreover, isolate GR is worst biofilm producer than three (2, GN and GI) of the four  
176 isolates positioned in quadrant II (Fig. 3 and Fig. 1).

177 A hierarchical clustering, based on the first principal component which explains the majority  
178 (43.9%) of the variance (Fig. 3) was performed and allowed to confirm the presence of the three  
179 clusters of isolates (C1, C2 and C3) (Fig. 4), as suggested by PCA. A two-way clustering was  
180 performed. According to the intensity of the response of each isolate in every experiment, the  
181 treatments with P3 for 5 min at 0.2%, in both clean and soiled conditions, were the ones that  
182 allowed the most differentiation of the isolates. It is also possible to observe that, persistent and  
183 non-persistent isolates from both species *L. innocua* and *L. monocytogenes* were allocated in both  
184 clusters, confirming that both species respond in a similar way to the conditions tested in this work.

#### 185 **4. DISCUSSION**

186 The use of *Listeria* spp. as an indicator of a possible contamination by *L. monocytogenes* has been  
187 suggested by some authors and guidelines (Food and Drug Administration [FDA], 2008;  
188 Pennsylvania State University [Penn State], 2003; Tompkin, Scott, Bernard, Sveum, & Gombas,  
189 1999). This may be a conservative approach since in Gorgonzola processing plant *L. innocua* was  
190 far more frequent than *L. monocytogenes* (Nucera et al., 2001). In fact, repeated positive testing for  
191 *Listeria* spp. requires more stringent cleaning and disinfecting procedures and indicates the need to  
192 elucidate the reasons for these positive results.

193 Meylheuc, Giovannacci, Briandet, and Bellon-Fontaine (2002) have compared the bioadhesive  
194 behavior of both species and concluded that the non-pathogenic strain exhibited a more marked  
195 electronegative character and a slightly more hydrophilic nature than *L. monocytogenes*.  
196 Nevertheless, to our knowledge, only two studies have previously compared the biofilm production  
197 of *L. monocytogenes* and *L. innocua*: Zhou et al. (2011) used the CV method and concluded that *L.*  
198 *innocua* is a weaker biofilm former compared to *L. monocytogenes*; Koo, Ndahetuye, O'Bryan,  
199 Ricke, and Crandall (2014) used the cell enumeration method on aluminum and stainless steel and  
200 concluded that after 24 h the attachment of *L. monocytogenes* was significantly higher than that of  
201 *L. innocua*, though no significant differences were observed between both species biofilms after 72  
202 h. In the present work, two methods were used (CV in polystyrene P96 microtiter plates and  
203 enumeration on SSC) to evaluate the biofilm forming ability of a set of *L. innocua* and *L.*  
204 *monocytogenes* isolates fairly representative of the contaminant microorganisms, collected from  
205 Gorgonzola processing plants. The comparison performed here, indicated that both species  
206 produced similar values, suggesting an equivalent biofilm production.

207 Regarding disinfectant susceptibility, Best, Kennedy, and Coates (1990) tested the efficacy of 14  
208 disinfectants against both species after been spotted onto the surface of stainless steel disks and  
209 dried for 30 minutes. The obtained results showed that the pathogenic species was slightly less  
210 susceptible to disinfection than *L. innocua*. A few other comparative studies have assessed  
211 differences in the susceptibility of planktonic cells of both species to disinfectants finding no  
212 differences between species (Margolles, Mayo & de los Reyes-Gavilán, 2000) or highlighting an  
213 higher resistance of planktonic *L. innocua* (Yeater, Kirsch, Taylor, Mitchell, & Osburn, 2015).

214 In this work, using *L. innocua* and *L. monocytogenes* collected from the same environment, a  
215 similar susceptibility to the tested disinfectants was found for biofilms of both species. Moreover,  
216 the susceptibility to P3 and to MS of biofilms grown in conditions mimicking clean and soiled  
217 environment, showed no significant differences in terms of log reduction between persistent and  
218 non-persistent isolates. According to these results, *L. innocua* could be used as a surrogate for *L.*

219 *monocytogenes*, not only regarding the biofilm production, but also the biofilm susceptibility to HP-  
220 based disinfectants.

221 *L. innocua* was the only *Listeria* species besides *L. monocytogenes* detected in the Gorgonzola  
222 processing plants from where the isolates analyzed in this work were collected (Nucera et al., 2011).  
223 Our results demonstrate that the presence of *L. innocua* could indicate a contamination by *L.*  
224 *monocytogenes*, since the disinfectant susceptibility of the two species was similar. In fact, as  
225 concluded in the review by Milillo et al. (2012), if the two species have adapted to fit different  
226 environmental niches, they may not always respond to stress the same way. Therefore, the absence  
227 of *Listeria* spp. (i.e. *L. innocua* and *L. monocytogenes*) on food contact surfaces, equipment and  
228 floors would suggest the effectiveness of the sanitation procedures in place in the processing plant.  
229 Conversely, the detection of positive samples for *Listeria* spp. would indicate a need of improving  
230 the procedures to keep the environmental contamination under control, as suggested by Tompkin et  
231 al. (1999). Moreover, Zitz, Zunabovic, Domig, Wilrich, and Kneifel (2011) also verified reduced  
232 detectability of *L. monocytogenes* in the presence of *L. innocua* mainly due to the overgrowth of *L.*  
233 *monocytogenes* by *L. innocua* during the selective enrichment, leading to false-negative results.  
234 Furthermore, the use of *Listeria* spp. as an indicator of a potential *L. monocytogenes* contamination  
235 represents lower costs for routine laboratory analysis, due to the higher cost of chromogenic media  
236 used for *L. monocytogenes* (Tomkin, 2002).

237 The presented work tried to mimic food industry conditions. Consequently, susceptibility testing  
238 was conducted with biofilms produced on stainless steel, in soiled and in clean conditions, and  
239 testing two commonly used HP-based disinfectants at the dairy plants where the isolates were  
240 collected from. Nevertheless, in the real industrial environment the strains will form multi-species  
241 biofilms and that may become relevant to the response of both *L. monocytogenes* and *L. innocua* to  
242 disinfectants.

243

244 **4. CONCLUSION**

245 This study has shown that biofilms of in-house *L. innocua* could be employed for the validation and  
246 monitoring of HP-based disinfectant efficacy and proper sanitation procedures in Gorgonzola  
247 processing plants. In fact, not only *L. innocua* susceptibility to HP-based disinfectants is similar to  
248 *L. monocytogenes* biofilms, but also both species, collected from the same food industry  
249 environment, showed no differences in biofilm forming ability. The common origin of the isolates  
250 is probably fundamental when looking for the adequacy of *L. innocua* as a surrogate of *L.*  
251 *monocytogenes*. The convenience and safety in using a non-pathogenic surrogate will certainly  
252 contribute to clarify the factors that contribute to *L. monocytogenes* persistent colonization not only  
253 in some Gorgonzola processing plants as in other food industry environments.

254 **ACKNOWLEDGMENTS**

255 The authors thank to Ana Carla Silva for the support with the work performed in Laboratory of  
256 Microbiology (ISA/ULisboa).

257 **FUNDING**

258 This work was supported by the Italian Ministry of University and Research (Progetto Giovani) and  
259 Piedmont Region (Progetto BIOPACK-175-1). It was also partially supported by funds allocated to  
260 the LEAF/ISA-University of Lisboa and to the University of Turin.

261 **Declarations of interest:** none.

262

263 **REFERENCES**

- 264 Anonymous (2001). European Standard EN 13697: Chemical disinfectants and antiseptics –  
265 qualitative non-porous surface test for the evaluation of bacteriocidal and/or fungicidal  
266 activity of chemical disinfectants used in food, industrial, domestic and institutional areas –  
267 test method and requirements without mechanical action (phase 2, step 2).
- 268 Best, M., Kennedy, M. E., & Coates, F. (1990). Efficacy of a variety of disinfectants against  
269 *Listeria* spp. *Applied and Environmental Microbiology*, 56, 377–380.
- 270 Borucki, M. K., Peppin, J. D., White, D., Loge, F., & Call, D. R. (2003). Variation in biofilm  
271 formation among strains of *Listeria monocytogenes*. *Applied and Environmental*  
272 *Microbiology*, 69, 7336–7342.
- 273 Carpentier, B., & Cerf, O. (2011). Review-Persistence of *Listeria monocytogenes* in food industry  
274 equipment and premises. *International Journal of Food Microbiology*, 145, 1–8.
- 275 Chambel, L., Sol, M., Fernandes, I., Barbosa, M., Zilhão, I., Barata, B., Jordan, S., Perni, S., Shama,  
276 G., Adrião, A., Faleiro, L., Requena, T., Peláez, C., Andrew, P.W., & Tenreiro, R. (2007).  
277 Occurrence and persistence of *Listeria* spp. in the environment of ewe and cow’s milk  
278 cheese dairies in Portugal unveiled by an integrated analysis of identification, typing and  
279 spatial-temporal mapping along production cycle. *International Journal of Food*  
280 *Microbiology*, 116, 52–63.
- 281 Costa, A., Bertolotti, L., Brito, L., & Civera, T. (2016). Biofilm formation and disinfectant  
282 susceptibility of persistent and nonpersistent *Listeria monocytogenes* isolates from  
283 Gorgonzola cheese processing plants. *Foodborne Pathogens and Disease*, 13, 602–609.
- 284 Delaquis, P., Stanich, K., & Toivonen, P. (2005). Effect of pH on the inhibition of *Listeria* spp. by 3  
285 vanillin and vanillic acid. *Journal of Food Protection*, 68, 1472–1476.
- 286 den Bakker, H. C., Cummings, C. A., Ferreira, V., Vatta, P., Orsi, R. H., Degoricija, L., Barker, M.,  
287 Petrauskene, O., Furtado, M. R., & Wiedmann, M. (2010). Comparative genomics of the

288 bacterial genus *Listeria*: genome evolution is characterized by limited gene acquisition and  
289 limited gene loss. *BMC Genomics*, 11, 688.

290 Fairchild, T. M., & Foegeding, P. M. (1993). A proposed nonpathogenic biological indicator for  
291 thermal inactivation of *Listeria monocytogenes*. *Applied and Environmental Microbiology*,  
292 59, 1247-1250.

293 Ferreira, V., Wiedmann, M., Teixeira, P., & Stasiewicz, M. J. (2014). *Listeria monocytogenes*  
294 persistence in food-associated environments: epidemiology, strain characteristics, and  
295 implications for public health. *Journal of Food Protection*, 77, 150–170.

296 Food and Drug Administration (FDA) (2008). Guidance for industry: control of *Listeria*  
297 *monocytogenes* in refrigerated or frozen ready-to-eat foods.

298 Friedly, E. C., Crandall, P. G., Ricke, S., O’Bryan, C. A., Martin, E. M., & Boyd, L. M. (2008).  
299 Identification of *Listeria innocua* surrogates for *Listeria monocytogenes* in hamburger  
300 patties. *Journal of Food Science*, 73, M174–M178.

301 Glaser, P., Frangeul, L., Buchrieser, C., Rusniok, C., Amend, A., Baquero, F., Berche, P., Bloecker,  
302 H., Brandt, P., Chakraborty, T., Charbit, A., Chetouani, F., Couve, E., de Daruvar, A.,  
303 Dehoux, P., Domann, E., Dominguez-Bernal, G., Duchaud, E., Durant, L., Dussurget, O.,  
304 Entian, K. D., Fsihi, H., Garcia-Del Portillo, F., Garrido, P., Gautier, L., Goebel, W.,  
305 Gomez-Lopez, N., Hain, T., Hauf, J., Jackson, D., Jones, L. M., Kaerst, U., Kreft, J., Kuhn,  
306 M., Kunst, F., Kurapkat, G., Madueno, E., Maitournam, A., Vicente, J. M., Ng, E., Ndjari,  
307 H., Nordsiek, G., Novella, S., de Pablos, B., Perez-Diaz, J.C., Purcell, R., Remmel, B.,  
308 Rose, M., Schlueter, T., Simoes, N., Tierrez, A., Vazquez-Boland, J. A., Voss, H., Wehland,  
309 J., & Cossart, P. (2001). Comparative genomics of *Listeria* species. *Science*, 294, 849–852.

310 Koo, O. K., Ndahetuye, J. B., O’Bryan, C. A., Ricke, S. C., & Crandall, P. G. (2014). Influence of  
311 *Listeria innocua* on the attachment of *Listeria monocytogenes* to stainless steel and  
312 aluminum surfaces. *Food Control*, 39, 135–138.

- 313 Lappi, V. R., Thimothe, J., Walker, J., Bell, J., Gall, K., Moody, M. W., & Wiedmann, M. (2004).  
314 Impact of intervention strategies on *Listeria* contamination patterns in crawfish processing  
315 plants: a longitudinal study. *Journal of Food Protection*, 67, 1163–1169.
- 316 Lomonaco, S., Decastelli, L., Nucera, D., Gallina, S., Bianchi, D. M., & Civera, T. (2009). *Listeria*  
317 *monocytogenes* in Gorgonzola: subtypes, diversity and persistence over time. *International*  
318 *Journal of Food Microbiology*, 128, 516–520.
- 319 Lourenco, A., Rego, F., Brito, L., & Frank, J. F. (2012). Evaluation of methods to assess the  
320 biofilm-forming ability of *Listeria monocytogenes*. *Journal of Food Protection*, 75, 1411–7.
- 321 Moshtaghi, H., & Mohamadpour, A. A. (2007). Incidence of *Listeria* spp. in raw milk in  
322 Shahrekord, Iran. *Foodborne Pathogens Disease*, 4, 107–110.
- 323 Meylheuc, T., Giovannacci, I., Briandet, R., & Bellon-Fontaine, M.-N. (2002). Comparison of the  
324 cell surface properties and growth characteristics of *Listeria monocytogenes* and *Listeria*  
325 *innocua*. *Journal of Food Protection*, 65, 786–793.
- 326 Milillo, S. R., Friedly, E. C., Saldivar, J. C., Muthaiyan, A., O’Bryan, C., Crandall, P. G., Johnson,  
327 M. G., & Ricke, S. C. (2012). A review of the ecology, genomics, and stress response of  
328 *Listeria innocua* and *Listeria monocytogenes*. *Critical Reviews in Food Science and*  
329 *Nutrition*, 52, 712–725.
- 330 Margolles, A., Mayo, B., & de los Reyes-Gavilán, C. G. (2000). Phenotypic characterization of  
331 *Listeria monocytogenes* and *Listeria innocua* strains isolated from short-ripened cheeses.  
332 *Food Microbiology*, 17, 461–467.
- 333 Nucera, D., Morra, P., & Grassi, M. A. (2011). Detection and characterisation of *Listeria* isolates  
334 collected along the Gorgonzola PDO production chain. *Italian Journal of Food Safety*, 1,  
335 25–31.
- 336 Nucera, D. M., Lomonaco, S., Costa, A., Morra, P., & Grassi, M. A. (2013). Diagnostic  
337 performance of rep-PCR as a rapid subtyping method for *Listeria monocytogenes*. *Food*  
338 *Analytical Methods*, 6, 868–871.

339 Pennsylvania State University (Penn State), College of Agricultural Sciences, Agricultural Research  
340 and Cooperative Extension (2003). Control of *Listeria monocytogenes* in Small Meat and  
341 Poultry Establishments.

342 Rørvik, L. M., Caugant, D. A., & Yndestad, M. (1995). Contamination pattern of *Listeria*  
343 *monocytogenes* and other *Listeria* spp. in a salmon slaughterhouse and smoked salmon  
344 processing plant. *International Journal of Food Microbiology*, 25, 19–27.

345 Sauders, B. D., Overdevest, J., Fortes, E., Windham, K., Schukken, Y., Lembo, A., & Wiedmann,  
346 M. (2012). Diversity of *Listeria* species in urban and natural environments. *Applied and*  
347 *Environmental Microbiology*, 78, 4420–4433.

348 Schmid, M. W., Ng, E. Y. W., Lampidis, R., Emmerth, M., Walcher, M., Kreft, J., Goebel, W.,  
349 Wagner, M., & Schleifer, K.-H. (2005). Evolutionary history of the genus *Listeria* and its  
350 virulence genes. *Systematic and Applied Microbiology*, 28, 1–18.

351 Silva-Angulo, A. B., Zanini, S. F., Rosenthal, A., Rodrigo, D., Klein, G., & Martínez, A. (2015).  
352 Comparative study of the effects of citral on the growth and injury of *Listeria innocua* and  
353 *Listeria monocytogenes* cells. *PLoS ONE*, 10, e0114026.

354 Simmons, C., Stasiewicz, M. J., Wright, E., Warchocki, S., Roof, S., Kause, J. R., Bauer, N.,  
355 Ibrahim, S., Wiedmann, M., & Oliver, H. F. (2014). *Listeria monocytogenes* and *Listeria*  
356 spp. contamination patterns in retail delicatessen establishments in three U.S. States. *Journal*  
357 *of Food Protection*, 77, 1929–1939.

358 Tompkin, R. B. (2002). Control of *Listeria monocytogenes* in the food-processing environment.  
359 *Journal of Food Protection*, 65, 709–725.

360 Tompkin, R. B., Scott, V. N., Bernard, D. T., Sveum, W. H., & Gombas, K. S. (1999). Guidelines  
361 to prevent post-processing contamination from *Listeria monocytogenes*. *Dairy, Food and*  
362 *Environmental Sanitation*, 19, 551–603.



363 Versalovic, J., Koeuth, T., & Lupski, J.R. (1991). Distribution of repetitive DNA sequences in  
364 Eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Research*,  
365 19, 6823–6831.


366 Vongkamjan, K., Fuangpaiboon, J., Turner, M. P., & Vuddhakul, V. (2016). Various ready-to-eat  
367 products from retail stores linked to occurrence of diverse *Listeria monocytogenes* and  
368 *Listeria* spp. isolates. *Journal of Food Protection*, 79, 239–245.

369 Yeater, M. C., Kirsch, K. R., Taylor, T. M., Mitchell, J., & Osburn, W. N. (2015). Effectiveness of  
370 sanitizing products on controlling selected pathogen surrogates on retail deli slicers. *Journal*  
371 *of Food Protection*, 78, 707–715.

372 Zhou, Q., Feng, F., Wang, L., Feng, X., Yin, X., & Luo, Q. (2011). Virulence regulator PrfA is  
373 essential for biofilm formation in *Listeria monocytogenes* but not in *Listeria innocua*.  
374 *Current Microbiology*, 63, 186–192.

375 Zitz, U., Zunabovic, M., Domig, K. J., Wilrich, P.-T., & Kneifel, W. (2011). Reduced detectability  
376 of *Listeria monocytogenes* in the presence of *Listeria innocua*. *Journal of Food Protection*,  
377 74, 1282–1287.

378



379

380 **Fig. 1 Biofilm forming ability of *Listeria monocytogenes* (grey bars) and *Listeria innocua***  
 381 **(white bars). Oblique line bars represent persistent isolates; A - assessed by cell enumeration on**  
 382 **SSC (grown in TSB-YE for 48 h at 25 °C). The isolate GR, only on region 1 is statistically different**  
 383 **from isolates 2, GI and GN, on region 3 (regions marked on the right), The isolates on region 2 are**  
 384 **neither statistically different from isolates on region 1 nor from region 3. Two biological replicates**  
 385 **with two technical replicates each were performed; B - assessed by crystal violet (CV) method in**  
 386 **polystyrene 96-well microtiter plates (grown in TSB-YE for 24 h at 25 °C). Isolates were not**  
 387 **statically different ( $P > 0.05$ ). *L. innocua* isolates 1 and 2 and *L. monocytogenes* isolates GR and GI**  
 388 **are persistent isolates (Table 1). Results from *L. monocytogenes* were previously published by**  
 389 **Costa et al. (2016) and are shown here only for comparison. Error bars represent standard**  
 390 **deviations. At least three biological replicates were performed, with six technical replicates, each.**

391

392

393

394 **Fig. 2 Log reductions (log CFU/cm<sup>2</sup>) obtained after disinfectant treatment of *L. innocua* (light**  
395 **dots) and *L. monocytogenes* (dark dots) biofilms grown for 48 h at 25 °C on SSC, in clean**  
396 **conditions (1/10 TSB-YE; x-axis) and soiled conditions (TSB-YE; y-axis), using P3 at 0.2% or**  
397 **MS at 0.5% and 1%. The isolates placed in quadrant III did not reach the 4-log reduction neither**  
398 **with biofilms formed in soiled conditions nor with biofilms formed in clean conditions and further**  
399 **treatments with extended contact time/increased disinfectant concentration were performed, as**  
400 **needed. The isolates placed in quadrant I reached the reduction threshold of 4 logs, with biofilms**  
401 **formed in both conditions and, for this reason, no other treatments were performed. The isolates**  
402 **placed in quadrant II and IV reached the 4 log reduction threshold, respectively, only in soiled or**  
403 **clean biofilm forming conditions. Further treatments respectively with biofilms grown in clean and**  
404 **soiled conditions, were carried out in order to achieve the 4-log reduction in both conditions.**  
405 **Isolates that underwent treatment only with biofilms produced under clean/soiled condition are**  
406 **marked with an asterisk (\*) and placed on x- or y-axis, respectively. Isolates' ID in bold represent**  
407 **persistent isolates. Results from *L. monocytogenes* were previously published by Costa et al. (2016)**  
408 **and are shown here only for comparison. Error bars represent standard deviations. For each**  
409 **treatment, two technical replicates were performed.**