

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Mechanochemical activation by ultrasound

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1663910> since 2018-03-26T22:24:07Z


*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

ISMech 2013



International  
Symposium  
on Mechanochemistry  
in Synthesis  
and Nanoscience

## PROGRAMME & ABSTRACTS

21-24 September 2013  
Warsaw-Ossa, Poland



## Mechanochemical activation by ultrasound

Giancarlo Cravotto,<sup>1\*</sup> Emanuela Calcio Gaudino,<sup>1</sup> Pedro Cintas.<sup>2</sup>

<sup>1</sup>Dipartimento di Scienza e Tecnologia del Farmaco, Università di Torino, I-10125 Torino, Italy.

<sup>2</sup>Departamento de Química Organica e Inorganica, UEX, E-06006 Badajoz, Spain.

\* giancarlo.cravotto@unito.it

The use of mechanical forces to induce chemical reactivity is the oldest and simplest technique used by mankind and one which has existed since before the dawn of science. Mechanical effects can take place in solution and heterogeneous systems, thus accounting for enhanced reactivity.<sup>1</sup> Acoustic cavitation may initiate chemical reactions via the action of mechanical pulses, in close similarity to other, solid-state reactions under mechanical loading. This would otherwise be consistent with the initial and broad concept of mechanochemistry by Ostwald, who used the term for the first time. Ostwald simply recognized that the discipline is a part of physical chemistry that is distinctive from thermochemistry, electrochemistry and photochemistry in terms of energy transmission.<sup>2</sup> Faraday had observed the rapid decomposition and dehydration of some salts under milling and grinding since the early 1820s. The creation of large tensile forces in liquids via simple friction or agitation is not a simple task. Sonication provides a viable exception by virtue of the mechanical and chemical events it causes via the unique phenomenon of acoustic cavitation.<sup>3</sup> The cavitation-based mechanical effects, arise from shear forces, microjets and shock waves that occur outside the bubble, resulting in profound physical changes when solids or metals are present.<sup>4</sup> These changes include improved mass transfer, particle size reduction, surface erosion and cleaning, and are often accompanied by changes in particle properties. Likewise, mechanochemistry can also generate radicals via the breaking of weak bonds and under extreme surface plasma conditions where covalent crystals are cracked by mechanical impact.<sup>5,6</sup> Clearly, both scenarios and strategies are different, although goals may be similar. This presentation is aimed at unveiling the features and paths where tribochemistry and sonochemistry meet and highlighting, whenever possible, comparative aspects.<sup>7,8</sup>

### References

- 1) Cravotto, G.; Calcio Gaudino, E.; Cintas, P. *Chem. Soc. Rev.* **2013**, DOI: 10.1039/c2cs35456j.
- 2) Ostwald, W. *Handbuch der allgemeine Chemie*, Band I, Academische Verlag., Leipzig, 1919, p. 70.
- 3) Cravotto, G.; Cintas, P. *Chem. Soc. Rev.* **2006**, 35, 180-196.
- 4) Cravotto, G.; Cintas, P. *Angew. Chem. Int. Ed.* **2007**, 46, 5476-5478.
- 5) Kaupp, G. in *Making Crystals by Design*, ed. Braga D. and Grepioni, F. Wiley-VCH, New York, 2007, pp. 87-148.
- 6) G. Kaupp, *J. Phys. Org. Chem.* **2008**, 21, 630-643;
- 7) Cravotto, G.; Cintas, P. *Chem. Sci.* **2012**, 3, 295-307.
- 8) Cintas, P.; Cravotto, G. *Nature Chem.* **2012**, 4, 77-78.