AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fungi as a toolbox for sustainable bioremediation of pesticides in soil and water

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1664622> since 2019-08-08T12:21:41Z

Published version:

DOI:10.1080/11263504.2018.1445130

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Fungi as a toolbox for sustainable bioremediation of pesticides in soil and water

Journal:	<i>Plant Biosystems</i>
Manuscript ID	TPLB-2017-0224.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	23-Jan-2018
Complete List of Authors:	<p>Spina, Federica; University of Torino, Department of Life Sciences and Systems Biology</p> <p>Cecchi, Grazia; University of Genoa, Department of Earth, Environment and Life Sciences</p> <p>Landinez-Torres, Angela; Fundación Universitaria Juan de Castellanos; University of Pavia, Department of Earth Science and Environmental Sciences</p> <p>Pecoraro, Lorenzo; Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences</p> <p>Russo, Fabiana; Sapienza University of Rome, Department of Environmental Biology</p> <p>Wu, Bing; Chinese Academy of Sciences, Institute of Microbiology</p> <p>Cai, Lei; Chinese Academy of Sciences, Institute of Microbiology</p> <p>Liu, Xingzhong; Chinese Academy of Sciences, Institute of Microbiology</p> <p>Tosi, Solveig; Università di Pavia,</p> <p>Varese, Giovanna; University of Turin, Department of Plant Biology</p> <p>Zotti, Mirca; Università di Genova, Polo Botanico Hanbury, DiSTAV</p> <p>Persiani, Annamaria; Sapienza Università di Roma,</p>
Keywords:	Pesticides, Agrochemicals, Antibiotics, Sustainable bioremediation, Fungi, Synthetic microbial community, Environmental risk assessment

Fungi as a toolbox for sustainable bioremediation of pesticides in soil and water

F. Spina¹, G. Cecchi², A. Landinez-Torres^{3,4}, L. Pecoraro⁵, F. Russo⁶, B. Wu⁵, L. Cai⁵, X. Z. Liu⁵, S. Tosi⁴, G.C. Varese¹, M. Zotti², A. M. Persiani⁶

AFFILIATION

1 *Mycotheca Universitatis Turinensis, Department of Life Sciences and Systems Biology, University of Torino, Viale Mattioli, 25, 1025 Torino, Italy.*

2 *Laboratory of Mycology, Department of Earth, Environment and Life Sciences, University of Genoa, Corso Europa, 26, 16136 Genoa, Italy*

3 *Facultad de Ciencias Agrarias y Ambientales, Fundación Universitaria Juan de Castellanos. Carrera 11 No. 11 - 44 Tunja, Boyacá - Colombia*

4 *Laboratory of Mycology, Department of Earth Science and Environmental Sciences, University of Pavia, Via S. Epifanio 14, 27100 Pavia, Italy*

5 *State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*

6 *Laboratory of Fungal Biodiversity, Department of Environmental Biology, Sapienza University of Rome, Piazzale A. Moro 5, 00185 Rome, Italy*

Corresponding Author: Anna Maria Persiani, Laboratory of Fungal Biodiversity, Department of Environmental Biology, Sapienza University of Rome, Piazzale A. Moro 5, 00185 Rome, Italy. Email: annamaria.persiani@uniroma1.it

Abstract

Pesticides can help reduce yield losses caused by pests, pathogens, and weeds, but their overuse causes serious environmental pollution. They are persistent in the environment and are biomagnified through the food chain, becoming a serious health hazard for humankind. Bioremediation, where microbes are used to degrade pesticides *in situ*, is a useful technology. This review summarizes data on the fungi involved in the biodegradation of chemical pesticides and their application in soil and water bioremediation. Indications for future studies in this field are given.

Keywords: Pesticides, Agrochemicals, Antibiotics, Sustainable bioremediation, Fungi, Synthetic microbial community, Environmental risk assessment.

Introduction

Because of their unique functions, fungi are involved in ecosystem services essential to human well-being. Among others, fungi also carry out the transformation and detoxification of pollutants. For this reason, learning from nature, they represent an effective toolbox for a sustainable bioremediation of pesticides in soil and water. Many researches have revealed the untapped potential of fungi, and recent years have witnessed very interesting developments regarding the application of fungi not only to improve environmental quality but also human health (e.g. Gargano et al. 2017).

Pesticides are a diverse group of inorganic and organic chemicals that include herbicides, insecticides, nematocides, fungicides, antibiotics and soil fumigants (Verger and Boobis 2013; Verma et al. 2014). They are employed in agriculture to enhance crop yield and quality, and to maximize economic returns by preventing pest or weed attack. They are bioactive, toxic substances, capable of directly or indirectly influencing soil fertility and health as well as agroecosystem quality (Pinto et al. 2012; Verma et al. 2014). Given that belowground biodiversity is closely linked to land management, agricultural intensification exerts many pressures that lead to loss of biodiversity.

Consequently, soil pollution is one of the main threats to the decline of taxonomic and functional biodiversity, and to agricultural soil sustainability (Harms et al. 2017). Most pesticide emission (99 %) in Europe is associated with agricultural practices, whereas industrial and urban sources such as the manufacturing of pesticides or the at-home use of insecticides have a minor impact (EEA 2016).

The extensive and massive use of pesticides in agricultural activities has a serious impact on the environment, compromising soil and water quality (Pinto et al. 2012; Zhang et al. 2015; Pinto et al. 2016). In addition to pesticides, large quantities of antibiotics are added to agricultural fields worldwide through the application of wastewater, manures and biosolids, also resulting in antibiotic contamination and elevated environmental risks (Jechalke et al. 2014; Zhang et al. 2015; Pan and Chu 2016). A clear correlation between agriculture and water contamination was observed in Mar Chiquita lake (Argentina), where large amounts of endosulfan residues were detected soon after application and post-application periods (Ballesteros et al. 2014). The presence of the fungicide thifluzamide in the water in rice paddies in China was maximal after application, with variation over time associated with the dilution effect of rainfalls in the area (Wei et al. 2015).

Preventive measures are required, to mitigate the impact of agriculture on the environment. These must take into account both the use of safe pesticides and the optimization of farmer procedures. Aravinna et al. (2017) found that most of the 32 studied pesticides leached off rice paddies following specific pathways. Since direct runoff and erosion from soil were the main vehicles of dispersion, authors suggested alternative strategies (high resident time for pesticides, holding ponds for rice drainage water, delayed filling of paddies after pesticide application, and the use of less mobile compounds) to reduce the movement of the pesticides.

The intensive use of organic agrochemicals (OACs) poses risks to both wild life and human health. Over 98% of sprayed insecticides and 95% of herbicides reach a destination other than their target species through air, water and soil (Miller 2004). Around 30% of pesticides marketed in developing countries do not meet internationally accepted quality standards, posing a serious threat to human health and the environment (Popp et al. 2013). They are persistent in the environment and are biomagnified through the food chain, and it has been estimated that millions of agricultural workers worldwide experience unintentional pesticide poisoning each year. The correlation between long-term exposure to pesticides in occupational settings and illness is known, but recently non-occupational exposures have also been associated with an elevated rate of chronic diseases (Parrón et al. 2014).

Varieties and consumption of pesticides worldwide have increased dramatically, by up to 4-fold since 40 years ago (Mnif et al. 2011). According to De et al. (2014), about 45 % are used in Europe, 25 % in the USA, and 25 % in the rest of the world. The main pesticide consumer is Spain (around 79,000 ton of active ingredients sold between 2011 and 2014), followed by France (~ 75,000), Italy (~ 64,000), Germany (~ 46,000) and United Kingdom (~ 23,000) (Eurostat 2016). The United States applies over 1 billion pounds annually (Alavanja 2009) with dramatic consequences for human beings and environment (Carvalho 2017). According to other authors (Huang McBeath and McBeath 2010), China is the world's largest pesticide user, with a pesticide output of around 3.7 million tons (National Bureau of Statistics of China - <http://data.stats.gov.cn>), and a consumption of about 1.8 million tons in 2014. More than 350 insecticides, herbicides, microbicides, nematicides and other pesticides are reported to be used. The average amount of pesticides used per hectare in China is roughly 1.5- to 4-fold higher than the world average (Qiu 2011), thus resulting in contamination of water bodies in the receiving areas and disturbance of ecological equilibrium (Hui et al. 2003). Overall, use of pesticides in China breaks down as herbicides 47.5 %, insecticides 29.5 %, fungicides 17.5 % and others 5.5 % (De et al. 2014).

The adverse effects of OAC pollution have been of concern for a long time and many highly toxic and persistent pesticides have been banned worldwide. Although relatively safer pesticides have

been developed and replaced the highly toxic ones, environmental pollution resulting from the long-term application of pesticides is far from being solved. Obsolete pesticides still represent a threat to environment, biodiversity, and human health for the region of Southeast Europe and their risk to the environment and to humans needs to be assessed in order to mitigate it. Many organochlorines, organophosphates and pyrethroids have been banned but this has not yet solved the problem (Aravinna et al. 2017). In Argentina, hexachlorocyclohexane pesticides have been limited since the late '90s and were definitely banned in 2011, but samples taken from a saline lake in 2014 showed levels to be more than 5-fold over the legal limit of 4 ng/l for lindane levels in the environment (Ballesteros et al. 2014). Likewise in China, although the use of organichlorine pesticides has been banned for over 20 years, they can still be found in the water and sediments of main drainage areas (Nakata et al. 2005; Xue et al. 2006; Zhou et al. 2006), due to run-off from aged and weathered agricultural soils and from anaerobic sediments (Zhou et al. 2006). Water bodies and sediments, the water, the soil and even the air in many cities in China are polluted by OACs, in both urban and suburban areas (Gong et al. 2004; Nakata et al. 2005; Yang et al. 2008).

OACs pose pivotal environmental problems, due to their high resistance in the environment and the consequent low natural attenuation. As an example, organochlorine pesticides were poorly affected by photochemical, chemical and biological processes, and more than 95% of them impacted on non-target organisms (Mrema et al. 2013). As a consequence, regulatory and risk assessment procedures have to be adopted against OACs. Driven by the carcinogenicity of pesticides, Directive 91/414/EEC aimed to regulate the authorization of pesticides marketing within the EU.

The particular attention given to pesticides is because, as confirmed in recent studies, even low doses might trigger adverse effects on wildlife and humans (EEA 2005). As groundwater is our primary source of drinking waters, both the Groundwater Directive 2006/118/EC and the Drinking Water Directive 98/83/EC deal with maximum pesticide exposure concentrations: 0.1 µg/l of a single pesticide and 0.5 µg/l total pesticide load. Risk assessment needs to consider not only the source of contamination, but also the multifaceted direct and indirect pathways of contact with human beings. Kim et al. (2017) reported a number of routes pesticides might follow to meet human beings; the resulting direct and indirect multi-pathway exposure may affect human health.

Experimental evidence of progress in natural restoration processes highlight that time is our ally, since the abandonment of disturbed/polluted agricultural land for long time can reduce contamination (Kardol and Wardle 2010). Studies by Morriën et al. (2017) reported that nature restoration on ex-arable land resulted in increased connettance of soil biota networks, as restoration progresses. Such results confirm that soil biota provide many and varied services, and that detoxification of pollutants and xenobiotics is one of the primary ones.

In this context, innovation involves the search for solutions inspired by nature, with the strategy being to accelerate the natural attenuation processes in contaminated sites. Bioremediation has arisen as a useful technology to degrade OACs (Singh 2008; Velázquez-Fernández et al. 2012), with several benefits over landfill disposal and incineration, such as the formation of non-toxic end products, lower costs of disposal, reduction of effects on health and ecology and on the long-term liabilities associated with destructive treatment methods, and the ability to perform the treatment *in situ* without unduly disturbing native ecosystems (Sarkar et al. 2005). Over the past decade, numerous microorganisms capable of degrading antibiotics and pesticides have been isolated, and detoxification processes for target pollutants have been analyzed. Fungi and especially ligninolytic fungi have been suggested as the most promising group of organisms, as they are able to transform recalcitrant compounds through a unique set of extracellular oxidative enzymes (Anastasi et al. 2013; Harms et al. 2017). Comparative genomic analysis of 49 fungi with different nutritional modes, such as saprotrophic fungi, white-rot fungi (WRF), brown-rot fungi, soft rot fungi and symbiotic fungi indicate that there is a relationship between nutrition models and the enzymes for lignocellulose degradation. Saprotrophic fungi have a greater number of enzymes than symbiotic fungi, and brown-rot fungi have a smaller number than WRF and soft rot fungi (Wu et al. 2015a). This might provide some insight into how to choose fungi in OACs degradation.

Finally and importantly, the metabolic activity of fungal or microbial consortia could potentially produce unknown reaction products that are more toxic than the parent compounds. García-Carmona et al. (2017) highlighted the importance of carrying out environmental monitoring activities ante- and post-operam phases, using bioassays to determine the success of the bioremediation process. Although it is fundamental to assess the quality of the environment to ensure it remains free of toxic residues, most of the analytical tests available for determining the concentration of toxic chemicals do not give the biological impacts of toxicants. For this reason, biotoxicity testing has grown steadily in recent years and is a useful tool in environmental risk assessment (Shen et al. 2016; Prokop et al. 2016).

Indeed, there is a clear need to develop and define decontamination of hazardous pollutants as a concept that will support sustainable remediation by involving a broader uptake of principles, approaches and tools that integrate environmental, social and economical dimensions into remediation processes (Ridsdale and Noble 2016). Several organizations, academia and standardization committees are currently assessing remediation process and evaluating the complexity of sustainability. Documents have been developed by many countries across Europe and globally, addressing sustainable indicators for remediation activities (Harclerode et al. 2015).

The present review summarizes the current state of scientific knowledge on research and application of fungi as effective bioresources, considering recent advances in understanding their capacity to face up the pesticide contamination.

Bioremediation of OACs by fungi in the soil system

Large quantities of OACs are being added to agricultural fields worldwide through the application of wastewater, manures and biosolids, resulting in pesticide and antibiotic contamination and elevated environmental risks in terrestrial environments (Jechalke et al. 2014; Zhang et al. 2015; Pan and Chu 2016). A large proportion of the OACs applied to soils with manure or biosolids are retained in surface soil, whereas those added through irrigation with wastewater can seep down to lower horizons or be diffused in surface run-off. Once present, OACs interact with the solid phase of soil and are prone to microbial transformation (Hammesfahr et al. 2008; Jechalke et al. 2014). In particular, veterinary antibiotics interact with the soil solid phase in sorption and desorption reactions. Sorption and desorption control not only their mobility and uptake by plants but also their biotransformation and biological effects. OACs, like microorganisms are not distributed homogeneously in soil but are concentrated in hotspots. The multiplicity of surfaces, voids, and pores provided by soil aggregates harbor a vast amount of biological diversity and chemical variability, and cause patchy distribution of natural organic matter, oxides, nutrients, and microorganisms on soil particle surfaces (Hammesfahr et al. 2008; Jones et al. 2012). Sorption, sequestration, and subsequent release of OACs likely also occur at and from hotspots. Little is known about the behavior of OACs at environmentally relevant concentrations in agricultural soil.

Recently, many studies have highlighted the ability of fungi to transform and degrade recalcitrant OACs. In particular, one of promising group is the ligninolytic fungi that possess a unique set of extracellular enzymes suitable to degrade lignin and are able to transform recalcitrant compounds, (Čvančarová et al. 2015) (Supplemental material Table I; Table I References). Nguyen et al. (2014) reported the removal of diverse trace organic contaminants (Dichloroethyl chloroformate (TrOC) including phenolic and non-phenolic compounds, pharmaceuticals, pesticides, steroid hormones, industrial precursors and products, and phytoestrogen by live (biosorption + biodegradation), intracellular, enzyme-inhibited and chemically inactivated (biosorption only) whole-cell preparations and the fungal extracellular enzyme extract (predominantly laccases) from *Trametes versicolor* (strain ATCC 7731). They showed how non-phenolic TrOC were readily biodegraded while the removal of hydrophilic TrOC was negligible. The whole-cell culture showed considerably higher degradation of the major compounds, indicating the importance of biosorption and subsequent degradation by intracellular and/or mycelium associated enzymes. However, there are

too few studies that examine both adsorption and degradation of antibiotics in agricultural soil, with most using unrealistically high concentrations (in mg/kg levels) to overcome limitations in measurement. In addition, no model has been developed to speculate about the adsorption and degradation of different types of antibiotics in agricultural soil and the environmental risks they may pose. Pan and Chu (2016) evaluated the adsorption and degradation of five antibiotics (tetracycline, sulfamethazine, norfloxacin, erythromycin, and chloramphenicol) by native microorganisms (bacteria and fungi) in non-sterilized (test) and sterilized (control) agricultural soils under aerobic and anaerobic conditions. They showed that all antibiotics were susceptible to microbial degradation under aerobic conditions, and most antibiotics were degraded by more than 92% in non-sterilized soil after 28 days of incubation. For all the antibiotics, a higher initial concentration was found to slow down degradation and prolong persistence in soil. The degradation pathway of antibiotics varied in relation to their physicochemical properties as well as the microbial activities and aeration of the recipient soil. In their study, Pan and Chu (1996) were the first to develop a model for the prediction of antibiotic persistence in soil.

Given the public concern for environmental pollution by OACs, there is increasing attention towards the development of biopurification systems for reducing the risk from point source contamination of soil resources. Various treatment methods (e.g. land filling, recycling, pyrolysis and incineration) have been used for the removal and remediation of these chemicals from the contaminated sites, but microbial degradation of pesticides is so far the most important and effective way to remove these compounds from the environment (Hai et al. 2012; Verma et al. 2014), (Supplemental material Table I; Table I References).

Microorganisms have the ability to interact both chemically and physically with substances, leading to structural changes or to complete degradation of the target molecule. In particular, fungi may transform pesticides and other xenobiotics by introducing minor structural changes to the molecule, producing nontoxic molecules that can be released into the soil for further degradation by microflora (Hai et al. 2012), (Supplemental material Table I; Table I References). Mir-Tutusaus et al. (2014) investigated the degradation of the insecticides imiprothrin and cypermethrin and the insecticide/nematicide carbofuran using the white-rot fungus *T. versicolor*. Experiments with fungal pellets demonstrated extensive degradation of the tested agrochemicals, while *in vivo* studies with inhibitors of cytochrome P450 revealed that this intracellular system plays an important role in the degradation of imiprothrin and carbofuran, but not of cypermethrin. The simultaneous degradation of the compounds successfully took place with minimal inhibition of fungal activity and resulted in reduction of global toxicity, thus supporting the potential use of *T. versicolor* for the treatment of several OACs.

To date, the number of studies investigating novel treatment techniques for the removal of OACs from contaminated agricultural soils is limited. The bacteria-dominated conventional activated sludge process has been proved to be ineffective for OAC removal. While the importance of a mixed microbial community to initiate and complete OAC removal in the soil environment has been convincingly demonstrated by several researchers, studies concerning the removal of OACs from soils have predominantly focused on selected bacterial or fungal species separately. Few studies have explored the bioaugmentation synergy of fungi together with bacteria (Hai et al. 2012; Zhang et al. 2015; Madrigal-Zúñiga et al. 2016). Combining cultures of bacteria and fungi could be key to the removal of toxic and recalcitrant organic substances from contaminated agricultural soils.

On-farm biopurification systems constitute a biotechnological approach to the mitigation of point source contamination by pesticides. The main component of biopurification systems is the biomixture, which acts as the biologically active core that accelerates the degradation of OACs. Madrigal-Zúñiga et al. (2016) studied the results of employing the ligninolytic fungus *T. versicolor* in the bioaugmentation of compost- (GCS) and peat-based (GTS) biomixtures for the removal of the insecticide-nematicide carbofuran (CFN). The transformation products of CFN were detected at the moment of CFN application, but their concentration decreased continuously until complete removal in both biomixtures. Mineralization of ^{14}C radiolabeled CFN was faster in GTS than in GCS. The authors demonstrated the complete elimination of toxicity in the matrices after 48 days. Overall data suggested that the bioaugmentation improved the performance of the GTS rather than the GCS biomixture.

Pinto et al. (2016) also studied the potential use of different substrates in biomixtures like cork, cork and straw, coat pine and LECA (Light Expanded Clay Aggregates) in the degradation of terbuthylazine, difenoconazole, diflufenican and pendimethalin pesticides. Bioaugmentation using the WRF *Lentinula edodes* inoculated into the CBX was also assessed. The results obtained from this study clearly demonstrated the relevance of using natural biosorbents such as cork residues to increase the capacity for pesticide dissipation in biomixtures for establishing biobeds. Furthermore, greater degradation of all the pesticides was achieved by the use of bioaugmented biomixtures. Indeed, biomixtures inoculated with *L. edodes* EL1 were able to mineralize the selected xenobiotics, revealing that this WRF might be a suitable fungus to be used as inoculum source to improve the degradation efficiency of sustainable on-farm biopurification systems.

Fungi isolated from biomixtures represent a biological source of potentially active bioremediation agents, and the adaptation skills developed by these microorganisms could make the difference in OAC removal (Supplemental material Table I; Table I References). This strategy was assessed by Pinto et al. (2012), who isolated fungi from a loamy sand soil and a biomixture contaminated with

terbuthylazine, difenoconazole and pendimethalin. The ability of autochthonous fungi (*Penicillium brevicompactum* and *Lecanicillium saksenae*) to degrade xenobiotics was compared with that of allochthonous strains taken from a culture collection (*Fusarium oxysporum*, *Aspergillus oryzae* and *L. edodes*). The best biodegradation yield was achieved with *P. brevicompactum*: its higher ability to metabolize terbuthylazine was presumably acquired through chronic exposure to contamination with the herbicide.

Bioremediation of OACs by fungi in aquatic ecosystems

Many OACs are common contaminants of fresh water due to their high water solubility associated with a low soil adsorption, and a high stability that assures them a long half-life. Contamination is heterogeneously distributed along watercourses as evidenced in several studies where pesticides were recurrently found in real water samples. In one accurate survey, more than 160 water samples taken in 23 European countries were assayed for the presence of pharmaceuticals, pesticides and recognised endocrine-disrupting chemicals (Loos et al. 2010). Among the most frequently detected compounds were the insecticide (DEET), and other pesticides (chloridazon-desphenyl, DMS, desethylatrazine, chloridazon-methyl-desphenyl, bentazone, desethylterbutylazine, dichlorprop) exceeded the European threshold of 0.1 µg/l. Overall, 29% of the water samples could not be considered safe (Loos et al. 2010). In a similar study in the USA, groundwater in 18 states was screened for 65 organic contaminants: along with plasticizers and detergent metabolites, 66% of the total pollutant load was ascribable to insect repellent (Barnes et al. 2008).

The extent of freshwater contamination and the actual risk to human life depend on several factors concerning the hydrogeological characteristics of the soil, weather conditions and the chemical-physical properties of the OACs. The environmental fate of a given compound is a critical issue in which the water/soil surface is the first barrier. For instance, the sorption kinetics of three widely used pesticides (simazine, imidacloprid, and boscalid) were found to be correlated with soil organic carbon content and the hydrophobicity of the pesticide, which ultimately affected soil retention behavior and bioavailability in waters (Salvestrini et al. 2014). Leaching into surface waters is also a matter of season, and a complex and unpredictable scenario is influenced by a variety of phenomena. A rainy period can cause massive run-off of OACs from the soil, contaminating the receiving basin (Sandin et al. 2018). The detection of high levels of OACs, however, is not exclusively coincident to their recent and massive use, but is ascribable to their persistency, their slow natural degradation and their accumulation in the various diffusion pathways (Aguilar et al. 2017). They could then travel long distances in surface or groundwaters and the contamination can last for several decades (Ballesteros et al. 2014; Aravinna et al. 2017).

The so-called ecological services may help to contain the diffusion of OACs. The adaptation of microflora (fungi, Gram-positive and negative bacteria, actinobacteria, and sulfate-reducing bacteria) to soil environmental conditions may attenuate the pesticides released into groundwater sources (Mattsson et al. 2015). Several factors such as soil composition, temperature, aeration due to soil weaving, and depth influence autochthonous microbial community activity; if this balance fails, OACs are free to move among different ecological niches (i.e. sediments and water), alter their functioning and ultimately directly affect their animal inhabitants. For instance, significant ecological risk was associated with the presence of the insecticide fipronil and its metabolites in water ponds: the concentrations measured (up to 200 ng/l) affected the proper development of larval insects and crustaceans (Wu et al. 2015b). Evidence of the pesticide's toxicity against fish has already been reported, and it clearly interferes in several metabolic pathways (Odukkathil and Vasudevan 2013; Ballesteros et al. 2014; Guerreño et al. 2016).

The preservation of water quality is a priority, but OAC removal cannot be based only on natural attenuation. Water treatment plants (WTPs) are the major barriers where OACs should be removed. Not being specifically designed for micropollutant removal, however, they are often only partially effective, with a strong impact on the receiving ecosystem. Pesticides such as atrazine, fluconazole, tebuconazole, diazinon and diuron are particularly resistant to commonly used treatments (Köck-Schulmeyer et al. 2013; Luo et al. 2014). There is plenty of evidence confirming the presence of OACs in WTP effluents at toxicologically and estrogenically relevant concentration, making them one of the most impactful sources of contamination (Bicchi et al. 2009; Campo et al. 2013; Jarošová et al. 2014).

Particular attention has been given to advanced biological oxidation. Novel cost-effective and eco-friendly processes based on fungi are an attractive option. Fungi are well-known for their physiological adaption skills, including the natural activation of tolerance mechanisms against pesticides (Talk et al. 2016). Some reports have already demonstrated that in comparison with bacteria, fungi can better tolerate the presence of organic contaminants. Although the insecticide endosulfan inhibited both fungi and bacteria, bacterial community structure significantly changed at concentrations as low as 0.1 mg/kg, while modifications to fungal community structures required 1 mg/kg of pollutant (Zhang et al. 2015). Linuron reduced the bacterial count, and especially total bacteria, N₂-fixing bacteria and nitrifiers, but not fungal numbers (Cycoń et al. 2010).

The provenance of isolated fungi is of unquestionable importance. Strains isolated from contaminated niches indeed seem to develop specific adaptation skills due to chronic exposure. Carles et al. (2017) demonstrated that the aquatic microflora found in association with submerged leaves exposed to nicosulfuron is more efficient in its degradation than are communities that come

from a less polluted site. The authors indicated fungi as the main constituents of this active microflora and as being responsible for herbicide degradation. In the literature, several fungi isolated from contaminated areas or WTPs have been identified as degraders of nicosulfuron, diuron, isoproturon, glyphosate, chlorpyrifos, chlorfenvinphos and atrazine (Song et al. 2013; Carranza et al. 2014; Oliveira et al. 2015).

Fungi can thus transform a broad range of recalcitrant organic compounds, including OACs (Gao et al., 2010). A number of fungi that are OAC degraders, mostly belonging to Basidiomycetes, such as *Trametes*, *Pleurotus*, *Phlebia*, *Cerrena*, *Coriolopsis*, etc., have been already investigated (Koroleva et al. 2002; Marco-Urrea et al. 2009; Xiao et al. 2011; Ulčnik et al. 2013; Chan-Cupul et al. 2014; Ceci et al. 2015). Several pesticides as lindane, atrazine, diuron, terbutylazine, metalaxyl, DDT, gamma-hexachlorocyclohexane (g-HCH), dieldrin, aldrin, heptachlor, chlordane, lindane, mirex, etc. were effectively transformed by fungal treatment based on mycelium or enzymes (Supplemental material Table II).

A bioremediation approach based on fungi may involve both biosorption and biodegradation processes; the latter combines biosorption, where the molecule binds to the fungal wall, and bioaccumulation with the pollutant being transported inside the cell in contact with intracellular enzymes (Kulshreshtha et al. 2014). Concentrations of the insecticide lindane decreased during time in the presence of two WRFs (*T. versicolor* and *Pleurotus ostreatus*) and one brown-rot fungus (*Gloeophyllum trabeum*), but the lack of any change in the chromatogram profile indicated that a fast adsorption process was mainly involved (Ulčnik et al. 2013). However, this phenomenon is often strain-dependent, and especially related to metabolic differences between Ascomycetes and Basidiomycetes. Belonging to the brown-rot fungi, *G. trabeum* lacks the ligninolytic enzymes, responsible for lignin degradation and likely for that of OACs as well: adsorption onto fungal mycelium was mainly involved in the removal of endosulfan. On the contrary, the white-rot fungi actively degraded, producing endosulfan sulphate via oxidative pathways (Ulčnik et al. 2013). Although biosorption is a phenomenon that cannot be ignored, it is often secondary or at least negligible compared to biodegradation (Carles et al. 2017). For instance, the removal of clofibric acid found for heat-killed mycelium was less than 10 %, but more than 97 % for active *T. versicolor* (Marco-Urrea et al. 2009).

Fungi have developed a specific mechanism that employs few enzymes and molecules with high oxidizing power, physiologically aimed at transforming lignocellulose structures. The same enzymatic pathway may play a pivotal role in transforming other aromatic molecules. White-rot fungi usually deploy extracellular lignocellulosic enzymes such as peroxidases (EC 1.11.1.x) and laccases (EC 1.10.3.2). The involvement of redox enzymes in fungal-mediated oxidation is

confirmed by the direct induction of enzyme production in the presence of OACs. The fungus *T. versicolor* responded to 17 pesticides by increasing laccase production in comparison with the control: particular attention was given to the transformation products of the herbicides diquat and monuron, capable of increasing fungal activity 10- and 17-fold, respectively (Mougin et al. 2002). The laccase production of *Pycnoporus sanguineus*, *Trametes maxima*, *Pleurotus* spp1, *Pleurotus* spp2, *Cymatoderma elegans*, and *Daedalea elegans* was stimulated by the presence of atrazine even at high concentrations of 3750 mg/l. Likewise, the manganese peroxidase activity of *Pleurotus* spp1 and *C. elegans* was positively correlated with the pesticide (Chan-Cupul et al. 2014). Oxidoreductase stimulation was also observed with picloram (Maciel et al. 2013), bentazon (Da Silva Coelho et al. 2010) and carbofuran (Mir-Tutusaus et al. 2014).

Although these oxidoreductases are probably the most-known enzymes for aromatic compound degradation, alternative pathways can be stimulated by the presence of OACs. Two clones (laccase-positive and laccase-negative) of *Mycelia sterilia* were used to treat atrazine (20 µg/ml): even though one clone was defective in laccase production, comparable transformation yields (70-80%) were reached, indicating that the fungus can deploy alternatives to laccase in the degradation process (Vasil'Chenko et al. 2002). This behavior is commonly found in brown-rot fungi, which can trigger both nonenzymatic and enzymatic mechanisms, i.e. the Fenton mechanism or cellobiose dehydrogenase (CDH) reactions (Fan and Song 2014). The degradation of atrazine (20 µg/l) by an unidentified mycelial fungus was associated with the presence in the liquid medium of OH radicals and CDH. Moreover, CDH secretion was induced by the presence of the herbicide itself (Khromonygina et al. 2004). In addition, some fungi may associate extracellular oxidoreductases with intracellular enzymes such as the cytochrome P450 system (cyt450). In an effort to better characterize the degradation skills of *T. versicolor*, cyt450 inhibitors were used: fungal performance against clofibric acid and fipronil decreased (Marco-Urrea et al. 2009; Wolfand et al. 2016). Mori et al. (2017), suggest that in *Phanerochaete sordida*, cyt450 is involved in the initial stage of reduction of the clothianidin N-nitro group, but that the enzymes responsible of the further urea derivatives production are unknown.

Fungal intra- and interspecies variability has long been recognized and has found confirmation in OAC treatment. Literature data about a given species cannot be taken for granted and preliminary screening is often required. Despite *Phanerochaete chrysosporium* often being indicated as the fungal model for organic degradation including pesticides (Wang et al. 2014), it was almost ineffective against clofibric acid (Marco-Urrea et al. 2009). Among five Basidiomycetes, only *T. versicolor* extensively degraded this herbicide (Marco-Urrea et al. 2009). Alvarenga et al. (2014) treated methyl parathion with several fungi, including 3 *Aspergillus sydowii*. Based on ability to

grow in the presence of the pesticide, only the isolate *A. sydowii* CBMAI 935 was selected for further studies. It indeed grew almost 4-fold more than the other *A. sydowii*. Bioremediation potential is often substrate-targeted, and the choice of fungus cannot be taken for granted. For instance, *A. sydowii* CBMAI 935, which totally converted methyl parathion (Alvarenga et al. 2014) was not the best performing one against the insecticide esfenvalerate. Among 6 fungi, *Microsphaeropsis* sp. *Acremonium* sp. and *Westerdykella* sp. gave better results than the *Aspergillus* strain (Birolli et al. 2016).

Although the majority of these strains are effective in OAC removal in model solutions, only few researchers have taken the next step, and assessed bioremediation potential in contaminated waters. The experimentation with model solutions (single-compound solutions, high concentrations, no interfering molecules, etc.) is the only way to acquire information about degradation pathways (Masaphy et al. 1993; Birolli et al. 2016), but it is less predictive of fungal performance in real environmental water samples. Each type of wastewater has its own critical issues, making it difficult to predict fungal behavior. Some data highlight the robustness of fungal systems, although detailed case-by-case investigation is needed. A partially diluted leachate was shown to disturb the growth of *T. versicolor* and *Stereum hirsutum*, but this did not prevent them totally degrading linuron and dimethoate at 10 mg/l. As regards dimethoate, the presence of adsorbents enhances final yields from 50% to 97%, because the adsorption action combines with and exalts fungal biodegradation processes (Castellana and Loffredo 2014). The immobilization of *Bjerkandera adusta* and *Irpex lacteus* on coffee grounds, almond shells and a biochar favored the removal of the non-phenolic herbicides fenuron and carbaryl from a municipal landfill leachate (Loffredo et al. 2016). Surface waters, ground waters and municipal wastewaters represent a very unique environment, characterized by extreme chemical and physical conditions, the presence of a heterogeneous and variable mixture of micropollutants and an active autochthonous microflora. When inoculated into real surface water, a fungal consortium (*Aspergillus fumigatus*, *Aspergillus terreus*, *Cladosporium tenuissimum*, *Cladosporium cladosporioides*, *Fusarium begoniae*, *Penicillium citrinum*, *Penicillium melanoconidium* and *Phoma glomerata*) was not stable over time, probably due to the presence of toxic pesticides and interaction with the natural microbial population: *P. citrinum*, *A. fumigatus* and *A. terreus* were the most robust to the environmental conditions and were found to degrade the spiked chlorfenvinphos (Oliveira et al. 2015).

The set-up of active microbial consortia offers the intriguing possibility of strengthening and combining the bioremediation potential of different organisms: the combination of *Bacillus subtilis* and *A. niger* led to higher degradation rates of nicosulfuron than those obtained by using each strain singly (Lu et al. 2012). The biodegradation of aldicarb, atrazine and alachlor by *Coriolus versicolor*

was strongly enhanced by combination with activated sludge. Along with modifications in fungal morphology, when the bacterial-fungal consortium was established, the bio-absorbed fraction of especially atrazine was reduced: over 98% of atrazine was removed by degradation processes in two weeks (Hai et al. 2012).

The fate of the treated OACs must be carefully considered. Residual toxicity is a critical issue. Interestingly fenuron and carbaryl degradation (up to 70%) catalyzed by *B. adusta* and *I. lacteus* led to significant abatement of the phytotoxicity (rapeseed and flax tests) (Loffredo et al. 2016). Mori et al. (2017) monitored the neurotoxicity of clothianidin and the main metabolite it produced during *P. sordida* treatment: following treatment the insecticide still altered the viability of the neuronal cell line, but the metabolite was no longer neurotoxic.

Despite their well-demonstrated properties, the application of whole cell systems has some drawbacks including the fact that a living organism needs controlled growing conditions in terms of nutrients, pH, O₂, etc. (Majeau et al. 2010). The addition of synthetic nutrients can strengthen fungal mycelium activity, but it should be carefully balanced to allow subsequent scale-up of the process. The fact that *T. versicolor* needed 1% of glucose as carbon source to degrade atrazine would ultimately interfere with its potential use in real WTPs (Khromonygina et al. 2004). Likewise several fungi such as *A. niger* and *Dacryopinax elegans*, etc. required both easily available carbon and nitrogen sources to efficiently act against nicosulfuron and diuron, respectively (Lu et al. 2012; Arakaki et al. 2013). Particular attention should be instead given to those fungi, like *A. sydowii* and *Penicillium decaturense*, that maintained the same performance without glucose addition, indicating potential for using methyl parathion or triclosan as sole carbon source (Alvarenga et al. 2014; Tian et al. 2016).

A promising alternative is offered by the direct use of fungal enzymes, capable of catalyzing strong, rapid oxidation reactions, with less technical drawbacks in comparison with fungal cultures. The potential of enzymes-based methods has been worldwide recognized; the Swiss Industrial Biocatalysis Consortium defined oxidative enzymes as the biocatalysts displaying the highest development potential for the next decades (Meyer and Munch 2005). Great importance is given to the discovery of novel enzymes with wide substrate specificity, stable and applicable to industrial uses. A number of articles have reported the ability of fungal enzymes to degrade OACs. The potential of laccase-mediator systems has been assessed for the degradation of isoproturon (Margot et al. 2015), imiprothrin (Mir-Tutusaus et al. 2014), chloroxuron (Palvannan et al. 2014), isoproturon (Zeng et al. 2017), atrazine (Chan-Cupul et al. 2016). Laccases cannot be considered a novelty, unlike a phytase of *A. niger* capable of degrading organophosphorus pesticides (Shah et al. 2017) or a cellulase of *Trichoderma longibrachiatum* active against dicofol (Wang et al. 2015).

Particular attention should be given to the use of crude enzyme extracts of ligninolytic enzymes with a lower economic impact on the process than that of purified enzymes (Matute et al. 2012; Kaur et al. 2016). A crude extract of *Trametes pubescens* laccases degraded up to 19 compounds in a model solution and confirmed its potential in a study on real municipal wastewater where the presence of suspended particles, colloids, solvents and xenobiotics as well as autochthonous microorganisms posed strong environmental pressure. The transformation of all the detected compounds determined also a strong reduction of the estrogenicity of the water sample (Spina et al. 2015).

Application of synthetic microbial communities in bioremediation

Bioremediation is a crucial way to eliminate OAC pollution in agricultural ecosystems. However, many factors affect the efficiency of bioremediation in pesticide pollution, such as the microbes applied, treatment sites, rhizosphere effects and soil chemical and physical properties (Zhou and Hua 2004). Bioremediation of soil or water pollution often cannot reach expected results in practice because the target contaminant cannot be degraded completely, and sometimes intermediate products occur that are more toxic than the original pesticides. Long-term application of various pesticides results in pollution with more than one type of chemical compound, which are unlikely to be degraded by a so-called microbe. Thus, attention has shifted to synthetic systems based on communication between cells, rather than on individual isolated cell functionality (Biliouris et al. 2012). A promising way to overcome the difficulties is to create artificial synthetic microbial communities that contain several microbes to retain the key features of their natural counterparts (Großkopf and Soyer 2014).

The so-called *synthetic microbial community* is created by a bottom-up approach where two or more defined microbial populations are put together in a well-characterized and controlled environment (De Roy et al. 2014). In synthetic communities, mixed populations can perform complex tasks, although in changing environmental conditions (Brenner et al. 2008). Synthetic communities have several potential advantages over monocultures or natural communities: 1) the species in a synthetic community are known and the community structure is relatively simple and controllable, while the natural community may contain many microorganisms with unknown functions; 2) synthetic communities can perform more complicated functions than individual organisms because members of microbial consortia communicate and differentiate (Brenner et al. 2008); 3) synthetic communities are often more robust to environmental fluctuations because they can resist invasion by other species and weather periods of nutrient limitation better than monocultures (Brenner et al. 2008); 4) synthetic communities can be described through mathematical models more easily than

natural systems, and they can be used to develop and validate models of more complex systems (Liu et al. 2017).

Liu et al. (2017) proposed three design principles to develop a cooperative, steady-state community that is performing a desirable biotechnological function. Firstly, safety should be prioritized by beginning with innocuous or commensal organisms (Brenner et al. 2008). Secondly, the community can converse a low-cost and/or recalcitrant waste material into a biotechnologically relevant product, partial or de-novo biosynthesize a compound via heterologous metabolic pathways, or bioconverse toxic substrates or products in a toxic milieu (Jagmann and Philipp 2014). Thirdly, the bioremediation process should be optimized and regularly monitored on the basis of the knowledge of stability and division of different microorganisms (Liu et al. 2017).

Bioremediation of polluted soils and water is one field of application synthetic microbial communities. Due to the complex structure of some pollutants, such as the diuron pesticides, adding synthetic microbial communities is much more effective than adding single microorganisms. The herbicide diuron is used in the control of broad-leaved weeds on agricultural land. Several fungal-bacterial consortia were investigated by combining three different diuron-degrading bacteria and two fungal strains. The fastest mineralization of diuron was obtained by the three-member consortium (*Mortierella* LEJ702, *Variovorax* SRS16, and *Arthrobacter globiformis* D47). As measured by evolved $^{14}\text{CO}_2$ it mineralized about 32 % of the added diuron within 54 days, whereas the single strains or other consortia achieved no more than 10% mineralization. In addition, the production of diuron metabolites by the consortium was minimal. This may be due to cooperative catabolism, where the first organism transforms the pollutant to products that are then used by the other organisms. In addition, fungal hyphae may function as transport vectors for bacteria, thereby facilitating the more effective spreading of degrader organisms in the soil (Ellegaard-Jensen et al. 2014).

Similarly, a fungal-bacterial consortium consisting of *Mortierella* sp. LEJ702 and the 2,6-dichlorobenzamide (BAM)-degrading *Aminobacter* sp. MSH1 achieved more rapid mineralisation of BAM than did the bacteria alone, especially at lower moisture contents (Knudsen et al. 2013). Methylophilic and hydrocarbon-utilizing yeasts and bacteria alone did not degrade PCBs significantly, but PCB degradation reached about 50% when WRFs were applied together (Šašek et al. 1993).

Evaluation of bioremediation effectiveness in contaminated matrices by means of ecotoxicological and genotoxic tests

In order to improve the effectiveness and performance of bioremediation processes it is important to pursue three essential goals at the same time. Focus should be not only on reducing chemical concentrations, but also on reducing chemical mobility between the environmental compartments and eventually lowering toxicity levels while ensuring that contaminants do not get into the natural biological cycle (Loehr and Webster 1997; Chakraborty et al. 2013).

Bioremediation is often monitored by following the concentration of targeted contaminants (Molina-Barahona et al. 2005). Numerous studies in recent years have shown that traditional chemical analyses are insufficient for a full assessment of the contaminated site because, for example, they do not provide any information about the interactions between chemicals and they do not consider the partition and the mobility of pollutants (Frische 2003; Molina-Barahona et al. 2005; Ma et al. 2005; Molnár et al. 2007). An integrated approach that links the various fields and levels of study involving contaminated sites has proven to be an efficient way to evaluate the effectiveness of bioremediation in contaminated sites (Chapman and Anderson 2005; Wernersson et al. 2015; Marziali et al. 2017). Consequently, to achieve the desired goals and implement a successful bioremediation program, given the chemical and biological complexity of the tasks involved, close collaboration between microbiologists, chemists and engineers is required (Van Gestel et al. 2001; Chakraborty et al. 2013).

Additionally, the use of ecotoxicological and genotoxic tests to evaluate the effectiveness of bioremediation may be a valid tool to partially overcome the existing gap between the reported successes of bioremediation on the laboratory scale, and that in the field.

Signals that bioremediation is going on should be monitored. Two important chemical compounds produced by microorganisms during their degradation activity are CO₂ and soluble phosphorus. Both increase notably in soil treated with insecticides and inoculated with fungi (Boyle 1995; Abd El-Ghany and Masmali 2016). However, it must be taken into consideration that during and after a bioremediation process the disappearance of the parent compounds or evidence of metabolic activity (e.g. CO₂ production) may not indicate detoxification. Although the fate of the toxicants may be followed by chemical analyses, many reaction products resulting from the bioremediation process and their potential toxicity are not known. The elimination of mother compounds does not necessarily result in toxicity removal, and evaluating the efficiency of the process is important to assess not only the removal of a specific compound, but also potential ecotoxicity. In fact, biodegradation of pesticides can proceed partially or totally due to the structure of the molecule itself or to unfavourable environmental or test conditions, or to the lack of 'acclimatized' microbial communities (De Henau 1997).

In some instances, it has been shown that an effective process of bioremediation corresponds with a decrease in the toxicity of the analysed matrix (Baud-Grasset et al. 1993; Dorn and Salanitro 2000). To acquire complete and useful information in an ecotoxicological assessment and to determine the effectiveness of bioremediation treatments, it is suggested that a battery of tests be used (Keddy et al. 1995; Van Gestel et al. 2001; Tigini et al. 2011). The battery should include a number of reference organisms that are representative of the different trophic levels, in order to select species with different roles in ecosystems, and different exposure conditions (Van Straalen and Van Gestel 1997). Moreover, environmental risk assessment must integrate chemical characterization, ecotoxicity and bioremediation data, in order to accurately assess the ecological hazard.

As emphasized by Shen et al. (2016), an increased level of ecotoxicity within the various bioindicators could either indicate incomplete decomposition of the substance or could result from the formation of intermediate products generated via the bioremediation process. For this reason, chronic tests are sometimes more appropriate in evaluating the toxicity caused by by-products (Lofrano et al. 2014).

In certain circumstances, there is a clear need to monitor the bioremediation process using different bioindicators. In Lizano-Fallas et al. (2017), for example, the ecotoxicity test with *Daphnia magna* showed clear detoxification, whilst the detoxification patterns remain unclear when applying the phytotoxicity test. Ecotoxicological tests can also be used to determine the most suitable bioremediation technique in a given case, as reported in Dudášová et al. (2016).

Without worldwide-recognized guidelines for water quality assessment, literature data are difficult to compare due to the variety of model organisms, end-points, etc. Synthetic indices summarizing the findings can help monitor the effectiveness of biological treatment. Such indices have already been applied for toxicity monitoring of wastewaters (Tigini et al. 2011) but municipal effluents containing AOCs have never been taken into consideration nor has estrogenic activity been included so far.

Several toxicity assays were included in the biodegradability study protocol to measure remediation efficiency. Assessing the toxicity of complex matrixes such as soil could acquire methods from bioassays used to test toxicity of chemical compounds, reported by the Organization for Economic Co-operation and Development (e.g. OECD 201 2006; OECD 211 2012). The OECD has published a series of standardized tests for determining the biodegradability of a given compound, based on the evaluation of overall parameters (such as COD, TOC and BOD) or metabolic tests, e.g. respirometry (OECD 209 1984) as Polo et al. (2011) used; or that reveal susceptibility of toxic compounds, comprising that of herbicides, to biological treatment. Standardized testing procedures using different organisms have been approved by various environmental organizations, including

the US Environmental Protection Agency, American Society for Testing and Materials, International Standardization Organization (Siciliano et al. 2015). Many scientists have explored the effects of polluted soil on the whole organism using various microorganisms, animals, and plants, or by means of cellular, and biochemical biomarkers, or by ecological scale up systems. Here below, tests at some different biological hierarchical levels of analysis are presented and discussed.

Organismal level

Concerning complex matrices such as soil, quality assessments are performed with organisms on extracts of the polluted matrix, generally applying short-term exposure periods (Van Gestel et al. 2001). Experimental models have included aquatic organisms such as *Daphnia magna*, *Raphidocelis subcapitata*, *Danio rerio*, *Myriophyllum aquaticum* and *Lemna minor* (Feiler et al. 2004). The use of freshwater and marine biota may be particularly useful in order to provide a more complete comprehension of the fate of pesticides and the environmental outcomes of agricultural activities (Guida et al. 2008). Terrestrial animals such as nematodes (*Caenorhabditis elegans*) (Traunspurger et al. 1997), oligochaetes (*Lumbriculus variegatus*) (Phipps et al. 1993), springtails such as *Folsomia candida* (Houx et al. 1996), and fish embryos (Hollert et al. 2003; Zielke et al. 2011) are considered among the most reliable models.

Among the higher plants, important experimental models include *Lepidium sativum*, *Cucumis sativus*, and *Sorghum saccharatum* (germination rate, inhibition of root elongation). Since assays based on animals, plants and algae are considered expensive, time consuming and require large sample volumes, recent studies have emphasized the benefits of rapid, reproducible and cost effective bacterial assays for toxicity screening and assessment. *Arthrobacter globiformis* (Neumann-Hensel and Melbye 2006), *Bacillus cereus* (Rönnpagel et al. 1995; Prokop et al. 2016), *Vibrio proteolyticus* (Ahlf and Heise 2005) and yeasts (*Saccharomyces cerevisiae*) (Weber et al. 2006) are often used. Among the bacterial bioassays, the *Vibrio fischeri* luminescence inhibition test is the most common. The review of Parvez et al. (2006) remarks that the *Vibrio fischeri* inhibition test is the most sensitive, cost effective, easy to operate and requires only 5–30 min for toxicity prediction.

Cellular and biomolecular level

Biomarkers signal the adaptative responses of organisms to xenobiotic exposure. Various studies have highlighted the cytotoxic and genotoxic effects on organisms of OACs and their metabolic products. The exposed organisms may exhibit histological, cellular, molecular, biochemical and/or physiological, or even behavioural changes (Depledge et al. 1993) that enable information to be

obtained on the biological effects of pollutants or their remains during or after a bioremediation process (Fontanetti et al. 2011).

Genetic endpoints and biomarkers. The most-used biomarkers are mitotic index, chromosome aberrations, micronuclei, sister chromatid exchanges and mutations.

Bacteria have been recommended for bioassays to evaluate genotoxicity in a variety of samples (Mortelmans and Zeiger 2000; White and Claxton 2004). The Ames test, one of the most famous and widely-used, is a short term bacterial reverse mutation assay especially designed to evaluate the mutagenic potential of a wide range of chemical substances (Mortelmans and Zeiger 2000). It was found to be very sensitive in tests with a wide range of mutagenic and carcinogenic chemicals, as reported in the review paper of Chahal et al. (2014).

With regards to plant models, higher plants are recognized as excellent genetic models to detect cytogenetic and mutagenic agents and are frequently used in environmental monitoring studies. The main organisms employed are *Allium cepa*, *Vicia faba* and *Tradescantia* spp. as reported in a review by De Souza et al. (2016). Their protocols were standardized under the International Program on Plant Bioassays (IPPB) conducted by the United Nations Environment Programme (UNEP) (Ma 1999). In addition, the US Environmental Protection Agency (USEPA) and the World Health Organization (WHO) validated plant bioindicators as an efficient model to detect environmental genotoxicity.

One of the most used higher plant models is *V. faba*. The main advantages are its year-round availability, that it is economical to use, and easy to grow and handle. Its use does not require sterile conditions and rate of cell division is fast. The *V. faba* test, meticulously reported and discussed in the review of Iqbal (2016), enables the assessment of a variety of endpoints, e.g., chromosomal aberration, mitotic index, micronuclei and nuclear aberration.

Enzymatic biomarkers. Enzyme activity inhibition has been widely evaluated as a biomarker to measure the toxicity of a matrix. Dehydrogenases, for example, are directly involved in many of the vital anabolic and catabolic processes of living organisms, and their activity is inhibited by chemical toxicants. Recently, many studies have reported the use of terrestrial organisms to obtain enzymatic biomarkers in response to residual pesticides (Henson-Ramsey et al. 2011; Radwan and Mohamed 2013; Stepić et al. 2013), and among these, earthworms' enzymes were widely used to understand the impacts of pesticides. In two earthworm species, *Eisenia fetida* and *Lumbricus terrestris*, multiple esterases, including acetylcholinesterase (AChE), butyrylcholinesterase, and carboxylesterase (CE), were assessed as biomarkers for malathion exposure (Henson-Ramsey et al. 2011). Several studies have also reported AChE, catalase (CAT), and glutathione-S-transferase as biochemical biomarkers in *Eisenia andrei* for the insecticides endosulfan, temephos, malathion, and

672 pirimiphos-methyl (Stepić et al. 2013), and AChE, CAT, CE, and the efflux pump as biomarkers in
673 *E. andrei* and *Octolasion lacteum* for dimethoa. Recently, surface-enhanced laser
674 desorption/ionization-time-of-flight (SELDI-TOF) mass spectrometry (MS) has strongly
675 contributed to the identification of more accurate, precise biomarkers, e.g. specific for human
676 cancers (Silsirivanit et al. 2014), or for endosulfan exposure in Japanese rice fish (*Oryzias latipes*)
677 (Lee et al. 2013). In a recent paper, selective protein biomarkers for 6 pesticides (captan, carbaryl,
678 carbofuran, and α -endosulfan chlorpyrifos, propoxur) were found in *E. fetida*, by means of SELDI-
679 TOF MS technology (Park et al. 2015).

680 Estrogen and androgen biomarkers. It is well-documented that several chemicals from agricultural,
681 industrial, and household sources possess endocrine-disrupting properties, which provide a potential
682 threat to human and wildlife reproduction (Colborn et al. 1993; Colborn 1995; Jensen et al. 1995).
683 A suggested mechanism is that environmental contaminants alter the normal functioning of the
684 endocrine and reproductive system by mimicking or inhibiting the action of endogenous hormones,
685 by modulating the production of endogenous hormones, or by altering hormone receptor
686 populations (Sonnenschein and Soto 1998). Several pesticides exert estrogenic and antiandrogenic
687 activities through interaction with estrogen and androgen receptors. The risks associated with OAC
688 exposure has been known for decades: many pesticides, such as p,p'-dichlorodiphenyl
689 trichloroethane (DDT) (Welch et al. 1969), methoxychlor (Bulger et al. 1978; Cummings 1997), β -
690 benzene hexachloride (BHC) (Coosen and van Velsen 1989), endosulfan, toxaphene, and dieldrin
691 (Soto et al. 1995), and fenvalerate (Garey and Wolff 1998) were the first to be signaled as
692 estrogenic. Despite increased institutional awareness and more compelling legislation pressure, the
693 most recent literature still reports the occurrence of pesticides in watercourses and in the trophic
694 chains, that show conspicuous estrogen or androgen levels (Saillenfait et al. 2016; Brander et al.
695 2016; Guo et al. 2017; Khalil et al. 2017; Scott et al. 2017; Miccoli et al. 2017; Marcoccia et al.
696 2017). Several bioassays have been developed and standardized in order to describe the estrogenic
697 potency of OACs. Andersen et al. (2002) indicated that several currently used OACs, such as
698 methiocarb, fenarimol, chlorpyrifos, deltamethrin, and tolclofos-methyl, possess estrogenic activity
699 on the basis of cell proliferation assays and transactivation assays using MCF-7 human breast
700 cancer cells. Kojima et al. (2004) tested 200 pesticides in vitro for agonism and antagonism to two
701 human estrogen receptor (hER) subtypes, hER α and hER β , and a human androgen receptor (hAR)
702 by means of highly sensitive transactivation assays, using Chinese hamster ovary cells. The results
703 demonstrated that many pesticides possess in vitro estrogenic and antiandrogenic action through
704 ERs and/or AR. Although it appears that various pesticides exert hormonal effects at concentrations
705 that are orders of magnitude higher than that required for physiologic hormones, wide exposure to

large numbers of OACs may have additive and synergistic effects. Bioassay with YES (yeast estrogen screen) and YAS (yeast androgen screen) can determine hormonally active compounds still present in the environment. Since the the first papers on this subject (Purvis et al. 1991), much more sophisticated bioassays have been developed, such as that proposed by Eldridge et al. (2007) in which a bioluminescent strain of *Saccharomyces cerevisiae* was genetically engineered to respond to androgenic chemicals.

Ecological level

The risk to natural systems of pollution with the chemical residues of bioremediation processes is underestimated. The ecological scaling-up experiment illustrated by Rodea-Palomares et al. (2016) underlined how real-world exposure to chemical pollution is often dominated by low-dose complex mixtures combined with other biotic and abiotic stressors. In the paper, a novel screening method (GSA-QHTS) was reported, that coupled the computational power of global sensitivity analysis (GSA) with the experimental efficiency of quantitative high-throughput screening (QHTS). In the study, they reported that GSA-QHTS allowed for the identification of the main pharmaceutical pollutants that were driving the biological effects of low-dose complex mixtures at the microbial population level. The target complex community was a river benthic microbial community inoculum obtained from an unpolluted stream. The effects of the toxic compounds in the mixture was evaluated together with other physico-chemical stressors, on a series of community-level metabolic end points. Photosynthetic parameters, the dark-adapted basal fluorescence, the light-adapted steady-state fluorescence, the maximum photosynthetic efficiency, as well as the extracellular enzymatic activities b-Glu and Phos were considered as both autotrophic and heterotrophic global fitness indicators suited to study the effects of chemical pollution on freshwater benthic microbial communities.

Prospect

Bioremediation is based on the idea that different organisms will work together to remove (biodegrade) the waste substances or pollutants (OACs) from the environment. Although there exist limitations to bioremediation practice, including the nature of organisms, the enzyme involved, the concentration and availability and final survival of microorganisms, as well as the cost/benefit ratio (i.e. cost versus overall environmental impact), these limitations can be solved to some extent by understanding the genetics and biochemistry of the desired microbe. The advent of synthetic communities has shown enormous potential to facilitate the bioremediation process, the degradative fungi appearing to be particularly effective.

Acknowledgement

B. Wu is funded by National Natural Science Foundations of China (No. 31701853). L. Pecoraro acknowledges CAS 153211KYSB20160029 for supporting his research at Chinese Academy of Sciences.

References

Abd El-Ghany T, Masmali IA. 2016. Fungal biodegradation of organophosphorus insecticides and their impact on soil microbial population. J Plant Pathol Microbiol. Available from: <https://www.omicsonline.org/open-access/fungal-biodegradation-of-organophosphorus-insecticides-and-their-impact-on-soil-microbial-population-2157-7471-1000349.php?aid=72754>

Aguilar JAP, Andreu V, Campo J, Picó Y, Masiá A. 2017. Pesticide occurrence in the waters of Júcar River, Spain from different farming landscapes. Sci Total Environ 607–608: 752–760.

Ahlf W, Heise S. 2005. Sediment toxicity assessment: rationale for effect classes. J Soils Sediments. 5:16–20.

Alavanja MCR. 2009. Introduction: pesticides use and exposure, extensive worldwide. Rev Environ Health. Available from: <https://www.degruyter.com/view/j/reveh.2009.24.4/reveh.2009.24.4.303/reveh.2009.24.4.303.xml>

Alvarenga N, Birolli WG, Selegim MHR, Porto ALM. 2014. Biodegradation of methyl parathion by whole cells of marine-derived fungi *Aspergillus sydowii* and *Penicillium decaturense*. Chemosphere. 117:47–52.

Anastasi A, Tigini V, Varese GC. 2013. The bioremediation potential of different ecophysiological groups of fungi. In: Goltapeh EM, Danesh YR, Varma A, editors. Fungi as Bioremediators. Springer Berlin Heidelberg; p. 29–49. Available from: <http://dx.doi.org/10.1007/978-3-642-33811>

Andersen HR, Vinggaard AM, Høj Rasmussen T, Gjermansen IM, Cecilie Bonefeld-Jørgensen E. 2002. Effects of currently used pesticides in assays for estrogenicity, androgenicity, and aromatase activity in vitro. Toxicol Appl Pharmacol. 179:1–12.

- 768 Arakaki RL, Monteiro DA, Boscolo M, Dasilva R, Gomes E. 2013. Halotolerance, ligninase
769 production and herbicide degradation ability of basidiomycetes strains. *Braz J Microbiol.* 44:1207–
770 1214.
- 771 Aravinna P, Priyantha N, Pitawala A, Yatigammana SK. 2017. Use pattern of pesticides and their
772 predicted mobility into shallow groundwater and surface water bodies of paddy lands in Mahaweli
773 river basin in Sri Lanka. *J Environ Sci Health Part B.* 52:37–47. Available from:
774 <http://dx.doi.org/10.1080/03601234.2016.1229445>
- 775 Ballesteros ML, Miglioranza KSB, Gonzalez M, Fillmann G, Wunderlin DA, Bistoni MA. 2014.
776 Multimatrix measurement of persistent organic pollutants in Mar Chiquita, a continental saline
777 shallow lake. *Sci Total Environ.* 490:73–80.
- 778 Barnes KK, Kolpin DW, Furlong ET, Zaugg SD, Meyer MT, Barber LB. 2008. A national
779 reconnaissance of pharmaceuticals and other organic wastewater contaminants in the United States
780 — I) Groundwater. *Sci Total Environ.* 402:192–200.
- 781 Baud-Grasset F, Baud-Grasset S, Safferman SI. 1993. Evaluation of the bioremediation of a
782 contaminated soil with phytotoxicity tests. *Chemosphere.* 26:1365–1374.
- 783 Bicchi C, Schilirò T, Pignata C, Fea E, Cordero C, Canale F, Gilli G. 2009. Analysis of
784 environmental endocrine disrupting chemicals using the E-screen method and stir bar sorptive
785 extraction in wastewater treatment plant effluents. *Sci Total Environ.* 407:1842–1851.
- 786 Biliouris K, Babson D, Schmidt-Dannert C, Kaznessis YN. 2012. Stochastic simulations of a
787 synthetic bacteria-yeast ecosystem. *BMC Syst Biol.* 6:58.
- 788 Birolli WG, Alvarenga N, Seleglim MHR, Porto ALM. 2016. Biodegradation of the pyrethroid
789 pesticide esfenvalerate by marine-derived fungi. *Mar Biotechnol.* 18:511–520.
- 790 Boyle CD. 1995. Development of a practical method for inducing white-rot fungi to grow into and
791 degrade organopollutants in soil. *Can J Microbiol.* 41:345–353.
- 792 Brander SM, Gabler MK, Fowler NL, Connon RE, Schlenk D. 2016. Pyrethroid pesticides as
793 endocrine disruptors: molecular mechanisms in vertebrates with a focus on fishes. *Environ Sci*
794 *Technol.* 50:8977–8992.
- 795 Brenner K, You L, Arnold FH. 2008. Engineering microbial consortia: a new frontier in synthetic
796 biology. *Trends Biotechnol.* 26:483–489.

- 797 Bulger WH, Muccitelli RM, Kupfer D. 1978. Studies on the in vivo and in vitro estrogenic
798 activities of methoxychlor and its metabolites. Role of hepatic mono-oxygenase in methoxychlor
799 activation. *Biochem Pharmacol.* 27:2417–2423.
- 800 Campo J, Masiá A, Blasco C, Picó Y. 2013. Occurrence and removal efficiency of pesticides in
801 sewage treatment plants of four Mediterranean River Basins. *J Hazard Mater.* 263:146–157.
- 802 Carles L, Rossi F, Joly M, Besse-Hoggan P, Batisson I, Artigas J. 2017. Biotransformation of
803 herbicides by aquatic microbial communities associated to submerged leaves. *Environ Sci Pollut*
804 *Res.* 24:3664–3674.
- 805 Carranza CS, Barberis CL, Chiacchiera SM, Magnoli CE. 2014. Influence of the pesticides
806 glyphosate, chlorpyrifos and atrazine on growth parameters of nonochratoxigenic *Aspergillus*
807 section *Nigri* strains isolated from agricultural soils. *J Environ Sci Health Part B.* 49:747–755.
- 808 Carvalho FP. 2017. Pesticides, environment, and food safety. *Food Energy Secur.* 6:48–60.
- 809 Castellana G, Loffredo E. 2014. Simultaneous removal of endocrine disruptors from a wastewater
810 using white rot fungi and various adsorbents. *Water Air Soil Pollut.* Available from:
811 <http://link.springer.com/10.1007/s11270-014-1872-6>
- 812 Ceci A, Pierro L, Riccardi C, Pinzari F, Maggi O, Persiani AM, Gadd GM, Petrangeli Papini M.
813 2015. Biotransformation of β -hexachlorocyclohexane by the saprotrophic soil fungus *Penicillium*
814 *griseofulvum*. *Chemosphere.* 137:101–107.
- 815 Chahal V, Nagpal A, Pakade Y, Katnoria J. 2014. Ecotoxicological studies of soil using analytical
816 and biological methods: a review. *World Acad Sci Eng Technol Int J Biol Biomol Agric Food*
817 *Biotechnol Eng.* 8:302–318.
- 818 Chakraborty S, Sikder J, Mukherjee D, Kanti Mandal M, Arockiasamy DL. 2013. Bioremediation
819 of water: a sustainable approach. In: Piemonte V, De Falco M, Basile A, editors. *Sustain Dev Chem*
820 *Eng Innov Technol.* Chichester, UK: John Wiley & Sons, Ltd; p. 241–266. Available from:
821 <http://doi.wiley.com/10.1002/9781118629703.ch10>
- 822 Chan-Cupul W, Abarca GH, Vázquez RR, Salmenes D, Hernández RG, Gutiérrez EA. 2014.
823 Response of ligninolytic macrofungi to the herbicide atrazine: dose-response bioassays. *Rev Argent*
824 *Microbiol.* 46:348–357.

- 825 Chan-Cupul W, Heredia-Abarca G, Rodríguez-Vázquez R. 2016. Atrazine degradation by fungal
826 co-culture enzyme extracts under different soil conditions. *J Environ Sci Health Part B*. 51:298–
827 308.
- 828 Chapman PM, Anderson J. 2005. A decision-making framework for sediment contamination. *Integr*
829 *Environ Assess Manag*. 1:163.
- 830 Colborn T. 1995. Environmental estrogens: health implications for humans and wildlife. *Environ*
831 *Health Perspect*. 103:135.
- 832 Colborn T, vom Saal FS, Soto AM. 1993. Developmental effects of endocrine-disrupting chemicals
833 in wildlife and humans. *Environ Health Perspect*. 101:378.
- 834 Coosen R, van Velsen FL. 1989. Effects of the β -isomer of hexachlorocyclohexane on estrogen-
835 sensitive human mammary tumor cells. *Toxicol Appl Pharmacol*. 101:310–318.
- 836 Cummings AM. 1997. Methoxychlor as a model for environmental estrogens. *Crit Rev Toxicol*.
837 27:367–379.
- 838 Čvančarová M, Moeder M, Filipová A, Cajthaml T. 2015. Biotransformation of fluoroquinolone
839 antibiotics by ligninolytic fungi - metabolites, enzymes and residual antibacterial activity.
840 *Chemosphere*. 136:311–320.
- 841 Cycoń M, Piotrowska-Seget Z, Kozdrój J. 2010. Linuron effects on microbiological characteristics
842 of sandy soils as determined in a pot study. *Ann Microbiol*. 60:439–449.
- 843 Da Silva Coelho J, de Oliveira AL, Marques de Souza CG, Bracht A, Peralta RM. 2010. Effect of
844 the herbicides bentazon and diuron on the production of ligninolytic enzymes by *Ganoderma*
845 *lucidum*. *Int Biodeterior Biodegrad*. 64:156–161.
- 846 De A, Bose R, Kumar A, Mozumdar S. 2014. Worldwide pesticide use. In: *Target Deliv Pestic*
847 *Using Biodegrad Polym Nanoparticles*. New Delhi: Springer India; p. 5–6. Available from:
848 http://link.springer.com/10.1007/978-81-322-1689-6_2
- 849 De Henau H. 1997. Biodegradation. In: Calow P, editor. *Handb Ecotoxicol*. Oxford, UK: Blackwell
850 Publishing Ltd; p. 355–377. Available from: <http://doi.wiley.com/10.1002/9781444313512.ch18>
- 851 Depledge MH, Amaral-Mendes JJ, Daniel B, Halbrook RS, Kloepper-Sams P, Moore MN, Peakall
852 DB. 1993. The conceptual basis of the biomarker approach. In: Peakall DB, Shugart LR, editors.

- 853 Biomarkers. Berlin, Heidelberg: Springer Berlin Heidelberg; p. 15–29. Available from:
854 http://www.springerlink.com/index/10.1007/978-3-642-84631-1_2
- 855 De Roy K, Marzorati M, Van den Abbeele P, Van de Wiele T, Boon N. 2014. Synthetic microbial
856 ecosystems: an exciting tool to understand and apply microbial communities. *Environ Microbiol.*
857 16:1472–1481.
- 858 De Souza CP, de Andrade Guedes T, Fontanetti CS. 2016. Evaluation of herbicides action on plant
859 bioindicators by genetic biomarkers: a review. *Environ Monit Assess.* 188:694.
- 860 Dorn PB, Salanitro JP. 2000. Temporal ecological assessment of oil contaminated soils before and
861 after bioremediation. *Chemosphere.* 40:419–426.
- 862 Dudášová H, Lászlová K, Lukáčová L, Balaščáková M, Murínová S, Dercová K. 2016.
863 Bioremediation of PCB-contaminated sediments and evaluation of their pre- and post-treatment
864 ecotoxicity. *Chem Pap.* Available from: [http://www.degruyter.com/view/j/chempap.2016.70.issue-](http://www.degruyter.com/view/j/chempap.2016.70.issue-8/chempap-2016-0041/chempap-2016-0041.xml)
865 [8/chempap-2016-0041/chempap-2016-0041.xml](http://www.degruyter.com/view/j/chempap.2016.70.issue-8/chempap-2016-0041/chempap-2016-0041.xml)
- 866 EEA. 2005. Environment and health. Luxembourg: Office for Official Publications of the European
867 Communities.
- 868 EEA. 2016. EMEP/EEA air pollutant emission inventory guidebook 2016: 3. Df, 3. I Agriculture
869 other including use of pesticides.
- 870 Eldridge ML, Sanseverino J, Layton AC, Easter JP, Schultz TW, Sayler GS. 2007. *Saccharomyces*
871 *cerevisiae* BLYAS, a new bioluminescent bioreporter for detection of androgenic compounds. *Appl*
872 *Environ Microbiol.* 73:6012-8.
- 873 Ellegaard-Jensen L, Knudsen BE, Johansen A, Albers CN, Aamand J, Rosendahl S. 2014. Fungal–
874 bacterial consortia increase diuron degradation in water-unsaturated systems. *Sci Total Environ.*
875 466-467:699–705.
- 876 Eurostat. 2016. Agri-environmental indicator - consumption of pesticides. Eurostat Statistics
877 Explained.
- 878 Fan X, Song F. 2014. Bioremediation of atrazine: recent advances and promises. *J Soils Sediments.*
879 14:1727–1737.

- 880 Feiler U, Kirchesch I, Heininger P. 2004. A new plant-based bioassay for aquatic sediments. *J Soils*
881 *Sediments*. 4:261–266.
- 882 Fontanetti CS, Nogarol LR, de Souza RB, Perez GD, Maziviero GT. 2011. Bioindicators and
883 biomarkers in the assessment of soil toxicity. In: Pascucci S, editor. *Soil Contam. InTech*. Available
884 from: [http://www.intechopen.com/books/soil-contamination/bioindicators-and-biomarkers-in-the-](http://www.intechopen.com/books/soil-contamination/bioindicators-and-biomarkers-in-the-assessment-of-soil-toxicity)
885 [assessment-of-soil-toxicity](http://www.intechopen.com/books/soil-contamination/bioindicators-and-biomarkers-in-the-assessment-of-soil-toxicity)
- 886 Frische T. 2003. Ecotoxicological evaluation of in situ bioremediation of soils contaminated by the
887 explosive 2,4,6-trinitrotoluene (TNT). *Environ Pollut*. 121:103–113.
- 888 Gao D, Du L, Yang J, Wu W-M, Liang H. 2010. A critical review of the application of white rot
889 fungus to environmental pollution control. *Crit Rev Biotechnol*. 30:70–77.
- 890 García-Carmona M, Romero-Freire A, Sierra Aragón M, Martínez Garzón FJ, Martín Peinado FJ.
891 2017. Evaluation of remediation techniques in soils affected by residual contamination with heavy
892 metals and arsenic. *J Environ Manage*. 191:228–236.
- 893 Garey J, Wolff MS. 1998. Estrogenic and antiprogesteragenic activities of pyrethroid insecticides.
894 *Biochem Biophys Res Commun*. 251:855–859.
- 895 Gargano ML, van Griensven LJLD, Isikhuemhen OS, Lindequist U, Venturella G, Wasser SP,
896 Zervakis GI. 2017. Medicinal mushrooms: valuable biological resources of high exploitation
897 potential. *Plant Biosyst - Int J Deal Asp Plant Biol*. 151:548–565.
- 898 Gong ZM, Xu FL, Dawson R, Cao J, Liu WX, Li BG, Shen WR, Zhang WJ, Qin BP, Sun R, Tao S.
899 2004. Residues of hexachlorocyclohexane isomers and their distribution characteristics in soils in
900 the Tianjin Area, China. *Arch Environ Contam Toxicol*. Available from:
901 <http://link.springer.com/10.1007/s00244-003-2301-9>
- 902 Großkopf T, Soyer OS. 2014. Synthetic microbial communities. *Curr Opin Microbiol*. 18:72–77.
- 903 Guerreño M, López Armengol MF, Luquet CM, Venturino A. 2016. Comparative study of toxicity
904 and biochemical responses induced by sublethal levels of the pesticide azinphosmethyl in two fish
905 species from North-Patagonia, Argentina. *Aquat Toxicol*. 177:365–372.
- 906 Guida M, Inglese M, Meriç S. 2008. A multi-battery toxicity investigation on fungicides.
907 *Desalination*. 226:262–270.

- 908 Guo D, Wang Y, Qian Y, Chen C, Jiao B, Cai L, Wang Q. 2017. Joint acute and endocrine
909 disruptive toxicities of malathion, cypermethrin and prochloraz to embryo-larval zebrafish, *Danio*
910 *rerio*. Chemosphere. 166:63–71.
- 911 Hai FI, Modin O, Yamamoto K, Fukushi K, Nakajima F, Nghiem LD. 2012. Pesticide removal by a
912 mixed culture of bacteria and white-rot fungi. J Taiwan Inst Chem Eng. 43:459–462.
- 913 Hammesfahr U, Heuer H, Manzke B, Smalla K, Thiele-Bruhn S. 2008. Impact of the antibiotic
914 sulfadiazine and pig manure on the microbial community structure in agricultural soils. Soil Biol
915 Biochem. 40:1583–1591.
- 916 Harclerode M, Ridsdale DR, Darmendrail D, Bardos P, Alexandrescu F, Nathanail P, Pizzol L,
917 Rizzo E. 2015. Integrating the social dimension in remediation decision-making: state of the
918 practice and way forward. Remediat J. 26:11–42.
- 919 Harms H, Wick LY, Schlosser D. 2017. The fungal community in organically polluted systems. In:
920 Dighton J, White JF, editors. Mycology. CRC Press Taylor & Francis Group 6000 Broken Sound
921 Parkway NW, Suite 300 Boca Raton, FL 33487-2742: CRC Press; p. 459–470. Available from:
922 <http://www.crcnetbase.com/doi/10.1201/9781315119496-32>
- 923 Henson-Ramsey H, Schneider A, Stoskopf MK. 2011. A comparison of multiple esterases as
924 biomarkers of organophosphate exposure and effect in two earthworm species. Bull Environ
925 Contam Toxicol. 86:373–378.
- 926 Hollert H, Keiter S, König N, Rudolf M, Ulrich M, Braunbeck T. 2003. A new sediment contact
927 assay to assess particle-bound pollutants using zebrafish (*Danio rerio*) embryos. J Soils Sediments.
928 3:197.
- 929 Houx NWH, Dekker A, Van Kammen-Polman AMM, Ronday R. 1996. Acute toxicity test for
930 terrestrial hazard assessment with exposure of *Folsomia candida* to pesticides in an aqueous
931 medium. Arch Environ Contam Toxicol. 30:9–14.
- 932 Huang McBeath J, McBeath J. 2010. Environmental change and food security in China. Dordrecht:
933 Springer Netherlands. Available from: <http://link.springer.com/10.1007/978-1-4020-9180-3>
- 934 Hui X, Yi Q, Bu-zhuo P, Xiliu J, Xiao-mei H. 2003. Environmental pesticide pollution and its
935 countermeasures in China. AMBIO J Hum Environ. 32:78–80.

- 936 Iqbal M. 2016. *Vicia faba* bioassay for environmental toxicity monitoring: A review. Chemosphere.
937 144:785–802.
- 938 Jagmann N, Philipp B. 2014. Reprint of design of synthetic microbial communities for
939 biotechnological production processes. J Biotechnol. 192:293–301.
- 940 Jarošová B, Erseková A, Hilscherová K, Loos R, Gawlik BM, Giesy JP, Bláha L. 2014. Europe-
941 wide survey of estrogenicity in wastewater treatment plant effluents: the need for the effect-based
942 monitoring. Environ Sci Pollut Res. 21:10970–10982.
- 943 Jechalke S, Heuer H, Siemens J, Amelung W, Smalla K. 2014. Fate and effects of veterinary
944 antibiotics in soil. Trends Microbiol. 22:536–545.
- 945 Jensen TK, Toppari J, Keiding N, Skakkebaek NE. 1995. Do environmental estrogens contribute to
946 the decline in male reproductive health? Clin Chem. 41:1896–1901.
- 947 Jones A, Panagos P, Barcelo S, Bouraoui F, Bosco C, Dewitte O, Gardi C, Erhard M, Hervás J,
948 Hiederer R. 2012. The state of soil in Europe. JRC Ref Rep Jt Res Cent Eur Comm Ispra.
- 949 Kardol P, Wardle DA. 2010. How understanding aboveground-belowground linkages can assist
950 restoration ecology. Trends Ecol Evol. 25:670–679.
- 951 Kaur H, Kapoor S, Kaur G. 2016. Application of ligninolytic potentials of a white-rot fungus
952 *Ganoderma lucidum* for degradation of lindane. Environ Monit Assess. 188:588.
- 953 Keddy CJ, Greene JC, Bonnell MA. 1995. Review of whole-organism bioassays: soil, freshwater
954 sediment, and freshwater assessment in Canada. Ecotoxicol Environ Saf. 30:221–251.
- 955 Khalil SR, Awad A, Mohammed HH. 2017. Behavioral response and gene expression changes in
956 fipronil-administered male Japanese quail (*Coturnix japonica*). Environ Pollut Barking Essex 1987.
957 223:51–61.
- 958 Khromonygina VV, Saltykova AI, Vasil'chenko LG, Kozlov IP, Rabinovich ML. 2004.
959 Degradation of the herbicide atrazine by the soil mycelial fungus INBI 2-26(-)--a producer of
960 cellobiose dehydrogenase. Prikl Biokhim Mikrobiol. 40:337–343.
- 961 Kim K-H, Kabir E, Jahan SA. 2017. Exposure to pesticides and the associated human health effects.
962 Sci Total Environ. 575:525–535.

- Knudsen BE, Ellegaard-Jensen L, Albers CN, Rosendahl S, Aamand J. 2013. Fungal hyphae stimulate bacterial degradation of 2,6-dichlorobenzamide (BAM). *Environ Pollut.* 181:122–127.
- Köck-Schulmeyer M, Villagrasa M, López de Alda M, Céspedes-Sánchez R, Ventura F, Barceló D. 2013. Occurrence and behavior of pesticides in wastewater treatment plants and their environmental impact. *Sci Total Environ.* 458-460:466–476.
- Kojima H, Katsura E, Takeuchi S, Niiyama K, Kobayashi K. 2004. Screening for estrogen and androgen receptor activities in 200 pesticides by in vitro reporter gene assays using Chinese hamster ovary cells. *Environ Health Perspect.* 112:524–531.
- Koroleva O, Stepanova E, Landesman E, Vasilchenko L, Khromonygina V, Zherdev A, Rabinovich M. 2002. Enzyme immunoassay of herbicide decomposition by soil and wood decay fungi. *Appl Biochem Microbiol.* 38:355–360.
- Kulshreshtha S, Mathur N, Bhatnagar P. 2014. Mushroom as a product and their role in mycoremediation. *AMB Express.* Available from: <http://www.amb-express.com/content/4/1/29>
- Lee S-E, Young-Woong C, Mo H, Son J, Park K, Cho K. 2013. Endosulfan-induced biomarkers in Japanese rice fish (*Oryzias latipes*) analyzed by SELDI-TOF-MS. *Int J Biol Sci.* 9:343–349.
- Liu WW, Wu B, Xiang M, Liu X. 2017. From microbiome to synthetic microbial community. *Microbiol China.* 44: 1-9 (in Chinese).
- Lizano-Fallas V, Masís-Mora M, Espinoza-Villalobos D, Lizano-Brenes M, Rodríguez-Rodríguez CE. 2017. Removal of pesticides and ecotoxicological changes during the simultaneous treatment of triazines and chlorpyrifos in biomixtures. *Chemosphere.* 182:106–113.
- Loehr RC, Webster MT. 1997. Changes in toxicity and mobility resulting from bioremediation processes. *Bioremediation J.* 1:149–163.
- Loffredo E, Castellana G, Taskin E. 2016. A two-step approach to eliminate pesticides and estrogens from a wastewater and reduce its phytotoxicity: adsorption onto plant-derived materials and fungal degradation. *Water Air Soil Pollut.* Available from: <http://link.springer.com/10.1007/s11270-016-2883-2>
- Lofrano G, Carotenuto M, Uyguner-Demirel CS, Vitagliano A, Siciliano A, Guida M. 2014. An integrated chemical and ecotoxicological assessment for the photocatalytic degradation of vancomycin. *Environ Technol.* 35:1234–1242.

- 992 Loos R, Locoro G, Comero S, Contini S, Schwesig D, Werres F, Balsaa P, Gans O, Weiss S, Blaha
993 L, et al. 2010. Pan-European survey on the occurrence of selected polar organic persistent pollutants
994 in ground water. *Water Res.* 44:4115–4126.
- 995 Luo Y, Guo W, Ngo HH, Nghiem LD, Hai FI, Zhang J, Liang S, Wang XC. 2014. A review on the
996 occurrence of micropollutants in the aquatic environment and their fate and removal during
997 wastewater treatment. *Sci Total Environ.* 473-474:619–641.
- 998 Lu X, Kang Z, Tao B, Wang Y, Dong J, Zhang J. 2012. Degradation of nicosulfuron by *Bacillus*
999 *subtilis* YB1 and *Aspergillus niger* YF1. *Appl Biochem Microbiol.* 48:460–466.
- 1000 Ma M, Li J, Wang Z. 2005. Assessing the detoxication efficiencies of wastewater treatment
1001 processes using a battery of bioassays/biomarkers. *Arch Environ Contam Toxicol.* 49:480–487.
- 1002 Ma T-H. 1999. Global environment monitoring for ecosystem health and human well being —
1003 report from a round table discussion on international collaborative studies. *JASSA J Appl Sci South*
1004 *Afr.* 5:89–94.
- 1005 Maciel GM, Inácio FD, de Sá-Nakanishi AB, Haminiuk CWI, Castoldi R, Comar JF, Bracht A,
1006 Peralta RM. 2013. Response of *Ganoderma lucidum* and *Trametes* sp. to the herbicide picloram:
1007 tolerance, antioxidants and production of ligninolytic enzymes. *Pestic Biochem Physiol.* 105:84–92.
- 1008 Madrigal-Zúñiga K, Ruiz-Hidalgo K, Chin-Pampillo JS, Masís-Mora M, Castro-Gutiérrez V,
1009 Rodríguez-Rodríguez CE. 2016. Fungal bioaugmentation of two rice husk-based biomixtures for
1010 the removal of carbofuran in on-farm biopurification systems. *Biol Fertil Soils.* 52:243–250.
- 1011 Majeau J-A, Brar SK, Tyagi RD. 2010. Laccases for removal of recalcitrant and emerging
1012 pollutants. *Bioresour Technol.* 101:2331–2350.
- 1013 Marcoccia D, Pellegrini M, Fiocchetti M, Lorenzetti S, Marino M. 2017. Food components and
1014 contaminants as (anti)androgenic molecules. *Genes Nutr.* Available from:
1015 <http://genesandnutrition.biomedcentral.com/articles/10.1186/s12263-017-0555-5>
- 1016 Marco-Urrea E, Pérez-Trujillo M, Vicent T, Caminal G. 2009. Ability of white-rot fungi to remove
1017 selected pharmaceuticals and identification of degradation products of ibuprofen by *Trametes*
1018 *versicolor*. *Chemosphere.* 74:765–772.

- 1019 Margot J, Copin P-J, von Gunten U, Barry DA, Holliger C. 2015. Sulfamethoxazole and
1020 isoproturon degradation and detoxification by a laccase-mediator system: Influence of treatment
1021 conditions and mechanistic aspects. *Biochem Eng J.* 103:47–59.
- 1022 Marziali L, Rosignoli F, Drago A, Pascariello S, Valsecchi L, Rossaro B, Guzzella L. 2017.
1023 Toxicity risk assessment of mercury, DDT and arsenic legacy pollution in sediments: A triad
1024 approach under low concentration conditions. *Sci Total Environ.* 593-594:809–821.
- 1025 Masaphy S, Levanon D, Vaya J, Henis Y. 1993. Isolation and characterization of a novel atrazine
1026 metabolite produced by the fungus *Pleurotus pulmonarius*, 2-chloro-4-ethylamino-6-(1-
1027 hydroxyisopropyl)amino-1,3,5-triazine. *Appl Environ Microbiol.* 59:4342–4346.
- 1028 Mattsson MK, Liu X, Yu D, Kontro MH. 2015. Depth, soil type, water table, and site effects on
1029 microbial community composition in sediments of pesticide-contaminated aquifer. *Environ Sci*
1030 *Pollut Res.* 22:10263–10279.
- 1031 Matute RG, Figlas D, Mockel G, Curvetto N. 2012. Degradation of metsulfuron methyl by *Agaricus*
1032 *blazei* Murrill spent compost enzymes. *Bioremediation J.* 16:31–37.
- 1033 Meyer H-P, Munch T. 2005. Swiss industrial biocatalysis consortium. *BioWorld Eur.* 1:14–16.
- 1034 Miccoli A, Maradonna F, De Felice A, Caputo Barucchi V, Estonba A, Genangeli M, Vittori S,
1035 Leonori I, Carnevali O. 2017. Detection of endocrine disrupting chemicals and evidence of their
1036 effects on the HPG axis of the European anchovy *Engraulis encrasicolus*. *Mar Environ Res.*
1037 127:137–147.
- 1038 Miller GT. 2004. *Living in the environment: principles, connections, and solutions*. 13. ed., internat.
1039 student ed. Pacific Grove, Calif.: Thomson/Brooks/Cole.
- 1040 Mir-Tutusa JA, Masís-Mora M, Corcellas C, Eljarrat E, Barceló D, Sarrà M, Caminal G, Vicent
1041 T, Rodríguez-Rodríguez CE. 2014. Degradation of selected agrochemicals by the white rot fungus
1042 *Trametes versicolor*. *Sci Total Environ.* 500-501:235–242.
- 1043 Mnif W, Hassine AIH, Bouaziz A, Bartegi A, Thomas O, Roig B. 2011. Effect of endocrine
1044 disruptor pesticides: a review. *Int J Environ Res Public Health.* 8:2265–2303.
- 1045 Molina-Barahona L, Vega-Loyo L, Guerrero M, Ramírez S, Romero I, Vega-Jarquín C, Albores A.
1046 2005. Ecotoxicological evaluation of diesel-contaminated soil before and after a bioremediation
1047 process. *Environ Toxicol.* 20:100–109.

- 1048 Molnár M, Gruiz K, Halász M. 2007. Integrated methodology to evaluate bioremediation potential
1049 of creosote-contaminated soils. *Period Polytech Chem Eng.* 51:23.
- 1050 Mori T, Wang J, Tanaka Y, Nagai K, Kawagishi H, Hirai H. 2017. Bioremediation of the
1051 neonicotinoid insecticide clothianidin by the white-rot fungus *Phanerochaete sordida*. *J Hazard*
1052 *Mater.* 321:586–590.
- 1053 Morriën E, Hannula SE, Snoek LB, Helmsing NR, Zweers H, de Hollander M, Soto RL, Bouffaud
1054 M-L, Buée M, Dimmers W, et al. 2017. Soil networks become more connected and take up more
1055 carbon as nature restoration progresses. *Nat Commun.* 8:14349.
- 1056 Mortelmans K, Zeiger E. 2000. The Ames *Salmonella*/microsome mutagenicity assay. *Mutat Res*
1057 *Mol Mech Mutagen.* 455:29–60.
- 1058 Mougin C, Kollmann A, Jolival C. 2002. Enhanced production of laccase in the fungus *Trametes*
1059 *versicolor* by the addition of xenobiotics. *Biotechnol Lett.* 24:139–142.
- 1060 Mrema EJ, Rubino FM, Colosio C. 2013. Obsolete pesticides – A threat to environment,
1061 biodiversity and human health. In: Simeonov LI, Macaev FZ, Simeonova BG, editors. *Environ*
1062 *Secur Assess Manag Obsolete Pestic Southeast Eur.* Dordrecht: Springer Netherlands; p. 1–21.
1063 Available from: http://link.springer.com/10.1007/978-94-007-6461-3_1
- 1064 Nakata H, Hirakawa Y, Kawazoe M, Nakabo T, Arizono K, Abe S-I, Kitano T, Shimada H,
1065 Watanabe I, Li W, Ding X. 2005. Concentrations and compositions of organochlorine contaminants
1066 in sediments, soils, crustaceans, fishes and birds collected from Lake Tai, Hangzhou Bay and
1067 Shanghai city region, China. *Environ Pollut.* 133:415–429.
- 1068 Neumann-Hensel H, Melbye K. 2006. Optimisation of the solid-contact test with *Arthrobacter*
1069 *globiformis*. *J Soils Sediments.* 6:201–207.
- 1070 Nguyen LN, Hai FI, Yang S, Kang J, Leusch FDL, Roddick F, Price WE, Nghiem LD. 2014.
1071 Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals
1072 and pesticides by *Trametes versicolor*. Role of biosorption and biodegradation. *Int Biodeterior*
1073 *Biodegrad.* 88:169–175.
- 1074 Odukkathil G, Vasudevan N. 2013. Toxicity and bioremediation of pesticides in agricultural soil.
1075 *Rev Environ Sci Biotechnol.* 12:421–444.

- 1076 OECD 201. 2006. Test No. 201: Alga, growth inhibition test. OECD Publishing. Available from:
1077 <http://www.oecd-ilibrary.org/environment/test-no-201-alga-growth-inhibition->
1078 [test_9789264069923-en](http://www.oecd-ilibrary.org/environment/test-no-201-alga-growth-inhibition-test_9789264069923-en)
- 1079 OECD 209. 1984. Test No. 209: Activated sludge, respiration inhibition test. OECD Publishing.
1080 Available from: <http://www.oecd-ilibrary.org/oecd/content/book/9789264070080-en>
- 1081 OECD 211. 2012. Test No. 211: Daphnia magna reproduction test. OECD Publishing. Available
1082 from: <http://www.oecd-ilibrary.org/environment/test-no-211-daphnia-magna-reproduction->
1083 [test_9789264185203-en](http://www.oecd-ilibrary.org/environment/test-no-211-daphnia-magna-reproduction-test_9789264185203-en)
- 1084 Oliveira BR, Penetra A, Cardoso VV, Benoliel MJ, Barreto Crespo MT, Samson RA, Pereira VJ.
1085 2015. Biodegradation of pesticides using fungi species found in the aquatic environment. Environ
1086 Sci Pollut Res Int. 22:11781–11791.
- 1087 Palvannan T, Saravanakumar T, Unnithan AR, Chung N-J, Kim D-H, Park S-M. 2014. Efficient
1088 transformation of phenyl urea herbicide chloroxuron by laccase immobilized on zein polyurethane
1089 nanofiber. J Mol Catal B Enzym. 99:156–162.
- 1090 Pan M, Chu LM. 2016. Adsorption and degradation of five selected antibiotics in agricultural soil.
1091 Sci Total Environ. 545-546:48–56.
- 1092 Park D-S, Jeon H-J, Park E-S, Bae IK, Kim Y-E, Lee S-E. 2015. Highly selective biomarkers for
1093 pesticides developed in *Eisenia fetida* using SELDI-TOF MS. Environ Toxicol Pharmacol. 39:635–
1094 642.
- 1095 Parrón T, Requena M, Hernández AF, Alarcón R. 2014. Environmental exposure to pesticides and
1096 cancer risk in multiple human organ systems. Toxicol Lett. 230:157–165.
- 1097 Parvez S, Venkataraman C, Mukherji S. 2006. A review on advantages of implementing
1098 luminescence inhibition test (*Vibrio fischeri*) for acute toxicity prediction of chemicals. Environ Int.
1099 32:265–268.
- 1100 Phipps GL, Ankley GT, Benoit DA, Mattson VR. 1993. Use of the aquatic oligochaete *Lumbriculus*
1101 *variegatus* for assessing the toxicity and bioaccumulation of sediment-associated contaminants.
1102 Environ Toxicol Chem. 12:269–279.

- 1103 Pinto AP, Rodrigues SC, Caldeira AT, Teixeira DM. 2016. Exploring the potential of novel
1104 biomixtures and *Lentinula edodes* fungus for the degradation of selected pesticides. Evaluation for
1105 use in biobed systems. *Sci Total Environ.* 541:1372–1381.
- 1106 Pinto AP, Serrano C, Pires T, Mestrinho E, Dias L, Teixeira DM, Caldeira AT. 2012. Degradation
1107 of terbuthylazine, difenoconazole and pendimethalin pesticides by selected fungi cultures. *Sci Total*
1108 *Environ.* 435-436:402–410.
- 1109 Polo AM, Tobajas M, Sanchis S, Mohedano AF, Rodríguez JJ. 2011. Comparison of experimental
1110 methods for determination of toxicity and biodegradability of xenobiotic compounds.
1111 *Biodegradation.* 22:751–761.
- 1112 Popp J, Pető K, Nagy J. 2013. Pesticide productivity and food security. A review. *Agron Sustain*
1113 *Dev.* 33:243–255.
- 1114 Prokop Z, Nečasová A, Klánová J, Čupr P. 2016. Bioavailability and mobility of organic
1115 contaminants in soil: new three-step ecotoxicological evaluation. *Environ Sci Pollut Res.* 23:4312–
1116 4319.
- 1117 Purvis IJ, Chotai D, Dykes CW, Lubahn DB, French FS, Wilson EM, Hobden AN. 1991. An
1118 androgen-inducible expression system for *Saccharomyces cerevisiae*. *Gene.* 106:35–42.
- 1119 Qiu D. 2011. Development strategy for bio-pesticide and biological control. *J Agric Sci Technol.*
1120 5:020.
- 1121 Radwan MA, Mohamed MS. 2013. Imidacloprid induced alterations in enzyme activities and
1122 energy reserves of the land snail, *Helix aspersa*. *Ecotoxicol Environ Saf.* 95:91–97.
- 1123 Ridsdale DR, Noble BF. 2016. Assessing sustainable remediation frameworks using sustainability
1124 principles. *J Environ Manage.* 184:36–44.
- 1125 Rodea-Palomares I, Gonzalez-Pleiter M, Gonzalo S, Rosal R, Leganes F, Sabater S, Casellas M,
1126 Munoz-Carpena R, Fernandez-Pinas F. 2016. Hidden drivers of low-dose pharmaceutical pollutant
1127 mixtures revealed by the novel GSA-QHTS screening method. *Sci Adv.* 2:e1601272–e1601272.
- 1128 Rönnpagel K, Liss W, Ahlf W. 1995. Microbial bioassays to assess the toxicity of solid-associated
1129 contaminants. *Ecotoxicol Environ Saf.* 31:99–103.

- 1130 Saillenfait A-M, Ndiaye D, Sabaté J-P. 2016. The estrogenic and androgenic potential of
1131 pyrethroids in vitro. Review. Toxicol Vitro. 34:321–332.
- 1132 Sandin M, Piikki K, Jarvis N, Larsbo M, Bishop K, Kreuger J. 2018. Spatial and temporal patterns
1133 of pesticide concentrations in streamflow, drainage and runoff in a small Swedish agricultural
1134 catchment. Sci Total Environ. 610–611: 623–634.
- 1135 Salvestrini S, Canzano S, Iovino P, Leone V, Capasso S. 2014. Modelling the biphasic sorption of
1136 simazine, imidacloprid, and boscalid in water/soil systems. J Environ Sci Health Part B. 49:578–
1137 590.
- 1138 Sarkar D, Ferguson M, Datta R, Birnbaum S. 2005. Bioremediation of petroleum hydrocarbons in
1139 contaminated soils: comparison of biosolids addition, carbon supplementation, and monitored
1140 natural attenuation. Environ Pollut. 136:187–195.
- 1141 Šašek V, Volfová O, Erbanová P, Vyas BRM, Matucha M. 1993. Degradation of PCBs by white rot
1142 fungi, methylotrophic and hydrocarbon utilizing yeasts and bacteria. Biotechnol Lett. 15:521–526.
- 1143 Scott PD, Coleman HM, Colville A, Lim R, Matthews B, McDonald JA, Miranda A, Neale PA,
1144 Nugegoda D, Tremblay LA, Leusch FDL. 2017. Assessing the potential for trace organic
1145 contaminants commonly found in Australian rivers to induce vitellogenin in the native rainbowfish
1146 (*Melanotaenia fluviatilis*) and the introduced mosquitofish (*Gambusia holbrooki*). Aquat Toxicol
1147 Amst Neth. 185:105–120.
- 1148 Shah PC, Kumar VR, Dastager SG, Khire JM. 2017. Phytase production by *Aspergillus niger*
1149 NCIM 563 for a novel application to degrade organophosphorus pesticides. AMB Express. 7:66.
- 1150 Shen W, Zhu N, Cui J, Wang H, Dang Z, Wu P, Luo Y, Shi C. 2016. Ecotoxicity monitoring and
1151 bioindicator screening of oil-contaminated soil during bioremediation. Ecotoxicol Environ Saf.
1152 124:120–128.
- 1153 Siciliano A, Gesuele R, Pagano G, Guida M. 2015. How Daphnia (Cladocera) Assays may be used
1154 as Bioindicators of Health Effects? J Biodivers Endanger Species. S1:005. Available from:
1155 [https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-](https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936)
1156 [bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936](https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936)
- 1157 Silsirivanit A, Sawanyawisuth K, Riggins GJ, Wongkham C. 2014. Cancer biomarker discovery for
1158 cholangiocarcinoma: the high-throughput approaches. J Hepato-Biliary-Pancreat Sci. 21:388–396.

- 1159 Singh DK. 2008. Biodegradation and bioremediation of pesticide in soil: concept, method and
1160 recent developments. *Indian J Microbiol.* 48:35–40.
- 1161 Song J, Gu J, Zhai Y, Wu W, Wang H, Ruan Z, Shi Y, Yan Y. 2013. Biodegradation of
1162 nicosulfuron by a *Talaromyces flavus* LZM1. *Bioresour Technol.* 140:243–248.
- 1163 Sonnenschein C, Soto AM. 1998. An updated review of environmental estrogen and androgen
1164 mimics and antagonists. *J Steroid Biochem Mol Biol.* 65:143–150.
- 1165 Soto AM, Sonnenschein C, Chung KL, Fernandez MF, Olea N, Serrano FO. 1995. The E-SCREEN
1166 assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ*
1167 *Health Perspect.* 103:113.
- 1168 Spina F, Cordero C, Schilirò T, Sgorbini B, Pignata C, Gilli G, Bicchi C, Varese GC. 2015.
1169 Removal of micropollutants by fungal laccases in model solution and municipal wastewater:
1170 evaluation of estrogenic activity and ecotoxicity. *J Clean Prod.* 100:185–194.
- 1171 Stepić S, Hackenberger BK, Velki M, Lončarić Ž, Hackenberger DK. 2013. Effects of individual
1172 and binary-combined commercial insecticides endosulfan, temephos, malathion and pirimiphos-
1173 methyl on biomarker responses in earthworm *Eisenia andrei*. *Environ Toxicol Pharmacol.* 36:715–
1174 723.
- 1175 Talk A, Kublik S, Uksa M, Engel M, Berghahn R, Welzl G, Schlöter M, Mohr S. 2016. Effects of
1176 multiple but low pesticide loads on aquatic fungal communities colonizing leaf litter. *J Environ Sci.*
1177 46:116–125.
- 1178 Tian J, Dong Q, Yu C, Zhao R, Wang J, Chen L. 2016. Biodegradation of the organophosphate
1179 trichlorfon and its major degradation products by a novel *Aspergillus sydowii* PA F-2. *J Agric Food*
1180 *Chem.* 64:4280–4287.
- 1181 Tigini V, Giansanti P, Mangiavillano A, Pannocchia A, Varese GC. 2011. Evaluation of toxicity,
1182 genotoxicity and environmental risk of simulated textile and tannery wastewaters with a battery of
1183 biotests. *Ecotoxicol Environ Saf.* 74:866–873.
- 1184 Traunsperger W, Haitzer M, Höss S, Beier S, Ahlf W, Steinberg C. 1997. Ecotoxicological
1185 assessment of aquatic sediments with *Caenorhabditis elegans* (nematoda) - a method for testing
1186 liquid medium and whole-sediment samples. *Environ Toxicol Chem.* 16:245–250.

- 1187 Ulčnik A, Kralj Cigić I, Pohleven F. 2013. Degradation of lindane and endosulfan by fungi, fungal
1188 and bacterial laccases. *World J Microbiol Biotechnol.* 29:2239–2247.
- 1189 Van Gestel CA, Van der Waarde JJ, Derksen JGM (Anja), Van der Hoek EE, Veul MFXW,
1190 Bouwens S, Rusch B, Kronenburg R, Stokman GNM. 2001. The use of acute and chronic bioassays
1191 to determine the ecological risk and bioremediation efficiency of oil-polluted soils. *Environ Toxicol*
1192 *Chem.* 20:1438–1449.
- 1193 Van Straalen NM, Van Gestel CAM. 1997. Soil invertebrates and micro-organisms. In: Calow P,
1194 editor. *Handb Ecotoxicol.* Oxford, UK: Blackwell Publishing Ltd; p. 251–277. Available from:
1195 <http://doi.wiley.com/10.1002/9781444313512.ch13>
- 1196 Vasil'Chenko L, Khromonygina V, Koroleva O, Landesman E, Gaponenko V, Kovaleva T, Kozlov
1197 YP, Rabinovich M. 2002. Consumption of triazine herbicide atrazine by laccase-positive and
1198 laccase-negative strains of soil fungus *Mycelia sterilia* INBI 2-26. *Appl Biochem Microbiol.*
1199 38:454–459.
- 1200 Velázquez-Fernández JB, Martínez-Rizo AB, Ramírez-Sandoval M, Domínguez-Ojeda D. 2012.
1201 Biodegradation and bioremediation of organic pesticides. In: Soundararajan R.P. editor.
1202 *Agricultural and Biological Sciences "Pesticides - Recent Trends in Pesticide Residue Assay"*; p.
1203 978-953.
- 1204 Verger PJP, Boobis AR. 2013. Reevaluate pesticides for food security and safety. *Science.*
1205 341:717–718.
- 1206 Verma JP, Jaiswal DK, Sagar R. 2014. Pesticide relevance and their microbial degradation: a-state-
1207 of-art. *Rev Environ Sci Biotechnol.* 13:429–466.
- 1208 Wang C, Yu L, Zhang Z, Wang B, Sun H. 2014. Tourmaline combined with *Phanerochaete*
1209 *chrysosporium* to remediate agricultural soil contaminated with PAHs and OCPs. *J Hazard Mater.*
1210 264:439–448.
- 1211 Wang Z, Yang T, Zhai Z, Zhang B, Zhang J. 2015. Reaction mechanism of dicofol removal by
1212 cellulase. *J Environ Sci China.* 36:22–28.
- 1213 Weber J, Kreutzmann J, Plantikow A, Claus E, Manz W, Heininger P. 2006. A novel particle
1214 contact assay with the yeast *Saccharomyces cerevisiae* for ecotoxicological assessment of
1215 freshwater sediments. *J Soils Sediments.* 6:84–91.

- 1216 Wei LN, Wu P, Wang FR, Yang H. 2015. Dissipation and degradation dynamics of thifluzamide in
1217 rice field. *Water Air Soil Pollut*. Available from: [http://link.springer.com/10.1007/s11270-015-](http://link.springer.com/10.1007/s11270-015-2387-5)
1218 2387-5
- 1219 Welch RM, Levin W, Conney AH. 1969. Estrogenic action of DDT and its analogs. *Toxicol Appl*
1220 *Pharmacol*. 14:358–367.
- 1221 Wernersson A-S, Carere M, Maggi C, Tusil P, Soldan P, James A, Sanchez W, Dulio V, Broeg K,
1222 Reifferscheid G, et al. 2015. The European technical report on aquatic effect-based monitoring tools
1223 under the water framework directive. *Environ Sci Eur*. Available from:
1224 <http://www.enveurope.com/content/27/1/7>
- 1225 White PA, Claxton LD. 2004. Mutagens in contaminated soil: a review. *Mutat Res Mutat Res*.
1226 567:227–345.
- 1227 Wolfand JM, LeFevre GH, Luthy RG. 2016. Metabolization and degradation kinetics of the urban-
1228 use pesticide fipronil by white rot fungus *Trametes versicolor*. *Environ Sci Process Impacts*.
1229 18:1256–1265.
- 1230 Wu B, Zhang XL, Cui BK, Dai YC. 2015a. Comparative genomic analysis of edible (medicinal)
1231 fungi reveals different ecological habitats. *Mycosystema* 34: 742-60 (in Chinese).
- 1232 Wu J, Lu J, Lu H, Lin Y, Wilson PC. 2015b. Occurrence and ecological risks from fipronil in
1233 aquatic environments located within residential landscapes. *Sci Total Environ*. 518-519:139–147.
- 1234 Xiao P, Mori T, Kamei I, Kiyota H, Takagi K, Kondo R. 2011. Novel metabolic pathways of
1235 organochlorine pesticides dieldrin and aldrin by the white rot fungi of the genus *Phlebia*.
1236 *Chemosphere*. 85:218–224.
- 1237 Xue N, Zhang D, Xu X. 2006. Organochlorinated pesticide multiresidues in surface sediments from
1238 Beijing Guanting reservoir. *Water Res*. 40:183–194.
- 1239 Yang Y, Li D, Mu D. 2008. Levels, seasonal variations and sources of organochlorine pesticides in
1240 ambient air of Guangzhou, China. *Atmos Environ*. 42:677–687.
- 1241 Zeng S, Qin X, Xia L. 2017. Degradation of the herbicide isoproturon by laccase-mediator systems.
1242 *Biochem Eng J*. 119:92–100.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1243 Zhang J, Qin J, Zhao C, Liu C, Xie H, Liang S. 2015. Response of bacteria and fungi in soil
1244 microcosm under the presence of pesticide endosulfan. Water Air Soil Pollut. Available from:
1245 <http://link.springer.com/10.1007/s11270-015-2309-6>

1246 Zhou Q, Hua T. 2004. Bioremediation: a review of applications and problems to be resolved. Prog
1247 Nat Sci. 14:937–944.

1248 Zhou R, Zhu L, Yang K, Chen Y. 2006. Distribution of organochlorine pesticides in surface water
1249 and sediments from Qiantang River, East China. J Hazard Mater. 137:68–75.

1250 Zielke H, Seiler T-B, Niebergall S, Leist E, Brinkmann M, Spira D, Streck G, Brack W, Feiler U,
1251 Braunbeck T. 2011. The impact of extraction methodologies on the toxicity of sediments in the
1252 zebrafish (*Danio rerio*) embryo test. J Soils Sediments. 11:352–363.

Fungi as a toolbox for a sustainable bioremediation of pesticides in soil and water

F. Spina¹, G. Cecchi², A. Landinez-Torres^{3,4}, L. Pecoraro⁵, F. Russo⁶, B. Wu⁵, L. Cai⁵, X. Z. Liu⁵, S. Tosi⁴, G.C. Varese¹, M. Zotti², A. M. Persiani⁶

AFFILIATION

¹ *Mycotheca Universitatis Turinensis, Department of Life Sciences and Systems Biology, University of Torino, Viale Mattioli, 25, 1025 Torino, Italy.*

² *Laboratory of Mycology, Department of Earth, Environment and Life Sciences, University of Genoa, Corso Europa, 26, 16136 Genoa, Italy*

³ *Facultad de Ciencias Agrarias y Ambientales, Fundación Universitaria Juan de Castellanos. Carrera 11 No. 11 - 44 Tunja, Boyacá - Colombia*

⁴ *Laboratory of Mycology, Department of Earth Science and Environmental Sciences, University of Pavia, Via S. Epifanio 14, 27100 Pavia, Italy*

⁵ *State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*

⁶ *Laboratory of Fungal Biodiversity, Department of Environmental Biology, Sapienza University of Rome, Piazzale A. Moro 5, 00185 Rome, Italy*

Corresponding Author: Anna Maria Persiani, Laboratory of Fungal Biodiversity, Department of Environmental Biology, Sapienza University of Rome, Piazzale A. Moro 5, 00185 Rome, Italy. Email: annamaria.persiani@uniroma1.it

Abstract

Pesticide can help reduce yield losses caused by pests, pathogens, and weeds, but its overuse causes serious environmental pollution. They are persistent in the environment and biomagnified through the food chain resulting a serious hazard for humankind. Bioremediation by microbes to degrade the pesticides *in situ* is a useful technology. This review mainly summarized the fungi associated with biodegradation of chemical pesticides and their application in the soil and water bioremediation. The future studies on this field were also prospected.

1
2
3
4
5
6
7 31 Keywords: Pesticides, Agrochemicals, Antibiotics, Sustainable bioremediation, Fungi,
8 32 Synthetic microbial community, Environmental risk assessment.
9
10 33
11
12

13 34 **Introduction**

14 35 Because of their unique functions, fungi are involved with important ecosystem services for human
15 36 well-being. Among others, fungi account for provisional services also through the activity of
16 37 transforming and detoxifying pollutants. For this reason, learning from nature, they represent an
18 38 effective toolbox for a sustainable bioremediation of pesticides in soil and water. Many researches
19 39 have unfolded the untapped potential of fungi, given that recent years have witnessed very
20 40 interesting developments regarding use of fungi not only to improve the environmental quality but
22 41 also human health (e.g. Gargano et al. 2017).
23 42 Pesticides are a diverse group of inorganic and organic chemicals like herbicides, insecticides,
24 43 nematicides, fungicides, antibiotics and soil fumigants, all belonging to the so-called organic
26 44 agrochemicals (OACs) (Verger and Boobis 2013; Verma et al. 2014). In agriculture, pesticides aim
27 45 to enhance crop yield and quality, and to maximize economic returns by prevention of pest or weed
29 46 attack. They are bioactive, toxic substances, capable of influencing, directly or indirectly, soil
30 47 fertility and health as well as agroecosystem quality (Pinto et al. 2012; Verma et al. 2014). Given
32 48 that belowground biodiversity is closely linked to land management, agricultural intensification
33 49 causes many pressures that leads to loss of biodiversity. Consequently, soil pollution is one of the
34 50 main threats related to the decline of taxonomic and functional biodiversity, and of agricultural soils
36 51 sustainability (Harms et al. 2017). Most of the pesticides emission (99 %) in Europe is associated to
37 52 agricultural practices whereas industrial and urban sources as the manufacturing of pesticides or the
38 53 at-home use of insecticides have a minor impact (EEA 2016). Thus, the extensive and massive
40 54 use of pesticides in agriculture activities has serious impacts on the environment, compromising soil
41 55 and water quality (Pinto et al. 2012; Zhang et al. 2015; Pinto et al. 2016). Besides,
43 56 In addition to pesticides, large quantities of antibiotics are added to agricultural fields
44 57 worldwide through the application of wastewater, manures and biosolids, resulting in antibiotic
45 58 contamination and elevated environmental risks (Jechalke et al. 2014; Zhang et al. 2015; Pan and
47 59 Chu 2016). A clear correlation between agriculture and water contamination was observed in Mar
48 60 Chiquita lake (Argentina), where high amount of endosulfan residues were detected soon after
49 61 application and post-application periods (Ballesteros et al. 2014). The presence of the fungicide
51 62 thiﬂuzamide in paddy water of rice fields in China was maximal after the application, and variation
52 63 during time was associated to the dilution effect of rainfalls in the area (Wei et al. 2015). Preventive

Formatted: English (U.S.)

measures to mitigate the impact of agriculture on the environment are required, taking into account both the use of safety pesticides and the optimization of farmer procedures. Aravinna et al. (2017) found that most of the 32 studied pesticides leached rice field following specific pathways. Since direct run off and erosion from soil were the main vehicles of dispersion, authors suggested alternative strategies (high resident time of pesticides, holding ponds of rice drainage water, delayed filling of paddies after pesticide application and use less mobile compounds) to reduce the movement of the pesticides.

The intensive use of these ~~organic agrochemicals (OACs)~~ has posed risks to both wild lives and human health. Over 98% of sprayed insecticides and 95% of herbicides reach a destination other than their target species, through air, water and soil (Miller 2004). Around 30% of pesticides marketed in developing countries do not meet internationally accepted quality standards, posing a serious threat to human health and environment (Popp et al. 2013). They are persistent in the environment and biomagnified through the food chain. Therefore, it has been estimated that millions of agricultural workers worldwide experience unintentional pesticide poisonings each year. The correlation between long-term exposures to pesticides in occupational settings is known but recently also non-occupational exposures have been associated to an elevated rate of chronic diseases (Parrón et al. 2014).

Varieties and consumption of pesticides worldwide are dramatically increasing, ~~up to, but literature reports conflicting data on overall use (2–4 million ton for year). 4-fold higher than 40 years ago (Mnif et al. 2011).~~ According to De et al. (2014), about 45 % is used by Europe, 25 % by USA, and 25 % in the rest of the world. The main pesticide consumer is Spain (around 79,000 ton of active ingredients sold between 2011 and 2014), followed by France (~ 75,000), Italy (~ 64,000), Germany (~ 46,000) and United Kingdom (~ 23,000) (Eurostat 2016). The United States is also a large consumer of pesticides, applying usually applies over 1 billion pounds annually (Alavanja 2009) with dramatical consequences for human beings and environment (Carvalho 2017). Overall, herbicides account for 47.5 %, insecticides for 29.5 %, fungicides for 17.5 % and others account for 5.5 %.

~~On the contrary, according to~~ According to other authors (Huang McBeath and McBeath 2010), China is the world's largest pesticide user, with an output of pesticide around 3.7 million ton (National Bureau of Statistics of China - <http://data.stats.gov.cn>), and a consumption volume of about 1.8 million ton in 2014. The average amount of pesticides used per hectare in China is roughly 1.5- to 4-fold higher than the world average (Qiu 2011), thus resulting in the contamination of water bodies in the receiving areas and disturbance of ecological equilibrium (Hui et al. 2003).

~~Overall, herbicides account for 47.5 %, insecticides for 29.5 %, fungicides for 17.5 % and others~~

1
2
3
4
5
6
7 98 account for 5.5 % (De et al. 2014). More than 350 insecticides, herbicides, microbicides,
8 99 nematocides and other pesticides are reported to be used (Huang McBeath and McBeath 2010). ~~The~~
9 ~~average amount of pesticides used per hectare in China is roughly 1.5 to 4 fold higher than the~~
10 ~~world average (Qiu 2011), thus resulting in the contamination of water bodies in the receiving areas~~
11 ~~and disturbance of ecological equilibrium (Hui et al. 2003).~~
12 ~~The United States is also a large the next largest consumer of pesticides, applying over 1 billion~~
13 ~~pounds annually (Alavanja 2009) with dramatical consequences for human beings and environment~~
14 ~~(Carvalho 2017).~~
15 ~~As regards Europe, according to the Eurostat (2016), the main pesticide consumer is Spain (around~~
16 ~~79,000 ton of active ingredients sold between 2011 and 2014), followed by France (~ 75,000), Italy~~
17 ~~(~ 64,000), Germany (~ 46,000) and United Kingdom (~ 23,000).~~
18 106 The adverse effects of ~~pesticide and antibiotics~~ OACs pollution have been concerned for a long time
19 107 and many highly toxic and persistent pesticides have been banned worldwide. Although relatively
20 108 safer pesticides have been developed and replaced the highly toxic ones, environmental pollution
21 109 resulted by the long-term application of pesticides is far from being solved. ~~Still now~~
22 112 ~~Obsolete~~ ~~obsolete~~ pesticides ~~widely used in agriculture in the past, still~~ represent a threat to
23 113 environment, biodiversity, and human health for the region of Southeast Europe and their
24 114 environmental and human risk need to be assessed in order to mitigate their current risk. Many
25 115 organochlorines, organophosphates and pyrethroids have been banned but this did not solved the
26 116 problem yet (Aravinna et al. 2017). In Argentina, the use of hexachlorocyclohexane pesticides have
27 117 been limited from the late '90 and definitely banned in 2011, but this did not prevent to find
28 118 concentration of lindane during recent samplings. Although the maximum level of lindane in saline
29 119 water was fixed at 4 ng/l, in 2014 lindane exceeded this value of more than 5-fold (Ballesteros et al.
30 120 2014). Although the use of organo-chlorine pesticide has been banned for over 20 years, they can
31 121 still be found in the water and the sediment of main drainage area in China (Nakata et al. 2005; Xue
32 122 et al. 2006; Zhou et al. 2006), due to run off from aged and weathered agricultural soils, or
33 123 anaerobic sediments (Zhou et al. 2006). ~~Except for~~ Besides water bodies and sediment, water, soil
34 124 and even air in many cities are polluted by OACs, including urban or suburban areas (Gong et al.
35 125 2004; Nakata et al. 2005; Yang et al. 2008).
36 126 ~~For that matter,~~ OACs pose pivotal environmental problems, due to their high reistance in the
37 127 environment and the consequent low natural attenuation. As an example, ; among them,
38 128 organochlorine pesticides ~~and their metabolites, are resistant towere poorly affected by~~
39 129 photochemical, chemical and biological ~~degradation processes for a long time as reported by and~~
40 130 more than 95% of them impacted on non-target organisms (Mrema et al. (2013). ~~The authors~~
41 131

Formatted: Font:

Formatted: Font:

Formatted: Font:

highlighted the impacts of pesticides, which become widely dispersed in the environment; it was estimated that more than 95% of applied pesticides impact non-target organisms. As a consequence, Kim et al. (2017) reported a consequence, number of routes pesticides might follow to meet human beings; the resulting multi-pathway direct and indirect exposure may affect human health. For instance during the last decade, one of the most studied issues is cancer occurrence related to pesticide exposure.

As persistent organopollutants (POPs), pesticides represent one of the major problems in both terrestrial and aquatic ecosystems. Regulatory and risk assessment procedures have to be adopted against those compounds that could be categorized as POPs/OACs. Since early '90, European Union started taking care of the problem. Driven from the carcinogenicity of pesticides, Directive 91/414/EEC aimed to control the authorization for pesticides marketing within the EU. The particular attention given to pesticides is because recent studies confirmed that even low dose and chronic exposure might trigger adverse effects on wildlife and humans (EEA 2005). Being groundwater the primary source of drinking waters, both the Groundwater Directive 2006/118/EC and the Drinking Water Directive 98/83/EC deal with pesticides maximal exposure concentrations: 0.1 µg/l of a single pesticide and 0.5 µg/l of total pesticides load. The protracted exposure to low amount of pesticides cannot be underestimated because critical exposure levels can be chronically reached. A Risk assessment has to consider the possible source of contamination but also the direct and indirect multifaceted pathways of contact with human beings. Kim et al. (2017) reported a number of routes pesticides might follow to meet human beings; the resulting multi-pathway direct and indirect exposure may affect human health. Most of the pesticides emission (99 %) in the environment in Europe is associated to agricultural practices whereas industrial and urban sources as the manufacturing of pesticides or the at home use of insecticides have a minor impact (EEA 2016). Kim et al. (2017) reported a number of routes pesticides might follow to meet human beings; the resulting multi-pathway direct and indirect exposure may affect human health.

Point discharges of pesticides used in agriculture may occur and are mainly associated to accidental causes as spillage, inappropriate storage and disposal, etc. Most of pesticides instead reach surface waters, through direct surface run-off or by leaching to groundwater and then subsequently follow different transport pathways. Once entered in the aquatic system, they could ultimately contaminate water for human consumption.

A clear correlation between agriculture and water contamination was observed in Mar Chiquita lake (Argentina), since high amount of endosulfan residues were detected soon after application and post-application periods (Ballesteros et al. 2014). The presence of the fungicide thifluzamide in paddy water of rice fields in China was maximal after the application, and variation during time was

1
2
3
4
5
6
7 166 associated to the dilution effect of rainfalls in the area (Wei et al. 2015). Preventive measures to
8 167 mitigate the impact of agriculture on the environment are required, taking into account both the use
9
10 168 of safety pesticides and the optimization of farmer procedures. Aravinna et al. (2017) found that
11 169 most of the 32 studied pesticides leached rice field following specific pathways. Since direct run off
12 170 and erosion from soil were the main vehicles of dispersion, authors suggested alternative strategies
13
14 171 (high resident time of pesticides, holding ponds of rice drainage water, delayed filling of paddies
15 172 after pesticide application and use less mobile compounds) to reduce the movement of the
16
17 173 pesticides.
18 174 Experimental evidences of advances in natural restoration processes highlight that time is our
19 175 friend, since the abandonment of disturbed/polluted agricultural land for long time could reduce
20
21 176 their contaminatin. In fact, at a global scale, one of the most frequently used strategies is long term
22 177 remediation, which is represented by the abandonment of disturbed/polluted agricultural land
23 178 (Kardol and Wardle 2010). Studies by Morrin et al. (2017) reported that nature restoration on ex-
24
25 179 arable land resulted in increased connettance of soil biota's networks, as restoration progresses.
26 180 Such results confirm that the functions played by the soil biota provide many and varied services,
27
28 181 and detoxification of pollutants and xenobiotic is one of the included primary services. In this
29 182 context, innovation is represented by the research of solutions inspired by nature, as strategy to
30 183 accelerate the natural attenuation processes in contaminated sites, optimizing bioremediation in real
31
32 184 environment. Given that OACs represent a potential risk to humans, water, ecosystems and other
33 185 receptors, fungi can play a pivotal role addressing their removal from contaminated sites and thus
34 186 mitigating environmental pollution.
35
36 187 So clean and safe water is a critical step that stands between the *status quo* and a sustainable world.
37 188 This concept is no longer idealistic and became a milestone for the United Nations, as clearly stated
38
39 189 in the World Water Development Report of 2015 (WWAP, 2015). Human lifestyle and the
40 190 increasing urbanization lead to a worsened scenario. For instance, the actual pesticides use is 4 fold
41 191 higher than 40 years ago (Mnif et al. 2011). EC compiled a watch list including, among others,
42
43 192 pharmaceuticals, pesticides and personal care products. Being groundwater the primary source of
44 193 drinking waters, both the Groundwater Directive 2006/118/EC and the Drinking Water Directive
45 194 98/83/EC deal with pesticides maximal exposure concentrations: 0.1 µg/l of a single pesticide and
46
47 195 0.5 µg/l of total pesticides load.
48 196 In this context, bioremediation has aroused as an-is a ~~usefulis~~ ~~buseful~~ technology to degrade
49
50 197 pesticides OACs by microbes (Singh 2008; Velquez-Fernndez et al. 2012), with several benefits
51 198 over landfill disposal and incineration, such as the ~~conversion of toxic wastes to~~ ~~formation of~~ non-
52 199 toxic end products, a-lower ~~costs~~ ~~cost~~ of disposal-~~(or no disposal at all)~~, reduced health and
53
54
55
56
57
58
59
60

ecological effects and long-term liabilities associated with non-destructive treatment methods, and the ability to perform the treatment *in situ* without unduly disturbing native ecosystems (Sarkar et al. 2005). ~~Therefore, there is a growing interest in developing bioremediation techniques to degrade OACs in polluted environments.~~ During the past decade, numerous microorganisms capable of degrading antibiotics and pesticides have been isolated, and detoxification processes for target pollutants have been analyzed. ~~As for many other POPs (BTEX, PHAs, PCB congeners, etc) with structural similarities with lignin, fungi~~ and especially ligninolytic fungi have been suggested as the most promising group of organisms able to transform recalcitrant compounds through a unique set of extracellular oxidative enzymes (e.g. Anastasi et al. 2013; Harms et al. 2017). Comparative genomic analysis of 49 fungi with different nutritional modes such as saprotrophic fungi, white-rot fungi (WRF), brown-rot fungi, ~~straw-soft~~ rot fungi and symbiotic fungi indicated that there is a relationship between nutrition models and the enzymes for lignocellulose degradation. Saprotrophic fungi have greater number of enzymes than symbiotic fungi, and brown-rot fungi have smaller number than ~~white-rot fungi~~WRF and ~~straw-soft~~ rot fungi (Wu et al. 2015a). This might gain some insights into how to choose fungi in OACs degradation.

~~Experimental evidences of advances in natural restoration processes highlight that time is our friend. In fact, at a global scale, one of the most frequently used strategies is long term remediation, which is represented by the abandonment of disturbed/polluted agricultural land (Kardol and Wardle 2010). Studies by Morrión et al. (2017) reported that nature restoration on ex arable land resulted in increased connectance of soil biota's networks, as restoration progresses. Such results confirm that the functions played by the soil biota provide many and varied services, and detoxification of pollutants and xenobiotic is one of the included primary services. In this context, innovation is represented by the research of solutions inspired by nature, as strategy to accelerate the natural attenuation processes in contaminated sites, optimizing bioremediation in real environment. Given that OACs represent a potential risk to humans, water, ecosystems and other receptors, fungi can play a pivotal role addressing their removal from contaminated sites and thus mitigating environmental pollution.~~

Finally yet importantly, metabolic activity of fungal or microbial consortia could produce not-known reaction products potentially with a major toxicity than parental compounds.

García-Carmona et al. (2017) highlighted the importance to carry out environmental monitoring activities ante and post operam phases, using bioassays to determine the success of the bioremediation process. Although it is substantial to assess the quality of the environment to ensure it remains free of toxic residues, most of the analytical tests available for determining the concentration of toxic chemicals do not give the biological impacts of toxicants. For this reason,

1
2
3
4
5
6
7 234 biotoxicity testing has grown steadily in recent years and is a useful tool in environmental risk
8 235 assessment (Shen et al. 2016; Prokop et al. 2016).
9
10 236 Indeed, there is a clear need to develop and define decontamination of hazardous pollutants as a
11 237 concept towards sustainable remediation through a broader uptake of principles, approaches and
12 238 tools to integrate environmental, social and economical dimension into the remediation processes
13
14 239 (Ridsdale and Noble 2016). Several organizations, academia, standardization committees are
15 240 currently assessing remediation process, evaluating the complexity of the concept of sustainability.
16 241 Several documents have been developed by many countries across Europe and at global scale,
17
18 242 addressing sustainable indicators of remediation activities (Harclerode et al. 2015).
19 243 The present review article summarizes the current state of scientific knowledge on research and
20
21 244 application of fungi as effective bioresources, considering the recent advances in understanding
22 245 their capacity to handle pesticide contamination.

23 246
24
25 247 **Bioremediation of OACs by fungi in soil system**

26 248 Large quantities of ~~OACs~~~~antibiotics~~ are being added to agricultural fields worldwide through the
27
28 249 application of wastewater, manures and biosolids, resulting in pesticide and antibiotic
29 250 contamination and elevated environmental risks in terrestrial environments (Jechalke et al. 2014;
30 251 Zhang et al. 2015; Pan and Chu 2016). The largest fraction of ~~antibiotics~~~~OACs~~ applied to soils
31
32 252 with manure or biosolids is usually retained in surface soil whereas the part added through irrigation
33 253 with wastewater can diffuse easily deep or by surface run-off. Once added to soil, ~~antibiotics~~~~OACs~~
34 254 interact with soil solid phase and are prone to microbial transformation (Hammesfahr et al. 2008;
35
36 255 Jechalke et al. 2014). In particular, veterinary antibiotics interact with soil solid phase in sorption
37 256 and desorption reactions. Sorption and desorption control not only their mobility and uptake by
38
39 257 plants but also their biotransformation and biological effects. ~~Antibiotics~~~~OACs~~ as well as
40 258 microorganisms are not distributed homogeneously in soil but are concentrated in hotspots. The
41 259 different surfaces, voids, and pores provided by soil aggregates harbor a vast amount of biological
42
43 260 diversity and chemical variability, and cause a patchy distribution of natural organic matter, oxides,
44 261 nutrients, and microorganisms on soil particle surfaces (Hammesfahr et al. 2008; Jones et al. 2012).
45 262 Sorption, sequestration, and subsequent release of ~~antibiotics~~~~OACs~~ likely also occur at and from
46
47 263 hotspots, and little is known about the behavior of ~~antibiotics~~~~OACs~~ at environmentally relevant
48 264 concentrations in agricultural soil.
49
50 265 Recently, many studies highlighted the fungal capability to transform and degrade recalcitrant
51 266 OACs. In particular, one of a promising group is the ligninolytic fungi that possess a unique set of
52 267 extracellular enzymes suitable to degrade lignin and are able to transform recalcitrant compounds.
53

~~In particular, a promising group of fungi that are able to transform recalcitrant compounds and possess a unique set of extracellular ligninolytic enzymes are ligninolytic fungi~~ (Čvančarová et al. 2015). ~~(Supplemental data Table I; Table I References) 2015~~. Nguyen et al. (2014) reported the removal of diverse trace organic contaminants (i. e. trichloroethyl chloroformate (TrOC), phenolic and non phenolic, pharmaceuticals, pesticides, steroid hormones, industrial precursors and products, phytoestrogens) by live (biosorption + biodegradation), intracellular enzyme-inhibited, and chemically inactivated (biosorption only) whole-cell preparations and the fungal extracellular enzyme extract (predominantly laccases) from *Trametes versicolor* (strain ATCC 7731). They showed how non-phenolic TrOC were readily biodegraded while the removal of hydrophilic TrOC was negligible. The whole-cell culture showed considerably higher degradation of the major compounds, indicating the importance of biosorption and subsequent degradation by intracellular and/or mycelium associated enzymes. However, studies that examined both adsorption and degradation of antibiotics in agricultural soil are too few, with most of them using unrealistically high concentrations (in mg/kg levels) to overcome limitations in measurement. In addition, no model has been developed for speculating the adsorption and degradation of different types of antibiotics in agricultural soil and the environmental risks they may pose. Pan and Chu (2016) evaluated the adsorption and degradation of five antibiotics (tetracycline, sulfamethazine, norfloxacin, erythromycin, and chloramphenicol) by native microorganisms (bacteria and fungi) in non sterilized (test) and sterilized (control) agricultural soils under aerobic and anaerobic conditions. They showed that all antibiotics were susceptible to microbial degradation under aerobic conditions, and most antibiotics were degraded by more than 92% in non-sterilized soil after 28 days of incubation. For all the antibiotics, a higher initial concentration was found to slow down degradation and prolong persistence in soil. The degradation pathway of antibiotics, in fact, varied in relation to their physicochemical properties as well as the microbial activities and aeration of the recipient soil. The authors were the first to develop a model for the prediction of antibiotic persistence in soil, which was valuable for the investigation of the fate of antibiotics in the terrestrial environment.

Given the public concern for environmental pollution by OACs, there is increasing attention towards the development of biopurification systems for reducing the risk from the point source contamination of soil resources. Various treatment methods (e.g. land filling of contaminated sites, recycling, pyrolysis and incineration) have been used for the removal and remediation of these chemicals from the contaminated sites, but ~~for example~~ microbial degradation of pesticides ~~is~~ ~~results~~ the most important and effective way to remove these compounds from the environment (Hai et al. 2012; Verma et al. 2014). ~~(Supplemental data Table I; Table I References) 2014~~.

Formatted: English (U.S.)

Formatted: English (U.K.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted

1
2
3
4
5
6
7 302 Microorganisms have the ability to interact, both chemically and physically, with substances leading
8 303 to structural changes or complete degradation of the target molecule. In particular, fungi may
9
10 304 transform pesticides and other xenobiotics by introducing minor structural changes to the molecule,
11 305 producing nontoxic molecules that could be released into the soil for further degradation by
12 306 microflora (Hai et al. 2012), (Supplemental data Table I; Table I References).
13
14 307 In this context, Mir-Tutusa et al. (2014) investigated the degradation of the insecticides
15 308 imiprothrin and cypermethrin, the insecticide/nematicide carbofuran using the white-rot fungus *T.*
16 309 *versicolor*. Their experiments with fungal pellets demonstrated extensive degradation of the tested
17
18 310 agrochemicals. In vivo studies with inhibitors of cytochrome P450 revealed that this intracellular
19 311 system plays an important role in the degradation of imiprothrin and carbofuran, but not for
20
21 312 cypermethrin. The simultaneous degradation of the compounds successfully took place with
22 313 minimal inhibition of fungal activity and resulted in the reduction of the global toxicity, thus
23 314 supporting the potential use of *T. versicolor* for the treatment of several OACs.
24
25 315 To date, the number of studies investigating novel treatment techniques for the removal of
26 316 ~~pesticides—OACs~~ from contaminated agricultural soils is limited. The bacteria-dominated
27
28 317 conventional activated sludge process has been proved to be ineffective for ~~pesticide~~-removal.
29 318 While the importance of a mixed microbial community to initiate and complete ~~pesticide—OACs~~
30 319 removal in the soil environment has been convincingly demonstrated by several researchers, studies
31
32 320 concerning the removal of ~~pesticides—OACs~~ from soils have been predominantly focused on
33 321 selected bacterial or fungal species separately. Few studies have explored the bioaugmentation
34 322 synergy of fungi and bacteria (Hai et al. 2012; Zhang et al. 2015; Madrigal-Zúñiga et al. 2016).
35
36 323 Combining culture of bacteria and fungi could constitute a relevant process for the removal of toxic
37 324 and recalcitrant organic substances from contaminated agricultural soils. On-farm biopurification
38
39 325 systems represent a biotechnological approach for the mitigation of point source contamination by
40 326 ~~pesticidesOACs~~. The main component of the biopurification systems is the biomixture, which acts
41 327 as the biologically active core that accelerates the degradation of ~~OACs-pesticidesOACs~~ Madrigal-
42
43 328 Zúñiga et al. (2016) studied the employment possibility of the ligninolytic fungus *T. versicolor* in
44 329 the bioaugmentation of compost- (GCS) and peat-based (GTS) biomixtures for the removal of the
45 330 insecticide-nematicide carbofuran (CFN). The CFN transformation products were detected at the
46
47 331 moment of CFN application, but their concentration continuously decreased to complete removal in
48 332 both biomixtures. Mineralization of ¹⁴C radiolabeled CFN was faster in GTS than in GCS. The
49
50 333 authors demonstrated the complete elimination of toxicity in the matrices after 48 days. Overall data
51 334 suggested that the bioaugmentation improved the performance of the GTS rather than the GCS
52 335 biomixture.
53
54
55
56
57
58
59
60

Formatted: Font: Times New Roman, 12 pt, English (U.K.)

Moreover, Pinto et al. (2016) studied the potential use of different substrates in biomixtures as cork, cork and straw, coat pine and LECA (Light Expanded Clay Aggregates) on the degradation of terbuthylazine, difenoconazole, diflufenican and pendimethalin pesticides. Bioaugmentation strategies using the WRF *Lentinula edodes* inoculated into the CBX was also assessed. The results obtained from this study clearly demonstrated the relevance of using natural biosorbents as cork residues to increase the capacity of pesticide dissipation in biomixtures for establishing biobeds. Furthermore, higher degradation of all the pesticides was achieved by the use of bioaugmented biomixtures. Indeed, biomixtures inoculated with *L. edodes* EL1 were able to mineralize the selected xenobiotics, revealing that this WRF might be a suitable fungus for being used as inoculum sources in on-farm sustainable biopurification systems, in order to increase its degradation efficiency.

Fungi isolated from biomixture represents a biological source of potentially active bioremediation agents; the adaptation skills developed by these microorganisms could make the difference for OACs removal [\(Supplemental data Table I: Table I References\)](#). This challenging strategy was assessed by Pinto et al. (2012), who isolated fungi from a loamy sand soil and a biomixture contaminated with terbuthylazine, difenoconazole and pendimethalin. The capability of degrading xenobiotics by autochthonous fungi (*Penicillium brevicompactum* and *Lecanicillium saksenae*) was compared with allochthonous strains taken from a Culture Collection (*Fusarium oxysporum*, *Aspergillus oryzae* and *L. edodes*). The major biodegradation yield was reached with *P. brevicompactum*: its higher ability to metabolize terbuthylazine was presumably acquired through chronic exposure to contamination with the herbicide.

Bioremediation of OACs by fungi in aquatic ecosystem

Many OACs are common contaminant of freshwater due to their high water solubility associated to a low soil adsorption, and their high stability that assure them a long half-life. ~~These properties explain the recurring evidences of pesticides found in real water samples.~~ The contamination is not heterogeneously distributed along watercourses [as evidenced in several studies where and extensive studies are necessary.](#) ~~These properties explain the recurring evidences of pesticides were recurringly found in real water samples.~~ For instance, an accurate survey took into consideration 23 European countries with more than 160 water samplings studying mainly pharmaceuticals, pesticides and ~~known-recognised~~ endocrine ~~disrupting chemicals~~ [disrupting chemicals](#) (Loos et al. 2010). Among the 59 compounds under study, the most frequently detected compound was 1 insecticide (DEET), and 7 pesticides (chloridazon-desphenyl, DMS, desethylatrazine, chloridazon-methyl-desphenyl, bentazone,

Formatted: English (U.S.)

Formatted: English (U.S.)

desethylterbutylazine, dichlorprop) exceeded the European threshold of 0.1 µg/l. ~~On the whole,~~
Overall, 29% of the water samples could not be considered safe (Loos et al. 2010) accordingly to
this value. ~~Similarly~~ Similarly, in US, 18 states were monitored, focusing the attention of 65 organic
contaminants: along with plasticizers and detergent metabolites, 66% of the total pollutants load
was ascribable to insect repellent (Barnes et al. 2008).

The extent of the freshwaters contamination and the actual risk for human life depend on several
factors concerning the hydrogeological characteristics of the soil, the weather conditions and the
chemical-physical properties of the ~~pesticide~~OACs. The environmental fate of a certain compound
is a critical issue in which the water/soil surface is the first barrier. For instance, the sorption
kinetics of three widely used pesticides (simazine, imidacloprid, and boscalid) have been correlated
to the soil organic carbon content and the hydrophobicity of the pesticide, ultimately affecting their
soil retention behavior and the actual bioavailability in waters (Salvestrini et al. 2014). The flow of
the leaching into surface waters is also a matter of season, in which opposite phenomena draw a
complex scenario to be predict. A rainy period could cause a massive run-off of ~~OACs~~the pesticides
from the soil contaminating the receiving basin (Sandin et al. 2018), ~~but during dry season, the high~~
~~load of contaminants could be associated to evaporation and low water flow.~~ Besides the detection
of high levels of ~~pesticides~~pesticideOACs is not exclusively coincident to their recent and massive
use, ~~but it is ascribable to their~~ ~~Due to their~~ persistency, ~~their slow natural degradation,~~ their
~~accumulation and~~ the various diffusion pathways, ~~they~~ (Aguilar et al. 2017), ~~They~~ could then
tread long distances in surface or groundwater waters and the contamination can last for several
decades (Ballesteros et al. 2014; Aravinna et al. 2017).

The so-called ecological services could help to contain the ~~pesticides~~pesticideOACs diffusion.
Adapted microflora (fungi, Gram-positive and ~~negative~~ bacteria, actinobacteria, and sulfate-
reducing bacteria) to the soil environmental conditions may reduce the pesticides released in
groundwater sources (Mattsson et al. 2015). Several factors as soil composition, temperature,
aeration due to soil weaving and depth influence the autochthonous microbial community activity;
if this balance fails, ~~pesticides~~pesticideOACs are free to move among different ecological niches
(i.e. sediment and water), ~~and~~ alter their functioning, ~~and ultimately directly affecting their animal~~
~~inhabitants.~~ For instance, sSignificant ecological risk was associated to the presence of the
insecticide fipronil and its metabolites in three water ponds: concentration up to 200 ng/l affected
the proper development of larval insects and crustaceans (Wu et al. 2015b). Evidences of the
pesticides toxicity against fish has been already reported, demonstrating their interference with
different metabolic pathways (Odukkathil and Vasudevan 2013; Ballesteros et al. 2014; Guerreño et
al. 2016).

The preservation of water quality is a priority but OACs removal could not be based only on natural attenuation. Water treatment plants (WTPs) are the major barrages where OACs should be removed. Not being specifically designed for micropollutants removal, they are often only partially effective, with a strong impact on the receiving ecosystem. Pesticides as atrazine, fluconazole, tebuconazole, diazinon and diuron are particularly resistant to commonly in use treatments (Köck-Schulmeyer et al. 2013; Luo et al. 2014). A number of evidences confirmed the presence of OACs in WTPs effluents at toxicologically and estrogenically relevant concentration, becoming one of the most effecting source of contamination (Bicchi et al. 2009; Campo et al. 2013; Jarošová et al. 2014).

Particular attention has been given to advanced biological oxidation. Novel cost-effective and eco-friendly processes based on fungi are an attractive option. [They-Fungi](#) are well-known for to their physiological adaption skills, including the natural activation of tolerance mechanisms against pesticides (Talk et al. 2016). [In comparison with bacteria, Some reports already demonstrated that in comparison with bacteria,](#) fungi can better tolerate the presence of organic contaminants. Although the insecticide endosulfan inhibited both fungi and bacteria, bacterial community structure significantly changed already at 0.1 mg/kg while modifications on the fungal community structures required 1 mg/kg of pollutant (Zhang et al. 2015). Linuron reduced bacterial count, and especially total bacteria, N₂-fixing bacteria and nitrifiers, but not fungal numbers (Cycoń et al. 2010).

The importance of the isolation origin of fungi is out of discussion. Strains isolated from contaminated niches could have indeed developed specific adaptation skills due to the chronically exposure over time. Carles et al. (2017) demonstrated that the aquatic microflora associated to submerged leaves exposed to nicosulfuron is more efficient in its degradation than communities belonging to a less polluted site. The authors indicated fungi as the main constituents of this active microflora and as responsible of the herbicide degradation. In literature, several fungi isolated from contaminated areas or WTPs have been identified as degraders of nicosulfuron, diuron, isoproturon, glyphosate, chlorpyrifos, chlorfenvinphos and atrazine (Song et al. 2013; Carranza et al. 2014; Oliveira et al. 2015).

Exploiting this oxidative cascade, fungi may transform a broad range of recalcitrant organic compounds, including OACs (Gao et al. 2010). A number of fungi are [pesticidespesticideOACs](#) degraders, mostly belonging to Basidiomycetes as *Trametes*, *Pleurotus*, *Phlebia*, *Cerrena*, *Coriolopsis*, etc. have been already investigated (Koroleva et al. 2002; Marco-Urrea et al. 2009; Xiao et al. 2011; Ulčnik et al. 2013; Chan-Cupul et al. 2014; Ceci et al. 2015) ([Table-II2. Supplementary Materials](#)). Several ~~classes of~~ pesticides as lindane, atrazine, diuron,

1
2
3
4
5
6
7 438 terbuthylazine, metalaxyl, DDT, gamma-hexachlorocyclohexane (g-HCH), dieldrin, aldrin,
8 439 heptachlor, chlordane, lindane, mirex, etc. were effectively transformed by fungal treatment- [based](#)
9 [on mycelium or enzymes \(Table II2, Supplementary Materials\).](#)
10 440
11 441 A bioremediation approach based on fungi may involve both biosorption and biodegradation
12 442 processes; the latter one combines biosorption where the molecule binds to the fungal wall, and
13 443 bioaccumulation with the pollutant being transported inside the cell in contact with intracellular
14 444 enzymes (Kulshreshtha et al. 2014). Concentration of insecticide lindane decreased during time in
15 445 the presence of two WRF (*T. versicolor* and *Pleurotus ostreatus*) and one brown-rot fungus
16 446 (*Gloeophyllum trabeum*), but the lack of any change in the chromatogram profile indicated the main
17 447 involvement of a fast adsorption process (Ulčnik et al. 2013). However, this phenomenon is often
18 448 strains dependent, and especially related to metabolic differences between Ascomycetes and
19 449 Basidiomycetes. Belonging to brown-rot fungi, *G. trabeum* lacks the ligninolytic enzymes,
20 450 responsible for lignin degradation and likely for OACs as well: adsorption onto fungal mycelium
21 451 was mainly involved for removal of endosulfan. On the contrary, the WRF actively degraded
22 452 producing endosulfan sulphate via oxidative pathways (Ulčnik et al. 2013). Although biosorption is
23 453 a phenomenon that could be [not](#) ignored, it is often secondary or at least negligible respect to
24 454 biodegradation (Carles et al. 2017). For instance, the removal of clofibric acid associated to heat-
25 455 killed mycelium was less than 10 %, but more than 97 % in the presence of active *T. versicolor*
26 456 (Marco-Urrea et al. 2009).
27 457 Fungi have developed a specific mechanism that employs few enzymes and molecules with high
28 458 oxidizing power, physiologically aimed to transform ligninocellulose structure. The same enzymatic
29 459 pathway may play a pivotal role in transforming other aromatic molecules. White-rot fungi usually
30 460 involve ligninocellulosic extracellular enzymes as peroxidases (EC 1.11.1.x) and laccases (EC
31 461 1.10.3.2). The involvement of redox enzymes in the fungal-mediated oxidation is confirmed by the
32 462 direct induction of enzyme production due to the presence of [pesticides.pesticideOACs](#). The fungus
33 463 *T. versicolor* responded to 17 pesticides by increasing laccases production in comparison with the
34 464 control: particular attention was given to transformation products of the herbicides diquat and
35 465 monuron, capable of increasing the activity of 10- and 17-fold, respectively (Mougin et al. 2002).
36 466 Laccase production of *Pycnoporus sanguineus*, *Trametes maxima*, *Pleurotus* spp1, *Pleurotus* spp2,
37 467 *Cymatoderma elegans*, *Daedalea elegans* was stimulated by the presence of atrazine even at high
38 468 concentration 3750 mg/l. Likewise the pesticide positively affected the manganese peroxidase
39 469 activity of *Pleurotus* spp1 and *C. elegans* (Chan-Cupul et al. 2014). Oxidoreductases stimulation
40 470 was also observed with picloram (Maciel et al. 2013), bentazon (Da Silva Coelho et al. 2010),
41 471 carbofuran (Mir-Tutusaus et al. 2014).
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Although these oxidoreductases are probably the most known enzymes for aromatic compounds degradation, alternative pathways can be promoted by the presence of [pesticidespesticideOACs](#). Two clones (laccase positive and negative producers) of *Mycelia sterilia* were used to treat atrazine (20 µg/ml): even though one clone was defective for laccase production, comparable transformation yields (70-80%) were reached indicating their minor role in the degradation process (Vasil'Chenko et al. 2002). This behavior is commonly found in brown-rot fungi that may trigger both on nonenzymatic and enzymatic mechanisms, i.e. Fenton mechanism or cellobiose dehydrogenase (CDH) reactions (Fan and Song 2014). The degradation of atrazine (20 µg/l) by an unidentified mycelial fungus was associated to the presence in the liquid medium of OH radicals and CDH. Moreover, the CDH secretion was induced by the presence of the herbicide itself (Khromonygina et al. 2004). In addition, some fungi could associate extracellular oxidoreductases with intracellular enzymes [such](#) as the cytochrome P450 system (cyt450). In the effort to better characterize the degradation skills of *T. versicolor*, cyt450 inhibitors were used: fungal performances against clofibric acid and fipronil decreased (Marco-Urrea et al. 2009; Wolfand et al. 2016). Mori et al. (2017) suggested that cyt450 of *Phanerochaete sordida* is involved in the first reduction of the clothianidin N-nitro group but the enzymes responsible of the further urea derivatives production are unknown.

The intra- and interspecies variability has long been recognized and found [confirmation](#) [alsoconfirmation](#) for [pesticidespesticideOACs](#) treatment. Literature data about a certain specie could not be taken for granted and the set-up of a preliminary screening is often required. Despite *Phanerochaete chrysosporium* is often indicated as fungal model for organic degradation including pesticides (Wang et al. 2014), it was almost ineffective against clofibric acid. Among five Basidiomycetes, only *T. versicolor* extensively degraded the herbicide (Marco-Urrea et al. 2009). Alvarenga et al. (2014) treated methyl parathion with several fungi, including 3 *Aspergillus sydowii*. Based on the growth capability in the presence of the pesticide, [only](#) the isolate *A. sydowii* CBMAI 935 was selected for further studies. It indeed grew almost 4-fold more than the other *A. sydowii*. The bioremediation potential is often substrate targeted, and the choice of fungus cannot be taken for granted. For instance, the exact same isolate (*A. sydowii* CBMAI 935) that totally converted methyl parathion (Alvarenga et al. 2014) was not the best performing one against the insecticide esfenvalerate. Among 6 fungi, *Microsphaeropsis* sp. *Acremonium* sp. and *Westerdykella* sp. gave better results than the *Aspergillus* strain (Birolli et al. 2016).

Although the majority of these strains are effective in [pesticidespesticideOACs](#) removal in model solution, only few researchers have made a step forward, assessing the bioremediation potential of contaminated waters. The acquired information using model solutions (single-compound solution,

high concentration, no interfering molecules, etc.) is the unique way to acquire information about the degradation pathway (Masaphy et al. 1993; Birolli et al. 2016), but is less predictive of the fungal performances on real environmental water samples. Each wastewater has its own critical issues, making difficult to predict the fungal behavior. Some data highlighted the robustness of a fungal system, although this needs detailed investigation case-by-case. A partially diluted leachate showed to disturb the growth of *T. versicolor* and *Stereum hirsutum*, but this did not prevent them to totally degrade linuron and dimethoate at 10 mg/l. As regards dimethoate, the presence of adsorbents enhance the final process yields (from 50% to 97%), combining and exalting the action of adsorption and biodegradation processes (Castellana and Loffredo 2014). The immobilization of *Bjerkandera adusta* and *Irpex lacteus* on coffee grounds, almond shells, a biochar favored the removal of non-phenolic herbicides as fenuron and carbaryl from a municipal landfill leachate (Loffredo et al. 2016). Surface waters, ground waters or municipal wastewaters represent a very unique environment, characterized by extreme chemical and physical conditions, the presence of heterogeneous and variable micropollutant mixture and an active autochthonous microflora. When inoculated in real surface water, a fungal consortium (*Aspergillus fumigatus*, *Aspergillus terreus*, *Cladosporium tenuissimum*, *Cladosporium cladosporioides*, *Fusarium begoniae*, *Penicillium citrinum*, *Penicillium melanoconidium* and *Phoma glomerata*) was not stable in time due probably to the presence of toxic pesticides and the interaction with the natural microbial population: *P. citrinum*, *A. fumigatus* and *A. terreus* were the most robust to the environmental conditions and actually capable of degrading the spiked chlorfenvinphos (Oliveira et al. 2015).

The set-up of active microbial consortia is an intriguing solution to strengthen and combine the bioremediation potential of different organisms. Interestingly the combination of *Bacillus subtilis* and *A. niger* led to higher degradation rate of nicosulfuron than those obtained by using singly each strain (Lu et al. 2012). The biodegradation of aldicarb, atrazine and alachlor by *Coriolus versicolor* was strongly enhanced by the combination with activated sludge. Along with modifications in the fungal morphology, when the bacterial-fungal consortium was established, the bio-absorbed fraction of especially atrazine was reduced: over 98% of atrazine was removed by degradation processes in two weeks (Hai et al. 2012).

The fate of the treated ~~pesticides~~pesticideOACs is major issue that has to be carefully considered. The residual toxicity is a critical issue. Interestingly fenuron and carbaryl (~~up to 70%~~) degradation (up to 70%) catalyzed by *B. adusta* and *I. lacteus* led to significant abatement of the phytotoxicity (rapeseed and flax tests) (Loffredo et al. 2016). Mori et al. (2017) followed the neurotoxicity of clothianidin and its main metabolite produced by *P. sordida* treatment: the insecticide altered the cell viability of the neuronal cell line, but the metabolite was no longer neurotoxic.

Despite the well demonstrated properties, the application of whole cell system has some drawbacks including the fact that a living organism needs controlled growing conditions, in terms of nutrients, pH, O₂, etc. (Majeau et al. 2010). The addition of synthetic nutrients can strengthen fungal mycelium activity, but it should be carefully balanced for a further scale-up of the process. The fact that *T. versicolor* need 1% of glucose as carbon source to degrade atrazine would ultimately interfere with its potential use in real WTPs (Khromonygina et al. 2004). Likewise several fungi as *A. niger* and *Dacryopinax elegans*, etc. required both easily available carbon and nitrogen sources to efficiently act against nicosulfuron and diuron, respectively (Lu et al. 2012; Arakaki et al. 2013). Particular attention should be instead given to those fungi as *A. sydowii* and *Penicillium decaturense* that maintained the same performances without glucose addition, indicating the potential of using methyl parathion or triclosan as sole carbon source (Alvarenga et al. 2014; Tian et al. 2016).

A promising alternative could be given by the direct use of fungal enzymes, capable of catalyzing strong and fast oxidation reactions, with less technical drawbacks in comparison with fungal cultures. The potential of enzymes-based methods has been worldwide recognized; the Swiss Industrial Biocatalysis Consortium defined oxidative enzymes as the biocatalysts displaying the highest development potential in the next decades (Meyer and Munch 2005). Great importance is given to the discovery of novel enzymes with wide substrate specificity, stable and applicable to industrial uses. A number of articles have reported the ability of fungal enzymes to degrade pesticides, pesticide OACs. The potential of laccase-mediator systems have been assessed for the degradation of isoproturon (Margot et al. 2015), imiprothrin (Mir-Tutusaus et al. 2014), chloroxuron (Palvannan et al. 2014), isoproturon (Zeng et al. 2017), atrazine (Chan-Cupul et al. 2016). Laccases cannot be consider a novelty, as instead a phytase of *A. niger* capable of degrading organophosphorus pesticides (Shah et al. 2017) or a cellulase of *Trichoderma longibrachiatum* active against dicofol (Wang et al. 2015). Particular attention should be given to the use of crude enzyme extracts of ligninolytic enzymes with a minor economic impact on the process than purified enzymes (Matute et al. 2012; Kaur et al. 2016). A crude extract of *Trametes pubescens* laccases degraded up to 19 compounds in model solution and confirmed its ~~potential also~~ potential- with a real municipal wastewater where the presence of suspended particles, colloids, solvents and xenobiotics as well as autochthonous microorganisms posed a strong environmental pressure. The transformation of all the detected compounds determined also a strong reduction of the estrogenicity of the water sample (Spina et al. 2015).

Application of synthetic microbial community on bioremediation

Formatted: Level 1

1
2
3
4
5
6
7 574 Bioremediation is crucial way to eliminate the OACs pollution in agriculture ecosystem. However,
8 575 many factors effect bioremediation efficiency for pesticide pollution, such as microbes applied,
9 576 treatment sites, rhizosphere effects, soil chemical and physical properties (Zhou and Hua 2004). The
10 577 practice in the bioremediation of soil or water pollution often cannot reach expected results because
11 578 the target contaminant could not be degraded completely in most cases, and sometimes intermediate
12 579 products were occurred with more toxin than original pesticides. Long-term application of various
13 580 pesticides resulted in the pollution of more than one type of chemical compounds, which is hard to
14 581 be degraded by a sole microbe. Thus, attention has been shifted to synthetic systems based on
15 582 communication between cells, rather than individual isolated cell functionality (Biliouris et al.
16 583 2012). A promising way to overcome the difficulties is to create artificial synthetic microbial
17 584 communities that contain several microbes to retain the key features of their natural counterparts
18 585 (Großkopf and Soyer 2014).

19 586 Synthetic microbial community is a collective term that is created by a bottom-up approach where
20 587 two or more defined microbial populations are assembled in a well-characterized and controlled
21 588 environment (De Roy et al. 2014). In synthetic communities, mixed populations can perform
22 589 complex tasks, although in changing environmental conditions ~~to be robust to changes in~~
23 590 ~~environment~~ (Brenner et al. 2008). There are several potential advantages of synthetic community
24 591 compared to monocultures or natural community: 1) the species in a synthetic community are
25 592 identified and the community structure is relatively simple and controllable, while the natural
26 593 community is mixed up by many microorganisms with unknown functions; 2) synthetic community
27 594 can perform more complicated functions than individual organism because members of microbial
28 595 consortia communicate and differentiate (Brenner et al. 2008); 3) synthetic community can be more
29 596 robust to environmental fluctuations because communities might be more capable of better resisting
30 597 invasion by other species and weather periods of nutrient limitation compared with monocultures
31 598 (Brenner et al. 2008); 4) synthetic community might be described through mathematical models
32 599 more easily than natural systems, and they can be used to develop and validate models of more
33 600 complex systems (Liu et al. 2017).

34 601 To develop a cooperative and steady-state community that is performing a desirable
35 602 biotechnological function, Liu et al. (2017) concluded three design principles for the construction of
36 603 synthetic community. Firstly, safety should be prioritized by beginning with innocuous or
37 604 commensal organisms (Brenner et al. 2008). Secondly, the community can converse a low-cost
38 605 and/or recalcitrant waste material into a biotechnologically relevant product, partial or de-novo
39 606 biosynthesize a compound via heterologous metabolic pathways, or bioconverse toxic substrates or
40 607 products in a toxic milieu ~~process with toxic substrates or products or substrate conversion in a~~
41 608
42 609
43 610
44 611
45 612
46 613
47 614
48 615
49 616
50 617
51 618
52 619
53 620
54
55
56
57
58
59
60

Formatted

~~toxic milieu~~ (Jagmann and Philipp 2014). Thirdly, the bioremediation process should be optimized and regularly controlled based on the knowledge of stability and division of different microorganisms (Liu et al. 2017).

Field Code Changed

Bioremediation of polluted soils and water is one application field of synthetic microbial community. As the complex structure of some pollutants, the effect of adding synthetic microbial community is much higher than single microorganism, such as the biodegradation of pesticides diuron. The herbicide diuron is used for control of broad-leaved weeds on agricultural land. Several fungal-bacterial consortia were investigated by combining three different diuron-degrading bacteria and two fungal strains. The fastest mineralization of diuron was obtained by the three member consortium (*Mortierella* LEJ702, *Variovorax* SRS16, and *Arthrobacter globiformis* D47) as measured by evolved $^{14}\text{CO}_2$, mineralizing about 32 % of the added diuron within 54 days, whereas the single strains or other consortia reached no more than 10% mineralization. In addition, the production of diuron metabolites by consortium was minimal. This may be due to cooperative catabolism, where the first organism transforms the pollutant to products that are then used by other organisms. In addition, fungal hyphae may function as transport vectors for bacteria, thereby facilitating a more effective spreading of degrader organisms in the soil (Ellegaard-Jensen et al. 2014).

Similarly, a fungal-bacterial consortium consisting of *Mortierella* sp. LEJ702 and the 2,6-dichlorobenzamide (BAM)-degrading *Aminobacter* sp. MSH1 reached a more rapid mineralisation of BAM than the bacterial alone, especially at lower moisture contents (Knudsen et al. 2013). Methylophilic and hydrocarbon utilizing yeasts and bacteria alone did not degrade PCBs significantly, but PCB degradation achieved about 50% when WRF were applied together (Šašek et al. 1993).

Evaluation of bioremediation effectiveness in contaminated matrices by performing ecotoxicological and genotoxic tests

In order to improve the effectiveness and performance of bioremediation processes it is important to pursue three essential goals at the same time. Focus should be not only on reducing chemical concentrations, but also on reducing chemical mobility ~~between—in~~ the environmental compartments and eventually lowering toxicity levels ensuring that contaminants do not get into the natural biological cycle (Loehr and Webster 1997; Chakraborty et al. 2013).

Bioremediation is often monitored by following the concentration of targeted contaminants (Molina-Barahona et al. 2005). Numerous studies in recent years showed that traditional chemical analyses are insufficient for a full assessment of the contaminated site as they, for example, does not

1
2
3
4
5
6
7 642 provide any information about the interaction of chemicals and does not consider the partition and
8 643 the mobility of pollutants (Frische 2003; Molina-Barahona et al. 2005; Ma et al. 2005; Molnár et al.
9
10 644 2007). An integrated approach linking the various fields and levels of study involving contaminated
11 645 sites has proven to be an efficient system of evaluating bioremediation effectiveness in
12 646 contaminated sites (Chapman and Anderson 2005; Wernersson et al. 2015; Marziali et al. 2017).
13
14 647 Consequently, to achieve the desired goals and implement a successful bioremediation program a
15 648 close collaboration of microbiologists, chemists and engineers is requested by the chemical and
16 649 biological complexity of the tasks (Van Gestel et al. 2001; Chakraborty et al. 2013).
17
18 650 Additionally, the use of ecotoxicological and genotoxic tests in order to evaluate the bioremediation
19 651 effectiveness can be a valid tool to partially overcome the existing gap between the reported
20
21 652 successes of bioremediation on the laboratory scale and the field scale.
22 653 Signals that bioremediation is going on could be important to be monitored. Two important
23 654 chemical compounds produced by microorganisms during their degradation activity are CO₂ and
24
25 655 soluble phosphorus. Both increase distinctly in the soil treated with insecticides and inoculated with
26 656 fungi (Boyle 1995; Abd El-Ghany and Masmali 2016). However, it must be taken into
27
28 657 consideration that during and after a bioremediation process the disappearance of the parent
29 658 compounds or evidence of the metabolic activity (e.g. CO₂ production) may not indicate
30 659 detoxification. Beside the fact that the fate of the toxicants may be followed by chemical analyses,
31
32 660 many reaction products resulting from a bioremediation process are not known and their potential
33 661 toxicity, as well. The elimination of mother compounds does not necessarily result in toxicity
34 662 removal, and evaluating the efficiency of the process is important to assess not only the removal of
35
36 663 a specific compound, but also the potential ecotoxicity. In fact, biodegradation of pesticides can
37 664 proceed partially or totally due to the molecular structure itself or unfavourable environmental or
38
39 665 test conditions and the lack of 'acclimatized' microbial communities (De Henau 1997). In some
40 666 instances, it has been shown that to an effective process of bioremediation corresponds to a decrease
41 667 in the toxicity of the analysed matrix (Baud-Grasset et al. 1993; Dorn and Salanitro 2000). To
42
43 668 acquire complete and useful information in an ecotoxicological assessment and to determine the
44 669 effectiveness of bioremediation treatments, it is suggested to use a battery of tests (Keddy et al.
45 670 1995; Van Gestel et al. 2001; Tigini et al. 2011). The battery should include a number of biological
46
47 671 reference organisms ~~test species~~ that are representative of the different trophic levels, in order to
48 672 select species with different roles in ecosystems, and different ~~routes of exposure~~ conditions (Van
49
50 673 Straalen and Van Gestel 1997). Moreover, the environmental risk assessment must integrate
51 674 chemical characterization, ecotoxicity and bioremediation data, in order to accurately assess the
52 675 ecological hazard.
53
54
55
56
57
58
59
60

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: Not Highlight

As emphasized by Shen et al. (2016), an increased level of ecotoxicity within the various bioindicators either could indicate an incomplete decomposition of the substance or could result from the formation of intermediate products generated via the bioremediation process. For this reason, sometimes chronic tests are more appropriate in evaluating the toxicity caused by by-products (Lofrano et al. 2014). ~~In other cases, however, also the toxicity of the by products is effectively removed (Lofrano et al. 2016).~~

Formatted: Strikethrough

In certain circumstances, there is a clear need to monitor the bioremediation process using different bioindicators. In Lizano-Fallas et al. (2017), for example, the ecotoxicity test with *Daphnia magna* shows a clear detoxification, whilst the detoxification patterns remain unclear when applying the phytotoxicity test. Ecotoxicological tests can also be used to determine the most suitable bioremediation technique in relation to the examined case study as reported in Dudášová et al. (2016). Without worldwide-recognized unique guidelines for water quality assessment, literature data are difficult to compare due to the variety of model organisms, end-points, etc. Synthetic indices capable of summarizing these findings could help to have an objective advice about the effectiveness of the biological treatment. They have been already applied for toxicity monitoring of wastewaters (Tigini et al. 2011) but municipal effluents containing AOCs have never been taken into consideration nor estrogenic activity has been included so far.

Several toxicity assays were included in the treatability study protocol to measure remediation efficiency. Assessing the toxicity of complex matrixes such soil could acquire methods from bioassays used to test toxicity of chemical compounds reported by the Organization for Economic Co-operation and Development (e.g. OECD 201 2006; OECD 211 2012). OECD has published a series of standardized tests for determining the biodegradability of a given compound, based on the evaluation of overall parameters (such as COD, TOC and BOD) or methabolic tests, e.g. respirometric (OECD 209 1984) as Polo et al. (2011) used for revealing susceptibility ~~to~~ of toxic compound comprising herbicide to biological treatment. Standardized testing procedures using different organisms have been approved by various environmental organizations, including the US Environmental Protection Agency, American Society for Testing and Materials, International Standardization Organization (Siciliano et al. 2015). Many scientists have explored the effects of polluted soil on the whole organism using various microorganisms, animals, and plants, or by means of cellular, and biochemical biomarkers, or by ecological scale up systems. Here below, tests at some different biological hierarchical levels of analyses are reported and discussed.

Formatted: No underline

Organismal level

Concerning complex matrixes as soil, quality assessments are performed with organisms on extracts of the polluted matrix, generally applying short-term exposure periods (Van Gestel et al. 2001). Experimental models are aquatic organisms such as *Daphnia magna*, *Raphidocelis subcapitata*, *Danio rerio*, *Myriophyllum aquaticum* or *Lemna minor* (Feiler et al. 2004). The use of freshwater and marine biota may be particular useful in order to provide a more complete comprehension on the environmental outcomes of agricultural activities evaluating the fate of pesticides (Guida et al. 2008). Terrestrial animals such as nematodes (*Caenorhabditis elegans*) (Traunspurger et al. 1997), oligochaetes (*Lumbriculus variegatus*) (Phipps et al. 1993), ~~springtails~~, springtails as *Folsomia candida* (Houx et al. 1996), and fish embryos (Hollert et al. 2003; Zielke et al. 2011) are well considered among the most reliable models.

Among higher plants important experimental models are *Lepidium sativum*, *Cucumis sativus*, and *Sorghum saccharatum* (germination rate, inhibition of root elongation). Since assays based on animals, plants and algae are considered expensive, time consuming and require large sample volume, recent studies have emphasized the benefits of rapid, reproducible and cost effective bacterial assays for toxicity screening and assessment. *Arthrobacter globiformi* (Neumann-Hensel and Melbye 2006), *Bacillus cereus* (Rönnpapel et al. 1995; Prokop et al. 2016), *Vibrio proteolyticus* (Ahlf and Heise 2005) yeasts (*Saccharomyces cerevisiae*) (Weber et al. 2006) are often used; otherwise, among bacterial bioassays, *Vibrio fischeri* luminescence inhibition test is the most common. The review of Parvez et al. (2006) remarks that *Vibrio fischeri* inhibition test is the most sensitive test, cost effective, easy to operate and requires only 5–30 min for toxicity prediction.

Cellular and biomolecular level

Biomarkers are adaptive responses by the organisms after exposure to xenobiotics. Various studies highlighted the cytotoxicity and genotoxicity effect of OACs and their metabolic products on the organisms. The exposed organisms may exhibit histological, cellular, molecular, biochemical and/or physiological, or even by behavioural changes (Depledge et al. 1993) that enable the obtaining of information on the biological effects of pollutants or their remains during or after a bioremediation process (Fontanetti et al. 2011).

Genetic endpoints and biomarkers. The most used biomarkers are mitotic index, chromosome aberrations, micronuclei, sister chromatid exchange and mutations.

Various scientists have recommended bacteria for bioassays evaluating genotoxicity in different samples (Mortelmans and Zeiger 2000; White and Claxton 2004). Ames test, one of the most famous and used, is a short term bacterial reverse mutation assay especially designed to evaluate the mutagenic potential of wide range of chemical substances (Mortelmans and Zeiger 2000) and was

Formatted: Font color: Auto, Not Expanded by / Condensed by , Pattern: Clear

Formatted: Default Paragraph Font, Font: +Body (Calibri), 11 pt, Font color: Auto, English (U.S.), Not Expanded by / Condensed by , Pattern: Clear

found to be very sensitive to wide range of mutagenic and carcinogenic chemicals as reported in the review paper of Chahal et al. (2014).

On the side of plant models, higher plants are recognized as excellent genetic models to detect cytogenetic and mutagenic agents and are frequently used in environmental monitoring studies. The main organisms are *Allium cepa*, *Vicia faba* and *Tradescantia* spp. as reported in a review by De Souza et al. (2016). Their protocols are standardized through a program under the International Program on Plant Bioassays (IPPB) conducted by the United Nations Environment Programme (UNEP) (Ma 1999). In addition, the US Environmental Protection Agency (USEPA) and the World Health Organization (WHO) validated the results obtained with plant bioindicators as an efficient model to detect environmental genotoxicity.

One of the most used higher plant model is *V. faba*. The main advantages are its availability round the year, economical to use, easy to grow and handle; its use does not require sterile conditions and rate of cell division is fast. The *V. faba* test, deeply reported and discussed in the review of Iqbal (2016), enables the assessment of different endpoints i.e., chromosomal aberration, mitotic index, micronuclei and nuclear aberration.

Enzymatic biomarkers. Enzymatic activity inhibition as biomarker has been widely evaluated to measure toxicity of a matrix. Dehydrogenases, for example, are directly involved in many of the vital anabolic and catabolic processes of living organisms, and their activity is inhibited by chemical toxicants. Recently, many studies have reported the use of terrestrial organisms for developing enzymatic biomarkers in response to residual pesticides (Henson-Ramsey et al. 2011; Radwan and Mohamed 2013; Stepić et al. 2013), and among these, earthworms were widely used to understand the impacts of pesticides. In two earthworm species, *Eisenia fetida* and *Lumbricus terrestris*, multiple esterases, including acetylcholinesterase (AChE), butyrylcholinesterase, and carboxylesterase (CE), have been assessed as biomarkers for malathion exposure (Henson-Ramsey et al. 2011). Several studies have also reported AChE, catalase (CAT), and glutathione-S-transferase as bio-chemical biomarkers in *Eisenia andrei* for the insecticides endosulfan, temephos, malathion, and pirimiphos-methyl (Stepić et al. 2013), and AChE, CAT, CE, and the efflux pump as biomarkers in *E. andrei* and *Octolasion lacteum* for dimethoa. Recently, surface-enhanced laser desorption/ionization-time-of-flight (SELDI-TOF) mass spectrometry (MS) has strongly contributed to the identification of more accurate, precise biomarkers e.g. specific for human cancers (Silsirivanit et al. 2014), or for endosulfan exposure in Japanese rice fish (*Oryzias latipes*) (Lee et al. 2013). In a recent paper, selective protein biomarkers for 6 pesticides (captan, carbaryl, carbofuran, and α -endosulfan chlorpyrifos, propoxur) were found in *E. fetida*, by means of SELDI-TOF MS technology (Park et al. 2015).

1
2
3
4
5
6
7 Estrogenic and androgenic biomarkers. It has been well documented that several chemicals from
8 agricultural, industrial, and household sources possess endocrine-disrupting properties, which
9 provide a potential threat to human and wildlife reproduction (Colborn et al. 1993; Colborn 1995;
10 Jensen et al. 1995). A suggested mechanism is that environmental contaminants alter the normal
11 functioning of the endocrine and reproductive system by mimicking or inhibiting endogenous
12 hormone action, modulating the production of endogenous hormones, or altering hormone receptor
13 populations (Sonnenschein and Soto 1998). Besides several pesticides exert estrogenic and
14 antiandrogenic activities through interaction with estrogen and androgen receptors. The risk
15 associated to OACs exposure has been known for decades: many pesticides, such as p,p'-
16 dichlorodiphenyl trichloroethane (DDT) (Welch et al. 1969), methoxychlor (Bulger et al. 1978;
17 Cummings 1997), β -benzene hexachloride (BHC) (Coosen and van Velsen 1989), endosulfan,
18 toxaphene, and dieldrin (Soto et al. 1995), and fenvalerate (Garey and Wolff 1998) have been firstly
19 signaled as estrogenic. Despite increased institutional awareness and more compelling legislation
20 pressure, the most recent literature still reports the occurrence of pesticides in watercourses and in
21 passing through the trophic chains, ing-showing remarkable estrogenic or androgenic (Saillenfait et
22 al. 2016; Brander et al. 2016; Guo et al. 2017; Khalil et al. 2017; Scott et al. 2017; Miccoli et al.
23 2017; Marcoccia et al. 2017). Several bioassays have been developed and standardized in order to
24 describe the estrogenic potency of OACs. Andersen et al. (2002) indicated that several currently
25 used OACs, such as methiocarb, fenarimol, chlorpyrifos, deltamethrin, and tolclofos-methyl,
26 possess estrogenic activity on the basis of cell proliferation assay and transactivation assay using
27 MCF-7 human breast cancer cells. Kojima et al. (2004) tested 200 pesticides in vitro for agonism
28 and antagonism to two human estrogen receptor (hER) subtypes, hER α and hER β , and a human
29 androgen receptor (hAR) by highly sensitive transactivation assays, using Chinese hamster ovary
30 cells. The results demonstrated that many pesticides possess in vitro estrogenic and antiandrogenic
31 activities through ERs and/or AR. Although it appears that various pesticides exert hormonal effects
32 at concentration orders of magnitude higher than that required for physiologic hormones, wide
33 exposure to large numbers of OACs may have additive and synergistic effects. Bioassay with YES
34 (yeast estrogen screen) and YAS (yeast androgen screen) can determine hormonally active
35 compounds still present in the environment. By the the first papers about this subject (Purvis et al.
36 1991), much more sophisticated bioassays have been developed such as that proposed by Eldridge
37 et al. (2007) in which a bioluminescence strain of *Saccharomyces cerevisiae* was genetically
38 engineered to respond to androgenic chemicals.

52 *Ecological level*

The actual risk of chemical residues pollution from bioremediation process is underestimated at the ecological level in natural systems. The ecological scaling-up experiment illustrated by Rodea-Palomares et al. (2016) underlined how real-world exposure to chemical pollution is often dominated by low-dose complex combined with other biotic and abiotic stressors. In the paper, a novel screening method (GSA-QHTS) was reported, that coupled the computational power of global sensitivity analysis (GSA) with the experimental efficiency of quantitative high-throughput screening (QHTS). In the case of study, they reported GSA-QHTS allowed for the identification of the main pharmaceutical pollutants, driving biological effects of low-dose complex mixtures at the microbial population level. The target complex community was a river benthic microbial community inocula obtained from an unpolluted stream. The effect of the toxic compounds in a mixture was evaluated together with other physico-chemical stressors, on a series of community level metabolic end points. Photosynthetic parameters, the dark-adapted basal fluorescence, the light-adapted steady-state fluorescence, the maximum photosynthetic efficiency, as well as the extracellular enzymatic activities b-Glu and Phos were considered as both autotrophic and heterotrophic global fitness indicators suited to study the effects of chemical pollution on freshwater benthic microbial communities.

Prospect

Bioremediation is based on the idea that different organisms will work together to remove (biodegrade) the waste substances or pollutants (OACs) from environment. Although limitations for bioremediation practice might be occurred, including the nature of organisms, the enzyme involved, the concentration and availability and finally survival of microorganisms, as well as cost/benefit ratio (i.e. cost versus overall environmental impact), to some extent, these limitations can be solved by understanding the genetics and biochemistry of desired microbe. The advent of synthetic community showed giant potential ability in facilitating the bioremediation process, especially the effective utility of degradative fungi.

Acknowledgement

~~B. Wu is funded by National Natural Science Foundations of China (No. 31701853). The research was jointly supported by Beijing Municipal Science and Technology Project (No. D151100003915002) and Science and Technology Service Network Initiative (No. KFJ-SW-STS-143-5).~~ L. Pecoraro acknowledges CAS 153211KYSB20160029 for supporting his research at Chinese Academy of Sciences.

References

Abd El-Ghany T, Masmali IA. 2016. Fungal biodegradation of organophosphorus insecticides and their impact on soil microbial population. J Plant Pathol Microbiol. Available from: <https://www.omicsonline.org/open-access/fungal-biodegradation-of-organophosphorus-insecticides-and-their-impact-on-soil-microbial-population-2157-7471-1000349.php?aid=72754>

Aguilar JAP, Andreu V, Campo J, Picó Y, Masiá A. 2017. Pesticide occurrence in the waters of Júcar River, Spain from different farming landscapes. Sci Total Environ 607–608: 752–760.

Ahlf W, Heise S. 2005. Sediment toxicity assessment: rationale for effect classes. J Soils Sediments. 5:16–20.

Alavanja MCR. 2009. Introduction: Pesticides Use and Exposure, Extensive Worldwide. Rev Environ Health. Available from: <https://www.degruyter.com/view/j/reveh.2009.24.4/reveh.2009.24.4.303/reveh.2009.24.4.303.xml>

Alvarenga N, Birolli WG, Seleglim MHR, Porto ALM. 2014. Biodegradation of methyl parathion by whole cells of marine-derived fungi *Aspergillus sydowii* and *Penicillium decaturense*. Chemosphere. 117:47–52.

Anastasi A, Tigini V, Varese GC. 2013. The bioremediation potential of different ecophysiological groups of fungi. In: Goltapeh EM, Danesh YR, Varma A, editors. Fungi as Bioremediators. Springer Berlin Heidelberg; p. 29–49. Available from: <http://dx.doi.org/10.1007/978-3-642-33811>

Andersen HR, Vinggaard AM, Høj Rasmussen T, Gjermansen IM, Cecilie Bonefeld-Jørgensen E. 2002. Effects of currently used pesticides in assays for estrogenicity, androgenicity, and aromatase activity in vitro. Toxicol Appl Pharmacol. 179:1–12.

Arakaki RL, Monteiro DA, Boscolo M, Dasilva R, Gomes E. 2013. Halotolerance, ligninase production and herbicide degradation ability of basidiomycetes strains. Braz J Microbiol. 44:1207–1214.

Aravinna P, Priyantha N, Pitawala A, Yatigammana SK. 2017. Use pattern of pesticides and their predicted mobility into shallow groundwater and surface water bodies of paddy lands in Mahaweli

- river basin in Sri Lanka. *J Environ Sci Health Part B*. 52:37–47. Available from: <http://dx.doi.org/10.1080/03601234.2016.1229445>
- Ballesteros ML, Miglioranza KSB, Gonzalez M, Fillmann G, Wunderlin DA, Bistoni MA. 2014. Multimatrix measurement of persistent organic pollutants in Mar Chiquita, a continental saline shallow lake. *Sci Total Environ*. 490:73–80.
- Barnes KK, Kolpin DW, Furlong ET, Zaugg SD, Meyer MT, Barber LB. 2008. A national reconnaissance of pharmaceuticals and other organic wastewater contaminants in the United States — I) Groundwater. *Sci Total Environ*. 402:192–200.
- Baud-Grasset F, Baud-Grasset S, Safferman SI. 1993. Evaluation of the bioremediation of a contaminated soil with phytotoxicity tests. *Chemosphere*. 26:1365–1374.
- Bicchi C, Schilirò T, Pignata C, Fea E, Cordero C, Canale F, Gilli G. 2009. Analysis of environmental endocrine disrupting chemicals using the E-screen method and stir bar sorptive extraction in wastewater treatment plant effluents. *Sci Total Environ*. 407:1842–1851.
- Biliouris K, Babson D, Schmidt-Dannert C, Kaznessis YN. 2012. Stochastic simulations of a synthetic bacteria-yeast ecosystem. *BMC Syst Biol*. 6:58.
- Birolli WG, Alvarenga N, Seleghim MHR, Porto ALM. 2016. Biodegradation of the pyrethroid pesticide esfenvalerate by marine-derived fungi. *Mar Biotechnol*. 18:511–520.
- Boyle CD. 1995. Development of a practical method for inducing white-rot fungi to grow into and degrade organopollutants in soil. *Can J Microbiol*. 41:345–353.
- Brander SM, Gabler MK, Fowler NL, Cannon RE, Schlenk D. 2016. Pyrethroid pesticides as endocrine disruptors: molecular mechanisms in vertebrates with a focus on fishes. *Environ Sci Technol*. 50:8977–8992.
- Brenner K, You L, Arnold FH. 2008. Engineering microbial consortia: a new frontier in synthetic biology. *Trends Biotechnol*. 26:483–489.
- Bulger WH, Muccitelli RM, Kupfer D. 1978. Studies on the in vivo and in vitro estrogenic activities of methoxychlor and its metabolites. Role of hepatic mono-oxygenase in methoxychlor activation. *Biochem Pharmacol*. 27:2417–2423.

1
2
3
4
5
6
7 900 Campo J, Masiá A, Blasco C, Picó Y. 2013. Occurrence and removal efficiency of pesticides in
8 901 sewage treatment plants of four Mediterranean River Basins. J Hazard Mater. 263:146–157.
9
10 902 Carles L, Rossi F, Joly M, Besse-Hoggan P, Batisson I, Artigas J. 2017. Biotransformation of
11 herbicides by aquatic microbial communities associated to submerged leaves. Environ Sci Pollut
12 903 Res. 24:3664–3674.
13 904
14
15 905 Carranza CS, Barberis CL, Chiacchiera SM, Magnoli CE. 2014. Influence of the pesticides
16 glyphosate, chlorpyrifos and atrazine on growth parameters of nonochratoxigenic *Aspergillus*
17 906 section *Nigri* strains isolated from agricultural soils. J Environ Sci Health Part B. 49:747–755.
18 907
19
20 908 Carvalho FP. 2017. Pesticides, environment, and food safety. Food Energy Secur. 6:48–60.
21
22 909 Castellana G, Loffredo E. 2014. Simultaneous removal of endocrine disruptors from a wastewater
23 using white rot fungi and various adsorbents. Water Air Soil Pollut. Available from:
24 910 <http://link.springer.com/10.1007/s11270-014-1872-6>
25 911
26
27 912 Ceci A, Pierro L, Riccardi C, Pinzari F, Maggi O, Persiani AM, Gadd GM, Petrangeli Papini M.
28 2015. Biotransformation of β -hexachlorocyclohexane by the saprotrophic soil fungus *Penicillium*
29 913 *griseofulvum*. Chemosphere. 137:101–107.
30 914
31
32 915 Chahal V, Nagpal A, Pakade Y, Katnoria J. 2014. Ecotoxicological studies of soil using analytical
33 and biological methods: a review. World Acad Sci Eng Technol Int J Biol Biomol Agric Food
34 916 Biotechnol Eng. 8:302–318.
35 917
36
37 918 Chakraborty S, Sikder J, Mukherjee D, Kanti Mandal M, Arockiasamy DL. 2013. Bioremediation
38 of Water: A Sustainable Approach. In: Piemonte V, De Falco M, Basile A, editors. Sustain Dev
39 919 Chem Eng Innov Technol. Chichester, UK: John Wiley & Sons, Ltd; p. 241–266. Available from:
40 920 <http://doi.wiley.com/10.1002/9781118629703.ch10>
41 921
42
43 922 Chan-Cupul W, Abarca GH, Vázquez RR, Salmones D, Hernández RG, Gutiérrez EA. 2014.
44 Response of ligninolytic macrofungi to the herbicide atrazine: dose-response bioassays. Rev Argent
45 923 Microbiol. 46:348–357.
46 924
47
48 925 Chan-Cupul W, Heredia-Abarca G, Rodríguez-Vázquez R. 2016. Atrazine degradation by fungal
49 co-culture enzyme extracts under different soil conditions. J Environ Sci Health Part B. 51:298–
50 926 308.
51 927
52
53
54
55
56
57
58
59
60

- Chapman PM, Anderson J. 2005. A decision-making framework for sediment contamination. *Integr Environ Assess Manag.* 1:163.
- Colborn T. 1995. Environmental estrogens: health implications for humans and wildlife. *Environ Health Perspect.* 103:135.
- Colborn T, vom Saal FS, Soto AM. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect.* 101:378.
- Coosen R, van Velsen FL. 1989. Effects of the β -isomer of hexachlorocyclohexane on estrogen-sensitive human mammary tumor cells. *Toxicol Appl Pharmacol.* 101:310–318.
- Cummings AM. 1997. Methoxychlor as a model for environmental estrogens. *Crit Rev Toxicol.* 27:367–379.
- Čvančarová M, Moeder M, Filipová A, Cajthaml T. 2015. Biotransformation of fluoroquinolone antibiotics by ligninolytic fungi - Metabolites, enzymes and residual antibacterial activity. *Chemosphere.* 136:311–320.
- Cycoń M, Piotrowska-Seget Z, Kozdrój J. 2010. Linuron effects on microbiological characteristics of sandy soils as determined in a pot study. *Ann Microbiol.* 60:439–449.
- Da Silva Coelho J, de Oliveira AL, Marques de Souza CG, Bracht A, Peralta RM. 2010. Effect of the herbicides bentazon and diuron on the production of ligninolytic enzymes by *Ganoderma lucidum*. *Int Biodeterior Biodegrad.* 64:156–161.
- De A, Bose R, Kumar A, Mozumdar S. 2014. Worldwide Pesticide Use. In: Target Deliv Pestic Using Biodegrad Polym Nanoparticles. New Delhi: Springer India; p. 5–6. Available from: http://link.springer.com/10.1007/978-81-322-1689-6_2
- De Henau H. 1997. Biodegradation. In: Calow P, editor. *Handb Ecotoxicol.* Oxford, UK: Blackwell Publishing Ltd; p. 355–377. Available from: <http://doi.wiley.com/10.1002/9781444313512.ch18>
- Depledge MH, Amaral-Mendes JJ, Daniel B, Halbrook RS, Kloepper-Sams P, Moore MN, Peakall DB. 1993. The conceptual basis of the biomarker approach. In: Peakall DB, Shugart LR, editors. *Biomarkers.* Berlin, Heidelberg: Springer Berlin Heidelberg; p. 15–29. Available from: http://www.springerlink.com/index/10.1007/978-3-642-84631-1_2

1
2
3
4
5
6
7 955 De Roy K, Marzorati M, Van den Abbeele P, Van de Wiele T, Boon N. 2014. Synthetic microbial
8 956 ecosystems: an exciting tool to understand and apply microbial communities. *Environ Microbiol.*
9 957 16:1472–1481.
10
11
12 958 De Souza CP, de Andrade Guedes T, Fontanetti CS. 2016. Evaluation of herbicides action on plant
13 959 bioindicators by genetic biomarkers: a review. *Environ Monit Assess.* 188:694.
14
15 960 Dorn PB, Salanitro JP. 2000. Temporal ecological assessment of oil contaminated soils before and
16 961 after bioremediation. *Chemosphere.* 40:419–426.
17
18
19 962 Dudášová H, Lászlóvá K, Lukáčová L, Balaščáková M, Murínová S, Dercová K. 2016.
20 963 Bioremediation of PCB-contaminated sediments and evaluation of their pre- and post-treatment
21 964 ecotoxicity. *Chem Pap.* Available from: [http://www.degruyter.com/view/j/chempap.2016.70.issue-](http://www.degruyter.com/view/j/chempap.2016.70.issue-8/chempap-2016-0041/chempap-2016-0041.xml)
22 965 8/chempap-2016-0041/chempap-2016-0041.xml
23
24
25 966 EEA. 2005. Environment and health. Luxembourg: Office for Official Publications of the European
26 967 Communities.
27
28
29 968 EEA. 2016. EMEP/EEA air pollutant emission inventory guidebook 2016: 3. Df, 3. I Agriculture
30 969 other including use of pesticides.
31
32 970 Eldridge ML, Sanseverino J, Layton AC, Easter JP, Schultz TW, Sayler GS. 2007. *Saccharomyces*
33 971 *cerevisiae* BLYAS, a new bioluminescent bioreporter for detection of androgenic compounds. *Appl*
34 972 *Environ Microbiol.* 73:6012-8.
35
36
37 973 Ellegaard-Jensen L, Knudsen BE, Johansen A, Albers CN, Aamand J, Rosendahl S. 2014. Fungal–
38 974 bacterial consortia increase diuron degradation in water-unsaturated systems. *Sci Total Environ.*
39 975 466-467:699–705.
40
41
42 976 Fan X, Song F. 2014. Bioremediation of atrazine: recent advances and promises. *J Soils Sediments.*
43 977 14:1727–1737.
44
45 978 Feiler U, Kirchesch I, Heininger P. 2004. A new plant-based bioassay for aquatic sediments. *J Soils*
46 979 *Sediments.* 4:261–266.
47
48
49 980 Fontanetti CS, Nogarol LR, de Souza RB, Perez GD, Maziviero GT. 2011. Bioindicators and
50 981 Biomarkers in the Assessment of Soil Toxicity. In: Pascucci S, editor. *Soil Contam. InTech.*
51 982 Available from: [http://www.intechopen.com/books/soil-contamination/bioindicators-and-](http://www.intechopen.com/books/soil-contamination/bioindicators-and-biomarkers-in-the-assessment-of-soil-toxicity)
52 983 [biomarkers-in-the-assessment-of-soil-toxicity](http://www.intechopen.com/books/soil-contamination/bioindicators-and-biomarkers-in-the-assessment-of-soil-toxicity)
53
54
55
56
57
58
59
60

- Frzsche T. 2003. Ecotoxicological evaluation of in situ bioremediation of soils contaminated by the explosive 2,4,6-trinitrotoluene (TNT). *Environ Pollut.* 121:103–113.
- Gao D, Du L, Yang J, Wu W-M, Liang H. 2010. A critical review of the application of white rot fungus to environmental pollution control. *Crit Rev Biotechnol.* 30:70–77.
- García-Carmona M, Romero-Freire A, Sierra Aragón M, Martínez Garzón FJ, Martín Peinado FJ. 2017. Evaluation of remediation techniques in soils affected by residual contamination with heavy metals and arsenic. *J Environ Manage.* 191:228–236.
- Garey J, Wolff MS. 1998. Estrogenic and antiprogesterogenic activities of pyrethroid insecticides. *Biochem Biophys Res Commun.* 251:855–859.
- Gargano ML, van Griensven LJLD, Isikhuehnen OS, Lindequist U, Venturella G, Wasser SP, Zervakis GI. 2017. Medicinal mushrooms: valuable biological resources of high exploitation potential. *Plant Biosyst - Int J Deal Asp Plant Biol.* 151:548–565.
- Gong ZM, Xu FL, Dawson R, Cao J, Liu WX, Li BG, Shen WR, Zhang WJ, Qin BP, Sun R, Tao S. 2004. Residues of hexachlorocyclohexane isomers and their distribution characteristics in soils in the Tianjin Area, China. *Arch Environ Contam Toxicol.* Available from: <http://link.springer.com/10.1007/s00244-003-2301-9>
- Großkopf T, Soyer OS. 2014. Synthetic microbial communities. *Curr Opin Microbiol.* 18:72–77.
- Guerreño M, López Armengol MF, Luquet CM, Venturino A. 2016. Comparative study of toxicity and biochemical responses induced by sublethal levels of the pesticide azinphosmethyl in two fish species from North-Patagonia, Argentina. *Aquat Toxicol.* 177:365–372.
- Guida M, Inglese M, Meriç S. 2008. A multi-battery toxicity investigation on fungicides. *Desalination.* 226:262–270.
- Guo D, Wang Y, Qian Y, Chen C, Jiao B, Cai L, Wang Q. 2017. Joint acute and endocrine disruptive toxicities of malathion, cypermethrin and prochloraz to embryo-larval zebrafish, *Danio rerio*. *Chemosphere.* 166:63–71.
- Hai FI, Modin O, Yamamoto K, Fukushi K, Nakajima F, Nghiem LD. 2012. Pesticide removal by a mixed culture of bacteria and white-rot fungi. *J Taiwan Inst Chem Eng.* 43:459–462.

Formatted: Font: (Default) +Body (Calibri), 11 pt

Formatted: Normal, Left, Line spacing: single

1
2
3
4
5
6
7 1012 Hammesfahr U, Heuer H, Manzke B, Smalla K, Thiele-Bruhn S. 2008. Impact of the antibiotic
8 1013 sulfadiazine and pig manure on the microbial community structure in agricultural soils. Soil Biol
9
10 1014 Biochem. 40:1583–1591.
11
12 1015 Harclerode M, Ridsdale DR, Darmendrail D, Bardos P, Alexandrescu F, Nathanail P, Pizzol L,
13 1016 Rizzo E. 2015. Integrating the social dimension in remediation decision-making: state of the
14
15 1017 practice and way forward. Remediat J. 26:11–42.
16
17 1018 Harms H, Wick LY, Schlosser D. 2017. The fungal community in organically polluted systems. In:
18 1019 Dighton J, White JF, editors. Mycology. CRC Press Taylor & Francis Group 6000 Broken Sound
19
20 1020 Parkway NW, Suite 300 Boca Raton, FL 33487-2742: CRC Press; p. 459–470. Available from:
21 1021 <http://www.crcnetbase.com/doi/10.1201/9781315119496-32>
22
23 1022 Henson-Ramsey H, Schneider A, Stoskopf MK. 2011. A comparison of multiple esterases as
24
25 1023 biomarkers of organophosphate exposure and effect in two earthworm species. Bull Environ
26 1024 Contam Toxicol. 86:373–378.
27
28 1025 Hollert H, Keiter S, König N, Rudolf M, Ulrich M, Braunbeck T. 2003. A new sediment contact
29
30 1026 assay to assess particle-bound pollutants using zebrafish (*Danio rerio*) embryos. J Soils Sediments.
31 1027 3:197.
32
33 1028 Houx NWH, Dekker A, Van Kammen-Polman AMM, Ronday R. 1996. Acute toxicity test for
34
35 1029 terrestrial hazard assessment with exposure of *Folsomia candida* to pesticides in an aqueous
36 1030 medium. Arch Environ Contam Toxicol. 30:9–14.
37
38 1031 Huang McBeath J, McBeath J. 2010. Environmental change and food security in China. Dordrecht:
39 1032 Springer Netherlands. Available from: <http://link.springer.com/10.1007/978-1-4020-9180-3>
40
41 1033 Hui X, Yi Q, Bu-zhuo P, Xiliu J, Xiao-mei H. 2003. Environmental pesticide pollution and its
42
43 1034 countermeasures in China. AMBIO J Hum Environ. 32:78–80.
44
45 1035 Iqbal M. 2016. *Vicia faba* bioassay for environmental toxicity monitoring: A review. Chemosphere.
46 1036 144:785–802.
47
48 1037 Jagmann N, Philipp B. 2014. Reprint of design of synthetic microbial communities for
49
50 1038 biotechnological production processes. J Biotechnol. 192:293–301.
51
52
53
54
55
56
57
58
59
60

- Jarošová B, Erseková A, Hilscherová K, Loos R, Gawlik BM, Giesy JP, Bláha L. 2014. Europe-wide survey of estrogenicity in wastewater treatment plant effluents: the need for the effect-based monitoring. *Environ Sci Pollut Res*. 21:10970–10982.
- Jechalke S, Heuer H, Siemens J, Amelung W, Smalla K. 2014. Fate and effects of veterinary antibiotics in soil. *Trends Microbiol*. 22:536–545.
- Jensen TK, Toppari J, Keiding N, Skakkebaek NE. 1995. Do environmental estrogens contribute to the decline in male reproductive health? *Clin Chem*. 41:1896–1901.
- Jones A, Panagos P, Barcelo S, Bouraoui F, Bosco C, Dewitte O, Gardi C, Erhard M, Hervás J, Hiederer R. 2012. The state of soil in Europe. *JRC Ref Rep Jt Res Cent Eur Comm Ispra*.
- Kardol P, Wardle DA. 2010. How understanding aboveground-belowground linkages can assist restoration ecology. *Trends Ecol Evol*. 25:670–679.
- Kaur H, Kapoor S, Kaur G. 2016. Application of ligninolytic potentials of a white-rot fungus *Ganoderma lucidum* for degradation of lindane. *Environ Monit Assess*. 188:588.
- Keddy CJ, Greene JC, Bonnell MA. 1995. Review of whole-organism bioassays: soil, freshwater sediment, and freshwater assessment in Canada. *Ecotoxicol Environ Saf*. 30:221–251.
- Khalil SR, Awad A, Mohammed HH. 2017. Behavioral response and gene expression changes in fipronil-administered male Japanese quail (*Coturnix japonica*). *Environ Pollut Barking Essex* 1987. 223:51–61.
- Khromonygina VV, Saltykova AI, Vasil'chenko LG, Kozlov IP, Rabinovich ML. 2004. Degradation of the herbicide atrazine by the soil mycelial fungus INBI 2-26(-)--a producer of cellobiose dehydrogenase. *Prikl Biokhim Mikrobiol*. 40:337–343.
- Kim K-H, Kabir E, Jahan SA. 2017. Exposure to pesticides and the associated human health effects. *Sci Total Environ*. 575:525–535.
- Knudsen BE, Ellegaard-Jensen L, Albers CN, Rosendahl S, Aamand J. 2013. Fungal hyphae stimulate bacterial degradation of 2,6-dichlorobenzamide (BAM). *Environ Pollut*. 181:122–127.
- Köck-Schulmeyer M, Villagrasa M, López de Alda M, Céspedes-Sánchez R, Ventura F, Barceló D. 2013. Occurrence and behavior of pesticides in wastewater treatment plants and their environmental impact. *Sci Total Environ*. 458-460:466–476.

1
2
3
4
5
6
71067 Kojima H, Katsura E, Takeuchi S, Niiyama K, Kobayashi K. 2004. Screening for estrogen and
81068 androgen receptor activities in 200 pesticides by in vitro reporter gene assays using Chinese hamster
91069 ovary cells. Environ Health Perspect. 112:524–531.
10
11
121070 Koroleva O, Stepanova E, Landesman E, Vasilchenko L, Khromonygina V, Zherdev A, Rabinovich
131071 M. 2002. Enzyme immunoassay of herbicide decomposition by soil and wood decay fungi. Appl
141072 Biochem Microbiol. 38:355–360.
15
16
171073 Kulshreshtha S, Mathur N, Bhatnagar P. 2014. Mushroom as a product and their role in
181074 mycoremediation. AMB Express. Available from: <http://www.amb-express.com/content/4/1/29>
19
201075 Lee S-E, Young-Woong C, Mo H, Son J, Park K, Cho K. 2013. Endosulfan-induced biomarkers in
211076 Japanese rice fish (*Oryzias latipes*) analyzed by SELDI-TOF-MS. Int J Biol Sci. 9:343–349.
22
23
241077 Liu WW, Wu B, Xiang M, Liu X. 2017. From microbiome to synthetic microbial community.
251078 Microbiol China. 44: 1-9 (in Chinese).
26
271079 Lizano-Fallas V, Masís-Mora M, Espinoza-Villalobos D, Lizano-Brenes M, Rodríguez-Rodríguez
281080 CE. 2017. Removal of pesticides and ecotoxicological changes during the simultaneous treatment of
291081 triazines and chlorpyrifos in biomixtures. Chemosphere. 182:106–113.
30
31
321082 Loehr RC, Webster MT. 1997. Changes in toxicity and mobility resulting from bioremediation
331083 processes. Bioremediation J. 1:149–163.
34
35
361084 Loffredo E, Castellana G, Taskin E. 2016. A two-step approach to eliminate pesticides and
371085 estrogens from a wastewater and reduce its phytotoxicity: adsorption onto plant-derived materials
381086 and fungal degradation. Water Air Soil Pollut. Available from:
391087 <http://link.springer.com/10.1007/s11270-016-2883-2>
40
41
421088 Lofrano G, Carotenuto M, Uyguner-Demirel CS, Vitagliano A, Siciliano A, Guida M. 2014. An
431089 integrated chemical and ecotoxicological assessment for the photocatalytic degradation of
441090 vancomycin. Environ Technol. 35:1234–1242.
45
46
471091 ~~Lofrano G, Libralato G, Adinolfi R, Siciliano A, Iannece P, Guida M, Giugni M, Volpi Ghirardini~~
481092 ~~A, Carotenuto M. 2016. Photocatalytic degradation of the antibiotic chloramphenicol and effluent~~
491093 ~~toxicity effects. Ecotoxicol Environ Saf. 123:65–71.~~

- Loos R, Locoro G, Comero S, Contini S, Schwesig D, Werres F, Balsaa P, Gans O, Weiss S, Blaha L, et al. 2010. Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Res.* 44:4115–4126.
- Luo Y, Guo W, Ngo HH, Nghiem LD, Hai FI, Zhang J, Liang S, Wang XC. 2014. A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. *Sci Total Environ.* 473-474:619–641.
- Lu X, Kang Z, Tao B, Wang Y, Dong J, Zhang J. 2012. Degradation of nicosulfuron by *Bacillus subtilis* YB1 and *Aspergillus niger* YF1. *Appl Biochem Microbiol.* 48:460–466.
- Ma M, Li J, Wang Z. 2005. Assessing the Detoxication Efficiencies of Wastewater Treatment Processes Using a Battery of Bioassays/Biomarkers. *Arch Environ Contam Toxicol.* 49:480–487.
- Ma T-H. 1999. Global environment monitoring for ecosystem health and human well being — report from a round table discussion on international collaborative studies. *JASSA J Appl Sci South Afr.* 5:89–94.
- Maciel GM, Inácio FD, de Sá-Nakanishi AB, Haminiuk CWL, Castoldi R, Comar JF, Bracht A, Peralta RM. 2013. Response of *Ganoderma lucidum* and *Trametes* sp. to the herbicide picloram: Tolerance, antioxidants and production of ligninolytic enzymes. *Pestic Biochem Physiol.* 105:84–92.
- Madrigal-Zúñiga K, Ruiz-Hidalgo K, Chin-Pampillo JS, Masís-Mora M, Castro-Gutiérrez V, Rodríguez-Rodríguez CE. 2016. Fungal bioaugmentation of two rice husk-based biomixtures for the removal of carbofuran in on-farm biopurification systems. *Biol Fertil Soils.* 52:243–250.
- Majeau J-A, Brar SK, Tyagi RD. 2010. Laccases for removal of recalcitrant and emerging pollutants. *Bioresour Technol.* 101:2331–2350.
- Marcoccia D, Pellegrini M, Fiocchetti M, Lorenzetti S, Marino M. 2017. Food components and contaminants as (anti)androgenic molecules. *Genes Nutr.* Available from: <http://genesandnutrition.biomedcentral.com/articles/10.1186/s12263-017-0555-5>
- Marco-Urrea E, Pérez-Trujillo M, Vicent T, Caminal G. 2009. Ability of white-rot fungi to remove selected pharmaceuticals and identification of degradation products of ibuprofen by *Trametes versicolor*. *Chemosphere.* 74:765–772.

1
2
3
4
5
6
71122 Margot J, Copin P-J, von Gunten U, Barry DA, Holliger C. 2015. Sulfamethoxazole and
81123 isoproturon degradation and detoxification by a laccase-mediator system: Influence of treatment
9 conditions and mechanistic aspects. *Biochem Eng J.* 103:47–59.
101124
11
121125 Marziali L, Rosignoli F, Drago A, Pascariello S, Valsecchi L, Rossaro B, Guzzella L. 2017.
131126 Toxicity risk assessment of mercury, DDT and arsenic legacy pollution in sediments: A triad
14 approach under low concentration conditions. *Sci Total Environ.* 593-594:809–821.
151127
16
171128 Masaphy S, Levanon D, Vaya J, Henis Y. 1993. Isolation and characterization of a novel atrazine
181129 metabolite produced by the fungus *Pleurotus pulmonarius*, 2-chloro-4-ethylamino-6-(1-
19 hydroxyisopropyl)amino-1,3,5-triazine. *Appl Environ Microbiol.* 59:4342–4346.
201130
21
221131 Mattsson MK, Liu X, Yu D, Kontro MH. 2015. Depth, soil type, water table, and site effects on
231132 microbial community composition in sediments of pesticide-contaminated aquifer. *Environ Sci*
24 *Pollut Res.* 22:10263–10279.
251133
26
271134 Matute RG, Figlas D, Mockel G, Curvetto N. 2012. Degradation of metsulfuron methyl by *Agaricus*
281135 *blazei* Murrill spent compost enzymes. *Bioremediation J.* 16:31–37.
29
301136 Meyer H-P, Munch T. 2005. Swiss Industrial Biocatalysis Consortium. *BioWorld Eur.* 1:14–16.
31
321137 Miccoli A, Maradonna F, De Felice A, Caputo Barucchi V, Estonba A, Genangeli M, Vittori S,
33 Leonori I, Carnevali O. 2017. Detection of endocrine disrupting chemicals and evidence of their
341138 effects on the HPG axis of the European anchovy *Engraulis encrasicolus*. *Mar Environ Res.*
351139 127:137–147.
361140
37
38
391141 Miller GT. 2004. *Living in the environment: principles, connections, and solutions.* 13. ed., internat.
401142 student ed. Pacific Grove, Calif.: Thomson/Brooks/Cole.
41
421143 Mir-Tutusa JA, Masís-Mora M, Corcellas C, Eljarrat E, Barceló D, Sarrà M, Caminal G, Vicent
43 T, Rodríguez-Rodríguez CE. 2014. Degradation of selected agrochemicals by the white rot fungus
441144 *Trametes versicolor*. *Sci Total Environ.* 500-501:235–242.
451145
46
471146 Mnif W, Hassine AIH, Bouaziz A, Bartegi A, Thomas O, Roig B. 2011. Effect of endocrine
481147 disruptor pesticides: a review. *Int J Environ Res Public Health.* 8:2265–2303.
49
50
511148 Molina-Barahona L, Vega-Loyo L, Guerrero M, Ramírez S, Romero I, Vega-Jarquín C, Albores A.
521149 2005. Ecotoxicological evaluation of diesel-contaminated soil before and after a bioremediation
531150 process. *Environ Toxicol.* 20:100–109.
54
55
56
57
58
59
60

- Molnár M, Gruiz K, Halász M. 2007. Integrated methodology to evaluate bioremediation potential of creosote-contaminated soils. *Period Polytech Chem Eng.* 51:23.
- Mori T, Wang J, Tanaka Y, Nagai K, Kawagishi H, Hirai H. 2017. Bioremediation of the neonicotinoid insecticide clothianidin by the white-rot fungus *Phanerochaete sordida*. *J Hazard Mater.* 321:586–590.
- Morriën E, Hannula SE, Snoek LB, Helmsing NR, Zweers H, de Hollander M, Soto RL, Bouffaud M-L, Buée M, Dimmers W, et al. 2017. Soil networks become more connected and take up more carbon as nature restoration progresses. *Nat Commun.* 8:14349.
- Mortelmans K, Zeiger E. 2000. The Ames *Salmonella*/microsome mutagenicity assay. *Mutat Res Mol Mech Mutagen.* 455:29–60.
- Mougin C, Kollmann A, Jolival C. 2002. Enhanced production of laccase in the fungus *Trametes versicolor* by the addition of xenobiotics. *Biotechnol Lett.* 24:139–142.
- Mrema EJ, Rubino FM, Colosio C. 2013. Obsolete pesticides – A threat to environment, biodiversity and human health. In: Simeonov LI, Macaev FZ, Simeonova BG, editors. *Environ Secur Assess Manag Obsolete Pestic Southeast Eur.* Dordrecht: Springer Netherlands; p. 1–21. Available from: http://link.springer.com/10.1007/978-94-007-6461-3_1
- Nakata H, Hirakawa Y, Kawazoe M, Nakabo T, Arizono K, Abe S-I, Kitano T, Shimada H, Watanabe I, Li W, Ding X. 2005. Concentrations and compositions of organochlorine contaminants in sediments, soils, crustaceans, fishes and birds collected from Lake Tai, Hangzhou Bay and Shanghai city region, China. *Environ Pollut.* 133:415–429.
- Neumann-Hensel H, Melbye K. 2006. Optimisation of the solid-contact test with *Arthrobacter globiformis*. *J Soils Sediments.* 6:201–207.
- Nguyen LN, Hai FI, Yang S, Kang J, Leusch FDL, Roddick F, Price WE, Nghiem LD. 2014. Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*. Role of biosorption and biodegradation. *Int Biodeterior Biodegrad.* 88:169–175.
- Odukkathil G, Vasudevan N. 2013. Toxicity and bioremediation of pesticides in agricultural soil. *Rev Environ Sci Biotechnol.* 12:421–444.

1
2
3
4
5
6
7 1179 OECD 201. 2006. Test No. 201: Alga, Growth Inhibition Test. OECD Publishing. Available from:
8 1180 <http://www.oecd-ilibrary.org/environment/test-no-201-alga-growth-inhibition->
9 1181 [test_9789264069923-en](http://www.oecd-ilibrary.org/environment/test-no-201-alga-growth-inhibition-)
10
11
12 1182 OECD 209. 1984. Test No. 209: Activated Sludge, Respiration Inhibition Test. OECD Publishing.
13 1183 Available from: <http://www.oecd-ilibrary.org/oecd/content/book/9789264070080-en>
14
15 1184 OECD 211. 2012. Test No. 211: Daphnia magna Reproduction Test. OECD Publishing. Available
16 1185 from: <http://www.oecd-ilibrary.org/environment/test-no-211-daphnia-magna-reproduction->
17 1186 [test_9789264185203-en](http://www.oecd-ilibrary.org/environment/test-no-211-daphnia-magna-reproduction-)
18
19
20 1187 Oliveira BR, Penetra A, Cardoso VV, Benoliel MJ, Barreto Crespo MT, Samson RA, Pereira VJ.
21 1188 2015. Biodegradation of pesticides using fungi species found in the aquatic environment. Environ
22 1189 Sci Pollut Res Int. 22:11781–11791.
23
24
25 1190 Palvannan T, Saravanakumar T, Unnithan AR, Chung N-J, Kim D-H, Park S-M. 2014. Efficient
26 1191 transformation of phenyl urea herbicide chloroxuron by laccase immobilized on zein polyurethane
27 1192 nanofiber. J Mol Catal B Enzym. 99:156–162.
28
29
30 1193 Pan M, Chu LM. 2016. Adsorption and degradation of five selected antibiotics in agricultural soil.
31 1194 Sci Total Environ. 545-546:48–56.
32
33
34 1195 Park D-S, Jeon H-J, Park E-S, Bae IK, Kim Y-E, Lee S-E. 2015. Highly selective biomarkers for
35 1196 pesticides developed in *Eisenia fetida* using SELDI-TOF MS. Environ Toxicol Pharmacol. 39:635–
36 1197 642.
37
38
39 1198 Parrón T, Requena M, Hernández AF, Alarcón R. 2014. Environmental exposure to pesticides and
40 1199 cancer risk in multiple human organ systems. Toxicol Lett. 230:157–165.
41
42 1200 Parvez S, Venkataraman C, Mukherji S. 2006. A review on advantages of implementing
43 1201 luminescence inhibition test (*Vibrio fischeri*) for acute toxicity prediction of chemicals. Environ Int.
44 1202 32:265–268.
45
46
47 1203 Phipps GL, Ankley GT, Benoit DA, Mattson VR. 1993. Use of the aquatic oligochaete *Lumbriculus*
48 1204 *variegatus* for assessing the toxicity and bioaccumulation of sediment-associated contaminants.
49 1205 Environ Toxicol Chem. 12:269–279.
50
51
52
53
54
55
56
57
58
59
60

- Pinto AP, Rodrigues SC, Caldeira AT, Teixeira DM. 2016. Exploring the potential of novel biomixtures and *Lentinula edodes* fungus for the degradation of selected pesticides. Evaluation for use in biobed systems. *Sci Total Environ*. 541:1372–1381.
- Pinto AP, Serrano C, Pires T, Mestrinho E, Dias L, Teixeira DM, Caldeira AT. 2012. Degradation of terbuthylazine, difenoconazole and pendimethalin pesticides by selected fungi cultures. *Sci Total Environ*. 435-436:402–410.
- Polo AM, Tobajas M, Sanchis S, Mohedano AF, Rodríguez JJ. 2011. Comparison of experimental methods for determination of toxicity and biodegradability of xenobiotic compounds. *Biodegradation*. 22:751–761.
- Popp J, Pető K, Nagy J. 2013. Pesticide productivity and food security. A review. *Agron Sustain Dev*. 33:243–255.
- Prokop Z, Nečasová A, Klánová J, Čupr P. 2016. Bioavailability and mobility of organic contaminants in soil: new three-step ecotoxicological evaluation. *Environ Sci Pollut Res*. 23:4312–4319.
- Purvis IJ, Chotai D, Dykes CW, Lubahn DB, French FS, Wilson EM, Hobden AN. 1991. An androgen-inducible expression system for *Saccharomyces cerevisiae*. *Gene*. 106:35–42.
- Qiu D. 2011. Development strategy for bio-pesticide and biological control. *J Agric Sci Technol*. 5:020.
- Radwan MA, Mohamed MS. 2013. Imidacloprid induced alterations in enzyme activities and energy reserves of the land snail, *Helix aspersa*. *Ecotoxicol Environ Saf*. 95:91–97.
- Ridsdale DR, Noble BF. 2016. Assessing sustainable remediation frameworks using sustainability principles. *J Environ Manage*. 184:36–44.
- Rodea-Palomares I, Gonzalez-Pleiter M, Gonzalo S, Rosal R, Leganes F, Sabater S, Casellas M, Munoz-Carpena R, Fernandez-Pinas F. 2016. Hidden drivers of low-dose pharmaceutical pollutant mixtures revealed by the novel GSA-QHTS screening method. *Sci Adv*. 2:e1601272–e1601272.
- Rönnpapel K, Liss W, Ahlf W. 1995. Microbial bioassays to assess the toxicity of solid-associated contaminants. *Ecotoxicol Environ Saf*. 31:99–103.

1
2
3
4
5
6
7 1233 Saillenfait A-M, Ndiaye D, Sabaté J-P. 2016. The estrogenic and androgenic potential of
8 1234 pyrethroids in vitro. Review. Toxicol Vitro. 34:321–332.
9
10 1235 [Sandin M, Piikki K, Jarvis N, Larsbo M, Bishop K, Kreuger J. 2018. Spatial and temporal patterns](#)
11 1236 [of pesticide concentrations in streamflow, drainage and runoff in a small Swedish agricultural](#)
12 1237 [catchment. Sci Total Environ. 610–611: 623–634.](#)
13
14 1238 Salvestrini S, Canzano S, Iovino P, Leone V, Capasso S. 2014. Modelling the biphasic sorption of
15 1239 simazine, imidacloprid, and boscalid in water/soil systems. J Environ Sci Health Part B. 49:578–
16 1240 590.
17
18 1241 Sarkar D, Ferguson M, Datta R, Birnbaum S. 2005. Bioremediation of petroleum hydrocarbons in
19 1242 contaminated soils: comparison of biosolids addition, carbon supplementation, and monitored
20 1243 natural attenuation. Environ Pollut. 136:187–195.
21
22 1244 Šašek V, Volfová O, Erbanová P, Vyas BRM, Matucha M. 1993. Degradation of PCBs by white rot
23 1245 fungi, methylotrophic and hydrocarbon utilizing yeasts and bacteria. Biotechnol Lett. 15:521–526.
24
25 1246 Scott PD, Coleman HM, Colville A, Lim R, Matthews B, McDonald JA, Miranda A, Neale PA,
26 1247 Nugegoda D, Tremblay LA, Leusch FDL. 2017. Assessing the potential for trace organic
27 1248 contaminants commonly found in Australian rivers to induce vitellogenin in the native rainbowfish
28 1249 (*Melanotaenia fluviatilis*) and the introduced mosquitofish (*Gambusia holbrooki*). Aquat Toxicol
29 1250 Amst Neth. 185:105–120.
30
31 1251 Shah PC, Kumar VR, Dastager SG, Khire JM. 2017. Phytase production by *Aspergillus niger*
32 1252 NCIM 563 for a novel application to degrade organophosphorus pesticides. AMB Express. 7:66.
33
34 1253 Shen W, Zhu N, Cui J, Wang H, Dang Z, Wu P, Luo Y, Shi C. 2016. Ecotoxicity monitoring and
35 1254 bioindicator screening of oil-contaminated soil during bioremediation. Ecotoxicol Environ Saf.
36 1255 124:120–128.
37
38 1256 Siciliano A, Gesuele R, Pagano G, Guida M. 2015. How Daphnia (Cladocera) Assays may be used
39 1257 as Bioindicators of Health Effects? J Biodivers Endanger Species. S1:005. Available from:
40 1258 [https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-](https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936)
41 1259 [bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936](https://www.omicsonline.org/open-access/how-daphnia-cladocera-assays-may-be-used-as-bioindicators-of-health-effects-2332-2543-S1-005.php?aid=63936)
42
43 1260 Silsirivanit A, Sawanyawisuth K, Riggins GJ, Wongkham C. 2014. Cancer biomarker discovery for
44 1261 cholangiocarcinoma: the high-throughput approaches. J Hepato-Biliary-Pancreat Sci. 21:388–396.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Singh DK. 2008. Biodegradation and bioremediation of pesticide in soil: concept, method and recent developments. *Indian J Microbiol.* 48:35–40.
- Song J, Gu J, Zhai Y, Wu W, Wang H, Ruan Z, Shi Y, Yan Y. 2013. Biodegradation of nicosulfuron by a *Talaromyces flavus* LZM1. *Bioresour Technol.* 140:243–248.
- Sonnenschein C, Soto AM. 1998. An updated review of environmental estrogen and androgen mimics and antagonists. *J Steroid Biochem Mol Biol.* 65:143–150.
- Soto AM, Sonnenschein C, Chung KL, Fernandez MF, Olea N, Serrano FO. 1995. The E-SCREEN assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ Health Perspect.* 103:113.
- Spina F, Cordero C, Schilirò T, Sgorbini B, Pignata C, Gilli G, Bicchi C, Varese GC. 2015. Removal of micropollutants by fungal laccases in model solution and municipal wastewater: evaluation of estrogenic activity and ecotoxicity. *J Clean Prod.* 100:185–194.
- Stepić S, Hackenberger BK, Velki M, Lončarić Ž, Hackenberger DK. 2013. Effects of individual and binary-combined commercial insecticides endosulfan, temephos, malathion and pirimiphos-methyl on biomarker responses in earthworm *Eisenia andrei*. *Environ Toxicol Pharmacol.* 36:715–723.
- Talk A, Kublik S, Uksa M, Engel M, Berghahn R, Welzl G, Schlöter M, Mohr S. 2016. Effects of multiple but low pesticide loads on aquatic fungal communities colonizing leaf litter. *J Environ Sci.* 46:116–125.
- Tian J, Dong Q, Yu C, Zhao R, Wang J, Chen L. 2016. Biodegradation of the organophosphate trichlorfon and its major degradation products by a novel *Aspergillus sydowii* PA F-2. *J Agric Food Chem.* 64:4280–4287.
- Tigini V, Giansanti P, Mangiavillano A, Pannocchia A, Varese GC. 2011. Evaluation of toxicity, genotoxicity and environmental risk of simulated textile and tannery wastewaters with a battery of biotests. *Ecotoxicol Environ Saf.* 74:866–873.
- Traunsperger W, Haitzer M, Höss S, Beier S, Ahlf W, Steinberg C. 1997. Ecotoxicological assessment of aquatic sediments with *Caenorhabditis elegans* (nematoda) - a method for testing liquid medium and whole-sediment samples. *Environ Toxicol Chem.* 16:245–250.

- Ulčnik A, Kralj Cigić I, Pohleven F. 2013. Degradation of lindane and endosulfan by fungi, fungal and bacterial laccases. *World J Microbiol Biotechnol.* 29:2239–2247.
- Van Gestel CA, Van der Waarde JJ, Derksen JGM (Anja), Van der Hoek EE, Veul MFXW, Bouwens S, Rusch B, Kronenburg R, Stokman GNM. 2001. The use of acute and chronic bioassays to determine the ecological risk and bioremediation efficiency of oil-polluted soils. *Environ Toxicol Chem.* 20:1438–1449.
- Van Straalen NM, Van Gestel CAM. 1997. Soil Invertebrates and Micro-Organisms. In: Calow P, editor. *Handb Ecotoxicol.* Oxford, UK: Blackwell Publishing Ltd; p. 251–277. Available from: <http://doi.wiley.com/10.1002/9781444313512.ch13>
- Vasil'Chenko L, Khromonygina V, Koroleva O, Landesman E, Gaponenko V, Kovaleva T, Kozlov YP, Rabinovich M. 2002. Consumption of triazine herbicide atrazine by laccase-positive and laccase-negative strains of soil fungus *Mycelia sterilia* INBI 2-26. *Appl Biochem Microbiol.* 38:454–459.
- Velázquez-Fernández JB, Martínez-Rizo AB, Ramírez-Sandoval M, Domínguez-Ojeda D. 2012. Biodegradation and bioremediation of organic pesticides. In: Soundararajan R.P. editor. *Agricultural and Biological Sciences "Pesticides - Recent Trends in Pesticide Residue Assay"*; p. 978-953.
- Verger PJP, Boobis AR. 2013. Reevaluate pesticides for food security and safety. *Science.* 341:717–718.
- Verma JP, Jaiswal DK, Sagar R. 2014. Pesticide relevance and their microbial degradation: a-state-of-art. *Rev Environ Sci Biotechnol.* 13:429–466.
- Wang C, Yu L, Zhang Z, Wang B, Sun H. 2014. Tourmaline combined with *Phanerochaete chrysosporium* to remediate agricultural soil contaminated with PAHs and OCPs. *J Hazard Mater.* 264:439–448.
- Wang Z, Yang T, Zhai Z, Zhang B, Zhang J. 2015. Reaction mechanism of dicofol removal by cellulase. *J Environ Sci China.* 36:22–28.
- Weber J, Kreutzmann J, Plantikow A, Claus E, Manz W, Heininger P. 2006. A novel particle contact assay with the yeast *Saccharomyces cerevisiae* for ecotoxicological assessment of freshwater sediments. *J Soils Sediments.* 6:84–91.

- Wei LN, Wu P, Wang FR, Yang H. 2015. Dissipation and degradation dynamics of thifluzamide in rice field. *Water Air Soil Pollut.* Available from: <http://link.springer.com/10.1007/s11270-015-2387-5>
- Welch RM, Levin W, Conney AH. 1969. Estrogenic action of DDT and its analogs. *Toxicol Appl Pharmacol.* 14:358–367.
- Wernersson A-S, Carere M, Maggi C, Tusil P, Soldan P, James A, Sanchez W, Dulio V, Broeg K, Reifferscheid G, et al. 2015. The European technical report on aquatic effect-based monitoring tools under the water framework directive. *Environ Sci Eur.* Available from: <http://www.enveurope.com/content/27/1/7>
- White PA, Claxton LD. 2004. Mutagens in contaminated soil: a review. *Mutat Res Mutat Res.* 567:227–345.
- Wolfand JM, LeFevre GH, Luthy RG. 2016. Metabolization and degradation kinetics of the urban-use pesticide fipronil by white rot fungus *Trametes versicolor*. *Environ Sci Process Impacts.* 18:1256–1265.
- Wu B, Zhang XL, Cui BK, Dai YC. 2015a. Comparative genomic analysis of edible (medicinal) fungi reveals different ecological habitats. *Mycosystema* 34: 742-60 (in Chinese).
- Wu J, Lu J, Lu H, Lin Y, Wilson PC. 2015b. Occurrence and ecological risks from fipronil in aquatic environments located within residential landscapes. *Sci Total Environ.* 518-519:139–147.
- WWAP (United Nations World Water Assessment Programme). 2015. The United Nations World Water Development Report 2015: Water for a Sustainable World. Paris, UNESCO."
- Xiao P, Mori T, Kamei I, Kiyota H, Takagi K, Kondo R. 2011. Novel metabolic pathways of organochlorine pesticides dieldrin and aldrin by the white rot fungi of the genus *Phlebia*. *Chemosphere.* 85:218–224.
- Xue N, Zhang D, Xu X. 2006. Organochlorinated pesticide multiresidues in surface sediments from Beijing Guanting reservoir. *Water Res.* 40:183–194.
- Yang Y, Li D, Mu D. 2008. Levels, seasonal variations and sources of organochlorine pesticides in ambient air of Guangzhou, China. *Atmos Environ.* 42:677–687.

1
2
3
4
5
6
71346 Zeng S, Qin X, Xia L. 2017. Degradation of the herbicide isoproturon by laccase-mediator systems.
81347 Biochem Eng J. 119:92–100.
9
101348 Zhang J, Qin J, Zhao C, Liu C, Xie H, Liang S. 2015. Response of bacteria and fungi in soil
111349 microcosm under the presence of pesticide endosulfan. Water Air Soil Pollut. Available from:
121350 <http://link.springer.com/10.1007/s11270-015-2309-6>
131351
141352 Zhou Q, Hua T. 2004. Bioremediation: a review of applications and problems to be resolved. Prog
151351 Nat Sci. 14:937–944.
161352
171353 Zhou R, Zhu L, Yang K, Chen Y. 2006. Distribution of organochlorine pesticides in surface water
181354 and sediments from Qiantang River, East China. J Hazard Mater. 137:68–75.
191355
201356 Zielke H, Seiler T-B, Niebergall S, Leist E, Brinkmann M, Spira D, Streck G, Brack W, Feiler U,
211357 Braunbeck T. 2011. The impact of extraction methodologies on the toxicity of sediments in the
221358 zebrafish (*Danio rerio*) embryo test. J Soils Sediments. 11:352–363.
231359
241360
251361
261362
271363
281364
291365
301366
311367
321368
331369
341370
351371
361372
371373
381374
391375
401376
411377
421378
431379
441380
451381
461382
471383
481384
491385
501386
511387
521388
531389
541390
551391
561392
571393
581394
591395
601396

Supplemental material: Table I

Table I. Fungal species list for biodegradation of pesticide pollutants

Pesticide types	target pesticide	Fungal species	Fungal habitats	Origin	Literature
organochlorine	aldrin	<i>Phanerochete chrysosporium</i>	white-rot		Kennedy et al 1990
	chlordan	<i>Phanerochete chrysosporium</i>	white-rot		Kennedy et al 1990
	DDT	<i>Phanerochete chrysosporium</i>	white-rot		Arisoy 1998
	DDT	<i>Pleurotus sajor-caju</i>	white-rot		Arisoy 1998
	DDT	<i>Pleurotus florida</i>	white-rot		Arisoy 1998
	DDT	<i>Pleurotus eryngi</i>	white-rot		Arisoy 1998
	DDT	<i>Gloeophyllum trabeum</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Gloeophyllum sepiarium</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Gloeophyllum unguatum</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Gloeophyllum striatum</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Daedalea malicola</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Daedalea albida</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Daedalea serialis</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Daedalea dickinsii</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Fomitopsis palustris</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Fomitopsis annosa</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Fomitopsis insularis</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Fomitopsis pinicola</i>	brown-rot		Purnomo et al 2008
	DDT	<i>Boletus edulis</i>	ectomycorrhizal		Huang et al 2007
	DDT	<i>Gomphidius viscidus</i>	ectomycorrhizal		Huang et al 2007
	DDT	<i>Laccaria bicolor</i>	ectomycorrhizal		Huang et al 2007
	DDT	<i>Leccinum scabrum</i>	ectomycorrhizal		Huang et al 2007
	DDT	<i>Trichoderma harzianum</i>	saprotrophic	field soil	Katayama and Matsumura 1993
	DDD	<i>Trichoderma sp.</i>	saprotrophic	marine sponges	Ortega et al 2011
	DDD	<i>Penicillium miczynskii</i>	saprotrophic	marine sponges	Ortega et al 2011
	dieldrin	<i>Trichoderma harzianum</i>	saprotrophic	field soil	Katayama and Matsumura 1993

dielldrin	<i>Phanerochete chrysosporium</i>	white-rot		Kennedy et al 1990
endosulfan	<i>Trichoderma harzianum</i>	saprotrophic	field soil	Katayama and Matsumura 1993
endosulfan	<i>Phanerochaete chrysosporium</i>	white-rot		Kullman and Matsumura 1996
heptachlor	<i>Phanerochete chrysosporium</i>	white-rot		Arisoy 1998
heptachlor	<i>Pleurotus sajor-caju</i>	white-rot		Arisoy 1998
heptachlor	<i>Pleurotus florida</i>	white-rot		Arisoy 1998
heptachlor	<i>Pleurotus eryngi</i>	white-rot		Arisoy 1998
pentachloronitrobenzene	<i>Trichoderma harzianum</i>	saprotrophic	field soil	Katayama and Matsumura 1993
pentachlorophenol(PCP)	<i>Trichoderma harzianum</i>	saprotrophic	field soil	Katayama and Matsumura 1993
pentachlorophenol(PCP)	<i>Phanerochaete chrysosporium</i>	white-rot		Kang and Stevens 1994
pentachlorophenol(PCP)	<i>Pleurotus ostreatus</i>	white-rot		Rüttimann-Johnson and Lamar 1997
pentachlorophenol(PCP)	<i>Irpex lacteus</i>	white-rot		Rüttimann-Johnson and Lamar 1997
pentachlorophenol(PCP)	<i>Trametes versicolor</i>	white-rot		Rüttimann-Johnson and Lamar 1997
pentachlorophenol(PCP)	<i>Bjerkandera adusta</i>	white-rot		Rüttimann-Johnson and Lamar 1997
pendimethalin	<i>Fusarium oxysporum</i>	saprotrophic	soil	Singh and Kulshreya 1991
pendimethalin	<i>Paecilomyces varioti</i>	saprotrophic	soil	Singh and Kulshreya 1991
pendimethalin	<i>Rhizoctonia bataticola</i>	saprotrophic	soil	Singh and Kulshreya 1991
lindane	<i>Rhizopus oryzae</i>	saprotrophic		Young and Banks 1998
lindane	<i>Phanerochete chrysosporium</i>	white-rot		Arisoy 1998
lindane	<i>Pleurotus sajor-caju</i>	white-rot		Arisoy 1998
lindane	<i>Pleurotus florida</i>	white-rot		Arisoy 1998
lindane	<i>Pleurotus eryngi</i>	white-rot		Arisoy 1998
mirex	<i>Phanerochete chrysosporium</i>	white-rot		Kennedy et al 1990
PCB 77	<i>Phanerochaete chrysosporium</i>	white-rot		Vyas et al 1994

	PCB 77	<i>Trametes versicolor</i>	white-rot		Vyas et al 1994
	PCB 77	<i>Coriolopsis polyzona</i>	white-rot		Vyas et al 1994
	Delor 106 (PCB)	<i>Phanerochaete chrysosporium</i>	white-rot		Novotný et al 1997
	Delor 106 (PCB)	<i>Trametes versicolor</i>	white-rot		Novotný et al 1997
	Delor 106 (PCB)	<i>Coriolopsis polyzona</i>	white-rot		Novotný et al 1997
	Six PCB congeners	<i>Trametes versicolor</i>	white-rot		Beaudette et al 2000
	Six PCB congeners	<i>Bjerkandera adusta</i>	white-rot		Beaudette et al 2000
	Six PCB congeners	<i>Phanerochaete chrysosporium</i>	white-rot		Beaudette et al 2000
organophosphate	chlorthaloprophos	<i>Phanerochaete chrysosporium</i>	white-rot		Bumpus et al 1993
	chlorthaloprophos	<i>Hypholoma fasciculare</i>	white-rot		Bending et al 2002
	chlorthaloprophos	<i>Coriolus versicolor</i>	white-rot		Bending et al 2002
	chlorthaloprophos	<i>Trichoderma harzianum</i>	saprotrophic	soil	Omar 1998
	chlorthaloprophos	<i>Penicillium brevicompactum</i>	saprotrophic	soil	Omar 1998
	fonofos	<i>Phanerochaete chrysosporium</i>	white-rot		Bumpus et al 1993
	glyphosate	<i>Penicillium citrium</i>	saprotrophic		Zboinska et al 1992
	methyl parathion	<i>Aspergillus sydowii</i>	saprotrophic	marine	Alvarenga et al 2014
	methyl parathion	<i>Penicillium decaturense</i>	saprotrophic	marine	Alvarenga et al 2014
	terbufos	<i>Phanerochaete chrysosporium</i>	white-rot		Bumpus et al 1993
herbicide	alachlor	<i>Phanerochaete chrysosporium</i>	white-rot		Ferrey et al 1994
	alachlor	<i>Ceriporiopsis subvermispora</i>	white-rot		Ferrey et al 1994
	alachlor	<i>Phlebia tremellosa</i>	white-rot		Ferrey et al 1994
	alachlor	<i>Cunninghamella elegans</i>			Pothuluri et al 1993
	aroachlor	<i>Pleurotus ostreatus</i>	white-rot		Zeddel et al 1993
	aroachlor	<i>Trametes versicolor</i>	white-rot		Zeddel et al 1993
	three aroclors	<i>Phanerochaete chrysosporium</i>	white-rot		Yadav et al 1995
	atrazine	<i>Phanerochaete chrysosporium</i>	white-rot		Mougin et al 1994
	atrazine	<i>Pleurotus pulmonarius</i>	white-rot		Masaphy 1993
	atrazine	<i>Agrocybe semiorbicularis</i>	white-rot		Bending et al 2002
	atrazine	<i>Auricularia auriccola</i>	white-rot		Bending et al 2002
	atrazine	<i>Coriolus versicolor</i>	white-rot		Bending et al 2002

		atrazine	<i>Dichotomitus squalens</i>	white-rot	Bending et al 2002
		atrazine	<i>Flammulina velupites</i>	white-rot	Bending et al 2002
		atrazine	<i>Hypholoma fasciculare</i>	white-rot	Bending et al 2002
		atrazine	<i>Phanerochaete velutina</i>	white-rot	Bending et al 2002
		atrazine	<i>Pleurotus ostreatus</i>	white-rot	Bending et al 2002
		atrazine	<i>Stereum hirsutum</i>	white-rot	Bending et al 2002
			<i>Agrocybe</i>		
		diuron	<i>semiorbicularis</i>	white-rot	Bending et al 2002
		diuron	<i>Hypholoma fasciculare</i>	white-rot	Bending et al 2002
		diuron	<i>Stereum hirsutum</i>	white-rot	Bending et al 2002
		diuron	<i>Coriolus versicolor</i>	white-rot	Bending et al 2002
	fungicide	carbendazim	<i>Trichoderma sp.</i>	saprotrophic	mutant strain Tian and Chen 2009
		metalaxyl	<i>Coriolus versicolor</i>	white-rot	Bending et al 2002
		metalaxyl	<i>Stereum hirsutum</i>	white-rot	Bending et al 2002
		iprodione	<i>Hypholoma fasciculare</i>	white-rot	Bending et al 2002
		iprodione	<i>Stereum hirsutum</i>	white-rot	Bending et al 2002
		iprodione	<i>Coriolus versicolor</i>	white-rot	Bending et al 2002
	PAH	five PAHs	<i>Bjerkandera adusta</i>	white-rot	soil and lignite Gramss et al 1995
		five PAHs	<i>Gymnophilus sapineus</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Hypholoma fasciculare</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Hypholoma frowardii</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Hypholoma sublateritium</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Kuehneromyces mutabilis</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Lenzites betulina</i>	Wood-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Pleurotus ostreatus</i>	white-rot	soil and lignite Gramss et al 1995
		five PAHs	<i>Agrocybe praecox</i>	Wood- and straw-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Stropharia coronilla</i>	Wood- and straw-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Stropharia rugoso-annulata</i>	Wood- and straw-degrading	soil and lignite Gramss et al 1995
		five PAHs	<i>Agaricus aestivalis</i>	Terricolous	soil and lignite Gramss et al 1995
		five PAHs	<i>Agaricus arvensis</i>	Terricolous	soil and lignite Gramss et al 1995

five PAHs	<i>Agaricus bisporus</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Agaricus campestris</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Agaricus porphyizon</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Agrocybe dura</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Bovisa nigrescens</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Clitocybe odora</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Collybia dyophila</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Collybia maculata</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Coprinus comatus</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Lepista nebularis</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Lepista nuda</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Lepista saeva</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Lycoperdon perlatum</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Marasmius oreades</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Megacollybia platyphylla</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Phallus impudicus</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Psathyrella velutina</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Stropharia aeruginosa</i>	Terricolous	soil and lignite	Gramss et al 1995
five PAHs	<i>Amanita muscaria</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Amanita rubescens</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Amanita spissa</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Hebeloma crustuliniforme</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Hebeloma hiemale</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995

five PAHs	<i>Hebeloma sinapizans</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Laccaria amethystina</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Lactarius deliciosus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Lactarius deterrimus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Lactarius rufus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Lactarius torminosus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Morchella conica</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Morchella elata</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Morchella esculenta</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Paxillus involutus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Russula aeruginea</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Russula foetens</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Suillus granulatus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Suillus variegatus</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Tricholoma lascivum</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Tricholoma terreum</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Xerocomus badius</i>	Ectomycorrhizal	soil and lignite	Gramss et al 1995
five PAHs	<i>Botrytis cinerea</i>	Mitosporic	soil and lignite	Gramss et al 1995
five PAHs	<i>Scytalidium lignicola</i>	saprotrophic	soil and lignite	Gramss et al 1995
five PAHs	<i>Trichoderma sp.</i>	saprotrophic	soil and lignite	Gramss et al 1995

Supplemental material – Table I: References

- Alvarenga N, Birolli WG, Selegim MHR, Porto ALM. 2014. Biodegradation of methyl parathion by whole cells of marine-derived fungi *Aspergillus sydowii* and *Penicillium decaturense*. Chemosphere. 117:47–52.
- Arisoy M. 1998. Biodegradation of chlorinated organic compounds by white-rot fungi. Bull Environ Contam Toxicol. 60:872–876.
- Beaudette LA, Ward OP, Pickard MA, Fedorak PM. 2000. Low surfactant concentration increases fungal mineralization of a polychlorinated biphenyl congener but has no effect on overall metabolism. Lett Appl Microbiol. 30:155–160.
- Bending GD, Friloux M, Walker A. 2002. Degradation of contrasting pesticides by white rot fungi and its relationship with ligninolytic potential. FEMS Microbiol Lett. 212:59–63.
- Bumpus JA, Kakar SN, Coleman RD. 1993. Fungal degradation of organophosphorous insecticides. Appl Biochem Biotechnol. 39-40:715–726.
- Ferrey ML, Koskinen WC, Blanchette RA, Burnes TA. 1994. Mineralization of alachlor by lignin-degrading fungi. Can J Microbiol. 40:795–798.
- Gramss G, Kirsche B, Voigt K-D, Günther T, Fritsche W. 1999. Conversion rates of five polycyclic aromatic hydrocarbons in liquid cultures of fifty-eight fungi and the concomitant production of oxidative enzymes. Mycol Res. 103:1009–1018.
- Huang Y, Zhao X, Luan S. 2007. Uptake and biodegradation of DDT by 4 ectomycorrhizal fungi. Sci Total Environ. 385:235–241.
- Kang G, Stevens DK. 1994. Degradation of pentachlorophenol in bench scale bioreactors using the white rot fungus *Phanerochaete chrysosporium*. Hazard Waste Hazard Mater. 11:397–410.
- Katayama A, Matsumura F. 1993. Degradation of organochlorine pesticides, particularly endosulfan, by *Trichoderma harzianum*. Environ Toxicol Chem. 12:1059–1065.
- Kennedy DW, Aust SD, Bumpus JA. 1990. Comparative biodegradation of alkyl halide insecticides by the white rot fungus, *Phanerochaete chrysosporium* (BKM-F-1767). Appl Environ Microbiol. 56:2347–2353.

- Kullman SW, Matsumura F. 1996. Metabolic pathways utilized by *Phanerochaete chrysosporium* for degradation of the cyclodiene pesticide endosulfan. *Appl Environ Microbiol.* 62:593–600.
- Masaphy S, Levanon D, Vaya J, Henis Y. 1993. Isolation and characterization of a novel atrazine metabolite produced by the fungus *Pleurotus pulmonarius*, 2-chloro-4-ethylamino-6-(1-hydroxyisopropyl)amino-1,3,5-triazine. *Appl Environ Microbiol.* 59:4342–4346.
- Mougin C, Laugero C, Asther M, Dubroca J, Frasse P, Asther M. 1994. Biotransformation of the herbicide atrazine by the white rot fungus *Phanerochaete chrysosporium*. *Appl Environ Microbiol.* 60:705–708.
- Novotný Č, Vyas BRM, Erbanová P, Kubátová A, Šašek V. 1997. Removal of PCBs by various white rot fungi in liquid cultures. *Folia Microbiol (Praha).* 42:136–140.
- Omar SA. 1998. Availability of phosphorus and sulfur of insecticide origin by fungi. *Biodegradation.* 9:327–336.
- Ortega SN, Nitschke M, Mouad AM, Landgraf MD, Rezende MOO, Seleglim MHR, Sette LD, Porto ALM. 2011. Isolation of Brazilian marine fungi capable of growing on DDD pesticide. *Biodegradation.* 22:43–50.
- Pothuluri JV, Freeman JP, Evans FE, Moorman TB, Cerniglia CE. 1993. Metabolism of alachlor by the fungus *Cunninghamella elegans*. *J Agric Food Chem.* 41:483–488.
- Purnomo AS, Kamei I, Kondo R. 2008. Degradation of 1,1,1-trichloro-2,2-bis(4-chlorophenyl) ethane (DDT) by brown-rot fungi. *J Biosci Bioeng.* 105:614–621.
- Rüttimann-Johnson C, Lamar RT. 1997. Binding of pentachlorophenol to humic substances in soil by the action of white rot fungi. *Soil Biol Biochem.* 29:1143–1148.
- Singh SB, Kulshrestha G. 1991. Microbial degradation of pendimethalin. *J Environ Sci Health Part B.* 26:309–321.
- Tian L, Chen F. 2009. Characterization of a carbendazim-degrading *Trichoderma* sp. T2-2 and its application in bioremediation. *Wei Sheng Wu Xue Bao.* 49:925–930.
- Vyas BRM, Šašek V, Matucha M, Bubner M. 1994. Degradation of 3,3',4,4'-tetrachlorobiphenyl by selected white rot fungi. *Chemosphere.* 28:1127–1134.

Yadav JS, Quensen JF, Tiedje JM, Reddy CA. 1995. Degradation of polychlorinated biphenyl mixtures (Aroclors 1242, 1254, and 1260) by the white rot fungus *Phanerochaete chrysosporium* as evidenced by congener-specific analysis. *Appl Environ Microbiol.* 61:2560–2565.

Young E, Banks CJ. 1998. The removal of lindane from aqueous solution using a fungal biosorbent: the influence of pH, temperature, biomass concentration, and culture Age. *Environ Technol.* 19:619–625.

Zbońska E, Maliszewska I, Lejczak B, Kafarski P. 1992. Degradation of organophosphonates by *Penicillium citrinum*. *Lett Appl Microbiol.* 15:269–272.

Zeddel A, Majcherczyk A, Hüttermann A. 1993. Degradation of polychlorinated biphenyls by white-rot fungi *Pleurotus ostreatus* and *Trametes versicolor* in a solid state system. *Toxicol Environ Chem.* 40:255–266.

Supplemental material: Table II

Table II. Fungi and their enzymes capable of transforming OACs; whole-cell and enzymatic treatments are reported

Whole-cell treatment			
Fungal species	Pesticide	Enzymes involved	Literature
<i>Aspergillus niger</i>	nicosulfuron		Lu et al. 2012
<i>Auricularia fuscusuccinea</i>	endosulfan	laccase, phenol oxidase	Yanez-Montalvo et al. 2016
<i>Aspergillus sydowii</i> , <i>Penicillium decaturense</i>	methyl parathion		Alvarenga et al. 2014
<i>Aspergillus sydowii</i> , <i>Penicillium raistrickii</i> , <i>Cladosporium sp.</i> , <i>Microspheeropsis sp.</i> , <i>Acremonium sp.</i> , <i>Westerdykella sp.</i> , <i>Cladosporium sp.</i>	esfenvalerate		Birrolli et al. 2016
<i>Aspergillus fumigatus</i> , <i>Aspergillus terreus</i> , <i>Penicillium citrinum</i> , <i>Trichoderma harzianum</i>	chlorfenvinphos		Oliveira et al. 2015
<i>Aspergillus oryzae</i>	3-phenoxybenzoic acid		Zhu et al. 2016

<i>Aspergillus oryzae</i> , <i>Fusarium oxysporum</i> , <i>Lentinula edodes</i> , <i>Penicillium brevicompactum</i> , <i>Lecanicillium saksenae</i>	terbuthylazine, difenoconazole and pendimethalin		Pinto et al. 2012
<i>Aspergillus sydowii</i>	trichlorfon		Tian et al. 2016
<i>Aspergillus versicolor</i>	triclosan		Taştan and Dönmez 2015
<i>Coriolus versicolor</i>	aldicarb, atrazine, alachlor		Hai et al. 2012
<i>Dacryopinax elegans</i>	diuron	laccase, manganese peroxidase, lignin peroxidase	Arakaki et al. 2013
<i>Ganoderma lucidum</i>	lindane	laccase, manganese peroxidase, lignin peroxidase	Kaur et al. 2016
<i>Ganoderma lucidum</i>	bentazon	laccase, manganese peroxidase	Da Silva Coelho et al. 2010
<i>Ganoderma lucidum</i> , <i>Trametes sp</i>	picloram	laccase	Maciel et al. 2013
<i>Gloeophyllum trabeum</i> , <i>Trametes versicolor</i> , <i>Pleurotus ostreatus</i>	lindane, endosulfan		Ulčnik et al. 2013
<i>Mycelia sterilia</i>	atrazine	laccase	Vasil'chenko et al. 2002
<i>Penicillium citrinum</i> , <i>P.citrinum</i> , <i>Fusarium proliferatum</i>	methylparathion		Rodrigues et al. 2016
<i>Penicillium griseofulvum</i>	b-hexachlorocyclohexane		Ceci et al. 2015
<i>Phanerochaete sordida</i>	clothianidin	cytochrome P450, manganese peroxidase	Mori et al. 2017
<i>Pleurotus pulmonarius</i>	atrazine		Masaphy et al. 1993
<i>Phlebia tremellosa</i> , <i>Phlebia brevispora</i> , <i>Phlebia acanthocystis</i>	Heptachlor, heptachlor epoxide		Xiao et al. 2011
<i>Saccharomyces cerevisiae</i>	diazinon		Ehrampoush et al. 2017
<i>Talaromyces flavus</i>	nicosulfuron		Song et al. 2013
<i>Trametes versicolor</i>	imiprothrin, cypermethrin, carbofuran, oxytetracycline	laccase, cytochrome P450	Mir-Tutusaus et al. 2014
<i>Trametes versicolor</i>	fipronil	cytochrome P450	Wolfand et al. 2016
<i>Trametes versicolor</i>	6 pesticides, 2 phytoestrogens		Nguyen et al. 2014
<i>Trametes versicolor</i> , <i>Stereum hirsutum</i>	linuron, dimethoate		Castellana and Loffredo 2014
nonsporulating mycelial fungus	atrazine	cellobiose dehydrogenase	Khromonygina et al. 2004
Enzymatic treatment			

Enzymes involved	Pesticide	Literature
laccases of <i>Agaricus blazei</i>	metsulfuron	González Matute et al. 2012
phytase of <i>Aspergillus niger</i>	chlorpyrifos	Shah et al. 2017
extracellular extract of <i>Auricularia fuscusuccinea</i>	endosulfan	Yanez-Montalvo et al. 2016
laccase of <i>Trametes versicolor</i>	sulfamethoxazole, isoproturon	Margot et al. 2015
laccase of <i>Trametes versicolor</i>	chloroxuron	Palvannan et al. 2014
laccase of <i>Trametes versicolor</i>	lindane, endosulfan	Ulčnik et al. 2013
cellulose of <i>Trichoderma longibrachiatum</i>	dicofol	Wang et al. 2015
laccase of <i>Trametes versicolor</i>	isoproturon	Zeng et al. 2017

Supplemental material – Table II: References

Alvarenga N, Birololi WG, Seleghim MHR, Porto ALM. 2014. Biodegradation of methyl parathion by whole cells of marine-derived fungi *Aspergillus sydowii* and *Penicillium decaturense*. Chemosphere. 117:47–52.

Arakaki RL, Monteiro DA, Boscolo M, Dasilva R, Gomes E. 2013. Halotolerance, ligninase production and herbicide degradation ability of basidiomycetes strains. Braz J Microbiol. 44:1207–1214.

Birololi WG, Alvarenga N, Seleghim MHR, Porto ALM. 2016. Biodegradation of the pyrethroid pesticide esfenvalerate by marine-derived fungi. Mar Biotechnol. 18:511–520.

Castellana G, Loffredo E. 2014. Simultaneous removal of endocrine disruptors from a wastewater using white rot fungi and various adsorbents. Water Air Soil Pollut. 225. Available from: <http://link.springer.com/10.1007/s11270-014-1872-6>

Ceci A, Pierro L, Riccardi C, Pinzari F, Maggi O, Persiani AM, Gadd GM, Petrangeli Papini M. 2015. Biotransformation of β -hexachlorocyclohexane by the saprotrophic soil fungus *Penicillium griseofulvum*. Chemosphere. 137:101–107.

Da Silva Coelho J, de Oliveira AL, Marques de Souza CG, Bracht A, Peralta RM. 2010. Effect of the herbicides bentazon and diuron on the production of ligninolytic enzymes by *Ganoderma lucidum*. Int Biodeterior Biodegrad. 64:156–161.

Ehrampoush MH, Sadeghi A, Ghaneian MT, Bonyadi Z. 2017. Optimization of diazinon biodegradation from aqueous solutions by *Saccharomyces cerevisiae* using response surface methodology. AMB Express. 7. Available from: <http://amb-express.springeropen.com/articles/10.1186/s13568-017-0366-5>

- González Matute R, Figlas D, Mockel G, Curvetto N. 2012. Degradation of metsulfuron methyl by *Agaricus blazei* Murrill spent compost enzymes. *Bioremediation J.* 16:31–37.
- Hai FI, Modin O, Yamamoto K, Fukushi K, Nakajima F, Nghiem LD. 2012. Pesticide removal by a mixed culture of bacteria and white-rot fungi. *J Taiwan Inst Chem Eng.* 43:459–462.
- Kaur H, Kapoor S, Kaur G. 2016. Application of ligninolytic potentials of a white-rot fungus *Ganoderma lucidum* for degradation of lindane. *Environ Monit Assess.* 188:588.
- Khromonygina VV, Saltykova AI, Vasil'chenko LG, Kozlov IP, Rabinovich ML. 2004. Degradation of the herbicide atrazine by the soil mycelial fungus INBI 2-26(-)--a producer of cellobiose dehydrogenase. *Prikl Biokhim Mikrobiol.* 40:337–343.
- Lu X, Kang Z, Tao B, Wang Y, Dong J, Zhang J. 2012. Degradation of nicosulfuron by *Bacillus subtilis* YB1 and *Aspergillus niger* YF1. *Appl Biochem Microbiol.* 48:460–466.
- Maciel GM, Inácio FD, de Sá-Nakanishi AB, Haminiuk CWI, Castoldi R, Comar JF, Bracht A, Peralta RM. 2013. Response of *Ganoderma lucidum* and *Trametes* sp. to the herbicide picloram: Tolerance, antioxidants and production of ligninolytic enzymes. *Pestic Biochem Physiol.* 105:84–92.
- Margot J, Copin P-J, von Gunten U, Barry DA, Holliger C. 2015. Sulfamethoxazole and isoproturon degradation and detoxification by a laccase-mediator system: Influence of treatment conditions and mechanistic aspects. *Biochem Eng J.* 103:47–59.
- Masaphy S, Levanon D, Vaya J, Henis Y. 1993. Isolation and characterization of a novel atrazine metabolite produced by the fungus *Pleurotus pulmonarius*, 2-chloro-4-ethylamino-6-(1-hydroxyisopropyl)amino-1,3,5-triazine. *Appl Environ Microbiol.* 59:4342–4346.
- Mir-Tutusa JA, Masís-Mora M, Corcellas C, Eljarrat E, Barceló D, Sarrà M, Caminal G, Vicent T, Rodríguez-Rodríguez CE. 2014. Degradation of selected agrochemicals by the white rot fungus *Trametes versicolor*. *Sci Total Environ.* 500–501:235–242.
- Mori T, Wang J, Tanaka Y, Nagai K, Kawagishi H, Hirai H. 2017. Bioremediation of the neonicotinoid insecticide clothianidin by the white-rot fungus *Phanerochaete sordida*. *J Hazard Mater.* 321:586–590.
- Nguyen LN, Hai FI, Yang S, Kang J, Leusch FDL, Roddick F, Price WE, Nghiem LD. 2014. Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*. Role of biosorption and biodegradation. *Int Biodeterior Biodegrad.* 88:169–175.
- Oliveira BR, Penetra A, Cardoso VV, Benoliel MJ, Barreto Crespo MT, Samson RA, Pereira VJ. 2015. Biodegradation of pesticides using fungi species found in the aquatic environment. *Environ Sci Pollut Res Int.* 22:11781–11791.
- Palvannan T, Saravanakumar T, Unnithan AR, Chung N-J, Kim D-H, Park S-M. 2014. Efficient transformation of phenyl urea herbicide chloroxuron by laccase immobilized on zein polyurethane nanofiber. *J Mol Catal B Enzym.* 99:156–162.
- Pinto AP, Serrano C, Pires T, Mestrinho E, Dias L, Teixeira DM, Caldeira AT. 2012. Degradation of terbutylazine, difenoconazole and pendimethalin pesticides by selected fungi cultures. *Sci Total Environ.* 435–436:402–410.

- Rodrigues GN, Alvarenga N, Vacondio B, de Vasconcellos SP, Passarini MRZ, Seleglim MHR, Porto ALM. 2016. Biotransformation of methyl parathion by marine-derived fungi isolated from ascidian *Didemnum ligulum*. *Biocatal Agric Biotechnol*. 7:24–30.
- Shah PC, Kumar VR, Dastager SG, Khire JM. 2017. Phytase production by *Aspergillus niger* NCIM 563 for a novel application to degrade organophosphorus pesticides. *AMB Express*. 7:66.
- Song J, Gu J, Zhai Y, Wu W, Wang H, Ruan Z, Shi Y, Yan Y. 2013. Biodegradation of nicosulfuron by a *Talaromyces flavus* LZM1. *Bioresour Technol*. 140:243–248.
- Taştan BE, Dönmez G. 2015. Biodegradation of pesticide triclosan by *A. versicolor* in simulated wastewater and semi-synthetic media. *Pestic Biochem Physiol*. 118:33–37.
- Tian J, Dong Q, Yu C, Zhao R, Wang J, Chen L. 2016. Biodegradation of the organophosphate trichlorfon and its major degradation products by a novel *Aspergillus sydowii* PA F-2. *J Agric Food Chem*. 64:4280–4287.
- Ulčnik A, Kralj Cigić I, Pohleven F. 2013. Degradation of lindane and endosulfan by fungi, fungal and bacterial laccases. *World J Microbiol Biotechnol*. 29:2239–2247.
- Vasil'chenko L, Khromonygina V, Koroleva O, Landesman E, Gaponenko V, Kovaleva T, Kozlov YP, Rabinovich M. 2002. Consumption of triazine herbicide atrazine by laccase-positive and laccase-negative strains of soil fungus *Mycelia sterilia* INBI 2-26. *Appl Biochem Microbiol*. 38:454–459.
- Wang Z, Yang T, Zhai Z, Zhang B, Zhang J. 2015. Reaction mechanism of dicofol removal by cellulase. *J Environ Sci China*. 36:22–28.
- Wolfand JM, LeFevre GH, Luthy RG. 2016. Metabolization and degradation kinetics of the urban-use pesticide fipronil by white rot fungus *Trametes versicolor*. *Environ Sci Process Impacts*. 18:1256–1265.
- Xiao P, Mori T, Kamei I, Kiyota H, Takagi K, Kondo R. 2011. Novel metabolic pathways of organochlorine pesticides dieldrin and aldrin by the white rot fungi of the genus *Phlebia*. *Chemosphere*. 85:218–224.
- Yanez-Montalvo AF, Sánchez JE, Rafael V-D. 2016. Degradation of endosulfan by strains of *Auricularia fuscusuccinea*. *Sydowia Int J Mycol*.:7–15.
- Zeng S, Qin X, Xia L. 2017. Degradation of the herbicide isoproturon by laccase-mediator systems. *Biochem Eng J*. 119:92–100.
- Zhu Y, Li J, Yao K, Zhao N, Zhou K, Hu X, Zou L, Han X, Liu A, Liu S. 2016. Degradation of 3-phenoxybenzoic acid by a filamentous fungus *Aspergillus oryzae* M-4 strain with self-protection transformation. *Appl Microbiol Biotechnol*. 100:9773–9786.