

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**A review and meta-analysis of the effects of climate change on Holarctic mountain and upland bird populations**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1666682> since 2018-11-06T13:05:21Z

*Published version:*

DOI:10.1111/ibi.12585

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[Ibis, 160: 484-519, 2018, DOI: 10.1111/ibi.12585]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*<https://onlinelibrary.wiley.com/doi/full/10.1111/ibi.12585>*

20 **The Effects of Climate Change on Holarctic Mountain and Upland Bird**  
21 **Populations: A Review and Meta-Analysis**

22  
23  
24 D. SCRIDEL,<sup>1,2,3</sup> M. BRAMBILLA,<sup>1,4</sup> K. MARTIN<sup>5</sup>, A. LEHIKONEN<sup>6</sup>, A. IEMMA<sup>1</sup>, M. ANDERLE<sup>1</sup>, S.  
25 JÄHNIG<sup>7</sup>, E. CAPRIO, E.<sup>7</sup>, G. BOGLIANI<sup>2</sup>, P. PEDRINI<sup>1</sup>, A. ROLANDO<sup>7</sup>, R. ARLETTAZ<sup>8,9</sup> & D. E.  
26 CHAMBERLAIN<sup>7</sup>  
27  
28  
29

30 <sup>1</sup> *Museo delle Scienze di Trento (MUSE), Sezione Zoologia dei Vertebrati, Corso del Lavoro e della*  
31 *Scienza 3, 38122 Trento, Italy*

32 <sup>2</sup> *Dipartimento di Scienze della Terra e dell'Ambiente, Università degli Studi di Pavia, Pavia, Italy*

33 <sup>3</sup> *Ente Parco Naturale Paneveggio Pale di San Martino, loc. Castelpietra, 2 - Tonadico (TN) – Italy*

34 <sup>4</sup> *Fondazione Lombardia per l'Ambiente, Settore Biodiversità e Aree Protette, Largo 10 Luglio 1976,*  
35 *1, 20822 Seveso, MB, Italy*

36 <sup>5</sup> *Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, BC,*  
37 *V6T 1Z4, Canada*

38 <sup>6</sup> *The Helsinki Lab of Ornithology (HeLO), Finnish Museum of Natural History, University of*  
39 *Helsinki, Helsinki FI-00014, Finland*

40 <sup>7</sup> *Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Via Accademia*  
41 *Albertina 13, 10123, Turin, Italy*

42 <sup>8</sup> *University of Bern, Institute of Ecology and Evolution, Division of Conservation Biology,*  
43 *Baltzerstrasse 6, CH-3012 Bern, Switzerland*

44 <sup>9</sup> *Swiss Ornithological Institute, Valais Field Station, Nature Centre, CH-3970 Salgesch, Switzerland*  
45  
46

47 Corresponding author

48 Email: [davide.scridel01@universitadipavia.it](mailto:davide.scridel01@universitadipavia.it)  
49  
50

51 Running head: Climate change and mountain birds  
52  
53  
54  
55  
56  
57  
58

59 **Keywords:** Avian physiology, biotic interactions, conservation, elevational shift, global warming, high-  
60 elevation species, interspecific competition, phenology, population dynamics, projections, snow, trophic  
61 mismatch.  
62

63 Mountain regions are globally important areas for biodiversity, but are subject to multiple human-induced  
64 threats, including climate change, which has been more severe at higher elevations. We reviewed evidence  
65 for impacts of climate change on Holarctic mountain bird populations in terms of physiology, phenology,  
66 trophic interactions, demography, and observed and projected distribution shifts, including effects of other  
67 factors that interact with climate change. We developed an objective classification of high-elevation,  
68 mountain specialist and generalist species, based on the proportion of their breeding range occurring in  
69 mountain regions. Our review found evidence of responses of mountain bird populations to climatic  
70 (extreme weather events, temperature, rainfall and snow) and environmental (i.e. land use) change, but we  
71 know little about either the underlying mechanisms or about the synergistic effects of climate and land use.  
72 Long-term studies assessing reproductive success or survival of mountain birds in relation to climate  
73 change were rare. Few studies have considered shifts in elevational distribution over time and a meta-  
74 analysis did not find a consistent direction in elevational change. A meta-analysis carried out on future  
75 projections of distribution shifts suggested that birds whose breeding distributions are largely restricted to  
76 mountains are likely to be more negatively impacted than other species. Adaptation responses to climate  
77 change rely mostly on managing and extending current protected areas for both species already present,  
78 and for expected colonising species that are losing habitat and climatic space at lower elevation. However,  
79 developing effective management actions requires an improvement in the current knowledge of mountain  
80 species ecology, in the quality of climatic data, and in understanding the role of interacting factors.  
81 Furthermore, the evidence was mostly based on widespread species rather than mountain specialists.  
82 Scientists should provide valuable tools to assess the status of mountain birds, for example through the  
83 development of a mountain bird population index, while policy-makers should influence legislation to  
84 develop efficient agri-environment schemes and forestry practices for mountain birds, as well as to  
85 regulate leisure activities at higher elevations.

86

87 Climate change has been recognised, alongside modifications in land-use, as a key driver of global change in  
88 biological diversity (e.g. IPCC 2007, Ameztegui *et al.* 2016), and there is now a large body of evidence that  
89 animals and plants are responding to climate change through shifts in distribution (e.g. Chen *et al.* 2011),  
90 changes in population size (e.g. Stephens *et al.* 2016), and changes in phenology leading to inter-linked  
91 effects at different trophic levels (e.g. Both *et al.* 2006, Thackeray *et al.* 2016). Such effects vary  
92 geographically, and biodiversity in temperate, boreal and arctic regions is considered particularly vulnerable,  
93 with greater warming at higher latitudes (e.g. Meehl *et al.* 2007). Furthermore, rates of warming and  
94 frequency of extreme cold events are more pronounced at higher elevations (Beniston & Rebetez 1996, Liu  
95 & Chen 2010, Pepin *et al.* 2015). As a result, high-elevation areas are particularly threatened as they are  
96 more susceptible to changes in climate (Diaz *et al.* 2003, Böhning-Gaese & Lemoine 2004, La Sorte & Jetz  
97 2010).

98 Mountain and high latitude upland regions (henceforth 'mountains') cover around 25% of the Earth's  
99 surface (Kapos *et al.* 2000). They support one quarter of terrestrial biodiversity (Körner & Ohsawa 2006) and  
100 contain nearly half the world's biodiversity hotspots (Myers *et al.* 2000). These are complex ecosystems of  
101 high conservation value as they encapsulate a high diversity of small-scale habitats dictated by different  
102 topoclimates within narrow elevational gradients (Körner & Ohsawa 2006). As a result, mountains  
103 accommodate high levels of species diversity with heterogeneous communities adapted to specific  
104 environmental conditions that change along the elevational gradient, including climate and other abiotic  
105 factors such as slope, exposure, solar radiation, wind direction and substrate (Körner & Spehn 2002, Nagy &  
106 Grabherr 2009, Viterbi *et al.* 2013, Boyle & Martin 2015). For example, marked changes occur over short  
107 distances, with temperature varying in temperate regions on average by 0.6°C every 100m in elevation  
108 (Dillon *et al.* 2006). Aspect can also influence temperature, with greater solar radiation on southern than  
109 northern slopes in the Northern Hemisphere (Nagy & Grabherr 2009). Global warming is causing changes to  
110 these environments, with documented responses including the elevational advance of the tree-line and a  
111 general increase in dominance of woody deciduous shrubs at high elevations (Gehrig-Fasel *et al.* 2007,  
112 Myers-Smith *et al.* 2011).

113 Many unique ecological features of temperate mountain systems also arise from the strong  
114 seasonality in temperatures which result in a very short growing and reproductive season, typically less than  
115 three months in alpine-arctic and boreal habitats (Nagy & Grabherr 2009). Strong inter-annual variations in  
116 temperature, precipitation, and snow cover regimes are also observed in these systems (IPCC 2013, Klien *et al.*  
117 *et al.* 2016) with changes in the timing, quantity and duration of precipitation likely to influence mountain  
118 habitats and biodiversity (Beniston *et al.* 2003, IPCC 2013, Martin *et al.* 2017). Snow cover has been shown  
119 to have insulating properties, protecting plants and invertebrates from frost during the coldest months of  
120 the year and thereby influencing survival rates of many slow-growing high-elevation plants, insects and  
121 mammals (Hågvar 2010, Wipf & Rixen 2010, Berteaux *et al.* 2016). Snow cover in the Northern Hemisphere  
122 has declined since the 1920s, particularly in spring and summer (IPCC 2007).

123 Although often perceived as true wildlands, mountain ecosystems typically have a long history of  
124 human activity, especially in Europe and Asia (FAO 2015). Twenty percent of the global human population  
125 inhabits mountain regions, with about 8% living above 2500m (Körner & Ohsawa 2006). However, mountains  
126 provide essential ecosystem services, including nearly half of the human population's water supply, carbon  
127 storage and sequestration (forests and peatlands), and natural resources (timber, productive soils and  
128 medicinal plants; Körner & Ohsawa 2006). Mountains are also very important in terms of leisure and tourism  
129 activities (skiing, snowboarding, hiking, biking, wildlife watching and hunting). Mountain systems are thus  
130 continuously subject to landscape changes due to human activities, which might have more severe  
131 consequences than climate change itself (Jetz *et al.* 2007), or which could potentially exacerbate climatic  
132 effects (Mantyka-Pringle & Rhodes 2012).

133 Amongst birds, changes in climate have been reported to influence migration timing (Hüppop &  
134 Hüppop 2003, Knudsen *et al.* 2011), breeding output (Crick *et al.* 1997, Laaksonen *et al.* 2006), population  
135 size (Sæther *et al.* 2000, Townsen *et al.* 2016), and changes in elevational (Reif & Flousek 2012) and  
136 latitudinal (Hickling *et al.* 2006, Zuckerberg *et al.* 2009) distributions. Because the severity of climatic change  
137 varies over the Earth's surface (e.g. Meehl *et al.* 2007), avian responses may also vary in intensity depending  
138 on their geographic distribution. Birds may exhibit rapid distributional responses to climatic fluctuations, for

139 example tracking changes in surface temperature latitudinally (Hickling *et al.* 2006, Zuckerberg *et al.* 2009).  
140 However, there is also evidence that range shifts in birds are lagging behind climate change (Devictor *et al.*  
141 2008, Ralston *et al.* 2017), potentially due to asynchronous phenology of birds and their prey (e.g. Mayor *et al.*  
142 *et al.* 2017). Species inhabiting high-elevation mountain systems often exhibit a high degree of habitat  
143 specialisation and unique ecological traits within narrow thermal ranges (Reif & Flousek 2012, Reif *et al.*  
144 2015, Mahon *et al.* 2016, Pacifici *et al.* 2017, Scridel *et al.* 2017a). Adapting to rapid climate change may be  
145 particularly challenging along the elevational gradients of many mountains where temperatures and area  
146 decrease monotonically with elevation (Elsen & Tingley 2015). As a result, species tracking rising  
147 temperatures in these systems are predicted to decline according to the species-area relationship (Preston  
148 1962) as populations become isolated, and thus increasingly vulnerable to stochastic events (Lande 1993,  
149 Bech *et al.* 2009, Fjeldså *et al.* 2012). A successful shift into a new area by a species is possible only when  
150 abiotic as well biotic requirements are fulfilled (Martin 2001a, Heikkinen *et al.* 2007, Wilson & Martin 2012).  
151 Given the fast rate of warming, species might have to track temperatures in areas where their associated  
152 habitat and resources require longer to establish (e.g. mature trees, alpine and sub-nival plants; Engler *et al.*  
153 2011, Reif & Flousek 2012, Brambilla & Gobbi 2014), or where suitable habitat formation cannot occur due  
154 to constraints of other factors such as soil processes or rock substrate (Freppaz *et al.* 2010), or by direct  
155 human activities (e.g. deforestation; Nogués-Bravo *et al.* 2008, Patthey *et al.* 2008, Kohler *et al.* 2014;  
156 disturbance via outdoor recreation; Arlettaz *et al.* 2007, 2015). Finally, climatic effects coupled with negative  
157 synergistic changes in land use might pose even more severe constraints on adaptation of mountain birds to  
158 future climatic conditions.

159           Due to the documented general responses of birds, and the more extreme climatic changes  
160 observed in mountains, it seems reasonable to expect that mountain birds may be particularly threatened by  
161 climate change. In this review, we assess the existing evidence for direct and indirect effects of climate  
162 change on mountain birds in the Holarctic region (Heilprin 1887), and we evaluate their future conservation  
163 prospects. We address six specific objectives: (i) to define mountain generalist and high-elevation specialist  
164 birds for the Holarctic region; (ii) to review the impacts of climate change on mountain birds through a

165 summary of the literature, and a quantification of general responses throughout the Holarctic, including a  
166 meta-analysis; (iii) to review and quantify projected impacts from future climate change scenarios using a  
167 meta-analysis; (iv) to assess stressors that are likely to interact with climate change in affecting birds living at  
168 high elevations; (v) to review proposed conservation actions; and (vi) to identify current gaps and future  
169 priorities for research.

170

## 171 **METHODS**

172

### 173 **Defining mountain birds**

174 Mountain systems and species inhabiting them are difficult to describe geographically and ecologically, and  
175 definitions may not apply consistently across the globe (Strahler 1946, Gerrard 1990, Körner 2012, Scridel  
176 2014). In order to assess the status of mountain birds, it was first necessary to define mountain areas and  
177 habitats. Using elevational thresholds to define these regions would immediately exclude older and lower  
178 mountain systems such as the Urals, Scottish Highlands and Appalachians, and include areas with little  
179 topographic relief and few environmental gradients (e.g. large, high-elevation plateaux). Using slope as a  
180 criterion on its own or in combination with elevation may resolve the latter problem, but not the former. For  
181 these reasons, we adopted the definition of Kapos *et al.* (2000), who classified mountain systems in seven  
182 classes on the basis of elevation, slope and local elevation range (Fig. 1). The latter criterion is particularly  
183 useful as it identifies lower elevation mountain ranges (300-999m) by defining a radius of interest (5km)  
184 around each grid cell (30 arc-second) and measuring the maximum and minimum elevation within a  
185 particular neighborhood, and their difference, allowing the identification of areas that occur in regions with  
186 significant relief, even though elevations may not be especially high (Kapos *et al.* 2000). This is a broad  
187 definition which includes high latitude 'upland' habitats at relatively lower elevations, as well as mountain  
188 forest, the alpine belt (the treeless region between the natural climatic forest limit and the snow line) and  
189 the nival belt (the terrain above the snowline, which is defined as the lowest elevation where snow is



190 commonly present all year round; Kapos *et al.* 2000, Körner & Ohsawa 2006). Hereafter, we refer to  
191 ‘mountain regions’ as those as defined by Kapos *et al.* (2000).

192 We developed a broad definition of Holarctic mountain birds based on the proportion of their  
193 Holarctic breeding range that was within the defined mountain regions in order to assess the evidence base  
194 for impacts of climate change on birds largely restricted to mountains as a breeding habitat. We stress that  
195 we are interested in all bird species occurring in Holarctic mountain regions, including species that also occur  
196 in a range of habitats, rather than only focusing on high-elevation specialist species. We used a geographic  
197 information system (GIS) software (QGIS, Quantum GIS Development Team 2016; GRASS, GRASS  
198 Development Team 2015) to restrict the map of Kapos *et al.* (2000) to the Holarctic realm, and imposed over  
199 it the breeding range of global bird species ( $n = 10280$  species; BirdLife International & NatureServe 2015).  
200 We defined as ‘high-elevation mountain specialist’ a species for which at least 50% of its range was in the  
201 higher elevation classes 1-4 of Kapos *et al.* (2000). We further defined a ‘mountain generalist’ as a species  
202 for which at least 50% of its entire breeding range was within the defined Holarctic mountain region (i.e.  
203 classes 1-7 of Kapos *et al.* 2000) and which was not classed as a high-elevation mountain specialist. These  
204 definitions therefore identify broadly which species are particularly associated with mountains over the  
205 whole Holarctic region. There are many species (e.g. Capercaillie *Tetrao urogallus*, Pygmy Owl *Glaucidium*  
206 *passerinum*, Rock Ptarmigan *Lagopus muta*), termed boreo-alpine taxa, that occur in mountains at low  
207 latitudes (e.g. European Alps) which are also present at higher latitudes, but at lower elevations (e.g.  
208 northern Europe). The definition adopted here seeks to identify species that are linked more closely with  
209 mountains *per se* (for example due to topography or particular habitat types) across a broad region. We use  
210 the terms ‘high-elevation mountain specialist’ and ‘mountain generalist’ when specifically referring to our  
211 classification. We use the term ‘mountain bird’ to refer to any species occurring in our defined Holarctic  
212 mountain region, which also includes species which potentially breed in a range of habitats and at a range of  
213 elevations across their geographic range.

214

215 **Literature survey**

216 The literature search was generated through ISI Web of Knowledge ([www.webofknowledge.com](http://www.webofknowledge.com)). In order to  
217 obtain relevant studies we used the following keywords: (bird\* OR avian\*) AND (mountain\* OR montane\*  
218 OR upland\* OR alpine\* OR moorland\* OR arctic\* OR polar\* OR altitude\* OR elevation\*) AND (climate  
219 change\* OR global warming\*) NOT tropic\*. The search period was from 1950 until the 31st December 2016.  
220 Papers identified from this search were subsequently included if they concerned research wholly or partly  
221 carried out within the defined mountain regions, or if the study species was/were defined as a high-elevation  
222 mountain specialist or mountain generalist (see Supporting Information, Table S1 & S4). The latter group of  
223 studies included some broad-scale analyses that were not specifically focused on mountains, but which  
224 considered some high-elevation mountain specialists (typically analyses covering large regions, for example  
225 based on national atlases). A total of 764 studies was initially identified. The abstract of each of these papers  
226 was read to determine whether they were relevant for the purposes of this review, and 591 studies were  
227 eliminated at this stage. The remaining 173 papers were also checked for other relevant references missed  
228 in the previous search. This resulted in a further 61 relevant papers, giving a total of 234 which were  
229 subsequently assigned to eight broad topics: i) climate change, physiological constraints and life history  
230 strategies; ii) links between climate and population dynamics; iii) changes in phenology; iv) trophic linkages;  
231 v) observed evidence of elevational shift; vi) projected elevational shifts; vii) interactions between climate  
232 change and other drivers (agriculture, grazing and forestry, leisure and other threats, interspecific  
233 interactions); and, viii) conservation and policy papers. We used the standardized literature search to  
234 summarise the main trends in the resulting database in terms of location and topic, and also in terms of  
235 analysing elevational shifts and future projections of species' geographic range and population size. In detail,  
236 we conducted two meta-analyses: one testing whether mountain birds have shifted in elevation to track  
237 suitable climate, and a second one to test whether mountain birds will be more negatively impacted by  
238 climate change than non-mountain species according to projected distribution range and population size.  
239 We also used the selected papers, in conjunction with the wider literature, as the basis of a qualitative  
240 review to highlight the key issues and findings.

241

242 **Current and future elevational shifts in bird populations**

243 Papers that presented estimates for shifts in species distributions over time in relation to elevation were  
244 considered for meta-analyses if they focussed, either wholly or mostly, on the defined mountain regions.  
245 Given that conditions may change rapidly over small distances in mountains due to the steep topography,  
246 smaller-scale studies are more appropriate than larger-scale atlas studies in tracking species distributions  
247 (Chamberlain *et al.* 2012), and thus we focused on studies with a maximum sampling unit area of 1-km<sup>2</sup>.  
248 Additional data were collected for each study on the period considered (in years), the elevational range (in  
249 metres), and the estimated mean annual rate of temperature change (°C/year) over the period considered.  
250 Similarly, papers that predicted future effects of climate change on mountain birds were considered if they  
251 were largely restricted to mountain areas, if they estimated a proportional change in geographic distribution  
252 or population size over time, and if the sample size of the underlying data set on which models were based  
253 was presented. Additional variables recorded were the period over which projections were made, and the  
254 climate change scenarios considered, which were classed as either 'severe' (scenarios A2 and A1F1 or  
255 RCP8.5) or 'moderate' (all other scenarios and RCPs; IPCC 2007, 2013).

256

257 **Statistical analysis**

258 For elevational shifts, papers were included in the meta-analysis only if sample sizes and test statistics were  
259 presented, or if parameter estimates (including mean shift) and standard errors, standard deviations or  
260 confidence limits, were reported. In cases where only estimates of change in elevation and errors were  
261 presented (i.e. without any test statistics), z-scores were derived, testing against a hypothesis of zero  
262 change. In common with standard meta-analytical approaches (e.g. Koricheva *et al.* 2013), the goal was to  
263 estimate standardized responses of elevational shifts in bird distributions over time from studies that used a  
264 diversity of methods for quantifying a potential shift, which in most cases was calculated as the change, in  
265 meters, of the distribution of a given species (sometimes a group of species) between two time periods.  
266 However, some papers also tested the effect of the interaction between elevation and time period on the

267 probability of species presence, a significant interaction indicating a significant shift over time (e.g. Reif &  
268 Flousek 2012, Mizel *et al.* 2016).

269 Shifts in species distributions were tested using a variety of methods in the above papers. Different  
270 test statistics (e.g.  $F$ ,  $t$ ,  $\chi^2$ ) presented in these papers were converted to Pearson's  $r$  using standard  
271 conversion formulas (Lajeunesse 2013) so that effect sizes (i.e. change in elevational distribution over time)  
272 could be compared across studies (further details are given in the Supplementary material, S2. Meta-analysis  
273 methods). Pearson's  $r$  varied between 1 and -1, a positive value indicating an upslope shift in elevation over  
274 time. Pearson's  $r$  values were not normally distributed, so prior to analysis, standardized Pearson's  $r$  values  
275 from each study were transformed using Fisher's  $Z$  transformation to derive both normalized estimates and  
276 their variance (as per Musitelli *et al.* 2016).

277 Parameter estimates of standardized elevational shifts were derived by analysing  $Z$ -transformed  
278 Pearson's  $r$  values (henceforth 'standardized effects') and 95% confidence intervals based on linear mixed  
279 effects models using the nlme package in R (Pinheiro *et al.* 2017). The analytical unit was the estimate for a  
280 given species or group of species (some papers estimated shifts for the whole community), hence 'study'  
281 was included as a random effect to account for multiple estimates derived from the same paper, and 'family'  
282 was included as a random effect to account for the potential phylogenetic dependence of closely related  
283 species (or multiple observations from the same species). Models were weighted according to the inverse of  
284 the variance of standardized effects. An effect was considered as significant if confidence intervals on the  
285 parameter estimate did not overlap zero. To derive a single overall estimate of shift, no fixed effect was  
286 included (i.e. an intercept-only model). A significant effect of the intercept in this case would indicate a  
287 consistent standardized effect in terms of elevational shift across studies and species. Study duration and  
288 rate of temperature change were then tested by including each as a fixed effect in the model.

289 Papers that made future projections of species distributions or abundances did not typically present  
290 significance tests, therefore standardized effect sizes could not be estimated. Instead, the mean percentage  
291 change in the response variable (either range size or a measure of population size) was analysed. The  
292 response variable was approximately normally distributed. The model structure was similar to that for

293 observed elevation shifts in that initially an intercept-only model was specified which included 'study' and  
294 'family' as random effects, and then further fixed effects (high-elevation specialist or generalist species,  
295 period over which projections were made, climate change scenario) were tested. The sample size of the  
296 initial input data was specified as a weight in the model statement, the assumption being that models based  
297 on a larger sample size are likely to be more reliable than those based on small sample sizes. Confidence  
298 intervals of estimates which did not overlap zero were taken as evidence of consistent effects of future  
299 projections of elevational shifts.

300

## 301 RESULTS

302

303 The literature review considered a total of 234 articles relevant to climate change across various mountain  
304 regions of the Holarctic (Table 1). In Europe, most studies occurred in the Alps and Pyrenees ( $n = 45$ ),  
305 followed by Fennoscandia ( $n = 25$ ) and the uplands of Britain and Ireland ( $n = 24$ ). Many studies were also  
306 carried out in North America ( $n = 75$ ), particularly in the Rocky ( $n = 14$ ), and Appalachian ( $n = 10$ ) Mountains,  
307 while only seven studies were carried out in Holarctic Asia. There were 26 papers included that investigated  
308 climate change impacts on bird communities at a global scale. The number of published studies according to  
309 our research criteria increased considerably over time from one study in 1991 to 48 studies published in  
310 2016 (Fig. 2).

311 The most commonly investigated climate change-related topic was the general ecology/physiology  
312 and ecology of mountain bird species ( $n = 61$ ; Fig. 3), followed by papers that tested for effects of climate  
313 change on changes in population trends, elevational/latitudinal shifts or changes in community composition  
314 ( $n = 57$ ). Papers investigating future prospects of species according to various climatic scenarios were also  
315 frequent ( $n = 47$ ). The least studied category involved studies that investigated interspecific and/or  
316 synergistic interactions between climatic changes and other environmental or ecological factors ( $n = 4$ ).

317

## 318 Mountain birds of the Holarctic region

319 We identified 2316 bird species breeding in the Holarctic realm, 818 (35.3 %) of which were defined as either  
320 high-elevation mountain specialists ( $n = 324$  species) or mountain generalists ( $n = 494$  species). The most  
321 frequent Order of birds in both groups was Passeriformes (generalist  $n = 333$  species; high-elevation  
322 specialist  $n = 256$  species), followed by Piciformes for generalists ( $n = 29$ ) and Galliformes for high-elevation  
323 specialists ( $n = 27$ ; a complete list of the 2316 species is provided in the Supporting information, Table S4). A  
324 great proportion of the high-elevation specialists breed almost exclusively on the Tibetan plateaux (i.e.  
325 Tibetan Babax *Garrulax koslowi*, Tibetan Rosefinch *Carpodacus roborowskii*) or have a large proportion of  
326 their breeding range confined to this region (i.e. Bearded Vulture *Gypaetus barbatus*, Wallcreeper  
327 *Tichodroma muraria*, Twite *Carduelis flavirostris*). Examples of non-Tibetan high-elevation specialists were  
328 few and generally displayed a restricted breeding distribution confined to the lowest class that defines high-  
329 elevation specialists (class 4; Fig.1) and at the southern-most range of the Holarctic realm (i.e. Maroon-  
330 fronted Parrot *Rhynchopsitta terrisi*, White-naped Swift *Streptoprocne semicollaris*, Black Rose Finch  
331 *Leucosticte atrata*). Generalist mountain birds occur across various Holarctic mountains, ranging from the  
332 Tibetan Plateau and European Alps to the Pacific Mountain System in North America.

333 Comparing the list of mountain birds across 232 relevant articles from the literature search (no  
334 information was available for two articles) revealed that almost all generalist (97%;  $n = 453/464$ ) and high-  
335 elevation specialist species (96%;  $n = 311/324$ ) have been investigated in the literature, with the three most  
336 frequent generalist species studied being Black Restart *Phoenicurus ochruros* ( $n = 32$  studies), Water Pipit  
337 *Anthus spinoletta* and Ring Ouzel *Turdus torquatus* ( $n = 31$  each), whilst for high-elevation specialists, the  
338 most frequent species were White-winged Snowfinch *Montifringilla nivalis* ( $n = 22$ ), Yellow-billed Chough  
339 *Pyrrhocorax graculus* ( $n = 20$ ) and Wallcreeper *Tichodroma muraria* ( $n = 13$ ). However, when excluding  
340 studies based on solely distributional data (e.g. species distribution models), meta-analysis and reviews, only  
341 2% ( $n = 7/324$ ) of high-elevation mountain specialists and only 14% ( $n = 67/494$ ) of mountain generalist

342 species were investigated. This suggests that fine-scale studies on species ecology are scarce for these  
343 species.

344

#### 345 **Climate change, physiological constraints and life history strategies**

346 Birds breeding in mountain systems have evolved complex physiological, behavioural and morphological  
347 adaptations (Dragon *et al.* 1999, Cheviron & Brumfield 2012). Adaptations to prevent heat loss rely  
348 particularly on insulation, for example by producing a denser coat of feathers (Broggi *et al.* 2011) and by  
349 exhibiting a greater body mass than lower-elevation conspecifics (Bergmann's rule; Ashton 2002).  
350 Physiological constraints are likely to be major determinants of how species respond to climate change. For  
351 example, Root *et al.* (2003) found that more than 80% of the species from various taxa and habitats that  
352 showed changes linked to global warming shifted geographically in the direction expected on the basis of  
353 known physiological constraints. Birds with physiological responses that are tightly coupled to specific  
354 environmental conditions (such as mountain species) are believed to be particularly sensitive to changes in  
355 climate, but little has been done to test whether these adaptations (especially morphological) are  
356 counterproductive in a warming climate. Anecdotal evidence and the limited literature available suggest  
357 there may be costs to higher temperatures for species like Rock Ptarmigan, Ring Ouzel and White-winged  
358 Snowfinch which have been observed panting and bathing in water or snow during hot sunny days in the  
359 Swiss Alps and Scottish Highlands (Glutz von Blotzheim *et al.* 1973; DS pers. obs.), whilst Johnson (1986)  
360 found that White-tailed Ptarmigan *Lagopus leucurus* started panting at 21° C. The above studies did not  
361 establish whether these behavioural changes were sufficient to prevent reduced survival or reproduction in  
362 warming conditions.

363 A species' life history strategy may be crucial in responding to climatic alterations. Patterns along  
364 elevational gradients have highlighted that populations of the same species confined to higher elevations  
365 have slower life-history strategies (fewer nesting attempts, lower clutch size) compared to populations at  
366 lower elevation (Boyle *et al.* 2016). Higher nest survival has been found for higher elevation populations that

367 may partially compensate for the reduction in potential fecundity. Boyle *et al.* (2016) did not record a  
368 pattern of significant differences in body mass, egg or nestling size, or survival between paired populations  
369 of bird species breeding at high and low elevation. Tingley *et al.* (2012), however, found that species were  
370 more likely to shift their elevational range in the Sierra Nevada (USA) if they had smaller clutches, defended  
371 all-purpose territories and were residents or short-distance migrants, although these involved both upslope  
372 and downslope shifts. It is therefore possible that higher-elevation species may indeed be more threatened  
373 by climate change than lower-elevation species due both to their morphological adaptations to cooler  
374 systems (e.g. insulation), and their life history strategies. However, future work is required to elucidate these  
375 ideas.

376

#### 377 **Links between climate and population dynamics**

378 Although not addressing climate change *per se*, several studies have indicated that fluctuations in climate do  
379 influence demographic rates in mountain birds, and hence that potential climate change effects can be  
380 inferred. In several cases, increasing temperatures may increase reproductive output. Sæther *et al.* (2000)  
381 demonstrated that increases in winter temperature (together with population density) positively affected  
382 White-throated Dipper *Cinclus cinclus* dynamics in the upland regions of southern Norway. Cold winters  
383 caused low recruitment and a decrease in population size associated with the amount of ice cover, which  
384 impaired foraging opportunities. Novoa *et al.* (2008) demonstrated that weather variables during both pre-  
385 laying and post-laying influenced reproductive success in Rock Ptarmigan in the French Pyrenees.  
386 Reproductive success was positively associated with early snow free patches, but rainfall had negative  
387 effects, particularly after hatching. Positive effects of snowmelt on Rock Ptarmigan were also confirmed in a  
388 later study by Novoa *et al.* (2016), but the intensity of the effect varied with respect to the geographical  
389 region considered (i.e. Alps vs Pyrenees). In Mountain Plover *Charadrius montanus*, nest survival was  
390 favoured by drier and cooler weather over a seven-year period (Dreitz *et al.* 2012).

391           There is also evidence for negative effects of climate on demographic parameters. Barnagaud *et al.*  
392 (2011) showed that winter and summer NAO (North Atlantic Oscillation) affects several indicators of



393 breeding success of Black Grouse *Tetrao tetrix* in the French Alps, particularly during years of extreme  
394 weather. Interestingly, birds showed some acclimatisation, being able to optimise their reproductive output  
395 in relation to the NAO index, but they performed particularly badly when extreme weather events occurred.  
396 Twenty-five years of prolonged spring warming caused low breeding success in a Scottish population of  
397 Capercaillie (Moss *et al.* 2001).

398 In mountain systems, bird response to temperature may vary at both small and large scales. For  
399 example, nest-site characteristics of Water Pipits are selected based on their accessibility to predators,  
400 snowfall and microclimate, with the latter two influencing nestling rearing periods and survival (Rauter *et al.*  
401 2002). Because snowfall and predation pressure vary over time and space, it is possible that large-scale  
402 factors also influence species choices. The relative importance of small and large scale weather effects is still  
403 unclear as these seem to vary depending between species, populations, seasons and time periods  
404 considered. Ptarmigan exemplify such complex responses: even though they have been considered as  
405 indicators of temperature-induced effects on mountain biodiversity (Novoa *et al.* 2008, Wilson & Martin  
406 2010, Imperio *et al.* 2013), some studies have shown little effect of climate change on their demography  
407 (Sandercock *et al.* 2005, Novoa *et al.* 2016). In one study by Wang *et al.* (2002), local minimum winter  
408 temperatures had a stronger effect on White-tailed Ptarmigan population dynamics than large-scale indices  
409 like NAO. On the other hand, Wann *et al.* (2014) found that the same species in the same study area  
410 responded to climatic effects over a longer period when a two-year lag time was considered. This nuanced  
411 evidence highlights the importance of testing both small and large-scale weather predictors, and in particular  
412 in focussing more studies on lagged effects of NAO on demographic parameters of mountain species.

413

#### 414 **Changes in phenology**

415 Amongst birds, climate change has affected the phenology of many species, leading to changed timing of  
416 breeding and migration (e.g. Crick *et al.* 1997, Rubolini *et al.* 2007), which in some cases has led to  
417 population declines when phenological trends are mismatched with those of their key food resources (e.g.  
418 Both *et al.* 2006). Such phenological mismatches are hypothesised to underpin the declines in many long-

419 distance migrant species in the western Palearctic (e.g. Møller *et al.* 2008). However, in the Fennoscandian  
420 mountains, Lehtikoinen *et al.* (2014) found that long-distance migrants declined less on average than  
421 residents and short-distance migrants, suggesting the latter were more sensitive to climate change impacts.

422         There are few studies that have shown changes in mountain bird phenology explicitly linked to  
423 climate change. Timing of breeding in Mexican Jays *Aphelocoma wollweberi* has advanced in line with  
424 climate changes in the Chiricahua Mountains of Arizona (Brown *et al.* 1999). Inouye *et al.* (2000) found that  
425 American Robins *Turdus migratorius* in the Colorado Rocky Mountains arrived 14 days earlier over a 19 year  
426 period. However, local conditions (e.g. the average date of snowmelt) did not change on the study site  
427 resulting in a 65 day gap between date of the first robin sighting and date of snowmelt, suggesting that  
428 American Robins may have to cope with an extended pre-breeding period at higher elevation. Indeed, the  
429 extent, duration and timing of snow cover is likely to be an important factor acting on the phenology of  
430 mountain birds in general. For birds in the European Alps and North American mountains, patterns in  
431 breeding season phenology are typically tied to the snow melt (e.g. Novoa *et al.* 2008, Imperio *et al.* 2013,  
432 García-González *et al.* 2016). There is evidence that responses to changes in snowmelt phenology vary  
433 between species and populations. Martin & Wiebe (2004) compared White-tailed Ptarmigan and Willow  
434 Ptarmigan *Lagopus lagopus* breeding in alpine and arctic environments respectively, and found that extreme  
435 weather events greatly reduced breeding success in both species. In average years, breeding parameters of  
436 White-tailed Ptarmigan were not correlated with snowmelt phenology, therefore suggesting a constraint in  
437 adjusting their reproductive phenology to a changing environment. Willow Ptarmigan, however, tracked  
438 local conditions, breeding earlier in years of early snowmelt (Hannon *et al.* 1988, Martin & Wiebe 2004).  
439 Similarly Novoa *et al.* (2016) found that the median hatching date for Rock Ptarmigan was significantly  
440 correlated with the date of snowmelt in the French Alps, but not in the Pyrenees. None of the above studies  
441 found trends over time linked to climate change, but it can be inferred that climate change acting on snow  
442 melt phenology could affect these species in the future, especially given that snow melt has become  
443 progressively earlier, and snow cover has declined in extent in the northern hemisphere (IPCC 2007).

444

## 445 Trophic linkages

446 Global warming may influence the distribution and abundance of invertebrate communities directly  
447 (Grigaltchik *et al.* 2012) or indirectly via the modification of suitable habitat conditions (i.e. soil desiccation,  
448 changes in vegetation communities; Carroll *et al.* 2011). However, the links between such changes and bird  
449 populations have received little investigation. Most of the evidence comes from the British uplands. Pearce-  
450 Higgins *et al.* (2010) demonstrated how abundance of adult craneflies (Diptera: Tipulidae), a keystone group  
451 in many mountain systems, was negatively correlated with August temperatures in the previous year, and in  
452 turn how changes in the Golden Plover *Pluvialis apricaria* populations were negatively correlated with August  
453 temperatures two years earlier. Furthermore, Fletcher *et al.* (2013) also concluded that low temperatures in  
454 May (a surrogate for late cranefly emergence; Pearce-Higgins *et al.* 2005) positively influenced Red Grouse  
455 *Lagopus lagopus scoticus* chick survival. These findings suggest that continued warming would have negative  
456 effects on these species.

457 We found only one study considering the role of climate change on plant food sources for mountain  
458 birds. Santisteban *et al.* (2012) correlated declines in adult survival of Cassia Crossbill *Loxia sinesciuris* with  
459 increasing temperatures in South Hills and Albion Mountains (USA). The most supported explanation was  
460 that Lodgepole Pine *Pinus contorta* seed availability varied with temperature, where with increasing  
461 temperatures, trees prematurely shed their seeds, reducing the carrying capacity for Cassia Crossbill  
462 breeding later in the year. The warmer springs and increased precipitation in Europe will also influence food  
463 availability and the future geographical distribution for European Crossbills (Common Crossbill *L. curvirostra*,  
464 Parrot Crossbill *L. pytyopsittacus*, Scottish Crossbill *L. scotica*) (Mezquida *et al.* 2017)

465 Snow patches can represent an important foraging habitat, providing both arthropod fallout and  
466 suitable sites at their margins for the collection of soil invertebrates, particularly during the nesting and  
467 rearing period of many mountain birds such as White-winged Snowfinch, Snow Bunting *Plectrophenax*  
468 *nivalis*, Horned Lark *Eremophila alpestris* and Alpine Accentor *Prunella collaris* (Antor 1995, Camfield *et al.*  
469 2010, Brambilla *et al.* 2016a,b, Rosvold 2016). In some extreme cases, birds may even choose to nest directly

470 in glaciers (White-winged Diuca Finch *Diuca speculifera*; Hardy & Hardy 2008) or in very close proximity  
471 (Grey-crowned Rosy Finch *Leucosticte tephrocotis*, Johnson 1965, Rosvold 2016; Brandt's Rosefinch  
472 *Leucosticte brandt*, Potatov 2004) to capitalize on abundant supplies of insects. Changes in amount and  
473 duration of snow may therefore affect these species via food resources.

474 Where trees and shrubs have expanded their distribution upslope in response to increasing  
475 temperatures (Harsch *et al.* 2009, Myers-Smith *et al.* 2011), changes in invertebrate communities are  
476 expected. Ground and canopy-dwelling arthropod communities have been assessed in the arctic foothills of  
477 Alaska in relation to the presence of two passerine predators, Gambel's White-Crowned Sparrow *Zonotrichia*  
478 *leucophrys gambelii* and Lapland Longspur *Calcarius lapponicus*. Predicted changes in shrub dominance are  
479 likely to favour White-crowned Sparrow nesting habitat and food (canopy-dwelling arthropods), whilst  
480 declines of Lapland Longspurs have been projected as a consequence of shrub encroachment and  
481 consequent reduced availability of ground-dwelling arthropods (Boelman *et al.* 2015).

482 Climate change may also have effects by affecting more complex linkages across different trophic  
483 levels. Martin & Maron (2012) conducted an experiment showing that climate change in the form of reduced  
484 snowfall in mountains, and leading to increased ungulate herbivory in winter, can negatively affect diverse  
485 species interactions. They experimentally tested the hypothesis that declining snowfall, which enables  
486 greater over-winter herbivory by Elk *Cervus canadensis*, indirectly influences plants and associated bird  
487 populations in montane forests. When they excluded Elk from one of two paired snowmelt drainages, and  
488 replicated this paired experiment across three distant canyons over a six year period, there was a reversal in  
489 the multi-decadal declines in plant and bird populations. These experimental results suggest that climate  
490 impacts can interact with other drivers of habitat change and strongly influence plant–animal and other  
491 ecological interactions.

492

### 493 **Observed evidence of elevational shifts**

494 Evidence exists, typically from broad-scale atlases of species distributions, that some species are shifting  
495 their geographic distributions to higher elevations in response to climate change, the distributions

496 presumably tracking more suitable climatic conditions (e.g. Auer & King 2014, Roth *et al.* 2014), although  
497 such effects are not universal (e.g. Zuckerberg *et al.* 2009, Tingley *et al.* 2012, Massimino *et al.* 2015).  
498 Furthermore, apparent elevational shifts may occur due to habitat deterioration or destruction at lower  
499 elevations (Archaux 2004, Bodin *et al.* 2013).

500           Few studies have considered elevational shifts in the distributions of Holarctic mountain birds. We  
501 found 10 relevant studies in our literature search that considered specifically elevational shifts in bird species  
502 distributions along elevational gradients, partly or wholly in mountains, over time (Table 2). Including papers  
503 that considered more than one study site ( $n = 13$  sites from 10 papers), the mean period considered was  $38$   
504  $\pm 2.5$  SE years (range 9 – 102) and the mean length of elevation gradients was  $1970 \pm 76$  SE m (range = 500 –  
505 3400m). There was little evidence of consistent patterns in elevational shifts across the studies, and there  
506 was a wide variation among species. In some cases, there were fairly consistent upward shifts in most  
507 species (e.g. Reif & Flousek 2012, Rocchia 2016), but other studies found that different species exhibited  
508 upward and downward shifts (Tingley *et al.* 2012, DeLuca & King 2017), or found shifts in only a small  
509 proportion of the species considered (Archaux 2004). Additionally, there was sometimes marked variation in  
510 species' responses between geographic locations within the same study (Tingley *et al.* 2012, Pernellet *et al.*  
511 2015).

512           The lack of consistent trends apparent in Table 2 was supported by the meta-analysis. There were  
513 203 estimates of elevational shift from seven published studies analysed, five from Europe and two from  
514 North America. Pooling all estimates across the studies, there was no strong support for a general shift  
515 towards higher elevations (parameter estimate  $\pm$  SE =  $0.083 \pm 0.052$ , 95% CLs = -0.018, 0.184). Shifts towards  
516 higher elevations were more positive when rates of temperature change were higher (estimate  $\pm$  se =  $0.543 \pm$   
517  $0.152$ , 95% CLs = 0.245, 0.841). Duration of study had an unexpected negative effect on shifts, studies over  
518 longer time spans resulting in more downward shifts (estimate  $\pm$  se =  $-0.026 \pm 0.004$ , 95% CLs = -0.034, -  
519 0.018). The above findings were robust to different model structures and different subgroups of species  
520 (Supplementary Material, Table S3).

521 Most studies in Table 2 also considered temperature variations over the same period, either  
522 modelling them in relation to bird distributions (Archaux 2004, Popy *et al.* 2010, Reif & Flousek 2012, Tingley  
523 *et al.* 2012, Pernollet *et al.* 2015, Rocchia 2016), or considering climate trends over the same periods  
524 (Maggini *et al.* 2011, Mizel *et al.* 2016). In most cases, trends in elevational shifts were temporally matched  
525 with temperature trends over the same period, with a few exceptions (Pernollet *et al.* 2015, Mizel *et al.*  
526 2016). Tingley *et al.* (2012) found a broad range of responses of bird species along elevation gradients in the  
527 Sierra Nevada, due in part to differential responses to increasing temperature (exerting a general positive  
528 upwards shift) and increasing precipitation (exerting a general downslope shift), although few other studies  
529 considered potential effects of precipitation (only Archaux 2004, Popy *et al.* 2010 and Pernollet *et al.* 2015).

530 Changes in bird population trends along elevation gradients over time are similarly inconsistent  
531 across studies. Some find positive changes in lower-elevation species and negative changes in higher-  
532 elevation species that are consistent with elevational shifts as lower-elevation species colonise mountains  
533 and higher- elevation species lose suitable habitat (Flousek *et al.* 2015). However, others have reported  
534 opposite (Archaux 2007) or inconsistent (Zamora & Barea-Azcón 2015, Furrer *et al.* 2016) patterns. Tingley  
535 and Beissinger (2013) found a decrease in total species richness, and in species richness of high elevation  
536 species, over time in the Sierra Nevada, despite heterogeneous shifts in individual species in the same area  
537 (Tingley *et al.* 2012). At wider scales, there is evidence that bird communities are shifting towards warm-  
538 dwelling species (Switzerland; Roth *et al.* 2014), but also that communities at higher elevations have lower  
539 'climate debt' (the spatio-temporal divergence between temperature changes and community changes) as  
540 elevation increases (France; Gaüzère *et al.* 2016).

541

#### 542 **Projected elevational shifts**

543 Extinction risks are expected to increase as a result of climate-induced elevational range shifts in the future  
544 (Sekercioglu *et al.* 2008, La Sorte & Jetz 2010). Shifting vegetation zones in mountains, and in particular an  
545 advance of the tree-line towards higher elevations, has been observed in many studies (e.g. Lenoir *et al.*

2008, Harsch *et al.* 2009). As a consequence, high-elevation specialists, and in particular those of open, treeless habitats, are expected to be most threatened due to habitat loss or fragmentation (e.g. Chamberlain *et al.* 2013, Goodenough & Hart 2013, Siegel *et al.* 2014, Brambilla *et al.* 2016a). Nevertheless, some studies have also projected overall range loss in higher-elevation forest specialists (Braunisch *et al.* 2014, Brambilla *et al.* 2015). There were 95 estimates derived from 12 studies that satisfied the criteria to be included in the analysis (see Supporting Information; Table S2). There was a net prediction of negative impacts on species populations or distributions, although there was a degree of variability and confidence limits overlapped zero (estimate  $\pm$  se =  $-28.9 \pm 17.0\%$ , 95% CLs = -62.4, 4.6). High-elevation mountain specialists and generalists were projected to be more negatively impacted than other species (mountain specialists and generalists =  $-76.1 \pm 27.1\%$ , 95% CLs = -129.2, -23.0; other =  $29.8 \pm 25.7\%$ , 95% CLs = -20.6, 80.2). There was a tendency for greater negative impacts in severe than moderate scenarios (moderate =  $-26.6 \pm 17.1\%$ , 95% CLs = -60.1, 6.9; severe =  $-33.6 \pm 17.5\%$ , 95% confidence limits = -67.9, 0.7). There was no evidence of an effect of the number of years over which projections were made ( $0.01 \pm 0.79$ , 95% CLs = -1.53, 1.55). Re-running the models without weighting for sample size showed the same patterns, although results were less conservative (i.e. it was less likely that confidence intervals overlapped zero).

561

## 562 **Interactions of climate change effects with other drivers of change**

### 563 *Land use*

564 Disentangling the relative importance of climatic effects and other drivers of environmental change that  
565 influence the persistence and maintenance of biodiversity has been a key issue across mountain regions  
566 (Mantyka-Pringle & Rhodes 2012, Cumming *et al.* 2014, Maggini *et al.* 2014, Elmhagen *et al.* 2015), and is  
567 central to produce efficient, adaptive conservation frameworks for threatened species (Gehrig-Fasel *et al.*  
568 2007, Gienapp *et al.* 2007, Eglinton & Pearce-Higgins 2012, Titeux *et al.* 2016). For example, climate change  
569 and land-use often interact in ways that influence biodiversity (Parmesan & Yohe 2003), and these  
570 interactions may amplify or reduce the magnitude of potential effects (Clavero & Brotons 2010, Dreitz *et al.*

571 2012, Chamberlain *et al.* 2013, Oliver *et al.* 2017). Lehtikoinen & Virkkala (2016) acknowledged a land-use and  
572 species trait effect due to the high level of unexplained variation in models predicting the change in density  
573 of birds in relation to temperature change. Jetz *et al.* (2007) attempted to assess the relative importance of  
574 climate and land use changes using future scenarios. We identified 617 birds defined as either high-elevation  
575 mountain specialists or mountain generalists present in the study of Jetz *et al.* (2007). For these species, the  
576 average percentage loss in geographic range due to land use change was estimated at 24.8% and 28.6% by  
577 the years 2050 and 2100 respectively. In comparison, loss due to climate change alone was predicted to be  
578 7.3% and 11.5% respectively.

579 The interaction between climate and land use is particularly relevant to mountain habitats that are  
580 experiencing a faster rate of climate change than the global average (Diaz *et al.* 2003, Nogués-Bravo *et al.*  
581 2007) and are subjected to various anthropogenic changes at a landscape level (Arlettaz *et al.* 2007, 2015,  
582 Gellrich & Zimmermann 2007, Nogués-Bravo *et al.* 2008, Patthey *et al.* 2008, Braunisch *et al.* 2011, 2013,  
583 2016, Douglas *et al.* 2015). However, land use change has only been rarely incorporated into analyses of  
584 distribution shifts: Reif & Flousek (2012) and Rocchia (2016) found that elevation shifts more closely  
585 matched temperature than habitat changes, Tryjanowski *et al.* (2005) found significant effects of both,  
586 whereas Popy *et al.* (2010) could not separate the effects of the two.

587 Agro-forestry and pastoral practices have shaped the landscape of Holarctic mountains in Europe  
588 and Asia, influencing the species composition and abundance of mountain birds (e.g. Gehrig-Fasel *et al.*  
589 2007, Caprio *et al.* 2011, Douglas *et al.* 2014, Wilson *et al.* 2014, Mollet *et al.* in press). Over time, forest  
590 management has changed in intensity (e.g. clear-felling vs single-tree selection), composition (planting of  
591 exotic conifers) and age dynamics (establishment of even-aged monocultures; Kirby & Watkins 2015). At the  
592 same time, climate change may be affecting forest bird assemblages either directly or indirectly by  
593 influencing cover, productivity, and composition of forest systems. However, it is generally unclear which of  
594 these two pressures (climate change or forestry practices) is the most important driver in changes in bird  
595 distribution. Changes in forest composition could cause opposite shifts (i.e. downhill) to those forecast due  
596 to effects of climate warming (uphill). For example, Archaux (2004) suggested that changes in forest



597 management that favoured coniferous at the expense of broadleaved trees might have caused forest birds  
598 to have shifted their mean elevation downwards. In other cases, there is evidence from boreal forests  
599 (including some mountain areas) that climate, in addition to vegetation type and management, is a crucial  
600 driver for determining passerine species distribution (Cumming *et al.* 2014, Frey *et al.* 2016). Virkkala (2016)  
601 found that forest management favoured passerine species benefitting from climate change, so that direct  
602 habitat alteration was connected to the indirect effects of climate change.

603           Climatic variables can also be important factors for non-passerine species. Brambilla *et al.* (2015)  
604 found in the Italian Alps an important effect of climate in addition to habitat composition at the landscape  
605 scale in dictating the distribution of the cold-adapted Pygmy Owl and Boreal Owl *Aegolius funereus*. Both of  
606 these forest species were predicted to undergo range contraction in the Alps as a consequence of climate  
607 change. Braunisch *et al.* (2014) evaluated the importance of climate, landscape and vegetation variables on  
608 the occurrence of indicator species (i.e. Capercaillie, Hazel Grouse *Tetrastes bonasia*, Three-toed  
609 Woodpecker *Picoides tridactylus* and Pygmy Owl) in central European mountain forests, and assessed future  
610 changes in habitat suitability of these species according to future climatic projections. Although climate  
611 variables were the most important factors for most species, the models predicted that *in situ* management  
612 actions, such as increasing the number of forest gaps (for Capercaillie), increasing bilberry *Vaccinium* spp.  
613 cover (for Hazel Grouse) and increasing the number of snags and/or the proportion of high (>15m) canopy  
614 forest (for Three-toed Woodpecker), could to some extent mitigate the detrimental impact of climatic events  
615 and sustain bird populations. However, such interventions may have to work against the natural forest  
616 dynamics and could be expensive.

617           Historically, agricultural expansion and changes in livestock management have had major impacts on  
618 mountain birds (Lundmark 2007, Elmhagen *et al.* 2015). In many mountain areas, traditional grazing  
619 practices are characterised by low stocking densities or transhumant pastoralism, i.e. the seasonal  
620 movement of livestock between high-elevation summer pastures and lowland winter pastures (Arnold &  
621 Greenfield 2006). These traditional grazing practices have been largely abandoned in some areas due to  
622 social and economic factors, especially in the European Alps. For example, in Italy, the number of farms has

623 decreased drastically and many have changed to indoor production systems (Battaglini *et al.* 2014), which  
624 has led to substantial changes within mountain vegetation zones through encroachment of formerly open  
625 grasslands by trees and shrubs and a loss of structural heterogeneity (Braunisch *et al.* 2016). Elevational  
626 shifts in vegetation may be therefore due both to climate change and land abandonment (Gehrig-Fasel *et al.*  
627 2007).

628           The reintroduction of grazing is an often recommended management solution to counteract tree  
629 and shrub encroachment in open areas (Gehrig-Fasel *et al.* 2007), and it has the potential to increase plant  
630 structural diversity and composition (Hoiss *et al.* 2013, Peringer *et al.* 2013) which is key to preserve  
631 emblematic birds of semi-open habitat (Patthey *et al.* 2012). However, the effects of grazing on mountain  
632 bird populations are still not well understood. Long-term grazing at high stocking densities is known to have  
633 negative impacts on soil fertility and consequently on the productivity of the whole system (McVean &  
634 Lockie 1969), although effects on mountain birds are not consistent and vary substantially among  
635 geographical regions, livestock types and stocking levels. Several studies have reported that grazing increases  
636 richness or densities of mountain grassland birds (Laiolo *et al.* 2004, Evans *et al.* 2006, Bazzi *et al.* 2015).  
637 Evans *et al.* (2006) found that mixed sheep and cattle grazing, at low intensity, improved the breeding  
638 abundance of Meadow Pipit *Anthus pratensis* compared to sites stocked with sheep only (at high or low  
639 density) or unstocked sites in the Scottish uplands, while Loe *et al.* (2007) reported the highest bird density  
640 on pastures with high sheep density in Norway. Other studies have shown no differences in bird abundance  
641 or species richness between grazed and ungrazed sites (Moser & Witmer 2000) or a negative influence of  
642 grazing animals on nesting success (Pavel 2004, Warren *et al.* 2008).

643           Climate change can also have direct impacts on grazing management, although this seems to be less  
644 well studied. In Nepal, where transhumance is a common practice, herders perceived the impact of climate  
645 change through personal experience. In several studies, where herders have been interviewed, they  
646 described a rise in temperature, a decline of rain- and snowfall, a scarcity of water resources (Aryal *et al.*  
647 2014, Wu *et al.* 2015) and the presence of invasive weeds, which are replacing the valuable grasses on  
648 farmlands (Gentle & Thwaites 2016). These perceptions were also in line with temperature and rainfall

649 trends in the studied region. As a result, herders tried to adjust their transhumance patterns to the changed  
650 conditions by altering the timing of seasonal livestock movements. The consequences of such management  
651 responses for mountain bird populations, however, remain unknown. Given the varieties of different effects  
652 of grazing on mountain birds, and the lack of research on likely responses of grazing management practices  
653 to future climate change, further investigations are needed to examine potential effects of grazing regimes  
654 on mountain bird populations before we can apply them as potential conservation tools.

655

#### 656 *Leisure and other potential threats*

657 Mountains are important ecosystems for biodiversity, but are also multi-functional sites for various human  
658 activities, including leisure. People seek mountain landscapes to practice a range of different sports and  
659 hobbies such as skiing, snowboarding, hiking, biking, birdwatching, rock-climbing, paragliding and hunting.  
660 Local communities benefit economically from tourism. The leisure industry in mountain areas is growing  
661 (Debarbieux *et al.* 2014), and the potential effects of these activities on avian communities have received  
662 increasing attention from conservationists (e.g. Patthey *et al.* 2008, Arlettaz *et al.* 2013, DeLuca & King  
663 2014). The impact of snowsports on biodiversity is a major topic studied in the European Alps, where there  
664 are c. 40,000 kilometres of ski-runs served by c. 14,000 ski-lifts that are capable of transporting c. 1.5 million  
665 skiers per hour (Weed & Bull 2004). As a consequence of this and other activities, the Alps receive nearly 100  
666 million visitors per year, spending \$60 billion annually (Giuliano 1994). In contrast, snowsport activities,  
667 including skiing operations in North America and the Eastern Holarctic remain at relatively low density, with  
668 most likely local effects on biodiversity (Martin 2001b).

669         There are several lines of evidence showing that ski-pistes have deleterious effects on both grassland  
670 and forest birds via loss and degradation of habitat, and a decrease in food availability (Laiolo & Rolando  
671 2005, Rolando *et al.* 2007, Caprio *et al.* 2011, Rixen & Rolando 2013). In addition, there is evidence that  
672 hormonal stress in birds generated by intensive human activities can negatively impact already vulnerable  
673 populations of Capercaillie (Thiel *et al.* 2011) and Black Grouse (Arlettaz *et al.* 2007, 2013). Anthropogenic  
674 disturbance has furthermore been shown to entail extra energetic costs that may negatively affect

675 population dynamics (Arlettaz *et al.* 2015). Effects may also operate through infrastructure associated with  
676 skiing, for example increased mortality due to collision with ski cables (Baines & Andrew 2003, Watson &  
677 Moss 2004), and reduced reproductive success of ground nesting birds associated with development of  
678 tourist resorts (Watson & Moss 2004, Patthey *et al.* 2008, Tolvanen & Kangas 2016), although negative  
679 effects are not universal (Rimmer *et al.* 2004).

680 Interactive effects of climate change and outdoor sports could increase the above negative impacts  
681 on bird populations in the future. Global warming is having important economic consequences for the skiing  
682 industry due to reduced snow cover and persistence. Compensatory mechanisms are targeted at prolonging  
683 the ski season by direct spraying of artificial snow, or by creating new ski pistes at higher elevations where  
684 snow conditions are more reliable. Brambilla *et al.* (2016a) modelled ski-pistes and mountain bird presence  
685 according to future climatic scenarios. Strong overlaps between areas climatically and topographically  
686 suitable for the development of ski-pistes and areas suitable for breeding alpine birds were predicted to  
687 occur, suggesting that the conservation of mountain bird communities will require careful planning in order  
688 to reduce potential increased future conflicts between outdoor winter sports and birds. Global warming is  
689 also causing the abandonment of ski-runs at lower elevations. Natural grassland revegetation at some  
690 abandoned sites resulted in a partial recovery of important alpine birds, but never back to the state of the  
691 'original' alpine grasslands (Caprio *et al.* 2016).

692

### 693 *Novel interspecific interactions*

694 Species may respond to climate change by shifting their distribution to track local climates (Tingley *et al.*  
695 2009, Jackson *et al.* 2015), which may result in novel interactions as species colonize new areas.  
696 Including such interactions has improved model predictions at different scales (Araújo & Luoto 2007).  
697 Heikkinen *et al.* (2007) and Brambilla *et al.* (2013) suggested that including the presence of woodpeckers  
698 that produce the cavities used by secondary cavity nesting raptors improved model performance in  
699 predicting cavity-nesting forest owl distributions. We found only one relevant example that tested the  
700 importance of biotic interactions among birds along elevational gradients. Freeman & Montgomery (2015)

701 assessed potential competition between Swainson's Thrush *Catharus ustulatus*, which generally inhabits  
702 lower elevations but which has shifted its distributions towards higher elevations, and the conspecific  
703 Bicknell's Thrush *Catharus ustulatus*, which is largely confined to mountaintops. Using playback techniques,  
704 the authors found that, where the species co-occurred, Swainson's Thrush responded aggressively to  
705 Bicknell's Thrush, but not *vice-versa*.

706

## 707 **Conservation and policy**

708 Our literature review has clearly highlighted the need for more detailed studies of mountain birds, with  
709 several papers stating that a valuable conservation framework can be achieved only if such knowledge gaps  
710 are bridged (see Research gaps and Conclusion; Fig. 4). Despite this, we found that most studies on this topic  
711 identified adaptation strategies for mountain and upland species threatened by climate change. Most of  
712 these studies ( $n = 21$ ; Fig. 4) focussed on the quality, quantity and geographical location of protected areas.  
713 Existing protected areas may have already functioned as important compensatory systems, increasing  
714 species resilience to climate change (Virkkala *et al.* 2014, Gaüzère *et al.* 2016, Santangeli *et al.* 2016), and in  
715 future scenarios of greenhouse gas emission, greater biodiversity losses have been predicted in unprotected  
716 than in protected areas (Virkkala *et al.* 2013). In Europe, nationally designed protected areas are likely to  
717 retain climatic suitability better than unprotected areas in the future, as they tend to occur at high elevations  
718 and hence act as climatic *refugia* for species, although this was not found to be the case for the European  
719 Union-wide Natura 2000 network (Araújo *et al.* 2011). The same authors also highlighted that nearly all (i.e.  
720 97.2%) alpine species and sub-species of vertebrates and plants of European concern are projected to lose  
721 suitable habitat due to their small ranges. However, it must be highlighted that the study did not consider  
722 species dispersal.

723 Some habitats may also be more prone to climate change than others. Montane forest species are  
724 predicted to be less impacted by climate change due to the stronger self-regulation of the forest  
725 microclimate compared to open habitats (Reif & Flousek 2012), and to native forest expansion that has

726 already occurred, and which is predicted to continue in many areas (European Alps; British uplands;  
727 Chamberlain *et al.* 2013, Scridel *et al.* 2017b), but they could be prone to other climate change related  
728 threats such as pests, disease and wild fires (Dale *et al.* 2001, Sturrock *et al.* 2011, Lesk *et al.* 2017).  
729 Furthermore, natural grasslands in the Alpine region may face serious challenges to shift their distributions,  
730 as they are being progressively colonised due to both forest encroachment following land abandonment or  
731 release of grazing pressure and advancing treelines at lower elevations, while facing constraints on  
732 colonizing higher elevations, for example due to slow rates of soil formation (Freppaz *et al.* 2010,  
733 Chamberlain *et al.* 2013, Jackson *et al.* 2015).

734 Targeted habitat management should be considered as an adaptive conservation tool for various  
735 species threatened by climate change (Fig. 4). Improving habitat structure and offering greater prey  
736 availability has been reported to increase mountain species' resilience and resistance for forest, semi-open  
737 and open-habitat species (Caroll *et al.* 2011, Braunisch *et al.* 2014, Scridel *et al.* 2017b). This might be  
738 achieved by targeted grazing to maintain open habitats and enhance invertebrate populations (Signorell *et*  
739 *al.* 2010, Patthey *et al.* 2012, Braunisch *et al.* 2016). Such intensive actions can be very costly and in conflict  
740 with many economic goals, hence management should be targeted in large areas projected to support viable  
741 wildlife populations. Increasing the quantity and quality of protected areas is not just important for mountain  
742 species *per se*, but also because these areas are likely to become stopover *refugia* for many migrant species  
743 tracking climate change (Loarie *et al.* 2009, Boyle & Martin 2015), and management action should also  
744 accommodate these species' requirements. When intensive management *in situ* does not compensate for  
745 climatic effects, potential captive programs ( $n = 2$ ; Fig. 4), translocation of species to new suitable areas ( $n =$   
746  $2$ ; Bech *et al.* 2009), or the creation of corridors to favour dispersal and colonization of new areas (Huntley *et*  
747 *al.* 2008, Conroy *et al.* 2011, Lu *et al.* 2012, Virkkala *et al.* 2013) have been proposed.

748 All of these adaptation responses for mountain species threatened by climate change can work only  
749 if scientists and policy makers collaborate to influence current legislation. Our classification of high-elevation  
750 mountain specialists and mountain generalists indicates initial steps for a joint common Holarctic mountain  
751 bird index, which so far has been developed for some regions in the world (Fennoscandia; Lehikoinen *et al.*

752 2014; North America and British Columbia, Canada; Boyle & Martin 2015) and which could be essential for  
753 scientists and policy-makers to measure progress in the conservation of mountain birds, especially if this  
754 index includes full life cycle avian use of mountain habitats. While in the long-term, global measures to  
755 contain and reverse anthropogenic emissions are important ( $n = 3$ ; Fig. 4), most authors admit that the  
756 persistence of mountain species also depends on immediate short-term national and local conservation  
757 actions and legislation ( $n = 9$ ; Fig. 4).

758

## 759 DISCUSSION

760

761 Our literature review has shown that there is a growing body of evidence that climate change is impacting  
762 mountain birds in terms of distribution and population trends, reproduction and survival. These changes may  
763 have been mediated through direct effects of climate on physiology, indirect effects of changes in habitat, or  
764 via interactions with other biotic and abiotic changes. However, patterns were often highly variable (e.g.  
765 both increases and decreases in population size, range changes towards both higher and lower elevations),  
766 between species, and between different study areas for the same species.

767 Defining a 'mountain bird' across a large region like the Holarctic is difficult because many species  
768 that are mountain birds in warmer climates are lowland species in colder climates. Our goal was to derive an  
769 objective definition that could be applicable over a large geographic area and which identified species  
770 associated with mountains *per se*, rather than occurring in mountains due to interactive effects of climate,  
771 elevation, latitude and land use. This is important when considering species distributions over large scales,  
772 and in particular when projecting future distributions. For example, the Water Pipit was identified as a  
773 generalist mountain breeding bird across various mountain slopes, even in the northern, colder, parts of its  
774 geographic range. Predictions based on climate alone may therefore be inaccurate for such species (e.g.  
775 Huntley *et al.* 2008). In general, the species identified as high-elevation specialists or mountain birds (see  
776 Supporting Information; Table S4) were in-line with the authors' expectations, although there were some  
777 surprising results. For example, Rock Ptarmigan is considered an archetypal mountain bird in many parts of

778 its range (the European Alps, the Pyrenees, British Columbia and Alberta), but not according to our  
779 definition. This to some extent may have been due to the coarse scale of the defined breeding range used in  
780 the method, but it is also reflective of the widespread populations of this species inhabiting lowland arctic  
781 tundra. As conservation policy is typically applied at national or regional level, a regional-based definition of  
782 mountain birds would also be useful, which would be readily achievable under the current methodology.  
783 Although this first classification of Holarctic high-elevation mountain specialist and generalist birds was not  
784 the primary aim of this review, we regard this exercise of considerable value for future work on this group of  
785 poorly studied species (e.g. baseline monitoring, development of a joint mountain bird index, and ecological  
786 and conservation research).

787           According to our meta-analysis, there was no evidence for consistent elevational shifts in mountain  
788 bird species. Although we failed to detect any direct and conclusive effects that climate change has caused  
789 widespread distribution shifts in Holarctic mountain birds, it is likely that we lack sufficient data to generate  
790 robust conclusions. The meta-analysis included a range of species encompassing a great variation in life  
791 history strategies, demographic parameters and geographical regions. All of these factors are likely to  
792 influence potential responses to climate change and hence cause a wide variation in patterns of elevational  
793 shift among mountain birds across the mountain ecosystems and avian taxa considered (Martin & Wiebe  
794 2004, Wilson & Martin 2010, Tingley *et al.* 2012, Novoa *et al.* 2016).

795           There was consistency in climate projections across studies that was somewhat at odds with the  
796 heterogeneity in responses of observed elevational shifts. This may in part have been due to more mountain  
797 high-elevation specialists being included in the projection papers. However, in many cases, there was a focus  
798 on climate (usually temperature and precipitation) as a driving factor, and only half of the studies considered  
799 alternative scenarios of climate change in tandem with land use change or other anthropogenic pressures. In  
800 general, species distribution models only rarely include scenarios of changes in land use and human  
801 disturbance alongside those of climate change (Sirami *et al.* 2016). It is clear that range shifts in mountains  
802 may be influenced by many factors, including temperature, but also precipitation, habitat and topography,



803 and that species may vary widely in their response. All of these factors should be considered when assessing  
804 elevational range shifts, and predicting future shifts, in mountain birds.

805           Adaptation responses for mountain species threatened by climate change rely on enhancing the  
806 quality and quantity of suitable habitat in particular via protected areas, but also the conservation of suitable  
807 ecological conditions at regional and wider levels, including improving landscape connectivity. We have  
808 shown that human activities can be beneficial for climate-sensitive species (i.e. some pastoral activities), and  
809 yet mechanisation, leisure and urbanisation may impede potential benefits. Major changes can occur if  
810 scientists and legislators work closely together, for example through the development of efficient agri-  
811 environmental schemes, forestry practices, regulation of leisure activities and sustainable urban planning in  
812 mountain areas, and explicit recognition of the general ecological requisites for wildlife persistence such as  
813 connectivity across their full life cycle.

814

#### 815 **Research gaps and conclusions**

816 From our literature review, it was evident that mountain species are little studied relative to species in  
817 lowland habitats of the Holarctic, such as farmland, forest and wetlands. Many common species in  
818 mountains are lacking even basic biological and ecological knowledge (e.g. Alpine Accentor, White-winged  
819 Snowfinch, Twite, Wallcreeper, North American rosy finches *Leucostiche* spp). Whilst both high-elevation  
820 mountain specialists and mountain generalist are well-represented in the literature in terms of large-scale  
821 distribution studies (e.g. species distribution models based on atlas data), they are very poorly represented  
822 when considering finer-scale, usually more intensive, studies which address ecological mechanisms. In  
823 particular, there were very few studies that investigated the ecology of high-elevation mountain specialists,  
824 yet these are the species that may be most likely to be impacted by climate change. Aside from broad-scale  
825 species distribution, the evidence base therefore largely concerns species that occur across a range of  
826 habitats and elevations, rather than species whose geographical range, at least in the Holarctic, is largely  
827 restricted to mountain areas.

828           Our understanding of physiological mechanisms underpinning avian responses to climate change is  
829 still limited, even if recent studies have emphasized the importance of specifying ecological traits, notably  
830 physiological tolerance, when predicting responses to climate change (Kearney & Porter 2009, Reif & Flousek  
831 2012, Auer & King 2014, Pacifici *et al.* 2017). This is particularly important in terms of developing  
832 conservation strategies. If a species responds directly to climate through a physiological effect, then there  
833 might be limited conservation action that could be implemented beyond the need to reduce our dependence  
834 on non-renewable fossil energy sources. There is more potential for developing conservation actions for  
835 species that are affected indirectly by climate change, although for mountain birds, we still lack information  
836 about species' basic ecological requirements, such as key trophic resources for reproduction, that are  
837 required to develop management strategies. We therefore emphasize the need for more basic studies of  
838 both physiological tolerance and ecological requirements of mountain birds, and in particular high-elevation  
839 mountain specialists, as well as for all those lower-elevation species that are predicted to colonize mountain  
840 regions in the near future (Loarie *et al.* 2009).

841           The importance of considering cross-ecosystem linkages such as trophic structure when identifying  
842 climate change effects has been shown to be crucial for a clear understanding of the underlying mechanisms  
843 affecting species and populations (Pearce-Higgins *et al.* 2010, Santisteban *et al.* 2012, Fletcher *et al.* 2013).  
844 Furthermore, a better understanding of energetic values in food sources (prey) and how these influence  
845 demographic rates in species is particularly important for future climate-related adaptation responses. There  
846 were relatively few studies that considered long-term trends over several years that could encompass a full  
847 range of climate variation, and hence assess climate trends (rather than year-to-year changes in weather  
848 over shorter periods). The low number of studies assessing elevational distribution shifts in particular  
849 suggests that monitoring in high mountains is inadequate, probably due to a combination of complex terrain  
850 and lack of field surveyors available in these sparsely populated areas. Targeted monitoring in mountain  
851 areas, with a focus on high-elevation mountain specialists, is therefore essential if we are to improve our  
852 assessments of current and future climatic effects on bird distributions.

853 Monitoring reproductive success and survival of mountain birds would be similarly useful. The  
854 demographic mechanisms that underpin species distributions and population changes are not well  
855 understood for mountain birds. There have been some short-term effects of climate demonstrated in several  
856 species, but longer-term studies are rare. More intensive long-term studies would enhance understanding of  
857 the key factors that determine population trends and distributions, and therefore would facilitate the  
858 predictions of future climate change impacts by elucidating more complex mechanisms, such as phenological  
859 effects. Many studies acknowledge that a valuable understanding of climatic impacts can only be achieved if  
860 key interacting factors are considered, such as land-use changes and biotic interactions, including  
861 interspecific competition. Given that projections of future mountain bird species distributions may be quite  
862 sensitive to assumptions about how land use will change in the future (e.g. Chamberlain *et al.* 2013), we urge  
863 a greater consideration of land use change in species distribution modelling in mountain environments.  
864 Finally, we invite scientists and policy-makers to further develop studies and related frameworks to  
865 efficiently develop habitat restoration plans in mountain areas, particularly where climate change and  
866 changes in land-use are likely to offer such opportunities in the near future (i.e. encroaching pastures after  
867 grazing/ski-pistes abandonment, afforestation of native woodland on moorlands). Indeed, conservation and  
868 restoration frameworks have already been developed for various birds species inhabiting mountain regions  
869 considered susceptible to changes in climate and land use (e.g. Carroll *et al.* 2011, Patthey *et al.* 2012,  
870 Signorell *et al.* 2010, Braunisch *et al.* 2016, Caprio *et al.* 2016, Scridel *et al.* 2017b).

871  
872 We are grateful to Jeremy Wilson, Paul Donald, James Pearce-Higgins, Tómas G. Gunnarsson and one anonymous  
873 reviewer for helpful comments and advice. We also acknowledge Alessandro Franzoi, Giacomo Assandri, Simone Tenan,  
874 Emanuel Rocchia & Frank La Sorte for general advice and comments. We thank Bill DeLuca, Fränzi Korner, Jeremy Mizel,  
875 Claire Pernollet, Veronika Braunisch, Jaime Resano Mayor and Morgan Tingley for help with the meta-analyses. The  
876 study was funded by Museo delle Scienze of Trento (MUSE - Italy) and by the Paneveggio-Pale di San Martino  
877 Natural Park (Italy) as part of Davide Scridel's doctorate program. Matteo Anderle's contribution was funded by the  
878 Autonomous Province of Trento (Italy).

879

880

881

882 REFERENCES

- 883 **Ameztegui, A., Coll, L., Brotons, L. & Ninot, J.M.** 2016. Land-use legacies rather than climate change are  
884 driving the recent upward shift of the mountain tree line in the Pyrenees. *Glob. Ecol. Biogeogr.* **25**: 263–  
885 273.
- 886 **Antor, R.J.** 1995. The importance of arthropod fallout on snow patches for the foraging of high-alpine birds. *J.*  
887 *Avian Biol.* **26**: 81–85.
- 888 **Araújo, M.B. & Luoto, M.** 2007. The importance of biotic interactions for modelling species distributions  
889 under climate change. *Global Ecol. Biogeogr.* **16**: 743–753.
- 890 **Araújo, M.B., Alagador, D., Cabeza, M., Nogués-Bravo, D. & Thuiller, W.** 2011. Climate change threatens  
891 European conservation areas. *Ecol. Lett.* **14**: 484–492.
- 892 **Archaux, F.** 2004. Breeding upwards when climate is becoming warmer: no bird response in the French Alps.  
893 *Ibis* **146**: 138-144.
- 894 **Archaux, F.** 2007. Are mountains refuges for farmland bird species? A case study in the northern French Alps.  
895 *Bird Study* **54**: 73–79.
- 896 **Arlettaz, R., Patthey, P., Baltic, M., Leu, T., Schaub, M., Palme, R. & Jenni-Eiermann, S.** 2007. Spreading free-  
897 riding snow sports represent a novel serious threat for wildlife. *Proc. R. Soc. B* **274**: 1219–1224.
- 898 **Arlettaz, R., P. Patthey & Braunisch, V.** 2013. Impacts of Outdoor Winter Recreation on Alpine Wildlife and  
899 Mitigation Approaches: A Case Study of the Black Grouse. In: *The Impacts of Skiing and Related Winter*  
900 *Recreational Activities on Mountain Environments* (eds C. Rixen & A. Rolando), pp. 137-154. Bentham  
901 eBooks, Bussum.
- 902 **Arlettaz, R., Nusslé, S., Baltic, M., Vogel, P., Palme, R., Jenni-Eiermann, S., Patthey, P. & Genoud, M.** 2015.  
903 Disturbance of wildlife by outdoor winter recreation: allostatic stress response and altered activity-energy  
904 budgets. *Ecol. App.* **25**: 1197-1212.
- 905 **Arnold, E.R. & Greenfield, H.J.** 2006. *The Origins of Transhumant Pastoralism in Temperate South Eastern*  
906 *Europe. A zooarchaeological perspective from the Central Balkans.* Oxford: BAR International Series 1538.
- 907 **Aryal, S., Maraseni, T. N. & Cockfield, G.** 2014. Sustainability of transhumance grazing systems under socio-  
908 economic threats in Langtang, Nepal. *J. Mt. Sci.* **11**: 1023-1034.
- 909 **Ashton, K. G.** 2002. Patterns of within-species body size variation of birds: strong evidence for Bergmann's  
910 rule. *Global Ecol. Biogeogr.* **11**: 505-523.
- 911 **Auer, S. & King, D.** 2014. Ecological and life-history traits explain recent boundary shifts in elevation and  
912 latitude of western North American songbirds. *Glob. Ecol. Biogeogr.* **23**: 867–875.
- 913 **Baines, D. & Andrew, M.** 2003. Marking of deer fences to reduce frequency of collisions by woodland grouse.  
914 *Biol. Conserv.* **110**: 169-176.
- 915 **Barnagaud, J.Y., Crochet, P.A., Magnani, Y., Bernard Laurent, A., Menoni, E., Novoa, C. & Gimenez, O.** 2011.  
916 Short-term response to the North Atlantic Oscillation but no long-term effects of climate change on the  
917 reproductive success of an alpine bird. *J. Ornithol.* **152**: 631–641.
- 918 **Battaglini, L., Bovolenta, S., Gusmeroli, F., Salvador, S. & Sturaro, E.** 2014. Environmental sustainability of  
919 Alpine livestock farms. *Ital. J. Anim. Sci.* **13**: 431-443.
- 920 **Bazzi, G., Foglini, C., Brambilla, M., Saino, N. & Rubolini, D.** 2015. Habitat management effects on Prealpine  
921 grassland bird communities. *Ital. J. Zool.* **82**: 251-261.

- 922 **Bech, N., Boissier, J., Drovetski, S. & Novoa, C.** 2009. Population genetic structure of rock ptarmigan in the 'sky  
923 islands' of French Pyrenees: implications for conservation. *Anim. Conserv.* **12**: 138–146.
- 924 **Beniston, M. & Rebetez, M.** 1996. Regional behavior of minimum temperatures in Switzerland for the period  
925 1979 - 1993. *Theor. Appl. Climatol.* **53**: 231 – 243.
- 926 **Beniston, M., Keller, F. & Goyette S.** 2003. Snow pack in the Swiss Alps under changing climatic conditions: an  
927 empirical approach for climate impact studies. *Theor. Appl. Climatol.* **74**: 19–31.
- 928 **Berteaux, D., Gauthier, G., Dominé, F., Ims, R.A., Lamoureux, S.F., Lévesque, E. & Yoccoz, N.** 2016. Effects of  
929 changing permafrost and snow conditions on tundra wildlife: critical places and times. *Arctic Science*,  
930 10.1139/as-2016-0023.
- 931 **BirdLife International & NatureServe.** 2015. *Bird species distribution maps of the world*. BirdLife International,  
932 Cambridge, UK and NatureServe, Arlington, USA.
- 933 **Bodin, J., Badeau, V., Bruno, E., Cluzeau, C., Moisselin, J.M., Walther, G.R. & Dupouey, J.L.** 2013. Shifts of forest  
934 species along an elevational gradient in Southeast France: climate change or stand maturation. *J. Veg. Sci.*  
935 **24**: 269–283.
- 936 **Boelman, N.T., Gough, L., Wingfield, J., Goetz, S., Asmus, A., Chmura, H. E., Krause, J. S., Perez, J. H., Sweet, S.  
937 K. & Guay, K. C.** 2015. Greater shrub dominance alters breeding habitat and food resources for migratory  
938 songbirds in Alaskan arctic tundra. *Glob. Change Biol.* **21**: 1508–1520.
- 939 **Böhning-Gaese, K., & Lemoine, N.** 2004. Importance of climate change for the ranges, communities and  
940 conservation of birds. *Adv. Ecol. Res.* **35**: 211-236.
- 941 **Both, C., Bouwhuis, S., Lessells, C.M. & Visser, M.E.** 2006. Climate change and population declines in a long-  
942 distance migratory bird. *Nature* **441**: 81–83.
- 943 **Boyle, A.W. & Martin, K.** 2015. The conservation value of high elevation habitats to North American migrant  
944 birds. *Biol. Conserv.* **192**: 461-476.
- 945 **Boyle, A.W., Sandercock, B.K. & Martin, K.** 2016. Patterns and drivers of intraspecific variation in avian life  
946 history along elevational gradients: a meta-analysis. *Biol. Rev.* **91**: 469-482.
- 947 **Brambilla, M., Bassi, E., Bergero, V., Casale, F., Chemollo, M., Falco, R., Longoni, V., Saporetti, F., Viganò, E. &  
948 Vitulano, S.** 2013. Modelling distribution and potential overlap between Boreal Owl *Aegolius funereus* and  
949 Black Woodpecker *Dryocopus martius*: implications for management and monitoring plans. *Bird Conserv.*  
950 *Int.* **23**: 502-511.
- 951 **Brambilla, M. & Gobbi, M.** 2014. A century of chasing the ice: delayed colonisation of ice-free sites by ground  
952 beetles along glacier forelands in the Alps. *Ecography* **37**: 33–42.
- 953 **Brambilla, M., Bergero, V., Bassi, E. & Falco, R.** 2015. Current and future effectiveness of Natura 2000 network  
954 in the central Alps for the conservation of mountain forest owl species in a warming climate. *Eur. J. Wildl.*  
955 *Res.* **61**: 35–44.
- 956 **Brambilla, M., Pedrini, P., Rolando, A. & Chamberlain, D.** 2016a. Climate change will increase the potential  
957 conflict between skiing and high-elevation bird species in the Alps. *J. Biogeogr.* **43**: 2299–2309.
- 958 **Brambilla, M., Cortesi, M., Capelli, F., Chamberlain, D., Pedrini, P., Rubolini, D.** 2016b. Foraging habitat  
959 selection by Alpine White-winged Snowfinches *Montifringilla nivalis* during the nestling rearing period. *J.*  
960 *Ornithol.* **158**: 277-286.
- 961 **Braunisch, V., P. Patthey & Arlettaz, R.** 2011. Spatially explicit modeling of conflict zones between wildlife and  
962 snow sports: prioritizing areas for winter refuges. *Ecol. App.* **21**: 955-967.
- 963 **Braunisch, V., Coppes, J., Arlettaz, R., Suchant, R., Schmid, H. & Bollmann, K.** 2013. Selecting from correlated  
964 climate variables: a major source of uncertainty for predicting species distributions under climate change.  
965 *Ecography* **36**: 971–983.
- 966 **Braunisch, V., Coppes, J., Arlettaz, R., Suchant, R., Zellweger, F. & Bollmann, K.** 2014. Temperate mountain  
967 forest biodiversity under climate change: compensating negative effects by increasing structural  
968 complexity. *PLoS ONE* **9**: e97718.
- 969 **Braunisch, V., P. Patthey & Arlettaz, R.** 2016. Where to Combat Shrub Encroachment in Alpine Timberline  
970 Ecosystems: Combining Remotely-Sensed Vegetation Information with Species Habitat Modelling. *PLoS*  
971 *ONE* **11**: e0164318.

- 972 Broggi, J., Gamero, A., Hohtola, E., Orell, M., Nilsson, J.Å. 2011. Interpopulation Variation in Contour Feather  
973 Structure Is Environmentally Determined in Great Tits. *PLoS ONE* **6**: e24942.
- 974 Brown, J.L., Li, S.H. & Bhagabati, N. 1999. Long-term trend toward earlier breeding in an American bird: a  
975 response to global warming? *Proc. Natl. Acad. Sci. U.S.A.* **96**: 5565–5569.
- 976 Camfield, A.F., Pearson, S. & Martin, K. 2010. Life history variation between high and low elevation  
977 subspecies of horned larks *Eremophila spp.* *J. Avian Biol.* **41**: 273-281.
- 978 Caprio, E., Chamberlain, D., Isaia, M. & Rolando, A. 2011. Landscape changes caused by high altitude ski-pistes  
979 affect bird species richness and distribution in the Alps. *Conserv. Biol.* **144**: 2958–2967.
- 980 Caprio, E., Chamberlain, D. & Rolando, A. 2016. Ski-piste revegetation promotes partial bird community  
981 recovery in the European Alps. *Bird study* **63**: 470-478.
- 982 Carroll, M., Dennis, P., Pearce-Higgins, J.W. & Thomas, C.D. 2011. Maintaining northern peatland ecosystems  
983 in a changing climate: effects of soil moisture, drainage and drain blocking on Craneflies. *Glob. Chang.*  
984 *Biol.* **17**: 2991–3001.
- 985 Carroll, M., Heinemeyer, A., Pearce-Higgins, J., Dennis, P., West, C., Holden, J., Wallage, Z. & Thomas, C. 2015.  
986 Hydrologically-driven ecosystem processes determine the distribution and persistence of ecosystem-  
987 specialist predators under climate change. *Nat. Commun.* doi: 10.1038/ncomms8851.
- 988 Chamberlain, D., Arlettaz, R., Caprio, E., Maggini, R., Pedrini, P., Rolando, A. & Zbinden, N. 2012. The altitudinal  
989 frontier in avian climate impact research. *Ibis* **154**: 205–209.
- 990 Chamberlain, D., Negro, M., Caprio, E. & Rolando, A. 2013. Assessing the sensitivity of alpine birds to potential  
991 future changes in habitat and climate to inform management strategies. *Biol. Conserv.* **167**: 127–135.
- 992 Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. 2011. Rapid range shifts of species associated  
993 with high levels of climate warming. *Science* **333**: 1024 – 1026.
- 994 Cheviron Z.A. & Brumfield, R.T. 2012. Genomic insights into adaptation to high-altitude environments.  
995 *Heredity* **108**: 354–361.
- 996 Clavero, M. & Brontons, L. 2010. Functional homogenization of bird communities along habitat gradients:  
997 accounting for niche multidimensionality. *Glob. Ecol. Biogeogr.* **19**: 684-696.
- 998 Conroy, M.J., Runge, M.C., Nichols, J.D., Stodola, K.W. & Cooper, R.J. 2011. Conservation in the face of climate  
999 change: the roles of alternative models, monitoring, and adaptation in confronting and reducing  
1000 uncertainty. *Biol. Conserv.* **144**: 1204–1213.
- 1001 Crick, H.Q.P., Dudley, C., Glue, D.E. & Thomson, D.L. 1997. UK birds are laying eggs earlier. *Nature* **388**: 526.
- 1002 Cumming, S.G., Stralberg, D., Lefevre, K.L., Solymos, P., Bayne, E.M., Fang, S., Fontaine, T., Mazerolle, D.,  
1003 Schmiegelow, F.K.A. & Song, S.J. 2014. Climate and vegetation hierarchically structure patterns of  
1004 songbird distribution in the Canadian boreal region. *Ecography* **37**: 137–151.
- 1005 Dale, V.H., Joyce, L.A., McNulty, S., Neilson, R.P., Ayres, M.P., Flannigan, M.D., Hanson, P.J., Irland, L.C., Lugo,  
1006 A.E., Peterson, C.J., Simberloff, D., Swanson, F.J., Stocks, B.J. & Wotton, B.M. 2001. Climate change and  
1007 forest disturbances. *Bioscience* **51**: 723-734
- 1008 DeLuca, W.V. & King, D. I. 2014. Influence of hiking trails on montane birds. *Jour. Wild. Mgmt.* **78**: 494–  
1009 502.
- 1010 DeLuca, W.V. & King, D.I. 2017. Montane birds shift downslope despite recent warming in the northern  
1011 Appalachian Mountains. *J. Ornithol.* **158**: 493.
- 1012 Debarbieux, B., Oiry Varacca, M., Rudaz, G, Maselli, D., Kohler, T. & Jurek, M. 2014. Tourism in Mountain  
1013 Regions: Hopes, Fears and Realities. Sustainable Mountain Development Series. Geneva, Switzerland:  
1014 UNIGE, CDE, SDC, pp. 108.
- 1015 Devictor, V., Julliard, R., Couvet, D. & Jiguet, F. 2008. Birds are tracking climate warming, but not fast enough.  
1016 *Proc. R. Soc. B* **275**: 2743–2748.
- 1017 Diaz, H.R., Grosjean, M. & Graumlich, L. 2003. Climate variability and change in high elevation regions: past,  
1018 present and future. *Clim. Chang.* **59**: 1-4.
- 1019 Dillon, M., Frazier, M. & Dudley, R. 2006. Into thin air: Physiology and evolution of alpine insects. *Integr.*  
1020 *Comp. Biol.* **46**: 49–61.



- 1021 Douglas, D.J.T. & Pearce-Higgins, J. W. 2014. Relative importance of prey abundance and habitat structure as  
1022 drivers of shorebird breeding success and abundance. *Anim. Conserv.* **17**: 535–543.
- 1023 Douglas, D.J.T., Buchanan, G.M., Thompson, P., Amar, A., Fielding, D.A., Redpath, S.M. & Wilson, J.D. 2015.  
1024 Vegetation burning for game management in the UK uplands is increasing and overlaps spatially with soil  
1025 carbon and protected areas. *Biol. Conserv.* **191**: 243–250.
- 1026 Dragon, S., Carey, C., Martin, K. & Baumann, R. 1999. Effect of high altitude and in vivo adenosine/b-  
1027 adrenergic receptor blockade on ATP and 2,3BPG concentrations in red blood cells of avian embryos. *J.*  
1028 *Exp. Biol.* **202**: 2787–2795.
- 1029 Dreitz, V. J., Yale Conrey, R. & Skagen, S. K. 2012. Drought and cooler temperatures are associated with higher  
1030 nest survival in Mountain Plovers. *Aian Conserv. Ecol.* **7**: 6.
- 1031 Eglinton, S.M. & Pearce-Higgins, J.W. 2012. Disentangling the relative importance of changes in climate and  
1032 land-use intensity in driving recent bird population trends. *PLoS ONE* **7**: e30407.
- 1033 Elmhagen, B., Kindberg, J., Hellström, P. & Angerbjörn, A. 2015. A boreal invasion in response to climate  
1034 change? Range shifts and community effects in the borderland between forest and tundra. *AMBIO* **44**:  
1035 39–50.
- 1036 Elsen, P.R., Tingley, M.W. 2015 Global mountain topography and the fate of montane species under climate  
1037 change. *Nat. Clim. Change* **5**: 772–776.
- 1038 Engler, R., Randin, C.F., Thuiller, W., Dullinger, S., Zimmermann, N. E., Araújo, M.B., Pearman, P.B.,  
1039 Gwenaëlle, I.I., Piedallu, C., Albert, C.H., Choler, P., Coldea, G., De Iamo, X., Dirnböck, T., Gégout, J.C., Gómez-  
1040 García, D., Grytnes, J.A., Heegaard, E., Høistad, F., Nogués-Bravo, D., Normand, S., Puşcaş, M., Sebastià,  
1041 M.T., Stanisci, A., Theurillat, J.P., Trivedi, M.R., Vittoz, P. & Guisan, A. 2011. 21st century climate change  
1042 threatens mountain flora unequally across Europe. *Glob. Chang. Biol.* **17**: 2330–2341.
- 1043 Evans, D. M., Redpath, S. M., Evans, S. A., Elston, D. A., Gardner, C. J., Dennis, P. & Pakeman, R. J. 2006. Low  
1044 intensity, mixed livestock grazing improves the breeding abundance of a common insectivorous  
1045 passerine. *Biology Letters* **2**: 636–638.
- 1046 FAO 2015. Mapping the vulnerability of mountain peoples to food insecurity. Romeo, R., Vita, A., Testolin, R.  
1047 & Hofer, T. Rome.
- 1048 Fjeldså, J., Bowie, R.C.K. & Rahbek, C. 2012. The role of mountain ranges in the diversification of birds. *Annu.*  
1049 *Rev. Ecol. Evol. Syst.* **43**: 249–265.
- 1050 Fletcher, K., Howarth, D., Kirby, A., Dunn, R. & Smith, A. 2013. Effect of climate change on breeding  
1051 phenology, clutch size and chick survival of an upland bird. *Ibis* **155**: 456–463.
- 1052 Flousek, J., Telensky, T., Hanzelka, J. & Reif, J. 2015. Population trends of central European montane birds  
1053 provide evidence for adverse impacts of climate change on high-altitude species. *PLoS ONE* **10**: e0139465.
- 1054 Freeman, B. & Montgomery, G. 2015. Interspecific aggression by Swainson’s Thrush (*Catharus ustulatus*) may  
1055 limit the distribution of the threatened Bicknell’s Thrush (*Catharus bicknelli*) in the Adirondack Mountains.  
1056 *Condor* **118**: 169–178.
- 1057 Freppaz, M., Filippa, G., Caimi, A., Buffa, G. & Zanini, E. 2010. Soil and plant characteristics in the alpine tundra  
1058 (NW Italy). In: Gutierrez, B., Pena, C. (Eds.). *Tundras: Vegetation, Wildlife and Climate Trends*. Nova  
1059 Science Publishers, New York: 81–110.
- 1060 Frey, S. J. K., Hadley, A. S. & Betts, M. G. 2016. Microclimate predicts within-season distribution dynamics of  
1061 montane forest birds. *Divers. Distrib.* **22**: 944–959.
- 1062 Furrer, R., Schaub, M., Bossert, A., Isler, R., Jenny, H., Jonas, T., Marti, C. & Jenni, L. 2016. Variable decline of  
1063 alpine Rock Ptarmigan (*Lagopus muta helvetica*) in Switzerland between regions and sites. *J. Ornithol.*  
1064 **157**: 787–796.
- 1065 García-González, R., Aldezabal, A., Laskurain, N.A., Margalida, A. & Novoa, C. 2016. Influence of snowmelt  
1066 timing on the diet quality of Pyrenean Rock Ptarmigan (*Lagopus muta pyrenaica*): implications for  
1067 reproductive success. *PLoS ONE* **11**: doi:10.1371/journal.pone.0148632.
- 1068 Gaüzère, P., Jiguet, F. & Devictor, V. 2016. Can protected areas mitigate the impacts of climate change on  
1069 bird's species and communities? *Divers. Distrib.* **22**: 625–637.

- 1070 Gehrig-Fasel, J., Guisan, A. & Zimmermann, N. 2007. Tree line shifts in the Swiss Alps: climate change or land  
1071 abandonment? *J. Veg. Sci.* **18**: 571–582.
- 1072 Gellrich, M. & Zimmermann, N.E. 2007. Land abandonment in the Swiss mountains: A spatial statistical  
1073 analysis at the regional scale. *Landscape Urban Plan.* **79**: 65–76.
- 1074 Gentle, P. & Thwaites, R. 2016. Transhumant pastoralism in the context of socioeconomic and climate change  
1075 in the mountains of Nepal. *Mt. Res. Dev.* **36**: 173–182.
- 1076 Gerrard, A.J. 1990. *Mountain environments: an examination of the physical geography of mountains*. MIT  
1077 Press.
- 1078 Gienapp, P., Leimu, R. & Merila, J. 2007. Responses to climate change in avian migration time – micro-  
1079 evolution versus phenotypic plasticity. *Clim. Res.* **35**: 25–35.
- 1080 Giuliano, W. 1994. The impact of hiking and rock climbing in mountain areas. *Environ. Conserv.* **21**: 287–279.
- 1081 Glutz von Blotzheim, U.N., Bauer, K.M., Bezzel, E. 1973. *Handbuch der Vögel Mitteleuropas*, **5**: Galliformes und  
1082 Gruiformes. Akademische Verlagsgesellschaft, Frankfurt.
- 1083 Goodenough A.E. & Hart A.G. 2013. Correlates of vulnerability to climate-induced distribution changes in  
1084 European avifauna: habitat, migration and endemism. *Clim. Change* **118**: 659–669.
- 1085 Grigaltchik, V.S., Ward, A.J.W. & Seebacher, F. 2012. Thermal acclimation of interactions: differential  
1086 responses to temperature change alter predator-prey relationship. *Proc. R. Soc. A* **279**: 4058–4064.
- 1087 Hågvar, S. 2010. A review of Fennoscandian arthropods living on and in snow. *Eur. J. Entomol.* **107**: 281–298.
- 1088 Hannon, S. J., Martin, K. & Schieck, J.A. 1988. Timing of reproduction in two populations of Willow Ptarmigan  
1089 in northern Canada. *Auk* **105**: 330–338.
- 1090 Hardy, D.R. & Hardy, S.P. 2008. White-winged Diuca Finch (*Diuca speculifera*) nesting on Quelccaya Ice Cap,  
1091 Perú. *Wilson J. Ornithol.* **120**: 613–617.
- 1092 Harsch, M.A., Hulme, P.E., McGlone, M.S. & Duncan, R.P. 2009. Are treelines advancing? A global meta-  
1093 analysis of treeline response to climate warming. *Ecol. Lett.* **12**: 1040–1049.
- 1094 Heikkinen, R. K., Luoto, M., Virkkala, R., & Pearson, R.G. & Körber, J.H. 2007. Biotic interactions improve  
1095 prediction of boreal bird distributions at macro-scales. *Glob. Ecol. Biogeogr.* **16**: 754 – 763.
- 1096 Heilprin, A. 1887. *The Geographical and Geological Distribution of Animals*. International Scientific Series:  
1097 London.
- 1098 Hickling, R., Roy, D.B., Hill, J.K., Fox, R. & Thomas, C.D. 2006. The distributions of a wide range of taxonomic  
1099 groups are expanding polewards. *Glob. Chang. Biol.* **12**: 450– 455.
- 1100 Hoiss, B., Gaviria, J., Leingartner, A., Krauss, J. & Steffan-Dewenter, I. 2013. Combined effects of climate and  
1101 management on plant diversity and pollination type in alpine grasslands. *Divers. Distrib.* **19**: 386–395.
- 1102 Huntley, B., Collingham, Y.C., Willis, S.G. & Green, R.E. 2008. Potential impacts of climate change on European  
1103 breeding birds. *PLoS ONE* **3**: e1439.
- 1104 Hüppop, O. & Hüppop, K. 2003. North Atlantic Oscillation and timing of spring migration in birds. *Proc. R. Soc.*  
1105 *A* **270**: 233–240.
- 1106 Imperio, S., Bionda, R., Viterbi, R. & Provenzale, A. 2013. Climate change and human disturbance can lead to  
1107 local extinction of Alpine rock ptarmigan: new insight from the western Italian Alps. *PLoS ONE* **8**: e81598.
- 1108 Inouye, D.W., Barr, B., Armitage, K.B. & Inouye, B.D. 2000. Climate change is affecting altitudinal migrants and  
1109 hibernating species. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 1630–1633.
- 1110 IPCC 2007. *Climate change 2007: Synthesis report*. Intergovernmental Panel on Climate Change, Paris.
- 1111 IPCC 2013 *Climate change 2013: the physical science basis*. In: Stocker TF et al. (eds) Working group I  
1112 contribution to the fifth assessment report of the intergovernmental panel on climate change. Cambridge  
1113 University Press, Cambridge, pp 1–14.
- 1114 Jackson, M.M., Gergel, S.E. & Martin, K. 2015. Effects of climate change on habitat availability and  
1115 configuration for an endemic coastal alpine bird. *PLoS ONE* **10**: e0142110.
- 1116 Jetz, W., Wilcove, D.S. & Dobson, A.P. 2007. Projected impacts of climate and land-use change on the global  
1117 diversity of birds. *PLoS Biol.* **5**: 1211–1219.



- 1118 **Johnson, R.** 1965. Reproductive activities of rosy finches, with special reference to Montana. *The Auk* **82**: 190  
 1119 –205.
- 1120 **Johnson, R.** 1968. Temperature regulation in the white-tailed ptarmigan, *Lagopus leucurus*. *Comp. Biochem.*  
 1121 *Physiol.* **24**:1003–1014.
- 1122 **Kapos, V., J. Rhind, J., Edwards, M., Price, M.F. & Ravilious, C.** 2000. *Developing a map of the world's mountain*  
 1123 *forests*. In: Forests in Sustainable Mountain Development: A State-of-Knowledge Report for 2000, M.F.  
 1124 Price and N. Butt (eds.), CAB International, Wallingford: 4–9.
- 1125 **Kearney, M. & Porter, W.** 2009. Mechanistic niche modelling: combining physiological and spatial data to  
 1126 predict species' ranges. *Ecol. Lett.* **12**: 334 – 350.
- 1127 **Kirby, K.J. & Watkins, C.** 2015. *Europe's changing woods and forests: from wildwood to managed landscapes*.  
 1128 CABI, Wallingford, UK.
- 1129 **Klein, G., Vitasse, Y., Rixen, C., Marty, C., & M. Rebetez, M.** 2016. Shorter snow cover duration since 1970 in  
 1130 the Swiss Alps due to earlier snowmelt more than to later snow onset. *Clim. Chang*, **139**: 637, doi:  
 1131 10.1007/s10584-016-1806-y.
- 1132 **Kohler, T., Wehrli, A. & Jurek, M.** 2014. *Mountains and climate change: a global concern. Sustainable*  
 1133 *mountain development series*. Bern, Switzerland, Centre for Development and Environment (CDE), Swiss  
 1134 Agency for Development and Cooperation (SDC) and Geographica Bernensia: 136.
- 1135 **Koricheva, J., Gurevitch, J. & Mengersen, K.** 2013. *Handbook of Meta-analysis in Ecology and Evolution*.  
 1136 Oxford: Princeton University Press.
- 1137 **Körner, C. & Spehn, E. M.** 2002. *Mountain biodiversity: a global assessment*. Boca Raton, FL: Parthenon  
 1138 Publication Group.
- 1139 **Körner, C. & Ohsawa, M.** 2006. *Mountain systems*. In: Hassan R, Scholes R, Ash N (eds) Ecosystem and human  
 1140 well-being: current state and trends. Millennium Ecosystem Assessment, Island Press, Washington **1**: 681-  
 1141 716.
- 1142 **Körner, C.** 2012. *Alpine treelines: Funct. Ecol. of the global high elevation tree limits*. Basel, Switzerland:  
 1143 Springer.
- 1144 **Knudsen, E., Linden, A., Both, C., Jonzen, N., Pulido, F., Saino, N., Sutherland, W.J., Bach, L.A., Coppack, T.,**  
 1145 **Ergon, T., Gienapp, P., Gill, J.A., Gordo, O., Hedenström, A., Lehikoinen, E., Marra, P.P., Møller, A.P., Nilsson,**  
 1146 **A.L.K., Péron, G., Ranta, E., Rubolini, D., Sparks, T.H., Spina, F., Studds, C.E., Sæther, S.A., Tryjanowski, P. &**  
 1147 **Stenseth, N.C.** 2011. Challenging claims in the study of migratory birds and climate change. *Biol. Rev.* **86**:  
 1148 928–946.
- 1149 **Laaksonen, T., Ahola, M., Eeva, T., Väisänen, R.A. & Lehikoinen, E.** 2006. Climate change, migratory  
 1150 connectivity and changes in laying date and clutch size of the Pied Flycatcher. *Oikos* **114**: 277–290.
- 1151 **Laiolo, P., Rolando, A. & Valsania, V.** 2004. Responses of birds to the natural re-establishment of wilderness in  
 1152 montane beechwoods of North-western Italy. *Acta Oecol.* **25**: 129–136.
- 1153 **Laiolo, P. & Rolando, A.** 2005. Forest bird diversity and ski-runs: a case of negative edge effect. *Anim. Conserv.*  
 1154 **7**: 9–16.
- 1155 **Lajeunesse, M.J.** 2013. Recovering missing or partial data from studies: A survey of conversions and  
 1156 imputations for meta-analysis. In Koricheva, J., Gurevitch, J. & Mengersen, K (eds) *Handbook of Meta-*  
 1157 *Analysis in Ecology and Evolution*. Princeton: Princeton University Press.
- 1158 **Lande, R.** 1993. Risks of population extinction from demographic and environmental stochasticity and  
 1159 random catastrophes. *Am. Nat.* **142**: 911–927.
- 1160 **La Sorte, F.A. & Jetz, W.** 2010. Projected range contractions of montane biodiversity under global warming.  
 1161 *Proc. R. Soc. A* **277**: 3401–3410.
- 1162 **Lehikoinen, A., Green, M., Husby, M., Kålås, J.E. & Lindström, Å.** 2014. Common montane birds are declining in  
 1163 Northern Europe. *J. Avian Biol.* **45**: 3–14.
- 1164 **Lehikoinen, A. & Virkkala, R.** 2016. North by north-west: climate change and directions of density shifts in  
 1165 birds. *Glob. Chang. Biol.* **22**: 1121–1129.

- 1166 Lenoir, J., Gégout, J.C., Marquet, P.A., de Ruffray, P. & Brisse, H. 2008 A significant upward shift in plant  
1167 species optimum elevation during the 20th century. *Science* **320**: 1768–1771.
- 1168 Lesk, C., Coffel, E., D’Amato, A.W., Dodds, K., Horton, R. 2017. Threats to North American forests from  
1169 southern pine beetle with warming winters. *Nat. Clim. Change* **7**: 713–717.
- 1170 Liu, X.D. & Chen, B.D. 2000. Climatic warming in the Tibetan Plateau during recent decades. *Int. J. Clim.* **20**:  
1171 1729–1742.
- 1172 Loarie, S., Duffy, P., Hamilton, H. & Asner, G. 2009. The velocity of climate change. *Nature* **462**: 1052–5.
- 1173 Loe, L. E., Mysterud, A., Stien, A., Steen, H., Evans, D. M. & Austrheim, G. 2007. Positive short-term effects of  
1174 sheep grazing on the alpine avifauna. *Biology Letters* **3**: 109–111.
- 1175 Lu, N., Jing, Y., Lloyd, H., Sun, Y. 2012. Assessing the distributions and potential risks from climate change for  
1176 the Sichuan Jay (*Perisoreus internigrans*). *The Condor* **114**: 365–376.
- 1177 Lundmark, L. 2007. Reindeer pastoralism in Sweden 1550–1950. *Rangifer* **12**: 9–16.
- 1178 Maggini, R., Lehmann, A., Kery, M., Schmid, H., Beniston, M., Jenni, L. & Zbinden, N. 2011. Are Swiss birds  
1179 tracking climate change? Detecting elevational shifts using response curve shapes. *Ecol. Model.* **222**: 21–  
1180 32.
- 1181 Maggini, R., Lehman, A., Zbinden, N., Zimmerman, N.E., Bollinger, J., Schröder, B., Foppen, R., Schmid, H.,  
1182 Beniston, M. & Jenni, L. 2014. Assessing species vulnerability to climate and land use change: the case of  
1183 the Swiss breeding birds. *Divers. Distrib.* **20**: 708–719.
- 1184 Mahon, C.L., Holloway, G., Sólymos, P., Cumming, S.G., Bayne, E.M. Schmiegelow F.K.A. & Song, S.J. 2016.  
1185 Community structure and niche characteristics of upland and lowland western boreal birds at multiple  
1186 spatial scales. *For. Ecol. Manage.* **361**: 99–116.
- 1187 Mantyka-Pringle, C. S. & Rhodes, J.R. 2012. Interactions between climate and habitat loss effects on  
1188 biodiversity: A systematic review and meta-analysis. *Glob. Chang. Biol.* **18**: 1239– 1252.
- 1189 Martin, T.E. 2001a. Abiotic vs biotic influences on habitat selection of coexisting species: climate change  
1190 impacts? *Ecology* **82**: 175–188.
- 1191 Martin, K. 2001b. *Wildlife communities in alpine and sub-alpine habitats*. In D. H. Johnson and T. A. O’Neil  
1192 (Managing Directors), *Wildlife-habitat relationships in Oregon and Washington*. Oregon State Univ. Press,  
1193 Corvallis, Oregon: 285–310.
- 1194 Martin, T.E. & Maron, J.L. 2012. Climate impacts on bird and plant communities from altered animal-plant  
1195 interactions. *Nat. Clim. Chang.* **2**: 195–200.
- 1196 Martin, K., & Wiebe, K. L. 2004. Coping mechanisms of alpine and arctic breeding birds: extreme weather and  
1197 limitations to reproductive resilience. *Integr. Comp. Biol.* **44**: 177–185.
- 1198 Martin, K., Wilson, S. D., MacDonald, E.C., Camfield, A. F., Martin, M. & Trefrey, S. A. 2017. Effects of severe  
1199 weather on reproduction for sympatric songbirds in an alpine environment: interactions of climate  
1200 extremes influence nesting success. *The Auk* **134**: 696–709.
- 1201 Massimino, D., Johnston, A. & Pearce-Higgins, J.W. 2015. The geographical range of British birds expands  
1202 during 15 years of warming. *Bird Study* **62**: 523–534.
- 1203 Mayor, S.J., Guralnick, R.P., Tingley, M. W., Outegui, J., Withey, J.C., Elmendorf, S. C., Andrew, M.E., Leyk, S.,  
1204 Pearse, I.S. & Schneider, D.C. 2017 Increasing phenological asynchrony between spring green-up and  
1205 arrival of migratory birds. *Sci. Rep.* **7**: 1902 doi:10.1038/s41598-017-02045-z
- 1206 McVean, D.; & Lockie, J.D. 1969. *Ecology and Land Use in Upland Scotland*. Edinburgh: University Press.
- 1207 Meehl, G.A., Covey, C., Delworth, T., Latif, M., McAvaney, B., Mitchell, J.F.B., Stouffer, R.J. & Taylor, K.E. 2007.  
1208 The WCRP CMIP3 multimodel dataset—a new era in climate change research. *B. Am. Meteorol. Soc.* **88**:  
1209 1383–1394.
- 1210 Mezquida, E.T., Svenning, J.C., Summers, R.W., Benkman, C.W. 2017. Higher spring temperatures increase  
1211 food scarcity and limit the current and future distributions of crossbills. *Divers. Distrib.*  
1212 doi.org/10.1111/ddi.12694

- 1213 Mizel, J. D., Schmidt, J. H., McIntyre, C. L. & Roland C. A. 2016. Rapidly shifting elevational distributions of  
 1214 passerine species parallel vegetation change in the subarctic. *Ecosphere* **7**: e01264.
- 1215 Moser, B. W. & Witmer, G. W. 2000. The effects of elk and cattle foraging on the vegetation, birds, and small  
 1216 mammals of the Bridge Creek Wildlife Area, Oregon. *Int. Biodeterior. Biodegradation* **45**: 151-157.
- 1217 Moss, R., Oswald, J. & Baines, D. 2001. Climate change and breeding success: decline of the Capercaillie in  
 1218 Scotland. *J. Anim. Ecol.* **70**: 47-61.
- 1219 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. 2000. Biodiversity hotspots for  
 1220 conservation priorities. *Nature* **403**: 853–858.
- 1221 Myers-Smith, I.H., Forbes, B.C., Wilmsking, M., Hallinger, M., Lantz, t., Blok, D., Tape, K.D., Macias-Fauria, M.,  
 1222 Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P., Hermanutz, L., Trant, A., Collier, L.S., Weijers, S.,  
 1223 Rozema, J., Rayback, S.S., Schmidt, N.M., Schaepman-Strub, G., Wipf, S., Rixen, C., Ménard, C.C., Venn, S.,  
 1224 Goetz, S., Andreu-Hayles, L., Elmendorf, S., Ravolainen, V., Welker, J., Grogan, P., Epstein, H.E. & Hik, D.S.  
 1225 2011. Shrub expansion in tundra eco- systems: dynamics, impacts and research priorities. *Environ. Res.*  
 1226 *Lett.* **6**: 045509.
- 1227 Møller, A.P., Rubolini, D. & Lehikoinen, E. 2008. Populations of migratory bird species that did not show a  
 1228 phenological response to climate change are declining. *Proc. Natl. Acad. Sci. U.S.A.* **105**: 16195–16200.
- 1229 Mollet, P., Bollmann, K., Braunisch, V. & Arlettaz, R. In press. Avian communities in European high altitude  
 1230 woodlands. In: Mikusinski, G., Roberge, J.-M. & Fuller, R.J. (Eds.). *Ecology and Conservation of Forest*  
 1231 *Birds*. Cambridge University Press. Cambridge, UK, in press.
- 1232 Musitelli, F., Romano, A., Møller, A.P. & Ambrosini, R. 2016. Effects of livestock farming on birds of rural areas  
 1233 in Europe. *Biodivers. Conserv.* **25**: 615-631.
- 1234 Nagy, L. & Grabherr, G. 2009. *The biology of alpine habitats*. Oxford University Press, Oxford.
- 1235 Nogués-Bravo, D., Araújo, M.B., Erread, M.P. & Martínez-Ricad, J.P. 2007. Exposure of global mountain  
 1236 systems to climate warming during the 21st Century. *Glob. Environ. Chang.* **17**: 420–428.
- 1237 Nogués-Bravo, D., Araujo, M. B., Romdal, T. S. & Rahbek, C. 2008. Scale effects and human impact on the  
 1238 elevational species richness gradients. *Nature* **453**: 216-220.
- 1239 Novoa, C., Besnard, A., Brenot, J.F. & Ellison, L.N. 2008. Effect of weather on the reproductive rate of Rock  
 1240 Ptarmigan *Lagopus muta* in the Eastern Pyrenees. *Ibis* **150**: 270–278.
- 1241 Novoa, C., Astruc, G., Desmet, J.F. & Besnard, A. 2016. No short-term effects of climate change on the  
 1242 breeding of Rock Ptarmigan in the French Alps and Pyrenees. *J. Ornithol.* **157**: 797-810.
- 1243 Oliver, T. H., Gillings, S., Pearce-Higgins, J. W., Brereton, T., Crick, H. Q. P., Duffield, S. J., Morecroft, M. D. & Roy,  
 1244 D. B. 2017. Large extents of intensive land use limit community reorganization during climate warming.  
 1245 *Glob. Change Biol.* **23**: 2272–2283.
- 1246 Pacifici, M., Visconti, P., Butchard, S.H.M., Watson, J.E.M., Cassola, F.M. & Rondinini, C. 2017. Species' traits  
 1247 influenced their response to recent climate change. *Nat. Clim. Chang.* doi: 10.1038/NCLIMATE3223.
- 1248 Parmesan, C. & Yohe, G. 2003. A globally coherent fingerprint of climate change impacts across natural  
 1249 systems. *Nature* **421**: 37 - 42.
- 1250 Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol.*  
 1251 *Syst.* **37**: 637–669.
- 1252 Patthey, P., Wirthner, S., Signorell, N. & Arlettaz, R. 2008. Impact of outdoor winter sports on the abundance  
 1253 of a key indicator species of alpine ecosystems. *J. Appl. Ecol.* **45**: 174–1711.
- 1254 Patthey, P., N. Signorell, L. Rotelli & Arlettaz, R. 2012. Vegetation structural and compositional heterogeneity  
 1255 as a key feature in Alpine black grouse microhabitat selection: conservation management implications.  
 1256 *Eur. J. Wildl. Res.* **58**: 59-70.
- 1257 Pavel, V. 2004. The impact of grazing animals on nesting success of grassland passerines in farmland and  
 1258 natural habitats: a field experiment. *Folia Zool.* **53**: 171-178.
- 1259 Pearce-Higgins, J.W., Yalden, D.W. & Whittingham, M.J. 2005. Warmer springs advance the breeding  
 1260 phenology of Golden Plovers *Pluvialis apricaria* and their prey (*Tipulidae*). *Oecologia* **143**: 470–476.

- 1261 Pearce-Higgins, J.W., Dennis, P., Whittingham, M.J. & Yalden, D.W. 2010. Impacts of climate on prey  
1262 abundance account for fluctuations in a population of a northern wader at the southern edge of its range.  
1263 *Glob. Chang. Biol.* **16**: 12–23.
- 1264 Pepin, N., Bradley, R. S., Diaz, H.F., Baraer, M., Caceres, E.B., Forsythe, N., Fowler, H., Greenwood, G., Hashmi,  
1265 M.Z., Liu, X.D., Miller, J.R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I., Schöner, W., Severskiy, I.,  
1266 Shahgedanova, M., Wang, M.B., Williamson, S.N. & Yang, D.Q. 2015. Elevation-dependent warming in  
1267 mountain regions of the world. *Nat. Clim. Chang.* **5**: 424–430.
- 1268 Peringer, A., Siehoff, S., Chetelat, J., Spiegelberger, T., Buttler, A. & Gillet, F. 2013. Past and future landscape  
1269 dynamics in pasture-woodlands of the Swiss Jura Mountains under climate change. *Ecol. Soc.* **18**: 11.
- 1270 Pernollet, C.A., Korner-Nievergelt, F. & Jenni, L. 2015. Regional changes in the elevational distribution of the  
1271 Alpine Rock Ptarmigan *Lagopus muta helvetica* in Switzerland. *Ibis* **157**: 823–836.
- 1272 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team 2017. *nlme: Linear and Nonlinear Mixed Effects*  
1273 *Models*. R package version 3.1-131.
- 1274 Popy, S., Bordignon, L. & Prodon, R. 2010. A weak upward elevational shift in the distributions of breeding  
1275 birds in the Italian Alps. *J. Biogeogr.* **37**: 57–67.
- 1276 Potatov, R. 2004. Adaptation of birds to life in high mountains in Eurasia. *Acta Zool. Sinica* **50**: 970–977.
- 1277 Preston, F.W. 1962. The canonical distribution of commonness and rarity. *Parts I, II. Ecology* **43**: 185–215,  
1278 410–432.
- 1279 Quantum GIS Development Team. 2015. *Quantum GIS Geographic Information System*. Open source  
1280 geospatial foundation project.
- 1281 Ralston, J., DeLuca, W. V., Feldman, R. E. & King, D. I. 2017. Population trends influence species ability to track  
1282 climate change. *Glob. Change Biol.*, **23**: 1390–1399. doi:10.1111/gcb.13478
- 1283 Rauter, C.M., Reyer, H.U. & Bollmann, K. 2002. Selection through predation, snowfall and microclimate on  
1284 nest-site preferences in the Water Pipit *Anthus spinoletta*. *Ibis* **144**: 433–444.
- 1285 Reif, J. & Flousek, J. 2012. The role of species' ecological traits in climatically driven altitudinal range shifts of  
1286 central European birds. *Oikos* **121**: 1053–1060.
- 1287 Reif, J., Hořák, D., Krištín, A., Kopsová, L. & Devictor, V. 2015. Linking habitat specialization with species' traits  
1288 in European birds. *Oikos* **125**: 405–413.
- 1289 Rimmer, C.C., McFarland, K.P., Lambert, J.D. & Renfrew, R.B. 2004. *Evaluating the use of Vermont ski areas by*  
1290 *Bicknell's thrush: applications for Whiteface mountain*. Vermont Institute of natural Science, Woodstock,  
1291 VT, New York.
- 1292 Rixen, C. & Rolando, A. (eds). 2013. *The impacts of skiing and related winter recreational activities on*  
1293 *mountain environments*. Bentham doi: 10.2174/97816080548861130101.
- 1294 Rocchia, E. 2016. *Temporal variation of species distribution and species morphological traits along altitude in*  
1295 *the Alps* (Doctoral thesis). University of Milano-Bicocca, Italy.
- 1296 Rolando, A., Caprio, E., Rinaldi, E. & Ellena, I. 2007. The impact of high-altitude ski-runs on alpine grassland  
1297 bird communities. *J. Appl. Ecol.* **44**: 210–219.
- 1298 Root, T.L., Price, J.T., Hall, K.R., Schneider, S.H., Rosenzweig, C. & Pounds, J.A. 2003. Fingerprints of global  
1299 warming on wild animals and plants. *Nature* **421**: 57–60.
- 1300 Rosvold, J. 2016. Perennial ice and snow-covered land as important ecosystems for birds and mammals. *J.*  
1301 *Biogeogr.* **43**: 3–12.
- 1302 Roth, T., Plattner, M. & Amrhein, V. 2014. Plants, birds and butterflies: short-term responses of species  
1303 communities to climate warming vary by taxon and with altitude. *PLoS ONE* **9**: e82490.
- 1304 Rubolini, D., Møller, A.P., Rainio, K. & Lehikoinen, E. 2007. Intraspecific consistency and geographic variability  
1305 in temporal trends of spring migration phenology among European bird species. *Clim. Res.* **35**: 135–146.
- 1306 Sæther, B., Tufto, J., Engen, S. & Jerstad, K. 2000. Population dynamical consequences of climate change for a  
1307 small temperate songbird. *Science* **287**: 854–856.
- 1308 Sandercock, B.K., Martin, K. & Hannon, S.J. 2005. Life history strategies in extreme environments:  
1309 comparative demography of arctic and alpine ptarmigan. *Ecology* **86**: 2176–2186.



1310 Santangeli, A., Rajasärkkä, A. & Lehikoinen, A. 2016. Effects of high latitude protected areas on bird  
1311 communities under rapid climate change. *Glob. Chang. Biol.* doi:10.1111/gcb.13518.

1312 Santisteban, L., Benkman, C.W., Fetz, T. & Smith, J.W. 2012. Survival and population size of a resident bird  
1313 species are declining as temperature increases. *J. Anim. Ecol.* **81**: 352–363.

1314 Scridel, D. 2014. Ecology and conservation of birds in upland and alpine habitats: a report on the BOU's  
1315 Annual Conference held at the University of Leicester, 1–3 April. *Ibis* **156**: 896–900.

1316 Scridel, D., Bogliani, G., Pedrini, P., Iemma, A., von Hardenberg, A., Brambilla, M. 2017a. Thermal niche  
1317 predicts recent changes in range size for bird species. *Clim. Res.* **73**:207-216.

1318 Scridel, D., Groom J.D. & Douglas, D.J.T. 2017b. Native woodland creation is associated with increase in a  
1319 Black Grouse *Lyrurus tetrix* population. *Bird Study* **64**: 70-83.

1320 Sekercioglu, C.H., Schneider, S.H., Fay, J.P. & Loarie, S.R. 2008. Climate change, elevational range shifts, and  
1321 bird extinctions. *Conserv. Biol.* **22**: 140–150.

1322 Siegel, R. B., Pyle, P., Thorne, J. H., Holguin, A. J., Howell, C. A., Stock, S. & Tingley, M.W. 2014. Vulnerability of  
1323 birds to climate change in California's Sierra Nevada. *Avian Conserv. Ecol.* **9**: 7  
1324 <http://dx.doi.org/10.5751/ACE-00658-090107>

1325 Signorell, N., S. Wirthner, P. Patthey, R. Schranz, L. Rotelli & Arlettaz, R. 2010. Concealment from predators  
1326 drives foraging habitat selection in brood-rearing Alpine black grouse *Tetrao tetrix* hens: habitat  
1327 management implications. *Wildlife Biol.* **16**: 249-257.

1328 Sirami, C., Caplat, P., Popy, S., Clamens, A., Arlettaz, R., Jiguet, F., Brotons, L. & Martin, J.L. (2017), Impacts of  
1329 global change on species distributions: obstacles and solutions to integrate climate and land use. *Global*  
1330 *Ecol. Biogeogr.*, **26**: 385–394.

1331 Stephens, P.A., Mason, L.R., Green, R.E., Gregory, R.D., Sauer, J.R., Alison, J., Aunins, A., Brotons, L., Butchart,  
1332 S.H., Campedelli, T., Chodkiewicz, T., Chylarecki, P., Crowe, O., Elts, J., Escandell, V., Foppen, R.P., Heldbjerg,  
1333 H., Herrando, S., Husby, M., Jiguet, F., Lehikoinen, A., Lindström, Å., Noble, D.G., Paquet, J.Y., Reif, J., Sattler,  
1334 T., Szép, T., Teufelbauer, N., Trautmann, S., van, Strien, A.J., van Turnhout, C.A., Vorisek, P. & Willis, S.G.  
1335 2016. Consistent response of bird populations to climate change on two continents. *Science* **352**: 84-87.

1336 Strahler, A.N. 1946. Geomorphic terminology and classification of land masses. *J. Geol.* **54**: 32-42.

1337 Sturrock, R. N., Frankel, S. J., Brown, A. V., Hennon, P. E., Kliejunas, J. T., Lewis, K. J., Worrall, J. J. and Woods, A.  
1338 J. 2011. Climate change and forest diseases. *Plant Pathology*, **60**: 133–149.

1339 Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S., Helaouet, P., Johns, D.G., Jones,  
1340 I.D., Leech, D.I., Mackay, E.B., Massimino, D., Atkinson, S., Bacon, P.J., Brereton, T.M., Carvalho, L., Clutton-  
1341 Brock, T.H., Duck, C., Edwards, M., Elliott, J.M., Hall, S.J.G., Harrington, R., Pearce-Higgins, J.W., Høye, T.T.,  
1342 Kruuk, L.E.B., Pemberton, J.M., Sparks, T.H., Thompson, P.M., White, I., Winfield, I.J. & Wanless, S.  
1343 2016. Phenological sensitivity to climate across taxa and trophic levels. *Nature* **535**: 241–245.

1344 Thiel, D., Jenni-Eierman, S., Palme, R. & Jenni, L. 2011. Winter tourism increases stress hormone levels in the  
1345 Capercaillie *Tetrao urogallus*. *Ibis* **153**: 122–133.

1346 Tingley, M.W., Monahan, W.B., Beissinger, S.R. & Moritz, C. 2009. Birds track their Grinnellian niche through a  
1347 century of climate change. *Proc. Natl. Acad. Sci. U.S.A.* **106**: 19637–19643.


1348 Tingley, M.W., Koo, M.S., Moritz, C., Rush, A.C. & Beissinger, S.R. 2012. The push and pull of climate change  
1349 causes heterogeneous shifts in avian elevational ranges. *Glob. Change Biol.* **18**: 3279–3290.

1350 Tingley, M.W. & Beissinger, S.R. 2013. Cryptic loss of montane avian richness and high community turnover  
1351 over 100 years. *Ecology* **94**: 598–609.

1352 Titeux, N., Henle, K., Mihoub, J.-B., Regos, A., Geijzendorffer, I. R., Cramer, W., Verburg, P. H. & Brotons, L.  
1353 2016. Biodiversity scenarios neglect future land-use changes. *Glob. Chang. Biol.* **22**: 2505–2515.

1354 Tolvanen, A. & Kangas, K. 2016. Tourism, biodiversity and protected areas - Review from northern  
1355 Fennoscandia. *J. Environ. Manage.* **166**: 58-66.

- 1356 Townsend, A.K., Cooch, E.G., Sillett, T.S., Rodenhouse, N.L., Holmes, R.T. & Webster, M.S. 2016. The  
 1357 interacting effects of food, spring temperature, and global climate cycles on population dynamics of a  
 1358 migratory songbird. *Glob. Chang. Biol.* **22**: 544–555.
- 1359 Tryjanowski, P., Sparks, T.H. & Profus, P. 2005. Uphill shifts in the distribution of the white stork *Ciconia*  
 1360 *ciconia* in southern Poland: the importance of nest quality. *Divers. Distrib.* **11**: 219–223.
- 1361 Viterbi, R., Cerrato, C., Bassano, B., Bionda, R., von Hardenberg, A., Provenzale, A. & Bogliani, G. 2013. Patterns  
 1362 of biodiversity in the northwestern Italian Alps: a multi-taxa approach. *Community Ecol.* **14**: 18–30.
- 1363 Virkkala, R., Heikkinen, R.K., Fronzek, S. & Leikola, N. 2013. Climate change, northern birds of conservation  
 1364 concern and matching the hotspots of habitat suitability with the reserve network. *PLoS ONE* **8**: e63376.
- 1365 Virkkala, R., Pöyry, J., Heikkinen, R.K., Lehikoinen, A. & Valkama, J. 2014. Protected areas alleviate Climate  
 1366 Change effects on northern bird species of conservation concern. *Ecol. Evol.* **15**: 2991–3003.
- 1367 Virkkala, R. 2016. Long-term decline of southern boreal forest birds: consequence of habitat alteration  
 1368 or climate change? *Biodivers. Conserv.* **25**: 151–167.
- 1369 Wang, G., Hobbs, T.N., Galbraith, H. & Giesen, K.M. 2002. Signatures of large-scale and local climates on the  
 1370 demography of white-tailed ptarmigan in Rocky Mountain National Park, Colorado, USA. *Int. J.*  
 1371 *Biometeorol.* **46**: 197–201.
- 1372 Wann, G.T., Aldridge, C.L. & Braun, C.E. 2014. Estimates of annual survival, growth, and recruitment of a  
 1373 white-tailed ptarmigan population in Colorado over 43 years. *Popul. Ecol.* **56**: 555–567.
- 1374 Warren, J.M., Rotella, J. & Thompson, J.E. 2008. Contrasting Effects of Cattle Grazing Intensity on Upland-  
 1375 Nesting Duck Production at Nest and Field Scales in the Aspen Parkland, Canada. *Avian Conserv. Ecol* **3**: 6.
- 1376 Watson, A. & Moss, R. 2004. Impacts of ski-development on Ptarmigan (*Lagopus mutus*) at Cairn Gorm,  
 1377 Scotland. *Biol. Conserv.* **116**: 267–275.
- 1378 Weed, M. & Bull, C. 2004. *Sports tourism: participants, policy and providers*. Elsevier Butterworth-  
 1379 Heinemann, Oxford.
- 1380 Wilson, J.D., Anderson, R., Bailey, S., Chetcuti, J., Cowie, N.R., Hancock, M.H., Quine, C., Russell, N., Stephen, L.  
 1381 & Thompson, D.B.A. 2014. Modelling edge effects of mature forest plantations on peatland waders  
 1382 informs landscape-scale conservation. *J. Appl. Ecol.* **51**: 204–213.
- 1383 Wilson S. & K. Martin. 2010. Variable reproductive effort for two ptarmigan species in response to spring  
 1384 weather in a northern alpine ecosystem. *J. Avian. Biol.* **41**: 319–326.
- 1385 Wilson, S. & Martin, K. 2012. Influence of life history strategies on sensitivity, population growth and  
 1386 response to climate for sympatric alpine birds. *BMC Ecology* **12**:9.
- 1387 Wipf, S. & Rixen, C. 2010 A review of snow manipulation experiments in Arctic and alpine ecosystems. *Polar*  
 1388 *Research* **29**: 95–109.
- 1389 Wu, X.Y., Zhang, X.F., Dong, S.K., Cai, H., Zhao, T.R., Yang, W.J., Jiang, R., Shi, Y.D. & Shao, J.L. 2015. Local  
 1390 perceptions of rangeland degradation and climate change in the pastoral society of Qinghai-Tibetan  
 1391 Plateau. *Rangeland J.* **37**: 11–19.
- 1392 Zamora, R., & Barea-Azcón, M. 2015. Long-term changes in mountain passerine bird communities in the  
 1393 Sierra Nevada (southern Spain): a 30-year case study. *Ardeola* **62**: 3–18.
- 1394 Zuckerberg, B., Woods, A.M. & Porter, W.F. 2009. Poleward shifts in breeding bird distributions in New York  
 1395 State. *Global Change Biol.* **15**: 1866–1883.



1397

1398

1399 **Figure 1.** Mountains systems classified by Kapos *et al.* (2000) and adapted to the Holarctic region (above the Tropic of  
 1400 Cancer – grey line). The upper three classes ('CL ') are delimited purely by elevation ( $\geq 2500\text{m}$ ). Areas below  $2500\text{m}$   
 1401 were classified additionally in terms of slope, terrain roughness and local elevation range (LER).

1401


1402

1403


**Table 1.** Frequency of studies of Holarctic mountain birds and climate change resulting from the systematic literature search across various regions and countries of the world. Reviews/commentaries and meta-analyses ( $n = 25$ ) were excluded.

<b>Geographical region</b>	<b>Frequency</b>
Eastern European countries (Poland, Czech Rep., Russia)	6
Western European countries (France, Germany)	6
Spain	8
UK/Ireland uplands	24
Nordic countries (Denmark, Finland, Iceland, Norway, Sweden & Iceland)	27
Alps & Pyrenees (Switzerland, France, Italy, Spain, Germany, Austria, Slovenia)	44
Pan-European	12
<b>Total European studies</b>	<b>127</b>
Greenland	4
Pacific North West Coastal Mtns (Alaska, Yukon, British Columbia Coast Mtns, Hudson Bay Mnts, Cascades)	18
South West Coastal ranges (California, Sierra Nevada, New Mexico)	7
Continental Ranges (Rocky Mnts, Colorado, Arizona, Montana, Dakota, Wyoming)	23
Appalachian Mtns (incl. NY State)	11
N. America wide region (Canada, USA also in combination)	12
<b>Total N. America studies</b>	<b>75</b>
China	5
Asia	2
<b>Total Asian studies</b>	<b>7</b>
<b>Global or nearly global</b>	<b>26</b>
<b>All studies</b>	<b>209</b>






**Figure 2.** Frequency of relevant published papers and reviews over time resulting from the systematic literature search.



**Figure 3.** Frequency of climate-change related studies on Holarctic mountain birds (mutually inclusive) resulting from the systematic literature search, classified according to general subjects addressed.



**Figure 4.** Number of conservation (mutually inclusive) actions suggested across all papers classified as ‘conservation & policy’ ( $n = 26$ ) in the systematic literature search.

**Table 2.** A summary of papers considering shifts in the elevation of bird species distributions in mountains over time. Papers were included if they were based on data carried out at relatively small scales (maximum 1-km<sup>2</sup>) and which were wholly or partly in mountainous regions (as per Kapos *et al.* 2000). MA indicates whether a given study was included in the meta-analysis (Y) or not (N).

Author	Location	Species	Sampling unit	Period	Temperature change (°C/year)	Elevation range (m)	MA	Key findings
Archaux 2004	French Alps	All	Point count	1973-2002	0.05	350-3099	Y	41 site/species comparisons: 6 showed significant downwards shifts, 4 significant upwards shifts
DeLuca & King 2017	Appalachian Mountains, USA	All	Point count	1993-2009	0.01 <sup>2</sup>	740-1470	Y	9 of 16 low-elevation species shifted upwards; 9 of 11 high-elevation species shifted downwards
Maggini <i>et al.</i> 2011	Switzerland <sup>1</sup>	All	1-km <sup>2</sup>	1999/2002 - 2004/07	0.09 <sup>2</sup>	210-2710	N	95 species: 33 species shifted upwards, 28 shifted downwards
Mizel <i>et al.</i> 2016	Denali National Park, Alaska	Passerines	Point count	1995-2013	0.04	500-1200	Y	Upwards shifts associated with shrub/tundra-nesting species; weaker evidence of upward shifts in forest species
Pernollet <i>et al.</i> 2015	Swiss Alps	Ptarmigan	1-km <sup>2</sup>	1984 - 2012	0.1	1700-3100	Y	Mean elevation of Ptarmigan presence shifted upwards in 3 of 4 regions
Popy <i>et al.</i> 2010	Italian Alps	All	1-km <sup>2</sup>	1992/94 - 2003/05	0.08	550-2556	Y	Weak overall upwards community shift; wide variation in the response of individual species
Reif & Flousek 2012	Giant Mountains, Czech Republic	All	Point count	1986/88 - 1996/98	0.12	400-1602	Y	Significant overall mean shifts to higher elevations; open-habitat shifted more than forest species
Rocchia 2016	Italian Alps	All	Point count	1982 - 2012	NA	600-4000	N	Woodland species tended to show range expansion, higher-elevation grassland species range retraction; regional variation
Tingley <i>et al.</i> 2012	Sierra Nevada, USA	All <sup>3</sup>	Point count	1911/28 - 2003/09	NA	61-3356	N	Shifts were heterogeneous within species and among regions; both temperature and precipitation likely to be important drivers
Tryjanowski <i>et al.</i> 2005	Tatra Mountains, Poland	White Stork	Nest location	1974 - 2003	0.08	400-900	Y	White Storks nested at progressively higher elevations.

<sup>1</sup> Included as the majority of the area of Switzerland is classed as mountainous by Kapos *et al.* (2000)

<sup>2</sup> Annual temperature – others are spring temperatures

<sup>3</sup> Passerines and five other families (Odontophoridae, Phasianidae, Columbidae, Trochilidae and Picidae)