

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Philaenus spumarius: when an old acquaintance becomes a new threat to European agriculture

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1667526> since 2018-05-07T13:04:57Z

Published version:

DOI:10.1007/s10340-018-0966-0

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

[Click here to view linked References](#)

1 ***Philaenus spumarius*: when an old acquaintance become a new threat to**
2 **European agriculture**

3 **¹Cornara D., ²Bosco D., ¹Fereres A.**

4 ¹ Instituto de Ciencias Agrarias. Consejo Superior de Investigaciones Cientificas. ICA-CSIC. Calle
5 Serrano 115 dpdo. 28006 Madrid

6 ² DISAFA – Entomologia. Largo Paolo Braccini, 2. 10095 Grugliasco (TO) – Italy.

7 **Abstract**

8 The unique color pattern polymorphism and the foamy nymphal case of the
9 meadow spittlebug *Philaenus spumarius*, have attracted the attention of
10 scientists for centuries. Nevertheless, since this species has never been
11 considered a major threat to agriculture, biological, ecological and ethological
12 data are missing and rather scattered. To date this knowledge has become of
13 paramount importance, in view of the discovery of *P. spumarius* main role in
14 the transmission of the bacterium *Xylella fastidiosa* in Italy, and possibly in
15 other European countries. The aim of this review is to provide a state of the
16 art about this species, with particular focus on those elements that could help
17 developing environmental-friendly and sustainable control programs to
18 prevent transmission of *X. fastidiosa*. Moreover, recent findings on the role
19 of the meadow spittlebug as vector of the fastidious bacterium within the first
20 reported European bacterium outbreak in Apulia (South Italy) will be
21 discussed.

22

23 **Key Message**

- 24
- 25 • The meadow spittlebug *Philaenus spumarius* plays a major role in the
26 spread of *Xylella fastidiosa* in the first European outbreak of the
27 bacterium in the Apulia region (Southern Italy).
 - 28 • Biological, ecological and ethological data about *P. spumarius* are rather
29 scattered and needs further investigations.
 - 30 • Here, we comprehensively collected scattering data and unpublished
31 information about the meadow spittlebug and its relationship with the
32 fastidious bacterium. Furthermore, we reviewed the known control
33 tactics and proposed new management strategies against this pest.
- 34
- 35

36 Introduction

37 Spittlebugs and their nymphal case have received attention from naturalists
38 for centuries. Starting from Saint Isidorous from Seville in the sixth century,
39 and later with Moffet and Linnaeus, many scientists devoted their attention
40 to these unique creatures coming from a “frothy sticky whitish dew” (Moffet
41 1685, cited in Weaver and King 1954). In the literature, spittle masses have
42 been called in many ways: Gowk’s spittle, frog spit, snake spit, witch’s spit
43 and wood sear, beside cuckoo spittle, since the Cuckoo bird migrate in
44 Europe at the same time the first masses appear (Svanberg 2016). It has also
45 been suggested that these masses generate small locust (Yurtsever 2000).
46 The meadow spittlebug *Philaenus spumarius* L. (1758) belongs to the order
47 Hemiptera, superfamily Cercopoidea, family Aphrophoridae. The name
48 spittlebug came from the shell built up by the nymphs mixing fluid voided
49 from the anus and a secretion produced by glands located between the 7th
50 and the 8th abdominal sternites. Air bubbles are introduced within the
51 spittle by mean of caudal appendages and a ventral tube formed by
52 abdominal tergites (4th to 9th) bent downward (Yurtsever 2000). Due to its
53 polymorphism, more than 50 synonyms had been given to *P. spumarius*, as
54 reported by Nast (1972). The meadow spittlebug was commonly called
55 *Philaenus leucophthalmus* in the early literature, as for example in Severin
56 (1950) and Weaver and King (1954). The taxonomical confusion was solved
57 when, in 1961, the International Commission of Zoological Nomenclature
58 decided for the only valid specific name of *P. spumarius* (Yurtsever 2000).
59 The large body of literature on *P. spumarius* deals meanly with the genetic
60 basis of adult color polymorphism, and the damage caused by nymphs to
61 strawberry and alfalfa, when the insect was introduced in USA (Weaver and
62 King, 1954). Now we know that this ubiquitous, common and locally very
63 abundant insect is the main vector of the bacterium *Xylella fastidiosa* in the
64 Apulia Region of Italy, and has the potential to spread it in all the other
65 European regions where the pathogen is present. Nevertheless, since the
66 meadow spittlebug has never been considered an agricultural pest in

67 Europe before the introduction of *X. fastidiosa*, its biology, ecology and
68 ethology have never been investigated continuously and in a comprehensive
69 way. Therefore, the main aim of this manuscript is to provide an updated
70 and critical state of the art about *P. spumarius*, mainly focusing on those
71 elements that could help developing an environmental friendly and
72 sustainable control strategy to prevent *X. fastidiosa* spread.

73

74 **Taxonomy and description**

75 Until 1980's, only three species belonging to the genus *Philaenus* were
76 known: the Holarctic *P. spumarius*; the Mediterranean species *P. signatus*
77 (inhabiting the Balkans and Middle East); and *P. tesselatus* (Southern Iberia
78 and Maghreb), this latter often considered a subspecies or a synonym of *P.*
79 *spumarius* (Nast 1972). Starting from the 1990's, thanks to in-depth studies
80 carried out across the Mediterranean, five further species of the genus have
81 been described: *P. loukasi* (southern Balkans), *P. arslani* (Middle East), *P.*
82 *maghresignus* (Maghreb and southern Spain), *P. italosignus* (southern Italy
83 and Sicily), and *P. tarifa* (southern Iberia). The eight species are sympatric
84 with *P. spumarius*, and partially allopatric with each other (Maryanska-
85 Nadachowska et al. 2012). The proteobacterium *Wolbachia* could have
86 played a role in the speciation of *P. spumarius*, since it is almost exclusively
87 present in Northeastern mitochondrial clade (Lis et al. 2015). Currently, the
88 species can be distinguished according to anal tube and male genitalia
89 morphology in two groups: the "spumarius" group (*P. spumarius*, *P.*
90 *tesselatus*, *P. loukasi* and *P. arslani*), and the "signatus" group (*P.*
91 *maghresignus*, *P. italosignus*, *P. signatus*, *P. tarifa*) (Drosopoulos and
92 Remane 2000). Another classification takes into account nymphal food
93 plants, and allows a differentiation in three main groups: *P. signatus*, *P.*
94 *italosignus*, *P. maghresignus* and *P. tarifa*, whose nymphs elect the lily
95 *Asphodelus aestivalis* L. (1753) as their main host plant; *P. loukasi* and *P.*
96 *arslani*, whose nymphs develop on xerophilic plants; and *P. spumarius* and

97 *P. tessellatus*, that thrive on monocotyledonous and dicotyledonous plants,
98 although the former is likely to prefer dicots (Drosopoulos 2003). According
99 to Maryanska-Nadachowska et al. (2012), the genus *Philaenus* is
100 monophyletic, this claim being supported by morphological, ecological and
101 chromosomal data. *P. spumarius* is extremely varying in color, going from
102 unicolorous yellowish white to unicolorous black, with several intermediate
103 morphs. Most of these were originally described as species. Furthermore,
104 recently two new species belonging to the genus *Philaenus*, namely *P.*
105 *elbusiarnus* and *P. iranicus*, have been described in Iran (Tishechkin 2013). A
106 detailed morphological and phylogenetic description of the species is out of
107 the purpose of this review; for papers regarding these issues, please refer to
108 Delong and Severin (1950), Ossiannilsson (1981), Berry and Willmer (1986),
109 Stewart and Lees (1996), Quartau and Borges (1997), Drosopoulos (2003),
110 Maryańska-Nadachowska et al. (2012), Rodrigues et al. (2014), and further
111 references.

112

113 **Geographical range**

114 *P. spumarius* is widely distributed, covering most of the Palearctic regions,
115 and extending to Nearctic, as well as most of the temperate regions of earth
116 and oceanic islands (Stewart and Lees 1996; Drosopoulos and Asche 1991;
117 Drosopoulos and Remane 2000). Its distribution ranges from north Lapland
118 to the Mediterranean in Europe, including Turkey. It has been reported for
119 North Africa, several parts of the former Soviet Union, Afghanistan, Japan,
120 USA, Canada, Azores, Hawaii, New Zealand (Yurtsever 2000). The meadow
121 spittlebug was probably introduced in new continents, as North America, as
122 overwintering eggs in straw stubble (Whittaker 1973). Its distribution in
123 Europe and world-wide has been summarized by EFSA (2015). In Greece,
124 Drosopoulos and Asche (1991) reported *P. spumarius* at an altitude ranging
125 from the sea level to more than 2000 m. Climate change may significantly
126 have affected the distribution of *P. spumarius*: Karban and Strauss (2004)

127 suggested that the species Northward shift in California since 1988 is related
128 to variations in humidity and temperature.

129

130 **Host plants and feeding behavior**

131 *P. spumarius* is highly polyphagous, and occurs in most of the terrestrial
132 habitats (Stewart and Lees 1996). According to Maryanska-Nadachowska et
133 al. (2012), the common ancestor of the species belonging to the genus
134 *Philaenus* may have used lily as its main host plant, a character that still
135 remains in *P. maghresignus*, *P. italosignus*, *P. tarifa* and *P. signatus*. On the
136 contrary, the exploitation of a wide range of hosts belonging to
137 monocotyledonous and dicotyledonous may have been the leading factor
138 promoting the geographical expansion of the species. *P. spumarius* is a
139 xylem feeder, either as nymph or adult: the spittlebug ingests considerable
140 amount of sap from the main transpiration stem without causing vessels
141 cavitation, overcoming dramatically high tension reaching -10 bars, and
142 showing a mean excretion rate of 280 times its body weight in 24 hours
143 (Wiegert 1964; Horsfield 1978; Crews et al. 1998; Malone et al. 1999;
144 Watson et al. 2001; Ponder et al. 2002). The association with symbionts
145 potentially relaxes the severe energy limitations related to xylem sap
146 feeding, being the xylem sap nutritionally poor and energetically costly to
147 extract (Thompson 2004; Koga et al. 2013). Nymphs and adults feed
148 preferentially on actively growing parts (Mundinger 1946; Wiegert 1964).
149 Nitrogen fixing legumes and other plants with high aminoacids
150 concentration in the xylem sap (*Medicago sativa* L. (1753), *Trifolium* sp. L.,
151 *Vicia* spp. L., and *Xanthium strumarium* L. (1753)) are the preferred hosts
152 (Horsfield 1977; Thompson 1994). Overall, *P. spumarius* seems to prefer
153 plants that transport fixed nitrogen as aminoacids and amides than those
154 that transport fixed nitrogen as ureides (Thompson 1994). Nymphal
155 excretion rate has been proven to be positively correlated with aminoacids
156 concentration in the xylem-sap (Horsfield 1977). Nymphs and adults thrive

157 on various plants in habitats moist enough to provide them with sufficient
158 humidity to keep them alive, such as meadows, abandoned fields, waste
159 grounds, roadsides, streamsides, hayfields, marshlands, parks, gardens, and
160 cultivated fields (Yurtsever 2000). Gulijeva (1961) reported cereals,
161 Asteraceae, legumes and Lamiaceae as the most favorable hosts.
162 Ossiannilsson (1981) states that *P.spumarius* is the most polyphagous insect
163 currently known, with a host lists that exceed 1000 plants. Dicotyledonous
164 plants tend to be used more often than monocotyledonous (Wiegert 1964;
165 Halkka et al. 1967; Halkka et al. 1977). Pasture mowing or a general
166 decrease of succulence of herbaceous hosts, cause a dispersal of the adults
167 that may settle in high numbers plants such as grapevine, olive, peach,
168 almond, besides several trees and shrubs as holm oak, myrtle, and lentisk
169 (Goidanich 1954; Pavan 2006; Cornara et al. 2016b). For a *P. spumarius*
170 complete host list, refer to Delong and Severin (1950) and Weaver and King
171 (1954).

172

173 **Biology and ecology**

174 **Life history and behavior**

175 *P. spumarius* is a univoltine species, overwintering as egg. First mature eggs
176 are found in the ovaries starting from the end of August, and then increase
177 until November (Weaver 1951). Females are polyandrous; the multiple
178 mating does not influence the number of progeny, but provide great genetic
179 and evolutionary benefits to the meadow spittlebug, as shown in many
180 polyandrous species (Smith 1984). Yurtsever (2000) hypothesized that *P.*
181 *spumarius* very diverse habitats is a consequence of the advantages derived
182 from multiple mating. Mating occurs readily after adult appearance, and
183 continues throughout the seasons; the spermatogenesis and release of
184 sperma in the spermatheca is designed so that delayed fertilization could
185 take place (Robertson and Gibbs 1937). Weaver and King (1954) observed a
186 peak of development for eggs not occurring until 2nd week of September,

187 with no significant difference due to geographical location. The failure in
188 spittlebug control with treatments in the first week of September, is a
189 further evidence that oviposition takes place after this period (King 1952). In
190 Apulia, oviposition was achieved in semi-artificial conditions in October on
191 *Sorghum halepense* L., concomitantly with a decrease of average daily
192 temperature below ca. 15°C; furthermore, the only eggs observed in the
193 field were laid on the same plant along the orchard edges (Cornara and
194 Porcelli 2014, FIGURE 1). Eggs are oviposited in stubble, herbs, dead parts of
195 plants, plant residue, cracks and tree trunk barks, or in the litter; the
196 majority of eggs are laid close to the ground between two apposed surfaces
197 (Barber and Ellis 1922; Weaver and King 1954; Yurtsever 2000).
198 Furthermore, Weaver and King (1954) reported that the presence of straw
199 within experimental cages caused an increase of 65% in egg deposition. Oat,
200 Johnson grass, dwarf broad bean, alfalfa, red clover, and timothy, were
201 reported as experimental hosts for oviposition (Weaver and King 1954;
202 Halkka et al. 1966; Stewart and Lees 1988; Cornara and Porcelli 2014). Eggs
203 are elongated, ovoid and tapering in shape, yellowish-white with a dark
204 pigmented orange spot at one end. If the egg is fertilized, the orange spot
205 gets bigger and a black lid-like formation develops on it (Yurtsever 2000).
206 Eggs are laid in masses of one to 30 elements, with an average value of
207 seven, held together by a hardened frothy cement (Weaver and King 1954;
208 Ossianilsson 1981). Munding (1946) and Weaver and King (1954) agreed
209 upon the number of eggs oviposited being around 18 to 51 per female,
210 although a lower estimate, about 10 to 20 per female, was reported by
211 Wiegert (1964). On the contrary, Yurtsever (2000) claims that an individual
212 female may produce up to 350-400 eggs. These conflicting data suggest that
213 experiments under controlled conditions aimed at estimating prolificacy are
214 needed to estimate this important biological parameter. The oviposition
215 continues until the female dies naturally or is killed by severe frost (Weaver
216 and King 1954). The *pre-imago* pass through five instars. Pre-imaginal
217 development takes 5-6 weeks, although cold weather considerably reduces

218 the speed of the cycle; consequently, nymphal period may take from 35 to
219 100 days approximately (Weaver and King 1954; Yurtsever 2000; Halkka et
220 al. 2006). The first instar nymph is approximately 1.35 mm long, orange, and
221 produces a tenuous spittle. During the development the color became
222 gradually green-yellow; the last two instars produce a great amount of
223 spittle (Yurtsever 2000). Once hatched, nymphs crawl to the closest green
224 succulent plant and began forming a spittle (Weaver and King 1954).
225 Selection of a feeding site on a plant may occur after the insect has ingested
226 and sampled xylem sap (Horsfield 1977). The first nymphs can be found on
227 low rosettening plants or in plants that offer closely apposed leaf and stem
228 surfaces; these hosts indeed provide a shelter from direct sun and drying
229 winds (Weaver and King 1954; Grant et al. 1998). Apparently, the nymph is
230 able to survive on almost any plant that provides them with sufficient
231 moisture to maintain their feeding habits (Weaver and King 1954). Plant
232 mechanical defences, as trichomes present on the stem of plants as
233 *Anaphalis margaritacea* Benth & Hook (1873), or tissue hardness, may
234 inhibit young nymphs from feeding, mechanically impeding stylet
235 penetration (Hoffman and McEvoy 1985a; 1985b). The range of feeding sites
236 exploited increases with nymphal development (Hoffman and McEvoy
237 1985a). Wiegert (1964) observed peak densities of 1280 nymphs/m² and
238 466 adults/m² in an alfalfa field. In Europe, nymphs density has been
239 reported not exceeding 1000 nymphs/m² (Zajac et al. 1984). Nymphs tend
240 to aggregate on the host plants, sharing the same spittle. The aggregation,
241 maintained within certain levels in order to avoid competition, ensures a
242 bottom-up effect, for example overcoming physical barriers to feeding on
243 xylem (Wise et al. 2006). Individuals of different Cercopid species may be
244 found embedded in the same spittle mass (Halkka et al. 1977). As reported
245 by Whittaker (1973), whereas nymphs mortality is inversely density
246 dependent, in adults a slight although significant density dependent
247 regulation exists. The same author found that, when *P. spumarius* is present

248 in the field concomitantly with other spittlebug species as *Neophilaenus*
249 *lineatus* L. (1758), both of the populations tend to be more stable.

250 After the spittle is formed, the nymph is able to maintain its own micro-
251 climate; evaporation rate gradients ensure that spittles are largest close to
252 the ground, where they are most available to non-flying predators, and
253 greater insulation from high temperature is required (Whittaker 1970). The
254 same author also inferred that the spittle is a form of protection from
255 predators. At the time of the last molt, the nymph ceases to form the spittle,
256 which progressively dries up, forming a chamber where the adult stage will
257 appear (Weaver and King 1954). Adults appear in April and live until fall
258 (Weaver and King 1954), although they may survive throughout the
259 successive spring in case of mild winters (Saponari et al. 2014). The callow
260 adult is nearly white with a slight greenish cast; it takes some minutes to
261 acquire its characteristic colored pattern (Weaver and King 1954). Industrial
262 melanism for *P. spumarius* has been suggested (Lees and Dent 1983).
263 Thompson (1973) claimed that *P. spumarius* color pattern warns the
264 predator about the insect's exceptional escape ability through leaping.
265 Therefore, a learned predator tends to avoid the meadow spittlebug
266 because it associates the color pattern to a wasted effort in preying, due to
267 the strong and rapid leaping of the prey (Gibson 1974). Males appear earlier
268 than females and, over the year, the number of males declines in
269 comparison to females (Edwards 1935; Halkka 1964; Drosopoulos and Asche
270 1991).

271

272 **Phenology, developmental thresholds and temperature-dependent** 273 **development**

274 Difficulties faced by researchers for decades in rearing *P. spumarius*
275 continuously in the lab, strongly suggests that the entire life cycle relies on a
276 specific combination of environmental variables still not fully understood. A
277 deep knowledge about phenology and developmental threshold is

278 mandatory in order to set up an effective forecasting model for *P. spumarius*
279 control. Two are the key factors regulating *P. spumarius* development:
280 humidity and temperature.

281 According to Weaver and King (1954), several evidences such as the
282 behavior of nymphs seeking sheltered places, the production of foam and
283 the necessary structure to produce it, the adult migration during the
284 summer period, the delay until cool weather for the deposition of the eggs,
285 the manner of placing and cementing the eggs between two apposed
286 surfaces so that water losses are minimized, suggest that the entire life cycle
287 depends on humidity and water availability. Even a cornea thicker in the
288 adults compared to the nymph, that reduces water losses, might be
289 considered a further evidence of the fact that water represents the key
290 element around which the meadow spittlebug biology spins (Keskinen and
291 Meyer-Rochow 2004). Weaver and King (1954) stated that the highest
292 concentration of spittlebugs are contained within the regions of highest
293 humidity. Humidity likely elicits hatching. Indeed, if eggs hatch in a high
294 humidity environment, first instar nymph would survive to dehydration, and
295 could find a suitable tissue to settle on. The first plants on which nymphs are
296 observed are those exhibiting dense lateral growth, thus limiting air
297 movements and having a higher RH (relative humidity). Furthermore,
298 nymphs tend to congregate on closely apposed surfaces where the humidity
299 can be maintained at high levels, as noticed both in field and lab conditions
300 using *Sonchus* sp. L. as a rearing plant (Morente et al. unpublished). As
301 reported by Weaver and King (1954), early in the morning nymphs can be
302 found at the tip of the plant, but as the temperature raises, the masses dry
303 and they leave them to move down on the plant. The foam secreted by
304 nymphs creates an excellent protection against dehydration and UV
305 radiation. Indeed *P. spumarius* foam case can block as much as 88% of the
306 UV incident radiation (in the 250 to 400 nm range) (Chen et al., 2017). In
307 spite of the indications on *P. spumarius* preference for moist environments,
308 other reports point out that the meadow spittlebug colonizes nearly all

309 habitats including wet or dry meadows and dry mediterranean forests
310 (Guglielmino et al. 2005). Consistently, *P. spumarius* can be very abundant
311 on herbaceous vegetation within and surrounding olive groves in the Apulia
312 Region of Italy, as well as in vineyards (Nicoli Aldini et al. 1998; Braccini and
313 Pavan 2000; Pavan 2006). Olive and grapevine are rain fed Mediterranean
314 crops that grow in dry environments. Thus, it can be concluded that *P.*
315 *spumarius* has the potential to live under different environmental
316 conditions, from moist to relatively dry, as long as the host plants are
317 actively growing and not subjected to severe water stress. Due to the
318 exceptionally wide area of distribution of this species, it cannot be excluded
319 that the spittlebug requirement for humidity depends upon the
320 geographical area the population lives in, or that different populations
321 within the species have different humidity requirements.

322 Along the years, several authors have tried to establish correlations
323 between the meadow spittlebug development and temperature. According
324 to Medler (1955), eggs hatch after an accumulation of 150 degree days (DD),
325 with a maximum daily accumulation of 10 degrees over ca. 4.4 °C. King
326 (1952) failed to speed up egg development by decreasing temperature to
327 10°C and diminishing day length to 13 hours/day. Stewart and Lees (1988)
328 succeeded in achieving oviposition in lab conditions, exposing eggs to 10°C
329 for 75 to 100 days, photoperiod 12/12 light/dark, 100% HR and then
330 increasing the temperature up to 15°C until hatching occurred. Chmiel and
331 Wilson (1979) stated that the 1st hatch can be predicted using an
332 accumulation of 120 HU (heating units calculated based on a threshold
333 temperature of 6.5°C from the 1st of January). Weaver and King (1954)
334 hypothesized that hatching occurs at temperatures of ca. 10 to 21°C, and
335 that cold temperatures may have a conditioning effect, that speeds up eggs
336 development. Nevertheless, the same authors reported that eggs never
337 exposed to less than ca. 15°C were able to hatch early in February. Masters
338 et al. (1998), reported that milder winters resulted in an early hatching, with
339 no significant effect on nymphal development. Weaver and King (1954)

340 stated that in areas where the spring weather is variable and short cold
341 periods are interspersed with warm periods, the hatching may be prolonged
342 over a long period. According to Zajac et al. (1989), upper and lower
343 threshold for nymphal development are 2.8°C and 26.7°C, respectively. The
344 first through the fifth instar nymphs and adults began appearing in the field
345 at 2, 154, 262, 364, 472, and 660 HU respectively, as calculated from the
346 first eggs hatching. The mean residence time of the five instars had been
347 calculated in 154, 103, 101, 113, 181 HUs, respectively (Zajac et al. 1989). All
348 this information, often based on substantially different estimations of the
349 lower temperature thresholds, reveals that no clear and consistent data on
350 the influence of temperature on spittlebug development are available and
351 new studies are needed to fill the gap.

352 Manipulation of the life cycle under controlled conditions in order to obtain
353 more than one generation per year, thus extending the period for biological
354 investigations of this species, does not seem an easy task, especially if we
355 consider that termination of egg diapause requires a prolonged period of
356 low temperatures, from 83 to 100 days (West and Lees, 1988; Yurtsever,
357 2000). Also, experimental data on the viability of eggs stored at low
358 temperature for several months in order to obtain nymphs later in the
359 season are lacking. This represents a constraint in the studies of biology and
360 behavior of *P. spumarius* under controlled conditions because the
361 experiments need to be carried out in a limited period of the year when
362 nymphs or adults are available.

363

364 **Movement**

365 Although the nymphs live inside a spittle, they can actively crawl on short
366 distances, thus moving from one herbaceous plant to another, as observed
367 by Bodino et al. (2017)., Adults are much more mobile, both actively and
368 passively. They can fly but, more often, they crawl or leap (Ossianilsson
369 1981). Hind legs, usually dragged while walking, are the structures

370 underlying the *P. spumarius* amazing jumping ability. Power muscles
371 contracting slowly and storing energy, plus a peculiar joint interlocking
372 mechanism, allow the insect to generate a force of 414 times its body
373 weight, with a jump acceleration of 2800-4000 m/s² (Burrows 2003). A
374 migratory behavior has been observed by several authors, with females
375 migrating further and more readily than males (Weaver 1951; Weaver and
376 King 1954; Lavigne 1959; Halkka 1962; Wiegert 1964; Halkka et al. 1967;
377 Drosopulos and Asche 1991; Grant et al. 1998). *P. spumarius* active dispersal
378 is probably made possible by its high polyphagy (Halkka et al. 1967). The
379 meadow spittlebug distribution largely depend on the distribution of
380 suitable host plants, which often occur aggregated, and frequently form a
381 discrete pattern (Biederman 2002). First dispersal is likely to happen when
382 the adults are still tender and immature, and can be related to harvest of
383 the host crop or to a general decline in succulence of the host plant (Weaver
384 and King 1954; Waloff 1973). Luxuriant foliage of newly seeded plants
385 gradually but constantly attracts the spittlebug from the surroundings
386 (Weaver 1951). The dissemination of the adults from the meadow is
387 concomitant with an increase in population in other crops (Putman 1953).
388 Migration continues until September, when adults gradually lessen their
389 migratory activity. The diminishing of this tendency could be associated
390 both with cooler temperatures, and with the fact that females devote their
391 energy to oviposition (Weaver and King 1954). An indirect evidence of the
392 migration behavior is also provided by Drosopoulos and Asche (1991), that
393 suggest the presence of a bivoltine *P. spumarius* population in Greece, with
394 two peaks in adults collection, in May and October. Since the author did not
395 find nymphs during summer, it can be speculated that the two peaks
396 coincide with the migrations driven by loss of succulence in host plants and
397 oviposition. The same author also observed a drastic reduction in population
398 densities likely caused by spring and summer drought, with *P. spumarius*
399 becoming a rare species in some years. Weaver and King (1954) observed
400 marked *P. spumarius* travelling more than 30 meters with a single flight, and

401 moving as much as 100 meters within 24 hours from the release point. The
402 same authors stated that the spittlebug may hop for several feet but are
403 poorly balanced, so that they land on their back. Adults mainly move at a
404 height of 15-70 cm from the ground; higher movements seems unlikely,
405 although observations of adults flying up to 6 meters high are reported
406 (Weaver and King 1954; Wilson and Shade 1967; Halkka et al. 1971).
407 However, according to Freeman (1945), adults of *P. spumarius* can actually
408 fly much higher than data reported by other authors. Indeed, Freeman
409 collected one individual of *P. spumarius* and eight individuals of
410 *Neophilaenus lineatus* with nets located 84 m above ground in the area of
411 Lincolnshire (UK). Although in the mentioned paper the author reported
412 these specimens generically as Cercopidae, actually it refers to *N. lineatus*
413 and *P. spumarius* (Don Reynolds, personal communication). Such
414 information suggests that *P. spumarius* can be transported by wind currents
415 and is potentially capable of long distance migration. Passive dispersal over
416 great distances is mediated by wind and human activities (Weaver and King
417 1954). Dispersal power is sufficient to colonize all the micro-habitat within
418 an island and to reach nearby islands (Halkka et al. 1971; Schultz and Meijer
419 1978). Passive dispersal due to transportation by cars has been observed
420 (Bosco, personal observation). A seasonal movement of adults from the
421 herbaceous vegetation of olive groves to the olive canopy and other
422 evergreen and deciduous trees/shrubs on late spring-early summer has
423 been observed in Northern and Southern Italy (Cornara et al. 2016b; Bodino
424 et al. 2017). This movement is likely not only due to drying of the
425 herbaceous hosts, as it can be observed also where the grass cover persists
426 over the summer. An opposite movement occurs at the end of summer-
427 beginning of autumn when adults, mostly females, re-colonize herbaceous
428 vegetation looking for suitable sites of oviposition.

429

430 **Direct damage**

431 The meadow spittlebug began to receive attention during 1940's, as a
432 consequence of the built up of large population and large infestations in
433 meadow crop in the USA. As reported by Weaver and King (1954), during late
434 40's in Ohio approximately every legume hay field was heavily infested, with
435 the complete loss of the first hay cutting. Moreover, the species has been
436 regarded as a pest of strawberry (Mundiger 1946; Zajac and Wilson 1984).
437 Outside the USA, *P. spumarius* had never been considered a pest). Direct
438 damages by adult meadow spittlebugs seem unlikely, especially in view of the
439 large number of adults congregating on a crop (Weaver and King 1954). On
440 the contrary, losses associated with large infestation by nymphs on alfalfa,
441 red clover, carrot, peas and strawberries in areas where *P. spumarius* was an
442 alien pest have been reported in the USA, with nymphal feeding causing
443 mainly dwarfing (Fisher and Allen 1946; Scholl and Medler 1947; Poos 1953;
444 Weaver and King 1954). No effects of the spittlebug feeding on white clover
445 (*Trifolium repens*) seed production was observed (Pearson 1991).

446

447 ***Philaenus spumarius* as a vector of plant pathogens**

448 While direct damages seem unlikely, transmission of plant pathogens
449 represents the most serious threat posed by the meadow spittlebug to
450 agriculture and landscape. *P. spumarius* has been erroneously reported as
451 vector of the peach yellow virus, while further tests disproved its involvement
452 in pathogen transmission (Severin 1950). Phytoplasmas have been detected
453 in *P. spumarius* by several authors (Pavan 2000; Landi et al. 2007; Ivanauskas
454 et al. 2014), and in one case the species was claimed to be a vector of ash
455 yellows phytoplasmas (Matteoni and Sinclair 1988). However, this latter
456 finding was not confirmed by further works (Sinclair and Griffith 1994; Hill and
457 Sinclair 2000) so that *P. spumarius* cannot be considered a vector of
458 phytoplasmas until new convincing evidences are provided. Moreover, the
459 spittlebug has been reported as a passive carrier of the plum mite (Mundinger
460 1946). *P. spumarius* was first reported as a vector of the bacterium *Xylella*

461 *fastidiosa* Wells (1987) by Severin (1950). The ability of the meadow
462 spittlebug in transmitting the bacterium was confirmed by further research,
463 although it was suggested that this insect might play only a marginal role in
464 *X. fastidiosa* epidemiology in the American outbreaks (Purcell 1980; Almeida
465 et al. 2005; Sanderlin and Melanson 2010). It was not until 2014 that *P.*
466 *spumarius* became a serious threat to European agriculture, when it was
467 reported as the major vector of *X. fastidiosa* in the Apulia region, Southern
468 Italy (Saponari et al. 2014; Cornara et al. 2016b)

469

470 **Role of *P. spumarius* in the first outbreak of *X. fastidiosa* in Europe, and** 471 **remarks on other potential vectors**

472 *X. fastidiosa* establishment in Europe is a clear example of the consequences
473 related to pathogen introduction and emergence in a new environment,
474 where the pathogen itself finds a suitable vector able to drive disease
475 epidemics (Almeida and Nunney 2015; Fereres 2015; Martelli et al. 2016). *X.*
476 *fastidiosa* is a gram-negative xylem limited gamma-proteobacterium, order
477 Xanthomonadales, family Xanthomonadaceae, present throughout America.
478 It causes diseases in many crops of economic importance such as grapevine,
479 citrus, almond, and others (Purcell 1997). According to EFSA (2015), its host
480 list embraces 309 plant species belonging to 63 families. In Europe the first
481 establishment of the bacterium was reported by Saponari et al. (2013) on
482 olive plants in Apulia showing severe symptoms of leaf scorch and dieback.
483 This first detection was followed by findings of several subspecies and strains
484 of *X. fastidiosa* in Corsica, mainland France, Germany and Spain (Denance et
485 al. 2017; Olmo et al. 2017). The introduction is supposed to be related with
486 trade of infected plant materials (Loconsole et al. 2016; Giampetruzzi et al.
487 2017). *X. fastidiosa* is transmitted exclusively by xylem-sap sucking insects
488 (Frazier 1965). All the members of superfamilies Cercopoidea (commonly
489 known as froghoppers or spittlebugs), Cicadoidea, and the subfamily
490 Cicadellinae within the family Cicadellidae (also known as sharpshooters), are

491 considered xylem-sap feeders (Novotny and Wilson 1997). Epidemiological
492 data suggestive of an insect involvement in pathogen spread in USA, resulted
493 in the identification of sharpshooters as vectors of *X. fastidiosa* to grapevine
494 (Hewitt et al. 1942; Frazier and Freitag 1946). Thereafter, Severin (1950)
495 discovered that, besides sharpshooters, also spittlebugs (Hemiptera:
496 Aphrophoridae) were able to transmit the bacterium. Nevertheless, the
497 epidemiological relevance of spittlebugs seems negligible in the Americans
498 outbreaks. Almeida et al. (2005) suggests that spittlebugs might maintain the
499 inoculum in pastures surrounding diseased vineyard. On the contrary,
500 spittlebugs seem to play an important role in pecan leaf scorch in Louisiana
501 (Sanderlin and Melanson, 2010). Furthermore, cicadas have been claimed to
502 transmit the bacterium, although only two reports with limited datasets are
503 available, and the level of uncertainties about cicadas role as vectors is
504 currently very high (Paião et al. 2002; Krell et al. 2007; EFSA 2015). Overall,
505 the amount of data about *X. fastidiosa* transmission by and interaction with
506 sharpshooters is much larger than the whole background about spittlebugs
507 and cicadas. Noteworthy, in Europe, only nine sharpshooter species are
508 present (Fauna Europaea 2016), and few of them are common and abundant.
509 Conversely, the widespread candidate vectors of *X. fastidiosa* in Europe seem
510 to be spittlebugs (or froghoppers) and, possibly, cicadas (EFSA 2015).

511 The first vector survey during 2013 in Apulia, and successive transmission
512 tests on periwinkle and olive plants, led to the identification of *P. spumarius*
513 as vector of *X. fastidiosa* within the first European bacterium outbreak
514 (Saponari et al. 2014). During the first tests carried out in October-November
515 2013, *P. spumarius* transmitted the bacterium only to periwinkle plants but
516 not to olive (Saponari et al. 2014). The role of *P. spumarius* in the transmission
517 of the fastidious bacterium from olive to olive was proven by successive tests
518 carried out during June-July 2014 (Cornara et al. 2016b). Furthermore, during
519 2014 it was observed that adults emerged in spring on ground cover within
520 olive orchards tested negative to *X. fastidiosa* by qPCR. First positive
521 individuals of 2014 were collected from infected olive canopies, with a great

522 population colonizing this host approximately from sprouting to fruit setting
523 (Cornara et al. 2016b, FIGURE 2). These elements, although not entirely
524 conclusive, strongly suggest: i) the role of olive plants as the main bacterium
525 reservoir within the olive orchard; ii) the implication of *P. spumarius* as the
526 main species involved in the secondary spread of *X. fastidiosa* from olive to
527 olive. *P. spumarius* movements within the olive orchard are still unclear: if the
528 spittlebug follows the general rules for xylem-sap feeders, movement would
529 be influenced by plants physiology and biochemistry, with *P. spumarius*
530 moving from plant to plant according to daily fluctuation of nutrient elements
531 into the xylem sap (Andersen et al. 1992). Another important factor
532 influencing spittlebugs movement is humidity (as previously discussed).
533 During summer, when ground cover dries up and temperature dramatically
534 increases, the spittlebugs find a perfect shelter in the olive canopies, where
535 they can acquire the bacterium. After *X. fastidiosa* acquisition, *P. spumarius*
536 would play an important role either in secondary transmission within the
537 olive orchards, or in primary transmission to plants surrounding the orchard
538 or several kilometers apart. Short-range bacterial dispersal after acquisition
539 seems to rely on active spittlebug movements, whereas anthropogenic
540 factors may have played a major role in long-range dispersal of infective
541 individuals in Apulia. This theory is consistent with the spotted distribution of
542 the outbreaks within Lecce's province (Martelli et al. 2016). Sumatra clove
543 disease, caused by *Pseudomonas syzigii*, transmitted by Machaerotidae,
544 sister taxon of Aphrophoridae, shows an analogous pattern, with sources of
545 primary spread several kilometers far from new hotspots (Eden-Green et al.
546 1992). Furthermore, the large number of spittlebugs present within the olive
547 canopy for several weeks/months, may dramatically increase the probability
548 of infection, and reduce the disease incubation period (Daugherty and
549 Almeida 2009). According to Purcell (1981), the probability of a plant being
550 infected with *X. fastidiosa*, is directly proportional to four factors: vector
551 infectivity (i), transmission efficiency (E), number of vectors (n), and time they
552 spent on the host (t). As shown by Daugherty and Almeida (2009), i and E are

553 proportional to n and t: practically speaking, even if the vector is relatively
554 inefficient, the infection is inevitable when large population settles on the
555 host plant for long time. Incubation time on olive is still unknown: Saponari
556 et al. (2016) stated symptoms appear 12 to 14 months after artificial
557 inoculation on young olive plants in greenhouse conditions. Nevertheless,
558 incubation period in the field may differ from lab results, mainly because of
559 the different age of the plants and their previous and concomitant exposition
560 to biotic and abiotic stresses, and number of inoculation events. Regarding *P.*
561 *spumarius* transmission efficiency, Cornara et al. (2016c) estimated the daily
562 value in ca. 20% using as a proxy grapevine plants and *X. fastidiosa* subsp.
563 *fastidiosa* strain STL. Within grapevine, *X. fastidiosa* reaches very high
564 population, likely 100 to 1000 times greater than in olive (Saponari et al.
565 2016): taking into account that the main factor influencing transmission is
566 bacterial population present within the source plant (Hill and Purcell 1997),
567 *P. spumarius* transmission efficiency to olive could be lower than the value
568 reported for grapevine. Altogether, data gathered from transmission
569 experiments with grapevine, showed that *X. fastidiosa* transmission by *P.*
570 *spumarius* does not differ from the dynamics reported for sharpshooters.
571 Nevertheless, whereas several authors reported no correlation between
572 bacterial population in the vector foregut and pathogen transmission (Hill and
573 Purcell 1995; Almeida and Purcell 2003; Rashed et al. 2011), Cornara et al.
574 (2016c) showed that for *P. spumarius* this correlation exists. Moreover, either
575 in the experiment with grapevine carried out with Californian population of
576 the spittlebug, or analyzing insects collected from infected olive canopies in
577 Apulia, the bacterial population found within the insect was ten to one
578 hundred times lower than that reported for sharpshooters (Cornara et al.,
579 2016a, Cornara et al, 2016c). Thus, despite similarities in overall transmission
580 dynamic, *P. spumarius* showed two novel unexplored characteristics in
581 relation to better studied sharpshooters: the insect hosts a relatively low
582 population of bacterial cells, around 100 to 1000 cells for individual;
583 moreover, the extent of the population is directly correlated with

584 transmission efficiency. Furthermore, Killiny and Almeida (2009) reported
585 that, once acquired, *X. fastidiosa* starts to multiply at a constant rate within
586 the foregut, saturating the available space in ca. 7 days, reaching a population
587 of 10000 to 50000 cells/insect. Quantitative PCR analyses of the individuals
588 used for transmission experiment on grapevine, revealed that the bacterium
589 within the foregut of *P. spumarius* reaches the population peak of ca. 1000
590 cells/insect in less than three days (Cornara et al. 2016c). Cornara et al.
591 (2016c) hypothesized two possible explanations underlying the observed
592 phenomenon: the first relies on cuticle chemistry and potential bacterial
593 receptors within the foregut; the second is related to insect probing behavior.
594 During acquisition, *X. fastidiosa* adhesins bind to insect cuticle, likely on the
595 part of the precibarium proximal to cibarium (Almeida and Purcell 2006;
596 Killiny and Almeida 2009); chitin, the main cuticle polysaccharide, is used by
597 the bacterial cells as a carbon source (Killiny et al. 2010). *P. spumarius* foregut
598 may host few bacterial cells because of differences in availability of
599 polysaccharides or sites where the first binding or the successive
600 multiplication take place. Alternatively or concomitantly, the observed
601 difference may be related to *P. spumarius* probing behavior: the meadow
602 spittlebug has been demonstrated to feed on the main xylem stream, where
603 tremendous tension even greater than -10 bars occurs (Malone et al. 1999).
604 To feed on xylem mainstream *P. spumarius* has to overcome this tension,
605 loading the cibarial muscles until balancing vessel negative pressure. As
606 shown with sharpshooters, the cibarial pump performs one up-and-down
607 movement every second, and the fluid flows within the foregut very rapidly
608 (Purcell et al. 1979; Dugravot et al. 2008). Under these conditions, bacterial
609 cells binding should not be straightforward, but the feeding behavior of *P.*
610 *spumarius* may make either binding or multiplication even more challenging.
611 Electrical penetration graph (EPG) is a technology devised by McLean and
612 Kinsey in 1964, then improved by Tjallingii in 1978, and to date considered an
613 essential tool in research on probing behavior and pathogen transmission by
614 piercing-sucking insects (Walker 2000). A detailed EPG-assisted study of *P.*

615 *spumarius* probing behavior in relation to *X. fastidiosa* transmission, may
616 shed the light on this phenomenon, providing useful data for blocking
617 pathogen transmission, following the approach illustrated by Killiny et al.
618 (2012) and Labroussaa et al. (2016).

619 Besides olive, Cornara et al. (2016a) showed that *P. spumarius* transmits *X.*
620 *fastidiosa* *pauca* ST53 to several host plants, namely oleander, periwinkle, the
621 stonefruit rootstock GF677, and sweet orange, but not grapevines. For
622 transmission tests, groups of five insects per plant were used; as expected,
623 the number of infective spittlebugs was directly correlated with transmission
624 probability. The same authors reported that systemic colonization does not
625 take place neither in GF677 nor in sweet orange, consistently with bacterial
626 artificial inoculation data reported by Saponari et al. (2016). Furthermore, *X.*
627 *fastidiosa* was never detected in hundreds of *Citrus* spp. plants monitored
628 within the infected area (Martelli et al. 2016). Eventually, the above reported
629 findings demonstrate the main role of *P. spumarius* as vector of *X. fastidiosa*
630 in the Apulian outbreak. Besides the meadow spittlebug, three other xylem
631 "specialist" feeders have been found in surveyed Apulia olive orchards: *N.*
632 *campestris*, *Cercopis sanguinolenta* Scopoli (1763), *Cicada orni* L. (1758)
633 (Cornara et al. 2016b). Whereas either *N. campestris* or *C. sanguinolenta*
634 seem not to play a significant role in the transmission of *X. fastidiosa* to olive,
635 the impact of these species as vector of the bacterium should be investigated
636 on other host plants and agro-ecosystems (Cornara et al. 2016b). Regarding
637 *C. orni*, in a recent natural infectivity test Cornara et al. (unpublished) found
638 three out of 160 cicadas positive to *X. fastidiosa* by qPCR, while no
639 transmission to olive recipient plants occurred. Eventually, more research
640 efforts are needed in order to understand the epidemiological relevance of
641 either *P. spumarius* or other candidate vectors in agro-ecosystems different
642 from olive orchards, and across others European epidemics.

643

644 **Control: integrated pest management and sustainable control perspectives**

645 Integrated pest management strongly relies on effective **sampling and**
646 **surveillance methods**. Unfortunately, to date, an effective method for *P.*
647 *spumarius* sampling is still missing. Sweep net is the most common method
648 used for adult collection; however, as remarked by Purcell et al. (1994),
649 sweep net is a poorly effective tool for sampling insects from a tree canopy,
650 in contrast with its high efficacy on the ground cover. Although sweep net is
651 the tool largely used to collect *P. spumarius*, other methods, namely
652 minicage (biocenometers), pitfall traps, sticky traps, aerial suction traps,
653 beat tray, and tanglefoot bands have been tested. However, all these
654 methods were proven to be less effective than sweep net (Weaver and King
655 1954; Lavigne 1959; Wilson and Shade 1967; Novotny 1992; Pavan 2000;
656 Bleicher et al. 2010). To the best of our knowledge, only one study focused
657 on effectiveness of different color sticky traps in collecting the meadow
658 spittlebug has been performed, with yellow resulting more attractive than
659 green, red, pink, blue, and white (Wilson and Shade 1967). Nevertheless,
660 preliminary results of observations conducted in Apulia and Spain, suggests
661 the low efficacy of yellow sticky traps in *P. spumarius* collection and other
662 colors need to be tested (Morente et al., unpublished data). Researches
663 carried out on vibrational signals produced by *Homalodisca vitripennis*
664 Germar (1821) opened new and interesting perspectives for the control of
665 this *X. fastidiosa* vector in US vineyard (Nieri et al. 2017). The occurrence of
666 communication through vibrations should be explored also for *P. spumarius*;
667 the outcomes of such researches could open new venues in order to set up
668 an effective monitoring tool. On the other hand, recent studies on the fine
669 structure of antennal sensilla of the spittlebug allowed to identify
670 chemoreceptors (Ranieri et al. 2016). Although the presence of olfactory
671 receptors among the antenna is limited, it is possible that *P. spumarius*
672 responds to olfactory attractants, e.g. plant attractants, thus providing new
673 tools for monitoring and control. Unfortunately, so far pheromones have
674 not been identified in spittlebugs, with the exception of an aggregation
675 pheromone of the rice spittlebug *Callitettix versicolor* nymphs (Chen and

676 Liang 2015) and therefore monitoring and control methods based on the use
677 of pheromones are very unlikely to be developed for *P. spumarius*.

678 To date, considering that *X. fastidiosa* eradication is no more feasible and
679 Apulia had become a reservoir of the bacterium, an effective disease
680 management strategy is mandatory for the survival of agriculture and
681 landscape (Strona et al. 2017). Strategies focused on disruption of only one
682 single aspect of the complex interaction vector-plant-pathogen has proven
683 many times to be unsuccessful (Almeida et al., 2005). *X. fastidiosa* provides
684 one of the best example of an arthropod-borne pathogen whose control
685 strongly depends on several interacting variables: crop; agricultural
686 practices; weather; vector biology, host range and behavior; pathogen host
687 range; transmission mode, primary or secondary (Almeida et al. 2005).
688 Therefore, management of *X. fastidiosa* epidemics should be based on a
689 combination of multiple tactics that partially interrupt more than one
690 interaction of the pathosystem (Almeida et al. 2005). At least in Salento
691 olive orchards, available data strongly suggests that *P. spumarius* transmits
692 *X. fastidiosa* from tree to tree, with olive being the primary source of the
693 bacterium. In such case of “secondary transmission”, the disease control
694 strategy should be based on exclusion of the pathogen from propagative
695 material, removal of infected plants, and **vector control** to reduce
696 transmission within and between the orchards (Lopes and Krugner 2016).
697 According to the funding theories of integrated pest management, an
698 effective pest control strategy should target the most vulnerable stages of
699 the insect life cycle, when the control tools can act on the residual pest
700 population already affected by, or exposed to, biotic and abiotic factors
701 (Lewis et al. 1997; Kogan 1998). Looking at *P. spumarius* life cycle and
702 behavior, two are the weakest point on which control measures could
703 achieve the best results: nymphal stage, and newly emerged non-infective
704 adults shifting to olive plants. **Nymphs** develop in natural vegetation within
705 and on the margins and hedgerows of olive groves during spring. Removal of
706 ground cover hosting the nymphs either by mowing, soil tillage or herbicides

707 within and surrounding olive orchards, could be effective in drastically
708 reducing resident vector population. Nevertheless, indiscriminate removal
709 of ground cover could be ecologically deleterious, with large-scale
710 environmental impact (Civitello et al. 2015). Another alternative approach
711 could be represented by the use of physical or chemical compounds that
712 remove and further affect the spittle production, since the ability of the
713 nymph to survive out of the spittle is very limited. Chemical control of
714 nymphs was not yet been extensively investigated so far, but could also
715 reduce resident vector population in olive groves.

716 **Adult control** is mainly hampered by migration tendency, that would soon
717 balance the amount of adults dead after insecticides application. King (1952)
718 observed that treatments in mid-summer were ineffective in spittlebug
719 control, since the population would be soon equalized by successive
720 migration from surrounding habitats. *P. spumarius* adults control in olive
721 orchards should be mainly focused on disrupting *X. fastidiosa* acquisition
722 from olive plants, that likely occur when non-infective recently molted adults
723 migrate from ground cover to tender olive sprouts. Carefully planned
724 insecticides application to olive and surrounding plants before adult shift to
725 olive would expose twice the spittlebug to the pesticide: once before and
726 when the insects alight on infected tree; secondly, when potentially infective
727 vectors move to healthy trees (Almeida et al. 2005). Currently, very few
728 reports on the activity of insecticides against *P. spumarius* are available,
729 because before the *X. fastidiosa* European outbreak the species was not
730 considered a pest and therefore was not targeted with insecticides. A recent
731 experiment carried out in Apulia on insecticide control of adults on olive
732 showed that the neonicotinoids acetamiprid and imidacloprid and
733 pyrethroids deltamethrin and lambda-cyhalothrin displayed a high mortality
734 rate. The insect growth regulators buprofezin, and spirotetramat showed no
735 acute lethal effect as well as the pyridine-azomethine pymetrozine. Among
736 botanical insecticides, citrus oil showed a good insect mortality when applied
737 at the volume of 2,000 L/ha (although its activity is not persistent at all), while

738 no toxic effect was recorded using azadirachtin (Dongiovanni et al. 2016).
739 Data on chronic effect or impact of the compounds in reducing *X. fastidiosa*
740 transmission are still missing. Neonicotinoids were successfully used in Brazil
741 against CVC-vectors, through roots and soil application on less than 3 years-
742 old citrus plant, and by spraying on elder plants (Lopes and Krugner 2016).
743 Nevertheless, treatments with Imidacloprid proved to be ineffective in
744 preventing grapevines infection with *X. fastidiosa* in areas with prevalent
745 sources of inoculum and high vector abundance (Krewer et al. 2002). Besides
746 the induced mortality, insecticides as neonicotinoids and repellent as the
747 aluminium silicate kaolin, could interfere with *X. fastidiosa*-vector interaction
748 by affecting vector orientation, host determination and feeding behaviour, as
749 shown in *H. vitripennis* (Tubajika et al. 2007). Kaolin particles, that protects
750 the hosts against the vector by camouflaging the plant with a white coating,
751 making them visually unperceivable, or by reflecting sunlight, might represent
752 a valid control tool especially for organic olive orchards (Puterka et al. 2003).
753 The negative effect of spittlebug migration from the surroundings on the
754 effectiveness of insecticides applications could be mitigated by coupling
755 insecticide treatments with installation around the olive orchard of screen
756 physical barriers, that proved to be efficient in reducing GWSS population
757 migrating from the surrounding citrus orchards into vineyards (Blua et al.
758 2005). Nevertheless, even if effective, the benefits coming from proper
759 control strategies would result in just a “hold back the tide” strategy, if the
760 measures will not be extended to the widest possible area.

761 **The control of the meadow spittlebug with parasitoids and predators** is still
762 far from its application, and more research efforts are needed to find a
763 suitable candidate to pursue this task. Indeed, detailed information about
764 the meadow spittlebug natural enemies are still scattered and missing.
765 Predation seems not to be an important source of mortality (Whittaker
766 1973). Birds, frogs, Arachnids Phalangiidae, Hymenoptera, Diptera and
767 Coleoptera Carabidae, prey *P. spumarius* (Phillipson 1960; Halkka et al.
768 1976; Harper and Whittaker 1976; Henderson et al. 1990). Westwood in

769 1840 (cited by Weaver and King 1954), and more recently Pagliano and Alma
770 (1997), observed *Argogorytes mystaceus* L. (1761) (Hymenoptera:
771 Sphecidae) (*Gorytes mystaceus* in Westwood 1840) dragging *P. spumarius*
772 nymphs from their spittle masses. Very recently, the Reduviidae bug *Zelus*
773 *renardii* Kolenati has been proposed as a biological control agent in olive
774 orchards of *P. spumarius* (Salerno et al., 2017). The dipteran parasitoid
775 *Verrallia aucta* Fallen (1817) (Diptera: Pipunculidae), found in Europe and
776 Central Siberia, is responsible for adults sterility bringing them to death just
777 in the last part of their cycle; parasitism rate is likely to be not greater than
778 1% (Whittaker 1969; Whittaker 1973; Meyer and Bruyn 1984; van Driesche
779 and Peters 1987). Furthermore, the nematode *Agamermis decaudata* Cobb,
780 Steiner & Christie (1923), and the entomopathogenic fungi *Entomophthora*
781 sp. Fresen (1856), attack the adults (Weaver and King 1954, Harper and
782 Whittaker, 1976; Ben-Ze'Ev and Kenneth, 1981). Eggs are parasitized by
783 Hymenoptera of the genus *Ooctonus* spp., *Tumidiscapus* sp., and *Centrodora*
784 sp., which were found parasitizing around 10% of field collected eggs in
785 1951 in Ohio (Weaver and King 1954).

786 Promising researches focused on **disrupting *X. fastidiosa*-vector interactions**
787 are ongoing. Deliver of lectins, carbohydrates and antibodies to a vector
788 through artificial diet, significantly impacted bacterial acquisition and
789 subsequent transmission (Killiny et al. 2012). Furthermore, recombinant
790 peptides efficiently blocked *X. fastidiosa* acquisition and initial bind to
791 foregut, while they did not interfere with successive steps of bacterial
792 multiplication once the bacterium had been acquired and was bound to the
793 cuticle (Labroussa et al., 2016). Nevertheless, such strategies, tested on
794 sharpshooters, should be further assessed for *P. spumarius*, whose intimate
795 relationship with the bacterium is different, to some extent, to the one of
796 Cicadellinae (Cornara et al. 2016c). Moreover, deepening our knowledge
797 about *P. spumarius* feeding behavior and *X. fastidiosa* transmission
798 mechanism through a real-time observation device as EPG could open new

799 venues in the discovery of an effective strategy to disrupt bacterium-
800 spittlebug interaction.

801

802 **Concluding remarks**

- 803 • The meadow spittlebug *P. spumarius*, never considered a pest in
804 Europe, raised the attention of scientists and stake-holders after the
805 discovery of its main role in the transmission of *X. fastidiosa* strain ST53
806 to olive in the first reported European outbreak of the bacterium,
807 occurred in Apulia (South Italy) in 2013.
- 808 • *P. spumarius* is widely distributed, covering most of the Palearctic
809 region, and extending to Nearctic. The spittlebug is highly polyphagous,
810 occurring in most of the terrestrial habitats; furthermore, *P. spumarius*
811 has the potential to live under different environmental conditions, from
812 moist to relatively dry, as long as the host plants are actively growing,
813 and not subjected to severe water stress.
- 814 • Lack of key information on *P. spumarius* urgently calls for research on
815 aspects considered fundamental for developing effective pest
816 management strategies: life history, ecology, phenology, population
817 dynamics, movement and dispersal, tri-trophic relationships, host plant
818 association and preference, reproductive biology, feeding behavior,
819 vibrational communication , effect of plant volatiles on host search and
820 recognition, insect microbiome, natural enemies.
- 821 • *X. fastidiosa*-associated disease control strategies should include
822 measures aimed at i) suppressing vector populations ii) suppressing
823 sources of inoculum for the vector. To achieve these goals, we should
824 consider the ecology and population dynamics of *P. spumarius* in
825 different sites and crop systems, as there are no universally applicable
826 solutions. As for suppressing *P. spumarius* population, control strategies
827 should target two stages of the insect life history: nymphs, and newly
828 emerged non-infective adults, that can move toward *X. fastidiosa*-

829 source plants. Moreover, as soon as research will provide new insights
830 on vector-plant-pathogen interactions, innovative control strategies
831 should be developed with the aim of targeting different aspects of these
832 interactions. Finally, control measures should be applied on the widest
833 possible area.

834 • Eventually, the more we learn about the vector-bacterium-plant
835 relationships, the faster we will find the way to cohabit with *X. fastidiosa*
836 associated diseases, reducing the impact of a bacterium that, to date,
837 represents one of the most frightening threat to European agriculture
838 and landscape.

839

840 **Acknowledgements**

841 Authors would like to acknowledge an EFSA procurement action on the
842 information on biology and control of vectors of *Xylella fastidiosa*, the
843 European Union Horizon 2020 research and innovation program under grant
844 agreement no. 635646 POnTE (Pest Organisms Threatening Europe), and
845 grant agreement no. 727987 XF-ACTORS.

846

847 **Compliance with ethical standard**

848 The authors declare no conflicts of interest. This article does not contain any
849 studies with human participants or animals performed by any of the
850 authors.

851

852 **Authors contribution**

853 Collected data and wrote the paper: DC, DB, AF

854

References

- 856 Almeida RP, Blua MJ, Lopes JR, Purcell AH (2005) Vector transmission of *Xylella fastidiosa*:
857 applying fundamental knowledge to generate disease management strategies. *Ann Entomol Soc*
858 *Am* 98: 775–786.
- 859 Almeida RP, Nunney L (2015) How Do Plant Diseases Caused by *Xylella fastidiosa* Emerge? *Plant Dis*
860 99: 1457–1467.
- 861 Almeida RP, Purcell AH (2006) Patterns of *Xylella fastidiosa* colonization on the precibarium of
862 sharpshooter vectors relative to transmission to plants. *Ann Entomol Soc Am* 99: 884–890.
- 863 Almeida R.PP, Purcell AH (2003) *Homalodisca coagulata* (Hemiptera, Cicadellidae) transmission of
864 *Xylella fastidiosa* to almond. *Plant Dis* 87: 1255–1259.
- 865 Andersen PC, Brodbeck BV, Mizell RF (1992) Feeding by the leafhopper, *Homalodisca coagulata*, in
866 relation to xylem fluid chemistry and tension. *J Insect Physiol* 38: 611–622.
- 867 Barber GW, Ellis WO (1922) Eggs of Tree Cercopidae. *Psyche* 29: 1–3.
- 868 Ben-Ze'ev I, Kenneth RG (1981) *Zoophthora radicans* and *Zoophthora petchi* sp. nov. [Zygomycetes:
869 Entomophthorales], two species of the “*Sphaerosperma* group” attacking leaf-hoppers and frog-
870 hoppers [Hom.]. *Entomophaga* 26: 131–142.
- 871 Berry A J, Willmer PG (1986). Temperature and the colour polymorphism of *Philaenus spumarius*
872 (Homoptera: Aphrophoridae). *Ecol Entomol* 11: 251–259.
- 873 Biedermann R (2002) Leafhoppers (Hemiptera, Auchenorrhyncha) in fragmented habitats. *Denisia* 176:
874 523–530.
- 875 Bleicher K, Orosz A, Cross J, Markó V (2010) Survey of leafhoppers, planthoppers and froghoppers
876 (Auchenorrhyncha) in apple orchards in South-East England. *Acta Phytopathol Hun* 45: 93–105.
- 877 Blua MJ, Campbell K, Morgan DJW, Redak RA (2005) Impact of a screen barrier on dispersion
878 behavior of *Homalodisca coagulata* (Hemiptera: Cicadellidae). *J Econ Entomol* 98: 1664–1668.
- 879 Bodino N, Plazio E, Cavalieri V, Dongiovanni E, Ripamonti M, Volani S, Gilioli G, Fumarola G, Di
880 Carolo M, Porcelli F, Bosco D (2017) Host-plant association and host-shifting of nymphs and
881 adults of *Philaenus spumarius* L. in Italian olive orchards. *Proceedings 3rd Hemipteran-Plant*
882 *Interactions Symposium (HPIS), Madrid, Spain, June 4–8, 2017*, 36.
- 883 Braccini P, Pavan F (2000) Auchenorrhyncha: potential vectors of phytoplasmas associated with vine
884 yellows. *Informatore Agrario* 56: 103–107.
- 885 Burrows M (2003) Biomechanics: Froghopper insects leap to new heights. *Nature* 424: 509–509.
- 886 Chen X, Liang AP (2015) Identification of a self-regulatory pheromone system that controls nymph
887 aggregation behavior of rice spittlebug *Callitettix versicolor*. *Front Zool* 12: 1–12.
- 888 Chen X, Meyer-Rochow VB, Fereres A, Morente A, Liang AP (2017) The role of biofoam in shielding
889 spittlebug nymphs (Insecta, Hemiptera, Cercopidae) against bright light. *Ecol Entomol*. doi:
890 10/1111/een.12496.
- 891 Chmiel SM, Wilson MC (1979) Estimation of the lower and upper developmental threshold
892 temperatures and duration of the nymphal stages of the meadow spittlebug, *Philaenus*
893 *spumarius*. *Environ Entomol* 8: 682–685.
- 894 Civitello DJ, Cohen J, Fatima H, Halstead NT, Liriano J, McMahon TA, Ortega CN, Sauer EL, Sehgal,
895 T, Young S (2015) Biodiversity inhibits parasites: broad evidence for the dilution effect. *P Natl*
896 *Acad Sci* 112: 8667–8671.
- 897 Cornara D, Cavalieri V, Dongiovanni C, Altamura G, Palmisano F, Bosco D, Porcelli F, Almeida RP,
898 Saponari M (2016a) Transmission of *Xylella fastidiosa* by naturally infected *Philaenus*
899 *spumarius* (Hemiptera, Aphrophoridae) to different host plants. *J Appl Entomol*.
900 doi:10.1111/jen.12365
- 901 Cornara D, Porcelli F (2014) Observations on the biology and ethology of Aphrophoridae: *Philaenus*

902 *spumarius* in the Salento peninsula, in: Proceedings “International Symposium on the
903 European Outbreak of *Xylella Fastidiosa* in Olive”, Gallipoli–Locorotondo, Italy, 21-24
904 October 2014.

905 Cornara D, Saponari M, Zeilinger AR, Stradis A de, Boscia D, Loconsole G, Bosco D, Martelli GP,
906 Almeida RPP, Porcelli F (2016b) Spittlebugs as vectors of *Xylella fastidiosa* in olive orchards
907 in Italy. J Pest Sci 1–10. doi:10.1007/s10340-016-0793-0

908 Cornara D, Sicard A, Zeilinger AR, Porcelli F, Purcell AH, Almeida RPP (2016c) Transmission of
909 *Xylella fastidiosa* to grapevine by the meadow spittlebug. Phytopathology 106: 1285–1290.

910 Crews L, McCully M, Canny M, Huang C, Ling L (1998) Xylem feeding by spittlebug nymphs: some
911 observations by optical and cryo-scanning electron microscopy. Am J Bot 85: 449–449.

912 Daugherty MP, Almeida RPP (2009) Estimating *Xylella fastidiosa* transmission parameters: decoupling
913 sharpshooter number and feeding period. Entomol Exp Appl 132: 84–92.

914 Delong DM, Severin HH (1950). Spittle-insect vectors of Pierce’s disease virus. Hilgardia 19 : 339–
915 376.

916 Denance N, Legendre B, Briand M, Olivier V, de Boisseson C, Poliakoff F, Jacques MA (2017)
917 Several subspecies and sequence types are associated with the emergence of *Xylella fastidiosa*
918 in natural settings in France. Plant Pathol. doi: 10.1111/ppa.12695.

919 Dongiovanni C, Cavalieri V, Altamura G, Di Carolo M, Fumarola G, Saponari M, Porcelli F (2016)
920 Preliminary results of comparative efficacy evaluation trials against *Philaenus spumarius* L.,
921 vector of *Xylella fastidiosa*. Options Méditerranéennes, A No.121, 2017 - *Xylella fastidiosa* &
922 the Olive Quick Decline Syndrome (OQDS). A serious worldwide challenge for the safeguard
923 of olive trees 79-80.

924 Drosopoulos S (2003) New data on the nature and origin of colour polymorphism in the spittlebug
925 genus *Philaenus* (Hemiptera: Aphrophoridae). Ann Soc Entomol Fr 31–42.

926 Drosopoulos S, Asche M (1991) Biosystematic studies on the spittlebug genus *Philaenus* with the
927 description of a new species. Zool J Linn Soc 101: 169–177.

928 Drosopoulos S, Remane R (2000) Biogeographic studies on the spittlebug *Philaenus signatus*
929 Melichar, 1896 species group (Hemiptera: Aphrophoridae) with the description of two new
930 allopatric species Ann Soc Entomol Fr 269–277.

931 Dugravot S, Backus EA, Reardon BJ, Miller TA (2008) Correlations of cibarial muscle activities of
932 *Homalodisca* spp. sharpshooters (Hemiptera: Cicadellidae) with EPG ingestion waveform and
933 excretion. J Insect Physiol 54: 1467–1478.

934 Eden-Green S, Balfas R, Sutarjo T (1992) Characteristics of the transmission of Sumatra disease of
935 cloves by tube-building cercopoids, *Hindola* spp. Plant Pathol 41: 702–712.

936 Edwards WD (1935) Strawberry pests including the spittlebug. Ann Rpt Oregon State Hort Soc 27: 58-
937 65.

938 EFSA (2015) Scientific Opinion on the risk to plant health posed by *Xylella fastidiosa* in the EU territory,
939 with the identification and evaluation of risk reduction options. EFSA J 13:3989.

940 Fauna Europaea (2016) Museum für Naturkunde Leibniz-Institut für Evolutions- und
941 Biodiversitätsforschung Invalidenstr. Berlin, Germany. <https://fauna-eu.org/>

942 Fereres A. (2015) Insect vectors as drivers of plant virus emergence. Curr Opin Virol 10: 42-46.

943 Fisher EH, Allen TC (1946) Alfalfa and Clover Severely Damaged by Spittlebugs. What’s New in
944 Farm Science. Wisconsin Agr Exp Sta Bul 469: 15–16.

945 Frazier NW (1965) Xylem viruses and their insect vectors, in: Proceedings of the International
946 Conference on Virus and Vectors on Perennial Hosts, with Special Reference to *Vitis* 91–99.

947 Frazier NW, Freitag JH (1946) 10 Additional Leafhopper Vectors Of The Virus Causing Pierces
948 Disease Of Grapes. Phytopathology 36: 634–637.

949 Freeman JA (1945) Studies in the distribution of insects by aerial currents. The Journal of Animal

- 950 Ecology 128–154.
- 951 Giampetruzzi A, Saponari M, Loconsole G, Boscia D, Savino VN, Almeida RP, Zicca S, Landa BB,
952 Chacón-Díaz C, Saldarelli P (2017) Genome-Wide Analysis Provides Evidence on the Genetic
953 Relatedness of the Emergent *Xylella fastidiosa* Genotype in Italy to Isolates from Central
954 America. *Phytopathology* PHYTO–12.
- 955 Gibson DO (1974) Batesian mimicry without distastefulness? *Nature* 250: 77–79.
- 956 Goidanich A (1954) *Enciclopedia Agraria Italiana*. Roma, REDA.
- 957 Grant JF, Lambdin PL, Follum RA (1998) Infestation levels and seasonal incidence of the meadow
958 spittlebug (Homoptera: Cercopidae) on musk thistle in Tennessee. *J Agr Entomol* 15: 83-91.
- 959 Guglielmino A, Bückle C, Remane R (2005). Contribution to the knowledge of the Auchenorrhyncha
960 fauna of Central Italy (Hemiptera, Fulgoromorpha et Cicadomorpha). *Marburger*
961 *Entomologische Publikationen Band 3 Heft 3: 13 – 98.*
- 962 Gulijeva EM (1961). Pennitsa *P.spumarius*-vreditel' Zernovykh lul'tur v Azerbaidzhane. *Izvest Akad*
963 *Navk SSR Ser Biol I Med Nauk* 5: 73-81.
- 964 Halkka A, Halkka L, Halkka O, Roukka K, Pokki J (2006) Lagged effects of North Atlantic Oscillation
965 on spittlebug *Philaenus spumarius* (Homoptera) abundance and survival. *Glab Change Biol* 12:
966 2250–2262.
- 967 Halkka O (1962) Equilibrium Populations of *Philaenus spumarius* L. *Nature* 193: 93–94.
- 968 Halkka O (1964) Geographical, spatial and temporal variability in the balanced polymorphism of
969 *Philaenus spumarius*. *Heredity* 19: 383–401.
- 970 Halkka O, Heinonen L, Raatikainen M, Vasarainen A (1966) Crossing experiments with *Philaenus*
971 *spumarius* (Homoptera). *Hereditas* 56: 306–312.
- 972 Halkka O, Kohila T, Komila T (1976) Persistence of visual polymorphism, despite a low rate of
973 predation, in *Philaenus spumarius* (L.)(Homoptera, Aphrophoridae). *Ann Zool Fenn* 185–188.
- 974 Halkka O, Raatikainen M, Halkka L, Lokki J (1971) Factors determining the size and composition of
975 island populations of *Philaenus spumarius* (L.)(Homoptera). *Acta Entomol Fenn* 28: 83-100.
- 976 Halkka O, Raatikainen M, Halkka L, Raatikainen T (1977) Coexistence of four species of spittle-
977 producing Homoptera, in: *Annales Zoologici Fennici*. JSTOR, pp. 228–231.
- 978 Halkka, O., Raatikainen, M., Vasarainen, A., Heinonen, L., 1967. Ecology and ecological genetics of
979 *Philaenus spumarius* (L.) (Homoptera). *Ann Zool Fenn* 1–18.
- 980 Harper G, Whittaker JB (1976) The role of natural enemies in the colour polymorphism of *Philaenus*
981 *spumarius* (L.). *J Anim Ecol* 91–104.
- 982 Henderson G, Hoffman GD, Jeanne RL (1990). Predation on cercopids and material use of the spittle in
983 aphid-tent construction by prairie ants. *Psyche* 97: 43–53.
- 984 Hewitt WB, Frazier NW, Jacob HE, Freitag JH (1942). Pierce's disease of grapevines. California
985 Agricultural Experimental Station Circular 1–32.
- 986 Hill B, Purcell AH (1995) Acquisition and Retention of *Xylella fastidiosa* by an Efficient Vector,
987 *Graphocephala atropunctata*. *Phytopathology* 85: 209-212. doi:10.1094/Phyto-85-209
- 988 Hill BL, Purcell AH (1997). Populations of *Xylella fastidiosa* in plants required for transmission by an
989 efficient vector. *Phytopathology* 87: 1197–1201.
- 990 Hill GT, Sinclair WA (2000) Taxa of leafhoppers carrying phytoplasmas at sites of ash yellows
991 occurrence in New York State. *Plant Dis* 84: 134-138.
- 992 Hoffman G, McEvoy PB (1985a) Mechanical limitations on feeding by meadow spittlebugs *Philaenus*
993 *spumarius* (Homoptera: Cercopidae) on wild and cultivated host plants. *Ecol Entomol* 10: 415–
994 426.
- 995 Hoffman GD, McEvoy PB (1985b) The mechanism of trichome resistance in *Anaphalis margaritacea*
996 to the meadow spittlebug *Philaenus spumarius*. *Entomol Exp Appl* 39: 123–129.
- 997 Horsfield D (1977) Relationships between feeding of *Philaenus spumarius* (L.) and the amino acid

- 998 concentration in the xylem sap. *Ecol Entomol* 2: 259–266.
- 999 Horsfield D (1978) Evidence for xylem feeding by *Philaenus spumarius* (L.)(Homoptera: Cercopidae).
1000 *Entomologia Exp Appl* 24: 95–99.
- 1001 Ivanauskas A, Valiūnas D, Jomantienė R, Picciau L, Davis RE (2014) Possible insect vectors of
1002 “*Candidatus Phytoplasma asteris*” and “*Ca. Phytoplasma pruni*”-related strains in Lithuania.
1003 *Žemdirbystė (Agriculture)* 101: 313–320.
- 1004 Karban R, Strauss SY (2004) Physiological tolerance, climate change, and a northward range shift in
1005 the spittlebug, *Philaenus spumarius*. *Ecol Entomol* 29: 251–254.
- 1006 Keskinen E, Meyer-Rochow VB (2004) Post-embryonic photoreceptor development and dark/light
1007 adaptation in the spittle bug *Philaenus spumarius* (L.)(Homoptera, Cercopidae). *Arthropod*
1008 *Struct Dev* 33: 405–417.
- 1009 Killiny N, Almeida RP (2009) *Xylella fastidiosa* afimbrial adhesins mediate cell transmission to plants
1010 by leafhopper vectors. *Appl Environ Microb* 75: 521–528.
- 1011 Killiny N, Prado SS, Almeida RP (2010) Chitin utilization by the insect-transmitted bacterium *Xylella*
1012 *fastidiosa*. *Appl Environ Microb* 76: 6134–6140.
- 1013 Killiny N, Rashed A, Almeida RP (2012) Disrupting the transmission of a vector-borne plant pathogen.
1014 *Appl Environ Microb* 78: 638–643.
- 1015 King DR (1952) The ecology of the Meadow Spittlebug *Philaenus leucophthalmus* (L.) L Family
1016 Cercopidae. PhD Thesis, The Ohio State University.
- 1017 Koga R, Bennett GM, Cryan JR, Moran NA (2013). Evolutionary replacement of obligate symbionts in
1018 an ancient and diverse insect lineage. *Environ Microbiol* 15: 2073–2081.
- 1019 Kogan M (1998) Integrated pest management: historical perspectives and contemporary developments.
1020 *Ann Rev Entomol* 43: 243–270.
- 1021 Krell RK, Boyd EA, Nay JE, Park YL, Perring TM (2007) Mechanical and Insect Transmission of
1022 *Xylella fastidiosa* to *Vitis vinifera*. *Am J Enol Vitic* 58: 211–216.
- 1023 Krewer G, Dutcher JD, Chang CJ (2002) Imidacloprid Insecticide Slows Development of Pierce’s
1024 Disease in Bunch Grapes. *J Entomol Sci* 37: 101–112. doi:10.18474/0749-8004-37.1.101.
- 1025 Labroussaa F, Zeilinger A, Almeida RP (2016) Blocking the Transmission of a Non-circulative Vector-
1026 borne Plant Pathogenic Bacterium. *Mol Plant Microbe In* 29: 535-544.
- 1027 Landi F, Prandini A, Paltrinieri S, Mori N, Bertaccini A (2007) Detection of different types of
1028 phytoplasmas in stone fruit orchards in northern Italy. *B Insectol* 60: 163.
- 1029 Lavigne B (1959) Biology of *Philaenus leucophthalmus* (L.), in Massachusetts. *J Econ Entomol* 52:
1030 904-907.
- 1031 Lees DR, Dent CS (1983) Industrial melanism in the spittlebug *Philaenus spumarius* (L) (Homoptera:
1032 Aphrophoridae). *Biol J Linn Soc* 19: 115–129.
- 1033 Lewis WJ, Van Lenteren JC, Phatak SC, Tumlinson JH (1997) A total system approach to sustainable
1034 pest management. *P Natl A Sci* 94: 12243–12248.
- 1035 Lis A, Maryńska-Nadachowska A, Kajtoch L (2015) Relations of *Wolbachia* infection with
1036 Phylogeography of *Philaenus spumarius* (Hemiptera: Aphrophoridae) populations within and
1037 beyond the Carpathian contact zone. *Microb Ecol* 70.2 (2015): 509-521.
- 1038 Loconsole G, Saponari M, Boscia D, D’Attoma G, Morelli M, Martelli GP, Almeida RPP (2016)
1039 Intercepted isolates of *Xylella fastidiosa* in Europe reveal novel genetic diversity. *Eur J Plant*
1040 *Pathol* 1–10.
- 1041 Lopes JRS, Krugner R (2016) Transmission ecology and epidemiology of the citrus variegated
1042 chlorosis strain of *Xylella fastidiosa*. *Am Phytopatol Soc* 195–208.
- 1043 Malone M, Watson R, Pritchard J (1999). The spittlebug *Philaenus spumarius* feeds from mature
1044 xylem at the full hydraulic tension of the transpiration stream. *New Phytologist* 143: 261–271.
- 1045 Martelli GP, Boscia D, Porcelli F, Saponari M (2016) The olive quick decline syndrome in south-east

1046 Italy: a threatening phytosanitary emergency. Eur J Plant Pathol 144: 235–243.

1047 Maryńska-Nadachowska A, Kuznetsova VG, Lachowska D, Drosopoulos S (2012) Mediterranean

1048 species of the spittlebug genus *Philaenus*: Modes of chromosome evolution. J Insect Sci 12: 1–

1049 17.

1050 Masters GJ, Brown VK, Clarke IP, Whittaker JB, Hollier JA (1998) Direct and indirect effects of

1051 climate change on insect herbivores: Auchenorrhyncha (Homoptera). Ecol Entomol 23: 45–52.

1052 Matteoni JA, Sinclair WA (1988) Elm yellows and ash yellows. Tree mycoplasmas and mycoplasma

1053 diseases 19–31.

1054 Medler JT (1955) Method of predicting the hatching Date of the Meadow Spittlebug. J Econ Entomol

1055 48: 204-205.

1056 Meyer M de, Bruyn L de (1984) On the phenology of some Pipunculidae (Diptera) in Belgium.

1057 Bulletin et Annales de La Société Royale Belge D'entomologie 123–131.

1058 Mundinger FG (1946) The control of spittle insects in strawberry plantings. J Econ Entomol 39: 299–

1059 305.

1060 Nast J (1972) Palaearctic Auchenorrhyncha (Homoptera) an annotated check list. Polish Science

1061 Publications, Warszawa.

1062 Nicoli Aldini R, Guardiani MC, Cravedi P (1998) Faunistical notes on the hoppers (Homoptera

1063 Auchenorrhyncha) in vineyards in the province of Piacenza. Bollettino di Zoologia Agraria e di

1064 Bachicoltura 30: 61-68.

1065 Nieri R, Mazzoni V, Gordon SD, Krugner R (2017) Mating behavior and vibrational mimicry in the

1066 glassy-winged sharpshooter, *Homalodisca vitripennis*. J Pest Sci 90: 887–899.

1067 Novotny V (1992) Vertical-distribution of leafhoppers (Hemiptera, Auchenorrhyncha) within a

1068 meadow community. Acta Entomol Bohemos 89: 13-20.

1069 Novotny V, Wilson MR (1997) Why are there no small species among xylem-sucking insects? Evol

1070 Ecol 11: 419–437.

1071 Olmo D, Nieto A, Androver F, Urbano A, Beidos O, Juan A et al (2017) First detection of *Xylella*

1072 *fastidiosa* on cherry (*Prunus avium*) and *Polygala myrtifolia* plants, in Mallorca Island,

1073 Spain." Plant Dis 101: 1820.

1074 Ossiannilsson F (1981) The Auchenorrhyncha (Homoptera) of Fennoscandia and Denmark. Part 2: the

1075 families Cicadidae, Cercopidae, Membracidae, and Cicadellidae (excl. Deltocephalinae). Fauna

1076 Entomologica Scandinavica 7: 223–593.

1077 Pagliano G, Alma A (1997) Ricerche etologiche su Gorytini e Alyssonini (Hymenoptera Sphecidae)

1078 parassitoidi di Auchenorrhyncha (Rhynchota Homoptera). Rivista Piemontese di Storia

1079 Naturale 18: 173-181.

1080 Paião FG, Meneguim AM, Casagrande EC, Leite RP (2002) Envolvimento de cigarras (Homoptera,

1081 Cicadidae) na transmissão de *Xylella fastidiosa* em cafeeiro. Fitopatol Bras 27: 67.

1082 Pavan F (2000) Occurrence on elm and phenology of Auchenorrhyncha potential vectors of the

1083 phytoplasma associated with elm yellows disease. Bollettino di Zoologia Agraria e di

1084 Bachicoltura 32: 59-68.

1085 Pavan F (2006) Xylem-feeding auchenorrhyncha potentially involved in Pierce's disease of grapevines

1086 in Europe. Bollettino di Zoologia Agraria e di Bachicoltura 38: 103-114.

1087 Pearson WD (1991) Effect of meadow spittlebug and Australian crop mirid on white clover seed

1088 production in small cages. New Zeal J Agr Res 34: 439–444.

1089 Phillipson J (1960) A contribution to the feeding biology of *Mitopus morio* (F) (Phalangida). The J

1090 Anim Ecol 35–43.

1091 Ponder, K.L., Watson, R.J., Malone, M., Pritchard, J., 2002. Mineral content of excreta from the

1092 spittlebug *Philaenus spumarius* closely matches that of xylem sap. New Phytologist 153: 237–

1093 242.

- 1094 Poos FW (1953) The Meadow Spittlebug: How to Control it. UDA Leaflet, 341.
- 1095 Purcell AH (1980) Almond leaf scorch: leafhopper and spittlebug vectors. *J Econ Entomol* 73: 834–
1096 838.
- 1097 Purcell AH (1981) Vector preference and inoculation efficiency as components of resistance to Pierce's
1098 disease in European grape cultivars. *Phytopathology* 71: 429–435.
- 1099 Purcell AH (1997) *Xylella fastidiosa*, a regional problem or global threat? *J Plant Pathol* 99–105.
- 1100 Purcell AH, Finlay AH, McLean DL (1979) Pierce's disease bacterium: Mechanism of transmission by
1101 leafhopper vectors. *Science* 206: 839–841.
- 1102 Purcell AH, Gravena S, Donadio LC (1994) Sharpshooter in citrus crops. Citrus-integrated
1103 management of insect and mite pests. *Bebedouro, Estação Experimental de Citricultura* 213–
1104 229.
- 1105 Puterka GJ, Glenn DM, Sekutowski DG, Unruh TR, Jones SK (2003) Particle film, Surround WP,
1106 effects on glassy-winged sharpshooter behavior and its utility as a barrier to sharpshooter
1107 infestation in grapes. *Plant Health Prog.* doi: 10.1094/PHP-2003-0321-RS.
- 1108 Putman WL (1953) Notes on the Bionomics of some Ontario Cercopids (Homoptera). *Can Entomol* 85:
1109 244-248.
- 1110 Quartau JA, Borges PA (1997). On the colour polymorphism of *Philaenus spumarius* (L.)(Homoptera,
1111 Cercopidae) in Portugal. *Miscellanea Zoologica* 20: 19-30.
- 1112 Ranieri E, Ruschioni S, Riolo P, Isidoro N, Romani R (2016) Fine structure of antennal sensilla of the
1113 spittlebug *Philaenus spumarius* L.(Insecta: Hemiptera: Aphrophoridae). I. Chemoreceptors and
1114 thermo-hygroreceptors. *Arthropod Struct Dev* 45: 432-439.
- 1115 Rashed A, Killiny N, Kwan J, Almeida RP (2011) Background matching behaviour and pathogen
1116 acquisition: plant site preference does not predict the bacterial acquisition efficiency of vectors.
1117 *Arthropod-Plant Inte* 5: 97–106.
- 1118 Robertson A, Gibbs AJ (1937) Spermatogenesis and fertilization in *Philaenus spumarius* Fallen. *J Trop*
1119 *Med Hyg* 40: 257–262.
- 1120 Rodrigues AS, Silva SE, Marabuto E, Silva DN, Wilson MR, Thompson V, et al (2014) New
1121 mitochondrial and nuclear evidences support recent demographic expansion and an atypical
1122 phylogeographic pattern in the spittlebug *Philaenus spumarius* (Hemiptera,
1123 Aphrophoridae). *PloS one* 9.6: e98375.
- 1124 Salerno M, Russo V, Sefa V, Lamaj F, Basher N. Verrastro V. Porcelli F (2017) *Zelus renardii* an
1125 assassin bug candidate for *Philaenus spumarius* biocontrol. European Conference on *Xylella*.
1126 Finding answer to a global problem. Palma de Mallorca, 13-15 November 2017, 22-23.
- 1127 Sanderlin RS, Melanson RA (2010) Insect transmission of *Xylella fastidiosa* to pecan. *Plant Dis* 94:
1128 465–470.
- 1129 Saponari M, Boscia D, Altamura G, D'Attoma G, Cavalieri V, Zicca S, Morelli M, Tavano D,
1130 Loconsole G, Susca L (2016) External Scientific Report.
- 1131 Saponari M, Boscia D, Nigro F, Martelli GP (2013) Identification of DNA sequences related to *Xylella*
1132 *fastidiosa* in oleander, almond and olive trees exhibiting leaf scorch symptoms in Apulia
1133 (southern Italy). *J Plant Pathol* 95.
- 1134 Saponari M, Loconsole G, Cornara D, Yokomi RK, De Stradis A, Boscia D, Bosco D, Martelli GP,
1135 Krugner R, Porcelli F (2014) Infectivity and transmission of *Xylella fastidiosa* by *Philaenus*
1136 *spumarius* (Hemiptera: Aphrophoridae) in Apulia, Italy. *J Econ Entomol* 107: 1316–1319.
- 1137 Scholl JM, Medler JT (1947) Spittle Bugs in relation to Alfalfa Seed Production in Wisconsin. *J Econ*
1138 *Entomol* 40: 446-448.
- 1139 Schulz CA, Meijer J (1978) Migration of leafhoppers (Homoptera: Auchenorrhyncha) into a new
1140 polder. *Ecography* 1: 73–78.
- 1141 Severin H (1950) Spittle-insect vectors of Pierce's disease virus. II. Life history and virus transmission.

- 1142 Hilgardia 19: 357–376.
- 1143 Sinclair WA, Griffiths HM (1994) Ash yellows and its relationship to dieback and decline of ash. Annu
1144 Rev Phytopathol 32: 49-60.
- 1145 Smith RL (1984) Human sperm competition. Sperm competition and the evolution of animal mating
1146 systems 602-652.
- 1147 Stewart AJ, Lees DR (1996) The colour/pattern polymorphism of *Philaenus spumarius* (L.)
1148 (Homoptera: Cercopidae) in England and Wales. Philos T Roy Soc B 351: 69–89.
- 1149 Stewart AJ, Lees DR (1988) Genetic control of colour/pattern polymorphism in British populations of
1150 the spittlebug *Philaenus spumarius* (L.)(Homoptera: Aphrophoridae). Biol J Linn Soc 34: 57–
1151 79.
- 1152 Strona G., Carstens CJ, Beck PS (2017) Network analysis reveals why *Xylella fastidiosa* will persist in
1153 Europe. Scientific Reports 7: 71.
- 1154 Svanberg I (2016) Cuckoo Spit in Northern European Folk Biology. SLA.
- 1155 Thompson V (1973) Spittlebug polymorphic for warning coloration. Nature 242: 126–128.
- 1156 Thompson V (1994) Spittlebug indicators of nitrogen-fixing plants. Ecol Entomol 19: 391–398.
- 1157 Thompson V (2004) Associative nitrogen fixation, C4 photosynthesis, and the evolution of spittlebugs
1158 (Hemiptera: Cercopidae) as major pests of neotropical sugarcane and forage grasses. B Entomol
1159 Res 94: 189–200.
- 1160 Tishechkin D. Yu (2013) Two new species of the genus *Philaenus* (Homoptera, Aphrophoridae) from
1161 Iran. Entomol Review 1: 73-76.
- 1162 Tubajika KM, Civerolo EL, Puterka GJ, Hashim JM, Luvisi DA (2007) The effects of kaolin, harpin,
1163 and imidacloprid on development of Pierce’s disease in grape. Crop Prot 26: 92–99.
- 1164 Van Driesche RG, Prokopy RJ, Coli WM (1987) Potential for increased use of biological control
1165 agents in Massachusetts apple orchards. Research Bulletin, Massachusetts Agricultural
1166 Experiment Station 6–21.
- 1167 Walker GP (2000) A beginner’s guide to electronic monitoring of homopteran probing behavior.
1168 Principles and applications of electronic monitoring and other techniques in the study of
1169 homopteran feeding behavior. Thomas Say Publications in Entomology, Entomological Society
1170 of America, Lanham, MD 14–40.
- 1171 Waloff N (1973) Dispersal by Flight of Leafhoppers (Auchenorrhyncha: Homoptera). J Appl Ecol 10:
1172 705-730.
- 1173 Watson R, Pritchard J, Malone M (2001) Direct measurement of sodium and potassium in the
1174 transpiration stream of salt-excluding and non-excluding varieties of wheat. J Exp Bot 52:
1175 1873–1881.
- 1176 Weaver CR (1951) The seasonal behaviour of meadow spittlebug and its relation to a control method. J
1177 Econ Entomol 44: 350-3.
- 1178 Weaver CR, King DR (1954) Meadow spittlebug, *Philaenus leucophthalmus* (L.). Ohio Agric Exp Stat
1179 Bull 741: 1-99.
- 1180 West J, Lees DR (1988) Temperature and egg development in the spittlebug *Philaenus spumarius*
1181 (L.)(Homoptera: Aphrophoridae). The Entomologist 13: 46-51.
- 1182 Whittaker JB (1969) The biology of Pipunculidae (Diptera) parasitising some British Cercopidae
1183 (Homoptera). Physiol Entomol 44: 17–24.
- 1184 Whittaker JB (1970) Cercopid spittle as a microhabitat. Oikos 59–64.
- 1185 Whittaker JB (1973) Density regulation in a population of *Philaenus spumarius* (L.)(Homoptera:
1186 Cercopidae). J Anim Ecol 163–172.
- 1187 Wiegert RG (1964) Population energetics of meadow spittlebugs (*Philaenus spumarius* L.) as affected
1188 by migration and habitat. Ecol Monogr 34: 217–241.
- 1189 Wilson MC, Shade RE (1967) Relative attractiveness of various luminescent colors to the cereal leaf

- 1190 beetle and the meadow spittle bug. J Econ Entomol 60: 578-580.
- 1191 Wise MJ, Kieffer DL, Abrahamson WG (2006) Costs and benefits of gregarious feeding in the meadow
1192 spittlebug, *Philaenus spumarius*. Ecol Entomol 31: 548–555.
- 1193 Yurtsever S (2000) On the polymorphic meadow spittlebug, *Philaenus spumarius* (L.)(Homoptera:
1194 Cercopidae). Turk J Zool 24: 447–460.
- 1195 Zajac MA, Hall FR, Wilson MC (1989). Heat unit model for the development of meadow spittlebug
1196 (Homoptera Cercopidae) on strawberry. Environ Entomol 18: 347-350.
- 1197 Zajac MA, Wilson MC (1984) The effects of nymphal feeding by the meadow spittlebug, *Philaenus*
1198 *spumarius* (L.) on strawberry yield and quality. Crop Prot 3: 167–175.
- 1199

1200 **Figure 1:** Biological cycle of *Philaenus spumarius* in Southern Apulia Region
1201 of Italy (photos by A. Fereres and D. Cornara).

1202

1203 **Figure 2:** *Philaenus spumarius* abundance on olive and ground cover during
1204 the year, and hosts shifting in infected olive orchards in Apulia (South Italy).
1205 Black line refers to *P. spumarius* adults abundance on olive plants, gray line
1206 refers to ground cover. Figure elaborated from Cornara et al. (2016b).

1207
1208



EGGS: October-February




ADULTS: April-December



NYMPHS: March-April





Philaenus spumarius hosts shifting
in infected olive orchards in Apulia

