AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Timing of food intake: sounding the alarm about metabolic impairments? A systematic review.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1671160> since 2018-07-22T11:03:31Z

Published version:

DOI:10.1016/j.phrs.2017.09.005

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Timing of food intake: sounding the alarm about metabolic impairments? A systematic review

Beccuti Guglielmo¹, Monagheddu Chiara², Evangelista Andrea², Ciccone Giovannino², Broglio Fabio¹, Soldati Laura³, Bo Simona¹

¹ Department of Medical Sciences, University of Turin, Turin, Italy

² Unit of Clinical Epidemiology, CPO, “Città della Salute e della Scienza” Hospital of Turin, Turin, Italy

³ Department of Health Sciences, University of Milan, Milan, Italy

Corresponding author:

Simona Bo, Department of Medical Sciences, University of Turin, Corso Dogliotti 14, 10126 Turin, Italy;

Telephone +39 (0)11 6336036 Fax +39(0)11 6335401

E-mail: simona.bo@unito.it

Keywords:

Diabetes mellitus, Metabolism, Obesity, Timing of meals

Words count: abstract 203, text 3585

Abstract

Growing evidence points to an association between timing of food intake and obesity in humans, raising the question if when to eat matters as much as what and how much to eat. Based on the new definition of obesity as a chronobiological disease, an unusual or late meal timing represent a circadian chronodisruption, leading to metabolic impairments.

Preliminary data from cross-sectional and experimental studies suggest that changes in meal timing can influence obesity and success of weight loss therapy, independently from total energy intake, dietary composition and estimated energy expenditure.

A systematic review of observational and experimental studies in humans was conducted to explore the link between time of food ingestion, obesity and metabolic alterations. Results confirm that eating time is relevant for obesity and metabolism: observational and experimental studies found an association between meal timing, weight gain, hyperglycemia and diabetes mellitus with benefits deriving from an early intake of food in the day in a wide range of individuals. Herein clinical, future perspectives of chronoprevention and chronotherapy of obesity and type 2 diabetes are also provided.

In conclusion, meal timing appears as a new potential target in weight control strategies, and therapeutic strategies should consider this contributor in the prevention of obesity.

1. Introduction

In the last decade, a new relevant question has arisen: when to eat [1-3]. In addition to what and how much to eat, food timing represents a novel issue in our 24-h modern society, characterized by more exposure to artificial light, later food intake and bedtimes. Food is a major synchronizer of peripheral circadian clocks, and delayed feeding due to prolonged night-time wakefulness leads to desynchrony between central circadian and peripheral clocks [4].

51 Growing evidence points to an association between timing of food intake and obesity in humans, suggesting
52 that changes in meal timing can influence obesity and success of weight loss therapy [1]. Also in animals,
53 weight regulation is affected by the timing of food ingestion [5, 6].

54 On this basis, obesity could now represent a “chronobiological disease” [7]. Differently from the time-
55 restricted feeding pattern unintentionally practiced by our ancestors for thousands of years, the current trend
56 is to shift most of the caloric intake later in the day [8]. In a few cross-sectional studies, an increased risk of
57 overweight and obesity was found when a greater daily caloric intake was consumed in the evening [9-12],
58 while a reduced risk was observed when consuming a larger proportion of calories at lunch or breakfast [9,
59 11, 13].

60 Even though the association between evening eating and body weight was not confirmed in a prospective US
61 cohort, it was present in specific subgroups (smoking men, physically active men, inactive women) [14].

62 Another prospective study showed that late-night eaters had an increased coronary heart disease risk [15].

63 The clinical relevance of meal timing appears to be supported by its role in weight loss strategies. In a 20-week
64 intervention study, as compared with early lunch eaters, late lunch eaters lost less weight independently from
65 self-reported 24-h caloric intakes [16]. In overweight and obese women with metabolic syndrome, a 12-week
66 weight-loss program with high caloric breakfast was more effective in reducing weight and waist
67 circumference than an isocaloric diet with high caloric intake at dinner [17].

68 Aside from body composition and weight regulation, timing of food intake seems to have a negative impact
69 also on metabolism. Eating lunch later in the day was associated with poorer insulin sensitivity assessed by
70 HOMA-IR (Homeostasis-Model Assessment-Insulin Resistance) index [16]. Experimental studies showed a
71 higher decrease in HOMA-IR after a high caloric breakfast vs dinner in women with metabolic syndrome [17]
72 and polycystic ovary syndrome [18]. Late lunch eating was associated with decreased pre-meal resting energy
73 expenditure, lower pre-meal carbohydrate utilization, and decreased glucose tolerance after mixed-meal test
74 [19]. In another study exploring food-induced thermogenesis in the morning and evening, the same meal
75 consumed in the evening determined a lower after-meal resting metabolic rate and increased, delayed
76 concentrations of glucose and insulin [20].

These preliminary data suggest that consuming a larger proportion of total daily energy in the morning, as opposed to later in the day, might be more beneficial for weight loss.

The aim of this study was to perform a systematic review of observational and experimental studies comparing the effect of different food timing on body weight and metabolic outcomes in adults. The possibility to undertake a meta-analysis of the effects of the interventions on at least some of the outcomes was evaluated too.

2. Material and Methods

This article is structured according to the preferred items for Systematic Reviews and Meta-Analyses (PRISMA) guideline [21].

2.1 Literature search strategy

The following electronic databases were queried using a combination of search terms: PubMed (National Library of Medicine), Trip database and The Cochrane Library, until 01 March 2017. The construction of the search strategy was performed using database specific subject headings and keywords. The search terms included combinations of “timing meal” or “timing meals” or “timing of food” or “food timing”, and Body Mass Index (BMI), obesity, weight, hyperglycemia, glycemia, insulin, insulin-resistance and type 2 diabetes mellitus (free-term and MESH as possible) (**Appendix 1**).

These search strategies were supplemented by hand searching the bibliographies of all the included studies. Searches were limited to randomized controlled trials, parallel or cross-over, and observational studies in healthy volunteers or patients (e.g. individuals with obesity/overweight, polycystic ovary syndrome, metabolic diseases or other underlying diseases). We excluded studies performed in children.

2.2 Study selection

We included studies reporting comparisons of different timing meal interventions or habits (early eaters/late eaters or different timing of daily energy intake distribution) to reduce weight, insulinemic and glycemic areas-under-the-curve values and other metabolic variables.

Two review authors (SB, CM) independently scanned the abstract, title, or both, of every record retrieved, to determine which studies should be assessed further. All potentially relevant articles were investigated as full text. Any discrepancy about inclusion was resolved by discussing with a third review author (GB).

2.3 Data collection and extraction

For the trials that fulfilled the inclusion criteria, two authors independently abstracted key participant characteristics and reported data on efficacy outcomes using standard data extraction templates.

From each included study, information was extracted on:

- Characteristics of study participants (type of population, age, BMI);
- Type of intervention;
- Outcomes:
 - Anthropometric variables (BMI, weight, waist circumference, total body fat, etc.);
 - Metabolic variables (blood glucose values, triglycerides, total cholesterol, HDL and LDL-cholesterol, etc.);
 - Hormonal variables (blood insulin, progesterone, testosterone, etc.);
 - Calorimetric variables (fasting or after-meal resting metabolic rate, fasting or after meal respiratory quotients, etc).

2.4 Risk of bias assessment

The validity of each study was independently assessed by two authors (SB, GB) using two tools: a) the 'Risk of bias' tool developed by The Cochrane Collaboration for RCT [22], and b) the 'Risk Of Bias in Non-randomized Studies of Interventions' (ROBINS-I) tool for evaluating risk of bias in estimates of the comparative effectiveness of interventions from studies not using randomization to allocate units (individuals or clusters of

individuals) to comparison groups [23]. As they were familiar with the literature, review authors were not blinded with respect to the study authors, institution or journal. We resolved possible disagreements by consensus, or with consultation with a third review author (AE).

We could not undertake a meta-analysis of the effects of the interventions due to the great variability in outcome assessment and reporting, and in the type of interventions.

3. Results

3.1 Flow and characteristics of included studies

With the initial literature search, 926 articles were found (**Figure 1**). Fifteen records were identified and carefully assessed for eligibility, after excluding non-original articles, duplicates, and articles not meeting the inclusion criteria. Only 10 studies satisfied all the inclusion criteria and were selected for the systematic review, including a total number of 6401 subjects (**Table 1**). The largest study recruited 4243 subjects [24] while the smallest one only 6 subjects [25]. Included studies were conducted in Spain [16, 19, 24, 26], Israel [17, 18], Japan [27], UK [25], and Italy [20, 28], between 2001-2014. Participants were: healthy individuals [19, 20, 25, 27], general population [24, 28], overweight/obese subjects [16, 17], post-bariatric surgery patients [26], women with polycystic ovary syndrome [18]. In one case participants were paid [27]. The demographic and clinical characteristics of the included studies are shown in Table 1.

Five of the included studies were trials: randomized cross-over [19, 20, 25, 27] and randomized controlled trials [17, 18], respectively, while four were observational prospective studies [16, 24, 26, 28]. The duration of the observational period was respectively: 6 years [26, 28], 3.5 years [24], and 20 weeks [16].

The interventions of the trials varied from the acute consumption of one or more meals a day at different hours [20, 25, 27] to early eating vs late eating the greater amount of kcal/day for 2 weeks [19] or 12 weeks [17, 18].

Observational studies divided participants according to the timing of the main meal (lunch before or after 15:00) [16, 26], the timing of the consumption of the larger amount of calories [24] or the tertiles of the percentage of total daily caloric intake from dinner [28].

The following outcomes were evaluated: variation on anthropometric variables [16-19, 24, 28], energy expenditure by indirect calorimetry [19, 20, 27] or equations [16, 26], metabolic parameters [16-20, 25-28], sleep pattern [16, 19, 26], body temperature [19], carbohydrate absorption [27], satiety [17], hormonal assessments [18, 19], and other blood variables, such as inflammatory parameters and liver enzymes [28].

3.2 Risk of bias assessment

Most of the analyzed trials provided insufficient information about randomization procedures (**Table 2**). If blinding of participants was not feasible owing to the nature of the interventions, data about blinding of the personnel who performed the laboratory or statistical analyses was often unknown. In one study about 20% of participants dropped out [17]. Most trials appeared to be free of selective outcome reporting and of other sources of bias.

The risk of bias for the observational studies is reported in **Table 3**. Most of the evaluated risks of bias were low/moderate. Ruiz-Lozano classified post-bariatric surgery patients by their weight-loss pattern after surgery and compared the timing of meals among groups [26]. The three groups, however, significantly differed for age and gender: older male patients were more frequently poor responders. Hermengildo studied the risk of weight gain by the distribution of energy intake throughout the day, but weight gain was self-reported both at baseline and at the end of the follow-up [24].

3.3 Effect of timing of food intake on changes in weight and other anthropometric parameters

Observational studies showed that late lunch-eaters (after 15:00) were 2-fold more frequent in poor-weight loss responders to bariatric surgery, independent of dietary macronutrient composition [26]; the OR of gaining weight (>3kg) was 0.79 (95% CI 0.63-0.99), 0.82 (95% CI 0.64-1.04) and 0.62 (95% CI 0.47-0.80) respectively in the second, third and the highest quartile of percent energy intake at lunch, when compared to the lowest quartile, in a multivariate logistic regression analysis (p for trend=0.001) [24]; being in the highest tertile of daily percent caloric intake at dinner was significantly associated with an increased risk of incident obesity

(OR=2.33, 95%CI 1.17-4.65) [28]; late lunch eaters lost less weight than early lunch eaters (7.7 vs 9.9kg) after a 20-week weight-loss intervention [16].

Apart from two observational studies [24, 28], the nutritional composition of meals was not different between the groups of early or later-eaters. This finding strongly reinforces the role of meal timing on the studied outcomes.

The evaluated randomized trials reported: a significantly higher weight loss in the “more calories at breakfast” group when compared to the “more calories at dinner” group (-8.7 ± 1.4 vs -3.6 ± 1.5 kg, $p < 0.001$) after 12-weeks of a hypocaloric diet in overweight/obese women [17]; no significant change in weight between two groups with the same distribution of calories as above reported, after 12 weeks of a maintenance diet in women with the polycystic ovary syndrome.

A few studies evaluated other indices of body fat [16-18]. No difference in waist circumference values and total body fat, as measured by bioelectrical impedance, were evident at baseline among early vs late lunch eaters, but these data were not available at follow-up [16]. Individual eating “more calories at breakfast” showed greater waist circumference reduction compared to the “more calories at dinner” group after a weight-loss 1400 kcal diet [17]; this difference was not confirmed after a 1800-kcal maintenance diet in women with polycystic ovary syndrome [18]. Only in 3 studies, waist circumference was described with the same methods and at the same time (baseline, follow-up); therefore, we could not undertake a meta-analysis because of the low number of individuals, not representative of the complete review [16-18].

3.4 Effects of timing of food intake on glucose and insulin blood values

At baseline, late lunch eaters when compared to early lunch eaters showed increased values of Homeostasis Model Assessment-Insulin Resistance (HOMA-IR) index, but fasting glucose and insulin blood concentrations were similar between the two groups; no data at follow-up were available [16]. Individuals in the highest tertile of percent daily caloric intake at dinner showed an increase risk of incident type 2 diabetes (2.26; 95%CI 0.89-5.75) [28].

The trials evaluating the acute consumption of one or more meals/day reported that: a) glucose responses were greater after consuming the majority of energy load in the evening than in the morning, while insulin responses and post-prandial insulin resistance seemed to be mainly affected by the quality of carbohydrates [25]; b) the efficiency of digestion and absorption of dietary carbohydrates consumed at breakfast was higher if the previous supper was later (at 23:00) than under usual conditions (at 18:00) and, accordingly, after-breakfast glucose values were increased until 3 hours in case of previous late suppertime [27]; c) the same meal consumed in the evening determined delayed and larger increases in glucose and insulin blood concentrations and significant increases in the corresponding areas-under-the curve [20].

The study protocol 1 of Bandin showed increased post-prandial glucose responses in late lunch eaters with a 46% higher glucose area-under-the curve than in early lunch eaters [19]; women eating “more calories at breakfast” showed greater reduction in fasting glucose, insulin and insulin resistance evaluated by the HOMA-IR index and, similarly, reduced glycemic and insulinemic responses both to the oral glucose tolerance test and to a meal challenge when compared to the “more calories at dinner”, after 12 weeks of isocaloric 1400-kcal diet [17]; in lean women with polycystic ovary syndrome, a high caloric intake at breakfast resulted in significantly reduced glucose and insulin areas-under-the-curve and an improvement in insulin sensitivity than consuming a high caloric intake at dinner [18].

4. Discussion

New, intriguing contributors to the epidemic of obesity have been lately recognized, such as meal frequency and patterns [29, 30], as well as sleep duration and quality [31]. Emerging evidence sounds the alarm on the role of meal timing and questions whether when to eat matters as much as what and how much to eat.

Few cross-sectional studies tried to answer this question, finding that later timing of meals or eating more calories later in the day has a negative impact on body weight and metabolism [9-13].

Our systematic review of observational and experimental studies, including both healthy individuals [19, 20, 24, 25, 27, 28] and patients with different dysmetabolic conditions [16-18, 26] confirms the health benefits of early eating, with positive effects on body weight, weight loss success, and glucose metabolism.

230

231 *4.1 Timing of food intake and obesity*

232 Experimental studies show that animal models fed at unusual feeding time develop obesity, even without
 233 change in activity or total energy intake [5, 32]. High-fat meal at the end of the active phase leads to increased
 234 weight gain [33]. When challenged with a high-fat diet, chronodisrupted mice were more likely to be obese
 235 [34].

236 The pathophysiologic basis of these findings relies on the new definition of obesity as a chronobiological
 237 disease [7]. Unusual feeding time can represent a circadian disruption leading to clock gene functional
 238 alterations and uncoupling between the central and peripheral oscillators, circadian variations of peripheral
 239 clocks, gene expression, satiety hormones, and digestive processes [1]. Among mechanisms promoting
 240 obesity, diet-induced thermogenesis is lower at night [35], and the reduced thermic effect of glucose in obesity
 241 is likely related to the nocturnal insulin resistance [36]. Additionally, reduced fat oxidation has been observed
 242 during nighttime eating [37, 38].

243 Also in humans, cross-sectional studies suggest that eating time is relevant for obesity. Particularly, consuming
 244 a greater daily caloric in the evening is associated with higher risk of overweight and obesity [9-12], while
 245 eating more calories at lunch or breakfast appears to be protective against overweight/obesity [9, 11, 13].

246 The observational studies included in the present systematic review showed a positive association between
 247 meal timing and body weight [16, 24, 26, 28], that remained significant also after controlling for many
 248 confounding factors involved in the obesity development, such as physical activity and sleep time [16, 24, 26,
 249 28]. Even though short sleep duration is a well know, independent risk factor for obesity, self-reported data
 250 on sleep time appear similar among the different weight loss patterns [16, 26] and therefore probably did not
 251 mediate the observed outcomes. The association between meal timing and body weight was supported also
 252 by a causal direction described in the included experimental studies [17-20, 25, 27]. Furthermore, the benefits
 253 were evident in a wide range of individuals: post-bariatric surgery patients [26], women with metabolic
 254 syndrome [17], overweight/obese subjects attending nutrition clinics [16, 26] and general population [24, 28].
 255 Specifically, in post-bariatric surgery patients, left ventricular mass was decreased one year after procedure;

this improvement correlated only with the decrease in leptin levels [39], postulating a cardiovascular protection from weight loss also mediated by hormonal changes.

This key message has a clear practice implication, and should be considered by clinicians when drawing up a nutritional scheme.

4.2 Timing of food intake and hyperglycemia

Circadian misalignment is known to result in adverse metabolic and cardiovascular consequences [40, 41]. Experimental studies explored the possible mechanisms supporting the circadian modulation of insulin secretion or action. Pathophysiological hypotheses of decreased insulin sensitivity later in the day [42] are represented by increased levels of triglycerides [43] and urinary epinephrine [44], fluctuation in cortisol serum concentrations [45] and higher morning ACTH plasma values [46], and/or a delayed peak in the counteracting activity of glucagon after evening meals [47]. Moreover, under late suppertime conditions, an increased efficacy of dietary carbohydrates absorption has been described [27]. Yet, increased evening meal emptying time seems to lead to evening delay in reaching peak plasma concentrations of the absorbed substances [48]. Metabolic consequences of experimental interventions occur rapidly and are already observed after an acute consumption of one or more meals a day at different hours [20, 25, 27].

4.3 Clinical prospective

The success of weight loss therapy seems to be predicted by food timing; evening preference has a negative impact on metabolism, too. Even though not specifically design for food timing investigation, later chronotype individuals with type 2 diabetes, more likely characterized by later food ingestion, were characterized by a poorer glycemic control [49-52]. This observation raises the question whether meal timing intervention, with or without circadian phase changes, might be helpful in type 2 diabetes management; future studies are needed to verify this hypothesis. Indeed, The Academy of Nutrition and Dietetics has recently pointed up meal timing as a new potential target in weight control strategies [53], stating that consuming most of an individual's energy earlier in the day may enhance weight loss and weight maintenance.

Among other clinical aspects of chronobiology in type 2 diabetes, it is worth considering that chronotherapy might apply not only to lifestyle but also to drug treatment. A prospective, randomized, open-label, blinded trial showed that blood pressure lowering drugs at bedtime reduced cardiovascular risk in type 2 diabetes patients with hypertension over a mean of 5.4 years, compared with the ingestion of drugs upon awakening [54]. Varying the time of day at which antihypertensive medications are taken is highly effective not only in diabetic but also non-diabetic subjects [55]. Like chronotherapy, also chronoprevention might apply to both lifestyle and drug treatment. In fact, in hypertensive patients without diabetes, administering ≥ 1 antihypertensive medications at bedtime, particularly angiotensin receptor blockers and ACE inhibitors, compared with medications taken after awakening, reduced risk of incident diabetes during a 5.9-year median follow-up and improved blood pressure control with significant decrease of asleep blood pressure [56].

Another clinical application of food timing intervention might be represented by type 1 diabetes, even though it was not mentioned in the studies included in our systematic review. As type 1 diabetes is affected by increased mortality [57], it would be interesting to see whether optimal food timing and daily caloric distribution may improve short-term glycemic, endothelial dysfunction, inflammation and oxidative stress outcomes as cardiovascular risk markers.

In consideration of the relevance of obesity- and type 2 diabetes- cardiovascular related diseases, it looks fundamental to search for efficient strategies for weight-loss and cardiovascular risk reduction. Future studies should verify whether well-known cardiovascular risk markers associated with obesity [39] and diabetes [58] may improve after chronotherapy intervention.

4.4 Limitations

The heterogeneity of the population studies and the evaluated outcomes has prevented us from performing a meta-analysis. The findings of the present systematic review do not allow to definitely prove the relationship between meal timing and the improvement of overweight and dysmetabolic conditions in humans. The heterogeneity of the included studies should be considered as a limitation, since either healthy individuals or patients with different dysmetabolic conditions have been enrolled. As another limitation, some studies

included in this analysis were not primarily designed to assess the effects of meal timing on weight or metabolic variables, suggesting a high risk of both publication and outcome reporting biases. The use of any drug was considered as exclusion criteria in most studies [16-20,27], but in 3 of the observational studies [24,26,28], data relative to pharmacological treatment were not reported; we therefore could not exclude that therapeutic regimens might have influenced weight loss dynamics or food intake timing in these studies. Finally, the number of trials and individuals included in the present review was small, which made it difficult to definitively assess the metabolic effect of meal timing, and required further investigations. Nevertheless, to the best of our knowledge, this is the first systematic review on this topic and could contribute to advancing knowledge and generating new studies in the field

4.5 Conclusions

Accumulating evidence summarized in this systematic review supports the negative impact of later meal timing and calories distribution on body weight and metabolism. High quality studies are needed to clarify the effectiveness of changes in eating time as an additional strategy for obesity and diabetes prevention and treatment in adults.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

FIGURE LEGEND

Figure 1. Flow chart of the number of studies identified and included in the systematic review.

5. References

- [1] Garaulet M, Gomez-Abellan P. Timing of food intake and obesity: a novel association. *Physiol Behav.* 2014;134:44-50. doi: 10.1016/j.physbeh.2014.01.001. PubMed PMID: 24467926.
- [2] McCrory MA, Shaw AC, Lee JA. Energy and Nutrient Timing for Weight Control: Does Timing of Ingestion Matter? *Endocrinol Metab Clin North Am.* 2016;45(3):689-718. doi: 10.1016/j.ecl.2016.04.017. PubMed PMID: 27519140.
- [3] Varady KA. Meal frequency and timing: impact on metabolic disease risk. *Curr Opin Endocrinol Diabetes Obes.* 2016;23(5):379-83. doi: 10.1097/MED.0000000000000280. PubMed PMID: 27455514.
- [4] Bass J, Takahashi JS. Circadian integration of metabolism and energetics. *Science.* 2010;330(6009):1349-54. doi: 10.1126/science.1195027. PubMed PMID: 21127246; PubMed Central PMCID: PMC3756146.
- [5] Arble DM, Bass J, Laposky AD, Vitaterna MH, Turek FW. Circadian timing of food intake contributes to weight gain. *Obesity (Silver Spring).* 2009;17(11):2100-2. doi: 10.1038/oby.2009.264. PubMed PMID: 19730426; PubMed Central PMCID: PMC3499064.
- [6] Sherman H, Genzer Y, Cohen R, Chapnik N, Madar Z, Froy O. Timed high-fat diet resets circadian metabolism and prevents obesity. *FASEB J.* 2012;26(8):3493-502. doi: 10.1096/fj.12-208868. PubMed PMID: 22593546.
- [7] Garaulet M, Ordovas JM, Madrid JA. The chronobiology, etiology and pathophysiology of obesity. *Int J Obes (Lond).* 2010;34(12):1667-83. doi: 10.1038/ijo.2010.118. PubMed PMID: 20567242; PubMed Central PMCID: PMC34428912.
- [8] Almoosawi S, Winter J, Prynn CJ, Hardy R, Stephen AM. Daily profiles of energy and nutrient intakes: are eating profiles changing over time? *Eur J Clin Nutr.* 2012;66(6):678-86. doi: 10.1038/ejcn.2011.210. PubMed PMID: 22190135; PubMed Central PMCID: PMC3389619.
- [9] Wang JB, Patterson RE, Ang A, Emond JA, Shetty N, Arab L. Timing of energy intake during the day is associated with the risk of obesity in adults. *J Hum Nutr Diet.* 2014;27 Suppl 2:255-62. doi: 10.1111/jhn.12141. PubMed PMID: 23808897.

- [10] Berteus Forslund H, Lindroos AK, Sjostrom L, Lissner L. Meal patterns and obesity in Swedish women- a simple instrument describing usual meal types, frequency and temporal distribution. *Eur J Clin Nutr.* 2002;56(8):740-7. doi: 10.1038/sj.ejcn.1601387. PubMed PMID: 12122550.
- [11] Berg C, Lappas G, Wolk A, Strandhagen E, Toren K, Rosengren A, et al. Eating patterns and portion size associated with obesity in a Swedish population. *Appetite.* 2009;52(1):21-6. doi: 10.1016/j.appet.2008.07.008. PubMed PMID: 18694791.
- [12] Baron KG, Reid KJ, Kern AS, Zee PC. Role of sleep timing in caloric intake and BMI. *Obesity (Silver Spring).* 2011;19(7):1374-81. doi: 10.1038/oby.2011.100. PubMed PMID: 21527892.
- [13] Summerbell CD, Moody RC, Shanks J, Stock MJ, Geissler C. Relationship between feeding pattern and body mass index in 220 free-living people in four age groups. *Eur J Clin Nutr.* 1996;50(8):513-9. PubMed PMID: 8863011.
- [14] Kant AK, Schatzkin A, Ballard-Barbash R. Evening eating and subsequent long-term weight change in a national cohort. *Int J Obes Relat Metab Disord.* 1997;21(5):407-12. PubMed PMID: 9152744.
- [15] Cahill LE, Chiuve SE, Mekary RA, Jensen MK, Flint AJ, Hu FB, et al. Prospective study of breakfast eating and incident coronary heart disease in a cohort of male US health professionals. *Circulation.* 2013;128(4):337-43. doi: 10.1161/CIRCULATIONAHA.113.001474. PubMed PMID: 23877060; PubMed Central PMCID: PMC3797523.
- [16] Garaulet M, Gomez-Abellan P, Alburquerque-Bejar JJ, Lee YC, Ordovas JM, Scheer FA. Timing of food intake predicts weight loss effectiveness. *Int J Obes (Lond).* 2013;37(4):604-11. doi: 10.1038/ijo.2012.229. PubMed PMID: 23357955; PubMed Central PMCID: PMC3756673.
- [17] Jakubowicz D, Barnea M, Wainstein J, Froy O. High caloric intake at breakfast vs. dinner differentially influences weight loss of overweight and obese women. *Obesity (Silver Spring).* 2013;21(12):2504-12. doi: 10.1002/oby.20460. PubMed PMID: 23512957.
- [18] Jakubowicz D, Barnea M, Wainstein J, Froy O. Effects of caloric intake timing on insulin resistance and hyperandrogenism in lean women with polycystic ovary syndrome. *Clin Sci (Lond).* 2013;125(9):423-32. doi: 10.1042/CS20130071. PubMed PMID: 23688334.

- [19] Bandin C, Scheer FA, Luque AJ, Avila-Gandia V, Zamora S, Madrid JA, et al. Meal timing affects glucose tolerance, substrate oxidation and circadian-related variables: A randomized, crossover trial. *Int J Obes (Lond)*. 2015;39(5):828-33. doi: 10.1038/ijo.2014.182. PubMed PMID: 25311083.
- [20] Bo S, Fadda M, Castiglione A, Ciccone G, De Francesco A, Fedele D, et al. Is the timing of caloric intake associated with variation in diet-induced thermogenesis and in the metabolic pattern? A randomized cross-over study. *Int J Obes (Lond)*. 2015;39(12):1689-95. doi: 10.1038/ijo.2015.138. PubMed PMID: 26219416.
- [21] Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi: 10.1371/journal.pmed.1000097. PubMed PMID: 19621072; PubMed Central PMCID: PMC2707599.
- [22] Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. doi: 10.1136/bmj.d5928. PubMed PMID: 22008217; PubMed Central PMCID: PMC3196245.
- [23] Sterne JA, Hernan MA, Reeves BC, Savovic J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. doi: 10.1136/bmj.i4919. PubMed PMID: 27733354; PubMed Central PMCID: PMC5062054.
- [24] Hermengildo Y, Lopez-Garcia E, Garcia-Esquinas E, Perez-Tasigchana RF, Rodriguez-Artalejo F, Guallar-Castillon P. Distribution of energy intake throughout the day and weight gain: a population-based cohort study in Spain. *Br J Nutr*. 2016;115(11):2003-10. doi: 10.1017/S0007114516000891. PubMed PMID: 27044416.
- [25] Morgan LM, Shi JW, Hampton SM, Frost G. Effect of meal timing and glycaemic index on glucose control and insulin secretion in healthy volunteers. *Br J Nutr*. 2012;108(7):1286-91. doi: 10.1017/S0007114511006507. PubMed PMID: 22176632.
- [26] Ruiz-Lozano T, Vidal J, de Hollanda A, Scheer FA, Garaulet M, Izquierdo-Pulido M. Timing of food intake is associated with weight loss evolution in severe obese patients after bariatric surgery. *Clin Nutr*. 2016;35(6):1308-14. doi: 10.1016/j.clnu.2016.02.007. PubMed PMID: 26948400.

- [27] Tsuchida Y, Hata S, Sone Y. Effects of a late supper on digestion and the absorption of dietary carbohydrates in the following morning. *J Physiol Anthropol*. 2013;32(1):9. doi: 10.1186/1880-6805-32-9. PubMed PMID: 23705984; PubMed Central PMCID: PMC3685573.
- [28] Bo S, Musso G, Beccuti G, Fadda M, Fedele D, Gambino R, et al. Consuming more of daily caloric intake at dinner predisposes to obesity. A 6-year population-based prospective cohort study. *PLoS One*. 2014;9(9):e108467. doi: 10.1371/journal.pone.0108467. PubMed PMID: 25250617; PubMed Central PMCID: PMC4177396.
- [29] Hutchison AT, Heilbronn LK. Metabolic impacts of altering meal frequency and timing - Does when we eat matter? *Biochimie*. 2016;124:187-97. doi: 10.1016/j.biochi.2015.07.025. PubMed PMID: 26226640.
- [30] Davis CS, Clarke RE, Coulter SN, Rounsefell KN, Walker RE, Rauch CE, et al. Intermittent energy restriction and weight loss: a systematic review. *Eur J Clin Nutr*. 2016;70(3):292-9. doi: 10.1038/ejcn.2015.195. PubMed PMID: 26603882.
- [31] Broussard JL, Van Cauter E. Disturbances of sleep and circadian rhythms: novel risk factors for obesity. *Curr Opin Endocrinol Diabetes Obes*. 2016;23(5):353-9. doi: 10.1097/MED.0000000000000276. PubMed PMID: 27584008; PubMed Central PMCID: PMC4507079.
- [32] Fonken LK, Workman JL, Walton JC, Weil ZM, Morris JS, Haim A, et al. Light at night increases body mass by shifting the time of food intake. *Proc Natl Acad Sci U S A*. 2010;107(43):18664-9. doi: 10.1073/pnas.1008734107. PubMed PMID: 20937863; PubMed Central PMCID: PMC2972983.
- [33] Bray MS, Tsai JY, Villegas-Montoya C, Boland BB, Blasier Z, Egbejimi O, et al. Time-of-day-dependent dietary fat consumption influences multiple cardiometabolic syndrome parameters in mice. *Int J Obes (Lond)*. 2010;34(11):1589-98. doi: 10.1038/ijo.2010.63. PubMed PMID: 20351731; PubMed Central PMCID: PMC3021134.
- [34] Shi SQ, Ansari TS, McGuinness OP, Wasserman DH, Johnson CH. Circadian disruption leads to insulin resistance and obesity. *Curr Biol*. 2013;23(5):372-81. doi: 10.1016/j.cub.2013.01.048. PubMed PMID: 23434278; PubMed Central PMCID: PMC3595381.
- [35] Romon M, Edme JL, Boulenguez C, Lescroart JL, Frimat P. Circadian variation of diet-induced thermogenesis. *Am J Clin Nutr*. 1993;57(4):476-80. PubMed PMID: 8460600.

- [36] Ravussin E, Acheson KJ, Vernet O, Danforth E, Jequier E. Evidence that insulin resistance is responsible for the decreased thermic effect of glucose in human obesity. *J Clin Invest.* 1985;76(3):1268-73. doi: 10.1172/JCI112083. PubMed PMID: 3900136; PubMed Central PMCID: PMCPMC424038.
- [37] Gluck ME, Venti CA, Salbe AD, Votruba SB, Krakoff J. Higher 24-h respiratory quotient and higher spontaneous physical activity in nighttime eaters. *Obesity (Silver Spring).* 2011;19(2):319-23. doi: 10.1038/oby.2010.206. PubMed PMID: 20864947.
- [38] Hibi M, Masumoto A, Naito Y, Kiuchi K, Yoshimoto Y, Matsumoto M, et al. Nighttime snacking reduces whole body fat oxidation and increases LDL cholesterol in healthy young women. *Am J Physiol Regul Integr Comp Physiol.* 2013;304(2):R94-R101. doi: 10.1152/ajpregu.00115.2012. PubMed PMID: 23174861.
- [39] Perego L, Pizzocri P, Corradi D, Maisano F, Paganelli M, Fiorina P, et al. Circulating leptin correlates with left ventricular mass in morbid (grade III) obesity before and after weight loss induced by bariatric surgery: a potential role for leptin in mediating human left ventricular hypertrophy. *J Clin Endocrinol Metab.* 2005;90(7):4087-93. doi: 10.1210/jc.2004-1963. PubMed PMID: 15855267.
- [40] Morris CJ, Purvis TE, Hu K, Scheer FA. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc Natl Acad Sci U S A.* 2016;113(10):E1402-11. doi: 10.1073/pnas.1516953113. PubMed PMID: 26858430; PubMed Central PMCID: PMCPMC4790999.
- [41] Morris CJ, Yang JN, Garcia JI, Myers S, Bozzi I, Wang W, et al. Endogenous circadian system and circadian misalignment impact glucose tolerance via separate mechanisms in humans. *Proc Natl Acad Sci U S A.* 2015;112(17):E2225-34. doi: 10.1073/pnas.1418955112. PubMed PMID: 25870289; PubMed Central PMCID: PMCPMC4418873.
- [42] Van Cauter E, Desir D, Decoster C, Fery F, Balasse EO. Nocturnal decrease in glucose tolerance during constant glucose infusion. *J Clin Endocrinol Metab.* 1989;69(3):604-11. doi: 10.1210/jcem-69-3-604. PubMed PMID: 2668321.
- [43] Al-Naimi S, Hampton SM, Richard P, Tzung C, Morgan LM. Postprandial metabolic profiles following meals and snacks eaten during simulated night and day shift work. *Chronobiol Int.* 2004;21(6):937-47. PubMed PMID: 15646240.

- [44] Lucassen EA, Zhao X, Rother KI, Mattingly MS, Courville AB, de Jonge L, et al. Evening chronotype is associated with changes in eating behavior, more sleep apnea, and increased stress hormones in short sleeping obese individuals. *PLoS One*. 2013;8(3):e56519. doi: 10.1371/journal.pone.0056519. PubMed PMID: 23483886; PubMed Central PMCID: PMC3590198.
- [45] Van Cauter E, Shapiro ET, Tillil H, Polonsky KS. Circadian modulation of glucose and insulin responses to meals: relationship to cortisol rhythm. *Am J Physiol*. 1992;262(4 Pt 1):E467-75. PubMed PMID: 1566835.
- [46] Van Cauter E, Refetoff S. Multifactorial control of the 24-hour secretory profiles of pituitary hormones. *J Endocrinol Invest*. 1985;8(4):381-91. doi: 10.1007/BF03348519. PubMed PMID: 2999212.
- [47] Goetz F, Bishop J, Halberg F, Sothorn RB, Brunning R, Senske B, et al. Timing of single daily meal influences relations among human circadian rhythms in urinary cyclic AMP and hemic glucagon, insulin and iron. *Experientia*. 1976;32(8):1081-4. PubMed PMID: 182528.
- [48] Goo RH, Moore JG, Greenberg E, Alazraki NP. Circadian variation in gastric emptying of meals in humans. *Gastroenterology*. 1987;93(3):515-8. PubMed PMID: 3609660.
- [49] Reutrakul S, Siwasaranond N, Nimitphong H, Saetung S, Chirakalwasan N, Ongphiphadhanakul B, et al. Relationships among sleep timing, sleep duration and glycemic control in Type 2 diabetes in Thailand. *Chronobiol Int*. 2015;32(10):1469-76. doi: 10.3109/07420528.2015.1105812. PubMed PMID: 26595483.
- [50] Reutrakul S, Hood MM, Crowley SJ, Morgan MK, Teodori M, Knutson KL, et al. Chronotype is independently associated with glycemic control in type 2 diabetes. *Diabetes Care*. 2013;36(9):2523-9. doi: 10.2337/dc12-2697. PubMed PMID: 23637357; PubMed Central PMCID: PMC3747872.
- [51] Iwasaki M, Hirose T, Mita T, Sato F, Ito C, Yamamoto R, et al. Morningness-eveningness questionnaire score correlates with glycated hemoglobin in middle-aged male workers with type 2 diabetes mellitus. *J Diabetes Investig*. 2013;4(4):376-81. doi: 10.1111/jdi.12047. PubMed PMID: 24843683; PubMed Central PMCID: PMC3402023.
- [52] Osonoi Y, Mita T, Osonoi T, Saito M, Tamasawa A, Nakayama S, et al. Morningness-eveningness questionnaire score and metabolic parameters in patients with type 2 diabetes mellitus. *Chronobiol Int*. 2014;31(9):1017-23. doi: 10.3109/07420528.2014.943843. PubMed PMID: 25102425.

- [53] Raynor HA, Champagne CM. Position of the Academy of Nutrition and Dietetics: Interventions for the Treatment of Overweight and Obesity in Adults. *J Acad Nutr Diet*. 2016;116(1):129-47. doi: 10.1016/j.jand.2015.10.031. PubMed PMID: 26718656.
- [54] Hermida RC, Ayala DE, Mojon A, Fernandez JR. Influence of time of day of blood pressure-lowering treatment on cardiovascular risk in hypertensive patients with type 2 diabetes. *Diabetes Care*. 2011;34(6):1270-6. doi: 10.2337/dc11-0297. PubMed PMID: 21617110; PubMed Central PMCID: PMC3114338.
- [55] Hermida RC, Ayala DE, Mojon A, Fernandez JR. Influence of circadian time of hypertension treatment on cardiovascular risk: results of the MAPEC study. *Chronobiol Int*. 2010;27(8):1629-51. doi: 10.3109/07420528.2010.510230. PubMed PMID: 20854139.
- [56] Hermida RC, Ayala DE, Mojon A, Fernandez JR. Bedtime ingestion of hypertension medications reduces the risk of new-onset type 2 diabetes: a randomised controlled trial. *Diabetologia*. 2016;59(2):255-65. doi: 10.1007/s00125-015-3749-7. PubMed PMID: 26399404.
- [57] Mameli C, Mazzantini S, Ben Nasr M, Fiorina P, Scaramuzza AE, Zuccotti GV. Explaining the increased mortality in type 1 diabetes. *World J Diabetes*. 2015;6(7):889-95. doi: 10.4239/wjd.v6.i7.889. PubMed PMID: 26185597; PubMed Central PMCID: PMC4499523.
- [58] Fiorina P, Lanfredini M, Montanari A, Peca MG, Veronelli A, Mello A, et al. Plasma homocysteine and folate are related to arterial blood pressure in type 2 diabetes mellitus. *Am J Hypertens*. 1998;11(9):1100-7. PubMed PMID: 9752896.

Table 1. Characteristics of the included studies

Observational studies					
Author(year) [ref]	Methods	Participants	Intervention	Outcomes	Changes in outcomes
Ruiz Lozano (2016) [26]	Observational prospective study (2006-2011)	N=270 patients treated with bariatric surgery (Roux-en-Y gastric bypass and sleeve gastrectomy)	Early eating vs Late eating according to the timing of main meal (before and after 15.00)	<i>Anthropometry measures:</i> weight, BMI, postoperative weight loss Energy and dietary intake before/during/after bariatric surgery: 4-days food record Morningness/eveningness questionnaire	<i>Weight change:</i> 70% of late eaters in poor weight-loss responders vs 42% in secondarily poor weight-loss responders and 37% in good weight-loss responders (p=0.01)
Hermenegildo Y (2016) [24]	Observational prospective study (2008-2012)	N=4243 adults from a population-based cohort Inclusion criteria: ≥18y, living in Spain, alive at follow-up Exclusion criteria: institutionalized, unable to give valid data about diet, cases with missing data on the evaluated variables	Quartile of energy intake by different meals (breakfast, mid-morning meal, lunch, mid-afternoon meal, dinner, snacking)	<i>Anthropometry measures:</i> weight gain (>3 kg)	<i>Weight change:</i> compared with those in the lowest quartile of % energy intake at lunch, the multivariate OR of gaining >3kg was 0.79 (95% CI 0.63-0.99) in the second quartile, 0.82 (0.64-1.04) in the third quartile and 0.62 (0.47-0.80) in the highest quartile (Ptrend=0.001)
Bo S (2014)	Observational	N=1245 adults from a population-based cohort	Tertiles of the percentage of total daily caloric intake from dinner	<i>Anthropometric measures:</i> weight, height, BMI, waist circumference	<i>Incidence of obesity:</i> from the lowest to the highest tertiles of total % daily caloric intake at dinner,

[28]	prospective study (2001-2008)	Inclusion criteria: age 45-64y from 6 general practitioners, Caucasian, living in Asti (North-Western Italy) Exclusion criteria: obesity and/or diabetes mellitus at baseline, died during follow-up		<i>Metabolic parameters:</i> blood glucose, glycated hemoglobin, cholesterol, HDL-cholesterol, LDL-cholesterol, triglycerides, HOMA-IR index Other blood parameters: C-reactive protein, alanine aminotransferase, γ -glutamyl transferase	the incidence rate of obesity increased (from 4.7 to 11.4%, $p<0.01$). The increased obesity risk for subjects in the highest tertile was confirmed in a multiple regression model (OR=2.33; 95% CI 1.17–4.65; $p=0.02$). <i>Incidence of diabetes:</i> individuals in the highest tertile of dinner % daily caloric intake showed an increase risk of incident type 2 diabetes (2.26; 0.89-5.75)
Garaulet M (2013) [16]	Observational prospective study (2007-2008)	N=420 obese/overweight individuals Exclusion criteria: special diet, weight-loss drugs, diabetes mellitus, chronic renal failure, hepatic diseases, cancers, any nutrition program within 2-y	Early eaters (lunch before 15:00) and late eaters (lunch after 15:00) All subjects received a 60-min educative program (once/week) with nutritional and exercise recommendations and a cognitive-behavioral approach	Anthropometric measures: weight, height, BMI, total body fat, waist circumference Metabolic parameters: blood glucose, cholesterol, HDL-cholesterol, LDL-cholesterol, triglycerides, HOMA-IR index, leptin, ghrelin Energy intake before/during treatment: 1-day dietary recall Energy expenditure: estimated by equations Morningness/eveningness questionnaire Sleep duration: evaluated by questionnaire	<i>Weight change:</i> early lunch eaters lost more weight than late eaters during the 20-weeks of intervention (9.9 ± 5.8 vs 7.7 ± 6.1 kg, $p=0.008$). The weight loss, expressed as % of initial weight was respectively: 11.3 ± 5.8 vs 9.0 ± 7.1 ($p=0.006$)
Randomized cross over/ controlled studies					

Author(year) [ref]	Methods	Participants	Intervention	Outcomes	Changes in outcomes
Bo S (2015) [20]	Cross-over randomized trial	<p>N=20 healthy volunteers</p> <p>Inclusion criteria: age 20-35y, BMI 19-26 kg/m², habitual moderate exercise level, <10 cigarettes/day.</p> <p>Exclusion criteria: Any acute or chronic diseases, menopause, any drugs or supplementations, any alimentary restrictions or specific diets, being a shift or night workers, unable to give a written informed consent</p>	<p>The same meal at 8:00 and, 7 days after at 20:00 or vice versa</p> <p>Each experiment lasted about 2-h</p>	<p><i>Calorimetric evaluation:</i> fasting RMR, after-meal RMR, DIT, fasting RQ, after meal RQ, RQ difference</p> <p><i>Metabolic parameters:</i> blood glucose, insulin, FFA, triglycerides every 30 min after each meal for 180 min</p>	<p><i>Metabolic variables:</i> delayed and larger increases in glucose and insulin concentrations were found after the evening meals.</p>

Bandin C (2015) [19]	Cross-over randomized trial Protocol 1 Each experiment lasted 2 weeks, after 1 week wash out Protocol 2 Each experiment lasted 2 weeks, after 1 week wash out	N=32 healthy women Exclusion criteria: endocrine, renal, hepatic, psychiatric disorders, any drugs (other than oral contraceptives) Protocol 1: N=10 Protocol 2: N=22	Early eating (lunch at 13:00) vs late eating (lunch at 16:30) for 2 weeks.	Specific measurements to protocols: Protocol 1 <i>Calorimetric evaluation:</i> fasting RMR, after-meal RMR, fasting RQ, after meal RQ, carbohydrate oxidation <i>Metabolic parameters:</i> Mixed meal test for glucose tolerance Protocol 2 <i>Wrist temperature</i> <i>Hormonal assessments:</i> salivary cortisol	<i>Metabolic variables:</i> late-eating lunch individuals showed significantly increased post-prandial glucose areas-under-the-curve than early eaters (102.6±30.8 vs 70.0±32.9 mmol/l×h; p=0.002)
Jacobowitz D,a (2013) [17]	Randomized controlled trial	N=93 overweight and obese women. Inclusion criteria: age 20-65y, BMI 25-37 kg/m ² , non-diabetic OGTT, presence of the metabolic syndrome Exclusion criteria: abnormal thyroid, liver or kidney function, cardiovascular disease,	Subjects were randomized to one of the following 1400 kcal weight-loss diet for 12 weeks: -breakfast group (700 kcal breakfast, 500 kcal lunch, 200 kcal dinner; N=46) -dinner group (200 kcal breakfast, 500 kcal lunch, 700 kcal dinner; N=47)	<i>Anthropometric measures:</i> height, weight, BMI, waist circumference <i>Metabolic parameters:</i> blood glucose and insulin after an OGTT, total cholesterol, LDL-cholesterol, HDL-cholesterol, triglycerides, ghrelin, HOMA-IR and HOMA-b, ISI <i>Appetite:</i> evaluated by questionnaires	<i>Weight change:</i> the breakfast group showed the greater weight loss after the intervention (-8.7±1.4 vs -3.6±1.5 kg; p<0.001) <i>Metabolic variables:</i> % changes in fasting glucose (-11.5 vs -4.2%), insulin (-51 vs -29%) and HOMA-IR (-57 vs -32.5%) significantly

		cancer, hypoglycemic drugs			decreased in the breakfast group. Similarly, OGTT test led to a greater decrease of glucose and insulin in the breakfast group
Jacobowitz D,b (2013) [18]	Randomized controlled trial	N=60 women with polycystic ovary syndrome Exclusion criteria: BMI>24.9 kg/m ² , on any diet, any drugs affecting weight, changing in weight >4.5 kg or in physical activity within the last 6 months	Subjects were randomized to one of the following 1800 kcal maintenance diet for 12 weeks: -breakfast diet (980 kcal breakfast, 640 kcal lunch, 190 kcal dinner; N=29) -dinner diet (190 kcal breakfast, 640 kcal lunch; 980 kcal dinner; N=31)	<i>Anthropometry measures:</i> BMI, waist circumference <i>Metabolic parameters:</i> blood glucose, insulin, HOMA-IR, HOMA-b <i>Hormonal assessments:</i> blood progesterone level, free and total testosterone, SHBG, 17-OH estradiol, DHEA-S, 17OHP, FAI, leuprolide stimulation test	<i>Weight change:</i> after the maintenance diets, weights did not change in the breakfast and dinner groups <i>Metabolic variables:</i> in the breakfast group, % changes of fasting glucose (-8 vs +2%), fasting insulin (-53 vs 0%), HOMA-IR (-56 vs +1%), HOMA-b (-35 vs -7%), ISI (+135 vs +2%), and glucose (-20 vs 0%) and insulin (-49 vs -7%) areas-under-the-curve were significantly higher

Morgan LM (2012) [25]	Cross-over randomized trial	N=6 healthy volunteers	<p>Subjects were randomized to one of the following 2000 kcal diets by cross-over:</p> <ul style="list-style-type: none"> -Low glycemic index with big breakfast (1200 kcal) and small dinner (400 kcal) -Low glycemic index with big dinner (1200 kcal) and small breakfast (400 kcal) -High glycemic index with big breakfast (1200 kcal) and small dinner (400 kcal) -High glycemic index with big dinner (1200 kcal) and small breakfast (400 kcal) 	<p><i>Metabolic parameters:</i> Blood glucose and insulin every 30 min after each meal for 120 min, post-prandial HOMA-IR, interstitial glucose by a continuous glucose monitoring system applied the day before each test</p>	<p><i>Metabolic variables:</i> interstitial glucose and insulin areas-under-the curve were significantly higher after consuming a big dinner rather than a big breakfast at the same glycemic index</p>
Tsuchida Y (2013) [27]	Cross-over random trial	<p>12 females (paid participants)</p> <p>Inclusion criteria: university students</p> <p>Exclusion criteria: smoking, current antibiotic use</p>	<p>Two experimental conditions:</p> <ul style="list-style-type: none"> a meal at usual suppertime (18:00) a meal at late suppertime (at 23:00), performed in different days 	<p><i>Metabolic parameters:</i> blood glucose every 30 min after each supper and after the breakfast of the next day for 180 min, unabsorbed carbohydrates by breath hydrogen test</p> <p><i>Calorimetric evaluation:</i> RQ</p>	<p><i>Metabolic variables:</i> a late suppertime meal determined significantly increased glucose values at 30, 60, 120, 150 and 180-min after the breakfast consumed the day after, with respect to the usual suppertime meal</p>

Abbreviations: 17-alpha Hydroxyprogesterone (17OHP), Body Mass Index (BMI), Dehydroepiandrosterone-Sulfate (DHEA-S), Diet-Induced Thermogenesis (DIT), Free Androgen Index (FAI), High Density Lipoprotein (HDL), Homeostasis model Assessment-Insulin resistance (HOMA-IR), Homeostasis model Assessment-beta cell function (HOMA-b), Insulin Sensitivity Index (ISI), Low Density Lipoprotein (LDL), Oral Glucose Tolerance Test (OGTT), Respiratory Quotient (RQ), Resting Metabolic Rate (RMR), Sex Hormone-Binding Globulin (SHBG).

Table 2 Risk of bias assessment in the trials included in the systematic review

Study	Random sequence generation	Allocation concealment	Blinding	Incomplete outcome data	Selective reporting	Free of other bias
Bo (2015)	L	U	L	L	L	L
Bandin (2015)	L	U	U	L	L	U
Jacubowitz ^a (2013)	U	U	H	H	L	L
Jacubowitz ^b (2013)	U	U	U	L	L	L
Tsuchida (2013)	U	U	U	U	L	L
Morgan (2012)	U	U	U	L	L	U

Criteria defined for quality assessment are based on the Cochrane guidelines.

Abbreviations: H, high risk of bias; L low risk of bias; U unclear or unrevealed risk of bias

Table 3 Risk of bias assessment in the observational studies included in the systematic review

Study	Ruiz Lozano (2016)	Hermenegildo (2016)	Bo (2014)	Garaulet (2013)
Domain				
Bias due to confounding	Serious	Moderate	Moderate	Moderate
Bias in selection of participants into study	Moderate	Low	Moderate	Moderate
Bias in classification of interventions	Moderate	Moderate	Moderate	Moderate
Bias due to departure from intended interventions	Low	Low	Low	Low
Bias due to missing data	Moderate	Low	Low	Low
Bias in measurement of outcomes	Low	Serious	Low	Low
Bias in selection of the reported results	Moderate	Low	Low	Moderate
Overall*	Serious	Serious	Moderate	Moderate

*Overall assessment derived from the seven domains of ROBINS-I (Risk Of Bias In Non-randomized Studies -of Intervention scale) tool