



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Bioactive Lipid Species and Metabolic Pathways in Progression and Resolution of Nonalcoholic Steatohepatitis

This is the author's manuscript Original Citation: Availability: This version is available http://hdl.handle.net/2318/1673657 since 2018-08-16T10:19:18Z Published version: DOI:10.1053/j.gastro.2018.06.031 Terms of use: Open Access Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *Gastroenterology*, *155*(2),282-302.e8, 2018, doi: 10.1053/j.gastro.2018.06.031.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), doi: 10.1053/j.gastro.2018.06.031

Bioactive lipid species and lipid metabolic pathways in NASH

progression and resolution

Running title: emerging lipid species in NASH

Giovanni Musso, M.D.¹, Maurizio Cassader, Ph.D.², Elena Paschetta M.D.¹, Roberto Gambino Ph.D.²

¹HUMANITAS Gradenigo, University of Turin, Turin, Italy

²Dept. of Medical Sciences, San Giovanni Battista Hospital, University of Turin, Turin, Italy

Correspondingauthor:

Giovanni Musso

HUMANITAS Gradenigo

C.^{so} Regina Margherita 8

10132 Turin, Italy

E-mail: giovanni.musso@gradenigo.it

Key words: eicosanoids, shpingolipids, proresolving, NAFLD

Word count: 5976

Grant support: this work received no funding

Disclosures: no author has any present or past conflict of interest to disclose

Author's contributions:

Giovanni Musso: designed research, conducted research, analyzed data, wrote paper, has primary responsibility for final content;

Maurizio Cassader: conducted research, analyzed and discussed data, approved final version of the paper;

Elena Paschetta: conducted research analyzed and discussed data, approved final version of the paper

Roberto Gambino: conducted research analyzed and discussed data, approved final version of the paper;

Abstract

The liver-related burden of NASH is growing as a consequence of increased disease prevalence and of the lack of an effective treatment.

A decade has passed since initial lipidomic analyses on NAFLD and our knowledge of NASH as a lipotoxic disease has considerbly expanded: identifying novel lipid species and metabolic pathways as potential pathogenic factors and therapeutic targets, including enzymes involved in fatty acid and triglyceride synthesis, polyunsaturated-derived eicosanoids, and bioactive sphingolipids, which are reviewed here. Furthermore, the concept of NASH as a pro-resolving defective disease and the role of specialized proresolving lipid molecules in the resolution of inflammation and regression of fibrosis in NASH are discussed.

Introduction

Nonalcoholic fatty liver disease (NAFLD) is the most common chronic liver disease in the world and an emerging risk factor for liver-related complications, including cirrhosis and hepatocellular carcinoma¹. NAFLD encompasses a histological spectrum ranging from simple steatosis (nonalcoholic fatty liver, NAFL) to non-alcoholic steatohepatitis (NASH)¹, with variable degrees of fibrosis, the latter having the potential to progress to cirrhosis².

The liver-related burden of NASH is growing and NASH is projected to be the leading indication for liver transplantation by 2020, as a consequence of increased disease prevalence and of the lack of an effective treatment³.

Lipotoxicity, defined as abnormal cellular lipid composition, leading to toxic lipid accumulation, organelle dysfunction, cell injury and chronic inflammation, is the hallmark of NASH^{2,4}, Our understanding of the main lipid species and pathways involved in lipotoxicity in NASH has susbtantially advanced in the last decade due to the progress in lipidomics, a subcategory of metabolomics that uses analytical chemistry techniques like mass spectrometry(MS) and chromatography to identify and quantify the diverse lipid species contained in biological samples². The analysis of data derived from from liver and blood samples of patients across the whole NAFLD spectrum generated specific lipid signatures which are associated with different stages of liver disease^{5,6,7,8,9,10,11,} (**Table 1**). The interpretation of these signatures not only disclosed novel metabolic pathways potentially responsible for liver disease progression or resolution, paving the way to new therapeutic approaches, but allowed also the identification of non-invasive diagnostic biomarkers of NASH and of fibrosis, which currently can be most reliably identified by invasive liver biopsy¹.

Lipidomic studies revealed that, although most hepatic lipids in NAFLD accumulate in the form of triglycerides (TGs), TGs are an inert form of lipid storage and protect against cell lipotoxicity, which is determined by the accumulation of toxic lipids, including intermediates in TG synthesis

(saturated fatty acids, SFAs, and free cholesterol, FC), ceramide and/or by a deficiency in lipid species that mantain cellular integrity including phospholipids, ω 3-polyunsaturated Fatty Acids(PUFAs) or PUFA-derived Specialized Proresolving Mediators (SPMs)^{10,11}. We will review recent advances regarding lipid species involved in NASH progression and resolution and discuss mechanistic insights and potential therapeutic targets¹¹. We will focus on different types of free fatty acids (FFAs), phospholipids, sphingolpids, and PUFA-derived eicosanoids and SPMs, The role of cholesterol accumulation in the pathogenesis of NASH has been recently reviewed elsewhere¹².

Role of saturated fatty acids in the pathogenesis and progression of NASH

Studies with stable isotopes demonstrated that in NAFLD, hepatic FFAs derive from lipolysis of adipose tissue TG (60% of hepatic FFAs), from *de novo* lipogenesis (25%), which is inappropriately upregulated, and from dietary $TG(15\%)^{13}$.

In NAFLD, as in other insulin resistant states, the liver is chronically exposed to high levels of circulating FFAs due to unrestricted lipolysis of adipose tissue TGs; additionally, hepatic uptake of circulating FFAs, which involves a tetrameric plasma membrane protein complex that comprises plasma membrane fatty acid-binding protein (FABP), caveolin-1, fatty acid translocase (FAT/CD36) and calcium independent membrane phospholipase A2 (iPLA2β), is uprregulated and contributes substantially to circulating FFA internalization¹⁴.

The saturated fatty acids (SFAs) palmitate (PA, C16:0) and stearate (C18:0) are major components of the diet, are synthesized via *de novo* lipogenesis and accumulate in the steatotic liver, paralleling liver disease severity⁸⁻¹⁰. These SFAs possess substantial lipotoxicity through a variety of mechanisms: they can bind to plasma membrane Toll-Like Receptor (TLR)-4 and death receptor TNF-related apoptosis-inducing ligand (TRAIL)-2, triggering caspase-dependent hepatocyte apoptosis and inflammasome-mediated pro-inflammatory cytokine secretion, or can enter the cell

and trigger endoplasmic reticulum (ER) stress, mitochondrial dysfunction, and c-Jun N-terminal kinase (JNK) activation.

JNK belongs to the mitogen-activated protein kinase (MAPK) family and is a major mediator of hepatic lipotoxicity: in hepatocytes JNK inactivates Insulin Receptor Substrate (IRS)-1, inducing insulin resistance, interacts with the outer membrane mitochondrial protein SH3BP5 (Sab) to impair respiration, and induces reactive oxygen species (ROS) generation¹⁵. Furthermore, JNK suppresses Peroxisome Proliferator-Activated Receptor (PPAR)-α-mediated Fibroblast Growth Factor (FGF)-21 expression and mitochondrial and peroxisomal β-oxidation, and activates the proapoptotic protein p53-upregulated modulator of apoptosis(PUMA)^{16,17}(Figure 1A) Notably, although hepatocytes are the primary target of SFA lipotoxicity, nonparenchimal cells are also involved: SFAs activate TLR-4 on hepatic stellate cells (HSCs) to secrete the chemokine monocyte chemoattractant protein (MCP)-1 and activate JNK in Kupffer cells and macrophages to induce proinflammatory M1 polarization, chemotaxis and secretion of profibrogenic factors Transforming Growth Factor(TGF)-β and Tissue Inhibitor of Metalloproteinase(TIMP)-1^{15,18}(Figure 1B).

On this basis, two strategies have been proposed to antagonizehepaticSFA lipotoxicity: the modulation of nuclear transcription factors involved in SFA metabolism, and selective inhibition of enzymes involved in key steps of lipid synthesis.

Modulation of nuclear transcription factors involved in lipid metabolism LXR and SREBP inhibition

Nuclear receptors are ligand-activated transcription factors that regulate the expression of target genes to affect diverse cellular processes, including lipid metabolism. A number of these nuclear receptors (reviewed in²), including liver X receptors (LXRs), Farnesoid X receptor (FXR), and

PPAR- α , PPAR- β (also called PPAR- δ) and PPAR- γ regulate hepatic lipid metabolism and are being targeted by emergingpharmacological therapies.

Liver X Receptor(LXR)-a modulation

There are two LXRs, LXR- α and LXR- β , with considerable sequence homology and the same ligand binding, but with different tissue distribution: LXR α is highly expressed in the liver, adipose tissue and macrophages, whereas LXR β is expressed in many tissues¹⁹.

LXR- α is a key regulator of whole-body FFA and cholesterol metabolism: upon activation, LXR α upregulates hepatic SREBP-1c-mediated *de novo* lipogenesis and inhibits VLDL catabolism, promoting hepatic steatosis, large VLDL triglyceride production and hyperlipidemia²⁰(**online Supplementary Table 1**). LXR- α activation promotes also net cholesterol loss from the body by upregulating ABCA1, ABCG5/G8 in macrophages, hepatocytes and enterocytes and downregulating intestinal NPC1L1 expression, thereby enhancing reverse cholesterol transport and intestinal excretion. Furthermore, LXR- α increases the hepatic trascription of cytochrome P450 7A1 (CYP7A1), the rate-limiting enzyme in bile acid synthesis, promoting cholesterol conversion to bile acids, and accelerates LDL-receptor degradation, thereby reducing hepatocyte cholesterol uptake¹².

25-Hydroxycholesterol-3-sulfate (25HC3S) is an endogenous sulfated oxysterol that suppresses LXR/SREBP-1 activation and NF- κ B-mediated pro-inflammatory response²¹, showing potent anti-steatotic, anti-inflammatory and anti-fibrotic activity in rodent models of NASH²².

25HC3S (Dur-928) is being developed as an oral agent for the treatment of NASH and proved safein a phase Ib study in patients with NASH (ANZCTR number:ACTRN12615001355561).

Farnesoid X Receptor agonists

Originally known for its function of bile acid sensor in enterohepatic tissues, farnesoid X receptor (FXR) has recently emerged as a master regulator of lipid metabolism and of inflammatory and fibrogenic processes. FXR is expressed mainly in the liver, intestine, kidney and adrenal glands, and at lower levels in adipose tissue. In the liver, FXR activation reverses lipotoxocity by increasing FA β-oxidation, suppressing SREBP-1c-mediated *de novo* lipogenesis and promoting cholesterol excretion by ABCG5/G8 transporters and has insulin-sensitizing, anti-inflammatory and antifibrotic properties (**online Supplementary Table 1**)². The semi-synthetic bile acid FXR agonist obetichoic acid (OCA) was evaluated in NASH, with favourable effects on NASH resolution and fibrosis improvement but unwanted changes in blood cholesterol levels (increased LDL-C and decreased HDL-C)²³. The impact of OCA on long-term clinical outcomes in noncirrhoticNASH will be clarified by the ongoing 5-year phase III REGENERATE trial (ClinicalTrials.gov ID: NCT02548351). Other non-steroidal FXR agonists (Px-102, Px-104,LMB763, Gs-9674) are being evaluated in phase I-IIa RCTs in NASH (ClinicalTrials.gov Identifier: NCT01998659, NCT0199910, NCT02854605).

PPAR agonists

PPARs belong to the nuclear receptor superfamily and they can be classified into 3 isotypes designated PPAR- α , PPAR- β (also called PPAR- δ) and PPAR- γ . PPARs form heterodimers with RXR and the PPAR:RXR heterodimer extensively regulates genes involved in metabolism inflammation and fibrogenesis², promptingdevelopment of PPAR agonistsfor NASH therapeutics²(online Supplementary Table 1).

PPAR- α and PPAR- δ have overlapping metabolic effects and complementary tissue distribution, which led to the development of dual PPAR α/δ agonists. These agents reversed hepatic toxic lipid accumulation by upregulating mitochondrial and peroxisomal fatty acid β -oxidation and microsomal ω -oxidation and ketogenesis in hepatic parenchimal and nonparenchymal cells,

improving NASH and fibrosis in preclinical models of NASH^{24,25}. Furthermore, the lipid-oxidizing effects of PPAR α/δ agonists in skeletal muscle and adipose tissue improved systemic insulin resistance and adipose tissue dysfunction (**online supplementary Table 1**).

Elafibranor, a dual PPAR α/δ agonist, reversed NASH without fibrosis worsening and improved blood lipid profileand glucose utilization in a phase IIb RCT²⁶, while the selective PPAR- δ agonist seladepar (MBX-8025)improved liver enzymes, inflammatory markers, insulin resistance and atherogenic dyslipidemia in dyslipidemic patients²⁷.

PPAR- γ is expressed by a dipocytes, immune cells, including macrophages Kupffer cells, and HSCs, and to a lower extent by hepatocytes and skeletal miocytes.

PPAR- γ activation reverses lipotoxocity by enhancing adipocyte differentiation and insulin sensitivity and promoting adiponectin secretion, thereby reducing lipotoxic SFA overflow to the liver; furthermore, PPAR- γ upregulation shifts SFA-induced macrophage and Kupffer cell polarization from an M1 to an M2 phenotype¹⁸, and reverses HSC transdifferentiation to myofibroblasts²⁸(**online supplementary Table 1**)['].

On this basis, dual PPAR- α/γ agonists and panPPAR agonists have been developed to combine the lipid-oxidizing properties of PPAR- α/δ agonists with the insulin sensitizing effects of PPAR- γ agonists.

Saroglitazar, a dual PPAR- α/γ agonist, and IVA337(Lanifibranor), a pan-PPAR agonist, increased the expression of β -oxidation-related and fatty acid desaturation-related enzymes and ameliorated diet-induced NASH and fibrosis with an efficacy superior to single class PPAR agonists^{29,30}. Saroglitazar improved liver enzymes in diabetic NAFLD patients³¹ and a small, phase IIa clinical trial on biopsy-proven NASH patients has been completed (CTRI registration no.: CTRI/2010/091/000108).

Inhibition of enzymes involved in *de novo* lipogenesis and triglyceride synthesis

Acetyl-CoA carboxylase (ACC) inhibition

ACC catalyzes the ATP-dependent carboxylation of acetyl-CoA to malonyl-CoA, the first, ratelimiting step in *de novo* lipogenesis¹⁹.

ACC exists as two tissue-specific isozymes that are encoded by separate genes and display distinct cellular distributions: ACC1 is a cytosolic enzyme present in lipogenic tissues (liver, adipose); ACC2 is a mitochondrial surface-associated isozyme present in oxidative tissues (liver, heart, skeletal muscle)¹⁹.

In the liver, cytosolic malonyl-CoA formed by ACC1 is used primarily for *de novo* lipogenesis, whereas mitochondrial malonyl-CoA formed by ACC2 acts primarily as an allosteric inhibitor of mitochondrial fatty acid uptake by carnitine palmitoyltransferase (CPT)-1, resulting in a subsequent reduction in mitochondrial fatty acid β-oxidation¹⁹. Thus, inhibition of the ACCs may reduce *de novo* lipogenesis and increase fatty acid β-oxidation, two key steps in fatty acid metabolism that are dysregulated in human and experimental NASH¹⁹: consistently, mice with constitutively activated ACC have enhanced lipogenesis, hepatic insulin resistance and develop NASH and fibrosis, which are prevented by genetic ACC ablation and reversed by pharmacological ACC inhibition³². Following these preclinical results, the orally available, liver-targeted, selective small molecule allosteric ACC1/2 inhibitor GS-0976 (formerly NDI-010976) suppressed *denovo* lipogenesis and improved MRI-assessed steatosis and serum fibrosis markers in a small phase 2a RCT enrolling noncirrhotic NASH patients³³.

Diacylglycerol acyltransferase (DGAT)-1 and DGAT-2 inhibition

Acyl-CoA:diacylglycerol acyltransferase (DGAT) is an ER-membrane-bound enzyme that catalyzes the final step in TG synthesis: diglyceride (DAG) esterification by long-chain acyl-CoA esters. DGAT exists in two isoforms encoded by two distinct genes. DGAT-1 is most highly expressed in the small intestine enterocytes, where it participates to the lipolysis-reesterification cycle by reassemblingTG from dietary FFAs in the process of intestinal fat absorption to form chylomicrons¹⁹. DGAT-2 is expressed primarily in the liver, adipose tissue and skin, where it synthesizes TG from *de novo* synthesized FFAs and newly formed diglycerides $(DAGs)^{34}$. DGAT-1 knockout $(DGAT1^{-/-})$ mice are viable and have modest reductions in tissue TG, whereas DGAT-2 knockout $(DGAT2^{-/-})$ mice have severe lipopenia(~90 % reduction in whole-body TG) and impaired skin barrier function (including the inability to retain moisture), and die shortly after birth³⁵.

Due to the data from DGAT knockout mouse models and to the finding that dietary SFA absorption is upregulated in NASH patients³⁶, most research focused on selective DGAT-1 inhibitors for the treatment of obesity, NASH and hyperlipidemia. Pradigastat, a potent selective DGAT-1 inhibitor, slightly reduced MRI-assessed liver fat after 24 weeks, but diarrhea and steatorrhea due to fat malabsorption affected up to 86% of patients on active treatment³⁷.

Growing data suggest that pharmacological DGAT-2 inhibition is an effective and safe therapeutic option, due to the central role of this isoenzyme in hepatic TG metabolism: unlike DGAT-1 inhibition, selective hepatic DGAT-2 inhibition with small molecule inhibitors improved high fat diet-induced hepatic steatosis, fibrosis and insulin resistanceby down-regulating SREBP-1c-mediated lipogenesis, protein kinase C(PKC)-mediated insulin resistance, and NF- κ B-mediated inflammation and fibrgenesis and upregulated fatty acid oxidative and thermogenic pathways, including CPT-1 and uncoupling protein(UCP)-2^{38,39,40}.

Role of glycerophospholipid lysophosphatidylcholine (LPC) in the pathogenesis and progression of NASH

LPC is a glycerophospholipid generated from phosphatidylcholine (PC) by partial hydrolysis of one fatty acid group. This reaction can be catalyzed intracellularly by phospholipase A2(PLA2) or extracellularly by plasma lecithincholesterol acyltransferase (LCAT)¹⁹.

The relative contribution of these two pathways to hepatic LPC accumulation *in vivo* remains to be elucidated. PLA2-generated LPC is an important mediator of SFA-induced lipotoxicity: in

cultured hepatocytes, incubation wih LPC triggered cell apoptosis, while PLA2 inhibition reduced intracellular LPC and palmitate-induced apoptosis^{14,41}. Consistently, hepatic LPC content is increased in rodent models and in human NASH and parallels liver disease severity^{5,8,10}. Beside lipoapoptosis, which is triggered by JNK and ER stress activation, other mechanisms for LPC lipotoxicity have been identified: LPC impairs hepatic mitochondrial oxidative phosphorylation, causing mitochndrial dysfunction⁴², induces pro-inflammatory and pro-fibrogenic extracellularvescicle (EV) release from hepatocytes^{43,44} and is converted into the potent profibrogenic phospholipid lysophosphatidic acid (LPA) by the enzyme autotaxin⁴⁵(**Figure 1**; **online supplementary Table 2**).

Most of these lipotoxic mechanisms overlap with those of SFA, suggesting that LPC generation may be a major downstream effector of SFA citotoxicity. An additional, important mechanism of lipotoxicity is the depletion of membrane PC caused by PLA2 activation^{14,46}. PC is the most abundant phospholipid in mammalian cells, is essential for cell membrane integrity and is also an important feedback inhibitor of SREBP1c-mediated *de novo* lipogenesis⁴⁷. Hepatocytes have a high demand for PC, which is used for the production of VLDL and is secreted in bile, making these cells particularly liable to PC depletion.

The NASH liver is characterized by hepatic PC depletion (**Table 1**)^{9,10} and disruption of hepatocyte membrane functional integrity, which results in release of lipotoxic lipids into the extracellular space, hepatocyte apoptosis, inflammation, and liver disease progression^{46,48}. Remarkably, this chain of events was interrupted by genetic or pharmacological PLA2 inhibition, which normalized intracellular PC pool, prevented diet-indued NAFLD and reversed established NASH and fibrosis^{14,49}.

Bioactive Sphingolipids: role of the "sphingolipid rheostat" in the pathogenesis and treatment of NASH

Sphingolipids are a lipid species containing a sphingosine (18-carbon amino alcohol) backbone that is N-acylated with various fatty acids. Sphingolipids are ubiquitous components of eukaryotic cell membranes and are both structural lipids and potent signaling molecules regulating key cellular functions, like cell growth and survival, proliferation, differentiation, migration and immune responses in health and disease states, including metabolic disorders and NASH⁵⁰. Furthermore, ceramide, sphingosine, and sphingosine-1-Phosphate (S1P) are readily interconvertible and the delicate balance of their relative levels has been termed the "sphingolipid rheostat", as it can direct cellular fate to opposing directions: as an example, ceramide and sphingosine induce cell apoptosis, senescence, and growth arrest while S1P promotes cell survival, growth, and proliferation⁵⁰.

Role of ceramide in NASH

Ceramide consists of an amino group of a sphingoid base, typically sphingosine, bound to a saturated or monounsaturated fatty acyl chain, and constitutes the hydrophobic core of all complex sphingolipids, including sphingomyelin, cerebrosides and gangliosides⁵⁰. Ceramide can be synthesized *de novo* from serine and palmitate by the sequential action of three ER-resident enzymes—serine palmitoyltransferase (SPT), ceramide synthase (CeS) and dihydroceramide desaturase (DES) — or generated from hydrolysis of plasma membrane sphingomyelin into ceramide and phosphocholine by the enzyme sphingomyelinase (SMase)(**Figure 2**). Lipidomic studies indicate hepatic ceramide content is increased in NAFLD and correlates with liver disease severity⁵⁻¹⁰. Hepatic ceramide overload results from both increased hydrolysis of sphingomyelin by acid sphingomyelinase (ASMase), one isoform of SMase which is upregulated in NASH by proinflammatory stimuli, including reactive oxygen species (ROS), TNF- α , death receptor ligands⁵¹, and from increased availability of SFAs, the limiting substrate for *de novo* ceramide synthesis. Additionally, recent data indicate the intestine is a relevant source of ceramide, whose synthesis in ileum and cecum is stimulated by the activation of a bile acid/intestinal FXR axis(**Figure 2**).

metabolic dysregulation in NASH, intestine-specific genetic or pharmacological FXR inhibition reduced circulating ceramide levels by 30-50%, enhanced adipose tissue browning, and improved hepatic insulin resistance and liver injury in HFD obese mice; remarkably, these effects were reversed by ceramide injection, confirming the benefits of intestinal FXR inhibition are mediated by ceramide reduction.

Accumulating evidence suggests ceramide is a key mediator of SFA lipotoxicity, which requires ceramide accumulation and is prevented by inhibiting ceramide formation: for example, studies using lipid infusion revealed that ceramide formation is required for SFA-induced inhibition of Akt-mediated insulin signaling and insulin resistance⁵³. The mitochondria are another major cellular target of ceramide, which impairs fatty acid β -oxidation through inactivation of complex II and IV of the electron transport chain and promotes ROS production, TG accumulation and insulin resistance^{54,55}(**Figure 2**). Further mechanisms of hepatic lipotoxicity of ceramide overload in NASH include disruption of calcium homeostasis in the ER, which leads to ER-stress-mediated apoptosis Nlrp3 inflammasome activation impairment of autophagy, and upregulation of hepatic hepcidin, which leads to hepatic iron overload^{56,57,58,59,60}. Additionally, ASMAse activation may promote liver injury independently of ceramide accumulation by disrupting methionine and phosphatidylcholine metabolism, which promotes lysosomal membrane permeabilization⁵⁸ and directly activates HSCs⁶¹(**Figure 2**).

On this basis, several strategies aiming at relieving hepatic ceramide accumulation have been investigated, including inhibiting ceramide hydrolysis from sphingomyelin or ceramide *de novo* biosynthesis or enhancing ceramide degradation. ASMase can be inhibited byfunctional inhibitors of acid sphingomyelinases (FIASMAs), a heterogeneous group of weakly basic and highly lipophilic molecules (**Table 2**), which accumulate into the lysosome, detach ASMase from the inner lysosomal membrane and induce its degradation by lysosomial proteases⁶².

Reduction of ceramide *de novo* synthesis can be achieved through inhibition of the enzymes involved in ceramide synthesis, i.e., SPT, CeS or DES or through inhibition of intestinal FXR activation (**Table 2**).

Treatment of genetically induced (ob/ob) and HFD–induced obese rodents with the natural fungal metabolite myriocin, a specific inhibitor of SPT, decreased circulating ceramides and body weight and improved glucose tolerance, hepatic insulin resistance, steatosis and fibrosis^{63,64}. However, complete whole-body inhibition of ceramide synthesis may have significant side effects, including neurodegeneration⁶⁵, because of the pivotal role of ceramide in the formation of other sphingolipid derivatives that are essential to cell membrane function and intracellular signaling pathways. Hence, selective intestinal FXR inhibition may overcome the limitation of whole-body ceramide synthesis inhibition: the bile acid derivative glycine-β-muricholic acid (Gly-MCA), reduced systemic ceramide levels and improved obesity and NAFLD without affecting CNS ceramide synthesis⁵². The challenge will be to restrict FXR inhibition to the intestine, as hepatic FXR activation actually improved NASH²³.

Another strategy to limit side effects of whole-body ceramide depletion derives from the finding that most lipotoxic effects of ceramide are mediated by C16:0 ceramide, which predominantly accumulates in the liver and adipose tissue of NASH patients^{10,55}, but not by very-long acyl-chain fatty acid (VLCFA) ceramides^{54,55}.

In mammals, CeS comproses six isoforms (CerS1–6), which differ in their acyl-chain specificities, tissue distribution, and transcriptional, and post-translational regulation. In the liver, CeS2 uses very long acyl-chain C22–26 fatty acids, while CeS5-CeS6 use long-chain C14–16 acyl-chain fatty acids⁶⁶. Therefore, selective inhibition of CeS5-6 may prevent accumulation of toxic C16:0 ceramide without depriving the organism of VLCFA ceramides.

Compound ST1072 inhibits preferentially CerS4 and CerS6⁶⁷, but to date has not been evaluated in NASH or associated metabolic disorders. CerS activity can also be modulated by phosphorylation or deacetylation, which opens up further therapeutic options for selective inhibition of C16:0

ceramide production⁶⁸. Finally, Dihydroceramide desaturase (DES)-1 inhibitor fenretinide has been shown to inhibit CeS5, downregulate the levels of long acyl-chain ceramides in favour of VLCFA ceramides⁶⁹ and to improve diet-induced obesity and NAFLD⁷⁰

Another therapeutic strategy is to enhance ceramide degradation by the enzyme acid ceraminidase, which is an important mediator of the biological actions of adiponectin⁷¹(**Figure 2**). Acid ceraminidase overexpression reduced hepatocyte apoptosis, liver inflammation and fibrosis and prevented high fat diet-induced NASH and insulin resistance in transgenic mice⁷².

Role of Sphingosine-1-Phosphate (S1P) in NASH

Once generated from ceramide deacylation, sphingosine can be phosphorylated by sphingosine kinases (SphKs) to form Sphingosine-1-Phosphate(S1P)(**Figure 2**). There are two pathways of S1P degradation: reversible dephosphorylation to sphingosine by nonspecific phosphatases, and by two S1P-specific phosphatases, SPP1 and SPP2; and irreversible cleavage by S1P lyase (SPL), which leads to the formation of phosphatidylethanolamine and hexadecenal. The latter is the only exit pathway for degradation of sphingoid bases in mammalian cells(**Figure 2**).

Although the ready interconvertibility of ceramide, sphingosine, and S1P and the opposing effects of these sphingolipidson cell growth and survival led tocoin the term "sphingolipid rheostat"⁵⁰, growing data support a more complex picture for S1P, which can exert diverse and even opposing actions, depending on the subcellular compartment of synthesis, on the cellular molecular target, and on the cell type.

In humans two sphingosine kinase isoforms (SphK1 and SphK2) exist, with different subcellular localization, resulting in cellular S1P compartmentalization⁷³. Upon activation, SphK1 translocates to the plasma membrane, where it catalyses the formation of S1P from plasma membrane-associated sphingosine: here, generated S1P is preferentially released extracellularly to activate a family of five cell surface G-protein coupled receptors (GPCRs), named S1PR1-5.

Conversely, SphK2 localizes in intracellular compartments, including the nucleus, the ER and the mitochondria, where newly generated S1P acts directly on intracellular targets in a receptorindependent mode: in the nucleus, S1P arrests DNA synthesis by inhibiting Histone Deacetylases (HDAC)-1/2; in the ER, S1P contributes to TNF receptor-associated factor 2 (TRAF2) activation, which is required for NF-kB activation⁷⁴, and in the mitochondria, S1P interacts with prohibitin 2 to regulate respiration and triggers BAK-dependent cell apoptosis⁷³.

The cellular compartmentalization of SphKs and S1P is therefore so critical for the regulation of cellular functions that artificially targeting SK1 to the ER or nucleus can allow the otherwise prosurvival enzyme to promote apoptosis⁷⁵.

The expression of the two SphK isoforms also varies across different tissues/organs, with SK1 most highly expressed in lung, spleen and leukocyte and SK2 being the predominant isoform in the kidney and liver⁷³⁻⁷⁵. However in NASH hepatic SphK1 expression is considerably upregulated^{76, 77} as a result of SFA overflow to the liver and of the stimulatory effect of pro-inflammatory cyokine TNF- α and IL-1 elevation, which may play a critical role in promoting liver injury.

The complex interaction of the aforementioned factors explains the results of studies evaluating the role of SpK1/SphK2 and of S1P receptor (S1PR) modulation in NASH.

The functional manipulation of SphK isoforms in diet-induced NASH models demonstrated that SphK1 activation promotes NASH, enhances secretion of pro-inflammatory cytokines TNF- α and IL-6 and inhibits secretion of the anti-inflammatory adiponectin and IL-10 adipokines⁷⁸, while SphK2 activation has opposite effects⁷⁹. However, since SphK activation uses ceramideas a substrate to synthesize S1P, intracellular levels of these two sphingolipids are often inversely related, making it is difficult to dissect the effects of changes in S1P levels from the changes in ceramide concentrationfollowing functional SphK manipulation^{78,79}.

Studies using whole-body genetic or pharmacological manipulation of SIPRs⁸⁰ indicate S1PR1 and S1PR3 are the crucial receptor subtypes mediating not only hepatic necro-inflammation but also fibrosis progression: in human fibrotic liver SphK1 S1P and S1PR(1,3) expression were increased

irrespective of the etiology of fibrosis; S1PR(1,3) were massively upregulated and exerted a powerful migratory action on human myofibroblasts, whereas S1PR2, which inhibited myofibroblast migration, was downregulated^{81, 82}(**Figure 2**). Furthermore, TGF- β is a potent inducer of SphK1, but not of SphK2 expression, and S1P is a crucial mediator of intracellular TGF- β signaling⁸³.

The therapeutic implications of S1PR antagonism in NASH are unexplored.

Administration of FTY720 (fingolimod), a functional S1PR1 antagonist approved for the treatment of multiple sclerosis, showed substantial antisteatotic, anti-inflammatory and anti-fibrotic activity in diet-induced rodent models of NASH, and S1PR(1,3)antagonist VPC23019 inhibited HSC activation in cell cultures^{79,81}.

However, it should be noted that the functional effects of S1P axis activation may be even opposing, depending on the cell type involved: as an example, activation of S1PR1 in endothelial sinusoidal cells promotes regeneration and suppresses fibrosis in the liver⁸⁴, and adipocyte-specific inhibition of S1P synthesis leads to lipodystrophy, NASH and insulin resistance⁸⁵. Ongoing nanotechnology approaches may develop liver-specific FTY720 nanoparticles and limit off-target effects of S1PR antagonists⁸⁶. A deeper knowledge of the down-stream cellular signaling pathways and functional effects of S1P in different organs and tissues would allow more targeted therapeutic interventions.

Monounsaturated Fatty Acid (MUFA) in NASH

The most abundant and well-studied MUFAs in NAFLD are palmitoleic acid, a 16 carbon length monounsaturated fatty acid (C16:1) and oleic acid, a 18 carbon length MUFA (C18:1), which are generated by the enzyme stearoyl CoA desaturase-1 (SCD1) from SFA palmitic and stearic acid, respectively¹⁹.

Despite the fact that they contribute to steatosis, these MUFAs are less lipotoxic than SFAs:

when individually studied, these MUFAs induce apoptosis, but this effect is minimal, compared with SFAs and they greatly attenuated palmitate-induced apoptosis in cultured hepatocytes⁸⁷. The cytoprotective effects of MUFAs may depend on their lower ability to trigger ER stress and PUMA activation and to a more efficient incorporation into TG: consistently, in cultured hepatocytes the combination of a MUFA with palmitate or SCD1 upregulation mitigated palmitate-induced apoptosis but enhanced TG accumulation while, in the setting of impaired TG synthesis, oleate accumulaion induced lipotoxicity⁸⁸.

Polyunsaturated Fatty Acid (PUFA) biosynthetic pathways in the pathogenesis and treatment of NASH

Twenty-carbon and more than 20-carbon polyunsaturated fatty acids (PUFAs) include two classes of PUFA, i.e., n-6 (ω -6) and n-3 (ω -3) series, named after the position of the first unsaturation counting from the methyl end of the fatty acid, the so-called omega-C.

ω-6 PUFAs include dihomo-γ-linolenic (DGLA, 20:3n-6) and arachidonic acid (AA, 20:4n-6),while ω-3 PUFAs include eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3)(**Figure 3**).

Mammals do not have the necessary fatty acid desaturases to convert oleic acid (18:1*n*-9) into linoleic acid (LA, 18:2n-6) and α -linolenic acid(ALA, 18:3n-3), which must therefore be introduced with the diet and are referred to as dietary essential fatty acid (**Figure 3**). The conversion rates of 18-carbon fatty acids to 20-carbon PUFAs are less than 5-10%⁸⁹: hence, fish and their oils, which contain high levels of these PUFAs, such as EPA and DHA, are the primary source of these FAs. The relevance of ω -6 and ω -3 PUFAs in numerous cellular biological functions has emerged in the latest years⁹⁰. PUFAs modulate cell membrane fluidity and permeability, membrane microdomain composition and hormone receptor binding, a critical step in receptor signaling: as an example, ω -3 PUFAs increase membrane fluidity, resulting in an enhanced number of membrane insulin receptors, an increased affinity of insulin to its receptors and an improved insulin sensitivity⁹¹. Furthermore, PUFAs have been found to modulate the expression of genes involved in lipid metabolism, redox balance, inflammation and fibrogenesis through interacting with nuclear receptors and transcription factors in different experimental models of NASH^{92,93,94, 95}(**online**

Supplementary Table 3).

Finally, PUFAs are released by PLA2 from cell membrane phospholipids to become substrates for the synthesis of eicosanoids and specialized proresolving mediators (SPMs) by the coordinated activities of cyclooxygenases (COX-1 a constitutive enzyme, or COX-2 an inducible enzyme), lipoxygenases (5-, 12-, or 15-LOX), or cytochrome P450 monoxygenases: while ω -6 PUFA arachidonic acid gives rise to the pro-inflammatory 2-series PGs and TXs and to the 4-series LTs, ω -3 PUFAs EPA and DHA give rise to the potent specialized pro-resolving molecules(SPMs) lipoxins, resolvins, maresins and protectins, which drive inflammatory response resolution(see below)¹¹(**Figure 4**).

The synthesis of highly unsaturated fatty acids such as the ω -3 EPA and DHA (synthesized from α linolenic acid), and the ω -6 AA (synthesized from linoleic acid) is a multi-stage process requiring successive actions by elongase and desaturase enzymes. In humans, three desaturases occur: Fatty Acid Desaturase 1 (FADS1, or Δ 5 desaturase), Fatty Acid Desaturase 2 (FADS2, or Δ 6 desaturase), and Stearoyl CoA Desaturase(SCD)-1 (or Δ 9 desaturase, where Δ indicates the number of the first carbon atom that forms the double bond relative to the carboxyl group)⁸⁹.

FADS1 and FADS2 are membrane-bound enzymes that catalyze the rate-limiting steps in the formation of long-chain PUFAs AA, EPA AND DHA(**Figure 3**). SCD-1 forms the MUFA oleic acid (18:1) from stearic acid (18:0), a saturated fatty acid either synthesized in the body from palmitic acid or ingested directly.

Lipidomic analyses coupled with enzymatic activity and gene expression profiling of human and mouse liver tissue highlighted a major dysregulation in hepatic long chain fatty acid (LCFA) desaturation and an unbalanced ω -6 to ω -3 ratio, with impaired fluxes toward the omega-3 fatty

acids pathway resulting in an increased omega-6 to omega-3 ratio as a key pathogenic factor in NASH^{5,9,10,96}.

Growing evidence support a central role for impaired FADS1 activity in dysregulated long chain fatty acid (LCFA) fluxes and liver injury in NASH^{9,10}. Impaired FADS1 activity acts as a bottleneck leading upstream to the accumulation of SFA and of preferential ω-6 PUFA-derived proinflammatory eicosanoids and downstream to the deficiency in VLCFA, which are substrates for the synthesis of phospholipids, essential components of cell membranes Studies using pharmacological FADS1 activity manipulation demonstrated that this enzyme is a key determinant of membrane phospholipid composition and proinflammatory versus proresolving lipid mediator balance⁹⁷. Hepatic FADS1 inhibition with Antisense Oligonucleotides (ASOs) yields two major pathophysiological consequences promoting NASH development: downstream, it alters the composition and quantity of membrane phospholipid leading to cell membrane phospholipid deficiency, hepatocyte membrane integrity disruption, cell necrosis and extracellular leak of lipotoxic lipids, which contribute to liver injury^{9,10}. Upstream, the impaired FADS1 activity leads to an imbalance in desaturation fluxes toward the preferential synthesis of ω -6-derived pro-inflammatory eicosanoids PGs, TXs and LTs (see below) at the expense of ω -3derived specialized pro-resolving lipid mediators(SPMs), further promoting chronic hepatic inflammation (Figure 3-4)'. Importantly, the proinflammatory effects of FADS1 deficiency are non limited to hepatocytes, but occur also in cultured macrophages, where FADS1 knockout induced classic M1 polarization and suppressed pro-resolving M2 activation programs⁹⁷. In contrast with these proinflammatory effects, FADS1 inhibition has anti-lipogennic effects by inhibiting LXR/SREBP-1c-mediated lipogenesis, which may explain why pharmacological FADS1 inhibition reduced hepatic steatosis body weight and insulin resistance^{97,98}.

It should be noted that the impact of FADS1 activity on livdr injury is diet-specific and is affected by the relative ω -3 and ω -6 PUFA intake:

Mechanisms underlying impaired FADS1 activity in NASH are various: dietary FAs regulate hepatic *FADS1* and *FADS2* gene expression and the activity is reduced by PUFAs in human HepG2 cells⁹⁹ and in rodents fed PUFA-enriched diets¹⁰⁰. A genetic component has also been proposed, as functional genetic polymorphisms in *FADS1* gene have been found to modulate human hepatic lipid composition and content¹⁰¹.

The dominant strategy adopted to restore a normal hepatic hepatic ω -6/ ω -3 PUFA ratio has been dietary supplementation with varying doses and formulations of ω -3 PUFAs with different biological activity, which showed consistent benefits on liver injury and metabolic parameters in diet-induced models of NASH^{93-95,97}. However, the analysis of RCTs with PUFAs conducted to date in NASH, demonstrated that ω -3 supplementation rescued hepatic ER stress, mitochondrial dysfunction and deregulated *de novo* lipogenesis¹⁰², improved steatosis and cardiometabolic profile, but did not affect histological necro-inflammation, NAFLD disease activity and fibrosis ^{90,103}. Potential reasons for the discrepant results of PUFA supplementation on steatosis versus those on necroinflammation and fibrosis include the heterogeneity of the type (with DHA showing more consistent anti-inflammatory and anti-fibrotic activity that EPA in preclinical models^{94, 95}), doses and formulation of ω -3 PUFAs used in different RCTs. Furthermore, while the antisteatogenic and metabolic effects of PUFAs are mediated directly by PUFA interaction with transcription factors involved in lipid metabolism, the anti-inflammatory effcts are mediated by PUFA-derived SPMs⁹³ and it may be more difficult to reach hepatic ω -3 PUFA concentrations high enough to enhance generation of anti-inflammatory/pro-resolving mediators, as the largest amount of dietary FAs undergoes β -oxidation (20–30%) or storage as TGs in adipocytes (15–80%)¹⁹. Furthermore, in NASH SMPs may be more rapidly inactivated by eicosanoid oxidoreductases and/or by the SPM receptor down-regulation observed in obesity-related condition(discussed below)¹⁰⁴. Other approaches to replace or complement dietary supplementation have been tested preclinically to re-equilibrate hepatic ω -6/ ω -3 ratio: the introduction of ω -3 desaturase (FAT-1) activity, which is

capable of generating ω -3 from ω -6 PUFAs and is absent in humans, restored a normal hepatic ω -3 PUFA content and reversed steatosis and necro-inflammatory changes in transgenic mice fed a HFD diet⁹³.

Another focus of research is the development of inhibitors of SCD-1, which is the critical control point regulating *de novo* lipogenesis, triglyceride synthesis and FA oxidation through the generation of MUFAs¹⁰⁵. A small molecule SCD-1 inhibitor ameliorated hepatic steatosis and necroinflammation in diverse nutritional models of NASH¹⁰⁶, and aramchol, a conjugate of cholic and arachidic acid that inhibits SCD-1 and *de novo* lipogenesis, given for 3 months significantly decreased hepatic fat content by 12.5% in a phase II RCT enrolling NAFLD patients¹⁰⁷. Treatment was not associated with improvement in liver enzymes, raising a concern that the reduction in hepatic fat was not accompanied by an improvement in inflammation or cellular injury. This is important as a potential drawback of SCD-1 inhibition is the increase of substrate SFAs, which are lipotoxic and may exacerbate inflammation¹⁰⁸. A phase IIb study with histological end-points is currently evaluating the effects of higher aramchol doses (400 and 600 mg/day) in non-cirrhotic biopsy-proven NASH (clinicaltrials.gov.ID: NCT02279524)(**Table 2**).

Eicosanoids and Specialized Proresolving Mediators in NASH

A hallmark of obesity-related disorders is a state of chronic low-grade inflammation, which in NASH recognizes the adipose tissue and the liver as major targets. In abdominal adipose tissue, the inflammatory state is fuelled by an expansion of adipose tissue pro-inflammatory macrophage pool², resulting in adipose tissue and systemic insulin resistance, unrestrained lipolysis and release of lipotoxic fatty acids, and pro-inflammatory cytokine and chemokine secretion^{2,109}. In the liver, chronic unresolved inflammation evokes a wound healing process, which includes fibrogenesis and persistent production of new extracellular matrix to replace damaged tissue, resulting in tissue scarring, fibrosis and eventually cirrhosis. Among the different inflammatory mediators, mounting experimental and human evidence supports a key role for eicosanoids and Specialized Proresolving

mediators (SPMs) in determining the immunometabolic milieu of NASH, with relevant therapeutic implications.

Eicosanoids (from the Greek term "*eicosa-*", "twenty") is the collective term for a family of signaling lipid molecules derived from the oxidation of straight-chain 20C-containing PUFAs, mainly AA, but also DGLA and EPA (**Figure 4**)¹¹⁰. Multiple subfamilies of eicosanoids exist, including prostaglandins(PGs), thromboxanes(TXs), leukotrienes(LTs), lipoxins(LXs) and resolvins(Rvs), with functionally heterogeneous, and even opposing activities on inflammation, immune and cardiovascular system, metabolism, cell growth and tisue remodeling. Eicosanoids act as autocrine, paracrine or endocrine signaling agents through binding to surface G-protein coupled receptors(GPCRs) on target cells. Unlike FAs, eicosanoids are not stored within cells but rather synthesized from ω -6 and ω -3 PUFAs contained in membrane phospholipids upon the activation of phospholipases A2's (PLA2s). Following their release from cell membranes, ω -6 and ω -3 PUFAs are further metabolized through various pathways involving the enzymes cyclooxygenases (COXs) and the thre major lipoxygenases 5-, 12- and 15-lipoxygenase(5-, 12-, 15-LOXs).

resolving lipid molecules, collectively named SPMs, in enhancing self-limitation and resolution of chronic inflammation and, more recently, of tissue fibrosis, is being recognized: SPMs act on key molecular pathways to restore a normal cell homeostasis and tissue architecture in different organs, including the steatotic liver¹¹(**Table 3**).

These SPMs are also synthesized from omega-3 PUFAs EPA and DHA by the coordinate action of 5-, 12- and 15-LOX, and include lipoxins (LXs, derived from lipoxygenase interaction products), resolvins(Rvs, derived from resolution phase interaction products), which were classified as either E-series Rvs(if generated from EPA) or D-series Rvs (if derived from DHA)), protectins(PDs), and maresins(MaRs, from macrophage mediators in resolving inflammation), which derive from DHA⁷ (**Figure 4**).

Data from diet-induced models of NASH and cell cultures disclosed an imbalance between proinflammatory eicosanoids and anti-inflammatory/pro-resolving SPMs in the liver and adipose tissue as a central pathogenic feature of liver injury in NASH, and suggest the restoration of a normal pro/anti-inflammatory balance as a feasible and effective therapeutic target11^{,111,112}.Current approaches aim either at antagonizing proinflammatory eicosanoids or at enhancing SPM actions.

Antagonizing proinflammatory eicosanoids

1) Group IVA phospholipase A2 inhibition

Phospholipases A2 (PLA2s) catalyzes the hydrolysis of the sn-2 fatty acid subsistent from membrane glycerophospholipids, generating lysophospholipids and arachidonic acid (AA), a substrate for the synthesis of eicosanoids prostaglandins (PGs) and leukotrienes (LTs) by cyclooxygenases and lipoxygenases¹¹³(**Figure 4**).

The over 20 isozymes of mammalian PLA2s identified to date have been classified into intracellular (groups IV and VI PLA2s) and secretory forms(groups I, II, III, V, X, and XII PLA2s). Among them, the groupIV calcium-dependent cytosolic PLA2 α (PLA2 α), received most attention because of its central role in AA generation, its broad expression, and its active participation in cell metabolism⁶³.

Although groupIV PLA2 α is constitutively expressed in most cells and tissues, several posttranslational mechanisms regulate its activity under different pathophysiological conditions: beside Ca(2+) binding to its C2 domain, groupIV PLA2 α is functionally linked to cell surface receptors, that modulate its activity through phosphorylation by protein kinases, including ERK1/2 and p38 MAPK, S-nitrosylation through interaction with nitric oxide (NO), and interaction with proteins[vimentin , cPLA2-interacting protein (PLIP)] and phospholipids⁶³. Therefore, diverse extracellular signaling molecules, including pro-inflammatory and pro-fibrotic cytokines (IL-1, TNF- α , Angiotensin II)and growth factors(EGF), converge on group IV PLA2 α , whch catalizes the first, rate-limiting step in the AA cascade. Mounting evidence suggest a critical role for group IVPLA2 α in the onset of obesity-related disorders. In the liver and adipose tissue, group IVPLA2 α is required to synthesize AA-derived PGs and LTs by COXs and LOXs (discussed below)(**Figure 4**).

A central role for group IV PLA2 α in hepatic lipid metabolism emerged from high fat diet(HFD)induced rodent models, where genetic group IV PLA2 α deletion prevented the development of NASH and, more intriguingly, pharmacological inhibition of groupIV PLA2 α with orally active compounds, including the ω 3-polyunsaturated fatty acid derivatives AVX001 and AVX002 and the indole derivative ASB14780, reversed established steatohepatitis and fibrosis^{49,117}. AVX001 was safe and effective in patients with mild-to-moderate psoriasis¹¹⁴ but has not been evaluated in NASH (**Table 2**).

2) Cyclooxygenase-2 (COX-2) pathway inhibition

COX is a membrane-bound byfunctional enzyme that catalyzes the first two committed steps in the pathwayleading to the formation of PGs and TX, namely cyclooxygenation and peroxidation There are 2 COX isoforms: COX-1 is ubiquitous and isconstitutively expressed throughout the gastrointestinalsystem, the kidneys, the vascular smooth muscle andplatelets, while COX-2 is undetectable in most tissues, but its expression can be induced by a variety of stimuli related to inflammatory response and is therefore referred to as the inducible COX isoform. However, this distinction is not completely true, as COX-1 can be induced undercertain conditions and COX-2 has been shownto be constitutively expressed in the brainand the kidneys⁴³.

In methionine-choline deficient (MCD) diet and HFD-induced NASH models, hepatic and adipose COX-2 expression is significantly increased, as a result of NF- κ B and IL-1 axis activation, and correlates with the severity of steatohepatitisIn^{115,116}.

The effects of COX-2 activation on liver disease appear largely mediated by the generation of PGs, including15-deoxy-PGI₂ and PGE₂, which enhance triglyceride storagenecro-inflammation and fibrosis by a variety of mechanisms: autophagy inhibition¹¹⁷, enhancement of lipid droplet

formationandadipogenesis¹¹⁸, increased chemokine MCP-1 secretion and TGF- β 1-induced HSC activation (**online supplementary Table 3**).

Intriguingly, liver histology, adipose tissue inflammation and metabolic abnormalities were improved to a similar extent by celecoxib, a selective COX-2 inhibitor, or by PGE receptor antagonists, indicating that the effects of COX-2 activation in NASH are largely PGE2mediated^{115,119}. PGE2 induces hepatocyte TG accumulation and apoptosis, promotes Kupffer cell activation and adipose tissue inflammation and dysfunction^{120, 121, 122, 123,124} (online supplementary **Table 3**); furthermore, growing evidence suggests COX-2-driven PGE-2 biosynthesis promotes obesity-associated hepatocellular carcinoma and breast cancer through Prostaglandin E receptor 4(PTGER4)-mediated suppression of antitumor immunity and enhanced transcription of CYP-19, CYP-181 and aromatase-catalyzed estrogen biosynthesis¹²⁵(online supplementary **Table 3**)). Several issues regarding the safety of COX-2 inhibition remain: PGE2 enhances macrophage polarization switch from a pro-inflammatory M1 to a pro-resolving M2 phenotype in adipose tissue, thereby initiating inflammation resolution^{126,127}.

Secondly, COX-2 is constitutively expressed in endothelium and kidney, where it contributes to the synthesis of the protective PGI2 and it is still debated if COX-2 inhibition increases CVD risk¹²⁸.

3) 5-lipoxygenase (5-LOX)/leukotriene pathway inhibition

Arachidonate 5-lipoxygenase (5-LOX) is a calcium-requiring, ATP-requiring, iron-requiring enzyme that catalyses the two-step lipoxygenation of arachidonic acid to formbioactiveproinflammatory lipids leukotriene (LT) LTB4 and cysteinyl-leukotrienes (Cys-LTs) LTC4, LTD4 and LTE4¹¹. Furthermore, 5-LOX is also involved with 15-LOX in the synthesis of Lipoxins LXA4 and LXB4 which have anti-inflammatory and pro-resolving properties (see below)⁷(**Figure 4**). 5-LOX is expressed primarily in inflammatory cells including polymorphonuclear leukocytes, eosinophils, monocytes, mast cells, and B-lymphocytes, where LTs exert potent pro-inflammatory actions bypromoting recruitment and chemotaxis, partly through NF-κB activation¹¹¹.

In the liver, Kupffer cells express 5-LOX and synthesize LTB4 and cysteinyl-LTs, the latter being also produced by hepatocytes through transcellular metabolism of LTA4 secreted by Kupffer cells¹²⁹. LTspromoteKupffer cell and HSC activation. A similar role for adipocyte 5-LOX in mediating adipose tissue inflammation has been found in experimental models of obesity¹²⁹. Notably, the adipose tissue and liver from mice with high-fat diet (HFD)-induced NASH showed increased expression of 5-LOX and its products¹³⁰, and in humans the progression from healthy liver to NASH is paralleled by an increased formation of 5-LOX products⁶. Furthermore, functional 5-LOX manipulation studies corroborate a causal role for 5-LOX in liver injury: genetic deletion of 5-LOX protected from HFD-induced obesity, insulin ressitance and NASH^{131, 132}, and pharmacological 5-LOX inhibition activated AMPK activation, reduced NF-kB activation and lipolysis and improved insulin resistance, steatosis and necroinflammatory changes in NASH^{129,131}. On this basis, an orally available small molecule 5-LOX inhibitor, MN-001 (tipelukast), which acts also as LTD4 receptor antagonist and phosphodiesterases (PDE) 3/4 inhibitor, reduced inflammation and fibrosis and proinflammatory and profibrogenic gene expression in an advanced NASH model¹³³ and was FDA-approved for a Phase IIa RCT in NASH patients with advanced fibrosis¹³⁴.

Beside 5-LOX inhibition, another strategy to antagonize the proinflammatory actions of LTs is receptor antagonism, and, intriguingly, inhibition of the LTB4 receptor 1 (Ltb4r1) achieved the same anti-inflammatory and insulin sensitizing effects in the liver and adipose tissue as 5-LOX inhibition¹³⁵, indicating that specific LT receptor antagonism could achieve the benefits of 5-LOX inhibition while preserving 5-LOX-mediated synthesis of pro-resolving mediators lipoxins.

Enhancing resolution of inflammation and fibrosis with specialized proresolving mediators (SPMs)

It is being increasingly recognized that the resolution of inflammation is not a passive phenomenon, deterimined by dilution and dissipation of inflammatory mediators, but an active process which is

orchestrated by a temporally regulated, sequential secretion of pro-resolving molecular signals which lead to self-limitation and resolution of the inflammatory response and restore tissue function^{7,136}. Tightly connected to inflammation, , liver fibrosis is currently considered a wound healing response to chronic, unresolved tissue injury, which is primarily driven by inflammatory and immune mediated mechanisms². Experimental models indicate liver fibrosis is not an unidirectional, irreversible process but can be reversed through four steps, which may coexist independently of each other: cessation of chronic damage, shifting the balance from inflammation to resolution , deactivation of myofibroblasts, matrix degradation (reflected by an altered balance between matrix stabilizing and matrix degrading factors)²(**Table 3**).

Unlike their precursors ω -3 PUFAs, which act at the micromolar to millimolar range, SPMs exert their biological actions in the picomolar to nanomolar range, with concentrations as low as10nM producing a 50 percent reduction in PMN transmigration in models systems¹³⁷.

SPMs not only attenuate the inflammatory response but also expedite its resolution by shifting the key cells of innate and adaptive immunity, i.e., macrophages and T cells, from a pro-inflammatory to a pro-resolving phenotype: SPMs enhance monocyte migration, macrophage polarization into a M2 phenotype¹³⁸, promoting macropage autophagy¹³⁹ and clearance of apoptotic cells (efferocytosis), and shift CD4⁺ T cell differentiation from a pro-inflammatory T_h1/T_h17 to a pro-resolving T_{reg} phenotype¹⁴⁰.

In HFD-induced rodent models of NASH, in precision-cut liver slices and in cultured primary hepatocytes Kupffer cells and adipocytes, Maresin 1(MaR1) administered at physiological, nanomolar concentrations prevented palmitate- and hypoxia-induced ER stress and apoptosis and enhanced Kupffer cell phagocytosis¹¹².

Recent studies indicate that in obesity-associated NASH the formation of SPM in the inflamed liver and adipose tissue is severely deregulated¹⁴¹, and that the administration of SPMs ameliorates inflammatory metabolic and histological features of obesity-related NASH through mechanisms strictly involved in NASH pathogenesis, inclusing insulin sensitivity, triglyceride accumulation, cell

apoptosis and fibrogenesis.More specifically, LXA4, RvD1 RvE1, Protectin D1 (PD1) and MaR1 improved adipose tissue inflammation and insulin resistance and hepatic fat infiltration and insulin resistance and reduced hepatocyte ER stress-induced apoptosis through Akt and AMPK activation and JNK inhibition; and PD1 and MaR1 increased expression of adiponectin, a key anisteatotic, anti-inflammatory and antifibrogenic adipokine, to a similar extent as thiazolidinediones^{142, 143, 144, 145, 146}

SPM demostrated also potent anti-fibrotic properties preclinically: Resolvin D1, Resolvin E1, Protectin DX and MaR1 attenuated fibrosis progression and, more intriguingly, reversed established fibrosis in preclinical models of hepatic, renal and pulmonary fibrosis^{147, 148}, at least in part through suppressing and reversing TGF- β 1/Smad2/3-induced epithelial-to-mesenchimal transition (EMT) of epithelial cells, which provide up to 40% of extracellular matrix-deposing myofibroblasts, and restoring a normal tissue levels of Matrix metalloproteinase(MMPs), which are crucial for ECM resorption^{149, 150,151,152}.

A key issue to the clinical development of SPMs is the achievement of therapeutically effective SPMs concentrations to activate resolution signals in target tissues, including the live and adipose tissue, as SPMs are rapidly inactivated by eicosanoid oxidoreductases and leukocyte receptors for SPMs appear to be down-regulated in obesity-related conditions¹⁰⁴.

Pharmacological strategies to enhance biological activity and selective delivery of SPMs to target organs, are being investigated: thesynthetic oxidoreductase-resistant RvD1 analogue benzodiacetylenic-17R-RvD1-methyl ester (BDA-RvD1) showed a 3.5-fold higher potency than the natural compound in resolving ischemia-reperfusion-induced lung injury¹⁵³, while the incorporation of SPMs into liposomes (e.g., Lipo-RvD1) should enhance selective clearance by macrophages¹⁵⁴.

A RCT evaluating safety and efficacy of RX10045, a synthetic RvE1 analog, on dry eye disease, has been completed (ClinicalTrials.gov ID: NCT00799552).

Concluding remarks and future directions

The health-related burden of NASH is increasing all over the world along with the obesity epidemic. There are no approved treatments for NASH, as lifestye intervention is often hampered by low patient compliance and requires long-term committent. Proposed pharmacological options do not seem to reverse more advanced stages of disease and therefore are unlikely to change the natural history of the disease or are encumbered by unwanted effects¹⁵⁵. Treatment of NASH is challenging, as progression from steatosis to NASH and fibrosis is likely a multi-factorial process, involving varied molecular pathways that may operate in different patient subsets at different stages of the disease. In the last decade our knowledge of NASH as a lipotoxic disease has considerably expanded the spectrum ofnew lipid species and pathways involved in NAFLD progression. These advances in basic science are expected to make up to clinical stage of development, posingatwofold challenge for future research: first, the discovery of safe and effective pharmacological tools to antagonize the diverse lipotoxic species. Secondly, once a variegated pharmacological armamentariumis available, will be the selection of the optimal therapeutic strategy for each patient: in this context recent advances inlipidomics, genetics and epigenetics systems biology technologies will likelyenable treatments tailored to individual genomic and lipidomic profiles¹⁵⁶. As a example, the I148M variant of patatin-like phospholipase domaincontaining 3 (PNPLA3)protein, an intracellular lipase involved in lipid droplet (LD) remodelling, ia a major genetic determinant of the full spectrum of NAFLD across different ethnicities and of response to PUFA treatment¹⁵⁷, independently of traditional metabolic risk factors¹⁵⁸. The mutant (I148M) variant protein, with loss of enzymatic activity, accumulates on LD surface by evading proteasomaldegradationandimpairs triglyceride mobilization from the LD by other lipases¹⁵⁹: in hepatic stellate cells this impaired the release of retinol and lipids from lipid droplets, resulting in a more fibrogenic phenotype¹⁶⁰, while the precisemolecular mechanisms underlying lipotoxicity of this variant in hepatocytes remain to be defined. A potentialtherapeuticimplication of recent advances in PNPLA3 biology could be the development of agents that decrease PNPLA3 protein

levels and improve liver disease by reducing the impairment in LD remodeling in NASH carriers of the PNPLA3 I148M mutant allele, in an individualized pharmacogenomic approach.

FIGURE LEGENDS

Figure 1. Mechanisms of saturated fatty acid (SFA) lipotoxicity.

Panel A. SFAs possess substantial lipotoxicity for hepatocytes through a variety of mechanisms: they can bind to plasma membrane Toll-Like Receptor (TLR)-4 and to death receptor TNF-related apoptosis-inducing ligand (TRAIL)-R2, inducing caspase 8 activation, which cleaves Bid into its truncated form (tBid) and triggers caspase-dependent hepatocyte apoptosis.

SFA can also enter hepatocyte and trigger ceramide-dependent inflammasome-mediated proinflammatory cytokine secretion, endoplasmic reticulum (ER) stress, mitochondrial dysfunction, and c-Jun N-terminal kinase (JNK) activation.

JNK inactivates Insulin Receptor Substrate (IRS)-1, inducing insulin resistance, interacts with the outer membrane mitochondrial protein SH3BP5 (Sab) to impair respiration, and induces reactive oxygen species (ROS) generation.

Furthermore, JNK suppresses Peroxisome Proliferator-Activated Receptor (PPAR)-α-mediated Fibroblast Growth Factor (FGF)-21 expression and mitochondrial and peroxisomal β-oxidation, and activates the proapoptotic proteins p53-upregulated modulator of apoptosis (PUMA) and Bim. PUMA and Bim belongs to the Bcl2-2 Homology(BH3)-only protein family: both proteins are major mediators of SFA-induced lipoapoptosis through Bax activation, resulting in mitochondrial outer membrane permeabilization (MOMP) with cytochrome c liberation and in lysosome permealization with cathepsin B release. SFA and and SFA-derived LPC and ceramide all lead to the release of Extracellular vescicles (EVs) from hepatocytes. Recent studies have defined multiple roles of lipotoxic EVs in lipotoxicity and NASH pathogenesis through cell-to-cell communication via various cargoes. CXCL10 and ceramide-enriched EVs mediate monocyte/macrophage chemotaxis to the liver, TRAIL-enriched EVs contribute to macrophage activation, and miR-128-3p-laden EVs enhance HSC proliferation and activation in vitro.

Panel B. Mechanisms of SFA lipotoxicity on nonparenchimal liver cells.

Nonparenchimal cells are also involved in SFA lipotoxicity: SFAs activate TLR-4 on hepatic stellate cells (HSCs) to secrete the chemokine monocyte chemoattractant protein(MCP)-1 and activate JNK in Kupffer cells and macrophages to induce proinflammatory M1 polarization, chemotaxis and secretion of profibrogenic factors Transforming Growth Factor(TGF)-ß and Tissue Inhibitor of Metalloproteinase(TIMP)-1

Abbreviations: CXCL10: C-X-C motif ligand; MOMP: mitochondrial outer membrane permeabilization; PUMA: p53-upregulated modulator of apoptosis; CHOP: CAAT/enhancer binding homologous protein; JNK: c-Jun N-terminal kinase; LPC: lysophosphatidylcholine; LPA: lisophosphatidic acid; TRAIL: TNF-related apoptosis-inducing ligand; BID, pro-apoptotic BCL-2 interacting domain; CHOP, CCAAT/enhancer-binding homologous protein; Bim, Bcl-2 protein family member; SFA, saturated fatty acid; FoxO3a, forkhead box-containing protein, class O member 3a; TNFa, tumor necrosis factor a; NF-kB: Nuclear Factor-kB; ROS, reactive oxygen species; ER, endoplasmic reticulum; MCP: monocyte chemoattractant protein.

Figure 2. Metabolic circuits of Ceramide (CER), sphingosine (Sph) and Sphingosine-1-Phosphate (S1P) synthesis and catabolism (the "sphingolipid rheostat") and cellular targets of sphingolipid lipotoxicity.

Panel A. Ceramide can be synthesized *de novo* from serine and palmitate by the sequential action of three ER-resident enzymes—serine palmitoyltransferase (SPT), ceramide synthase (CeS) and dihydroceramide desaturase (DES). — or generated from hydrolysis of plasma membrane sphingomyelin into ceramide and phosphocholineby the enzyme sphingomyelinase. Once generated from ceramide deacylation, sphingosine can be phosphorylated by sphingosine kinases (SphKs) to form S1P. There are two pathways of S1P degradation: reversible dephosphorylation to sphingosine by nonspecific phosphatases, and by two S1P-specific phosphatases, SPP1 and SPP2; and irreversible cleavage by S1P lyase (SPL), which leads to the formation of phosphatidylethanolamine and hexadecenal. The latter is the only exit pathway for degradation of sphingoid bases in mammalian cells.

Ceramide inhibits Akt phosphorylation and activation, thereby impairing Akt-mediated insulin signaling and promoting insulin resistance, an effect that is antagonized by S1P. The mitochondria are another major cellular target of ceramide, which impairs fatty acid β-oxidation through inactivation of electron transport chain (ETC) complex II and IV and promotes ROS production and TG accumulation. Furthermore, ceramide triggers BAX-dependent mitochondrial membrane permeabilization and cytochrome c release, leading to apoptosis Further mechanisms of hepatic lipotoxicity of ceramide overload in NASH include disruption of calcium homeostasis in the ER, which leads to ER-stress-mediated apoptosis Ceramide synergizes with S1P to activate SREBP-1c and SREBP-2 to enhance *de novo* lipogenesis and cholesterol synthesis, and NF-kB and Nlrp3 inflammasome to induce proinflammatory cytokine and chemokine secretion.

Further mechanisms of lipotoxicity include impairment of autophagy, and upregulation of hepatic hepcidin, which leads to hepatic iron overload. Additionally, ASMAse activation may promote liver injury independently of ceramide accumulation by disrupting methionine and phosphatidylcholine metabolism, which promotes lysosomal membrane permeabilization and directly activates HSCs.

Panel B.

Recent data indicate the intestine is a relevant source of ceramide, whose synthesis in ileum and cecum is stimulated by the activation of a bile acid/intestinal FXR axis: the activity of FXR in the epithelial cells of the ileum is inhibited by different intestinal bile acids, including cholic acid (CA), chenodeoxycholic acid (CDCA) and β -muricholic acid β -MCA), while FXR activation upregulates ceramide synthesis.

In adipocytes, ceramide impairs adipose function through increase ER stress, decreasing the ration of beige to white adipocytes SphK1 activation induces an increase in secretion of TNF- α and IL-6 and inhibits secretion of adiponectin and IL-10, promoting inflammation and insulin ressitance. Furthermore, in adipocytes and macrophages, ceramide activates Nlrp3 inflammasome to induce proinflammatory cytokine and chemokine secretion.

In HSCs, SIPR(1, 3) activation induces activation, migration and HSC transdifferentiation into myofibroblasts

Finally, SIPR1 activation in endothelial cells enhances integrity and barrier function

Figure 3. Metabolism of. Scheme of short-, long- and n-6 and n- 3 PUFA biosynthesis leading to membrane phospholipids synthesis.

The metabolism of PUFA is a complex process involving several enzymes of desaturation, elongation, and β -oxidation. Shown here is the pathway of both n-6 and n-3 PUFA metabolism to more unsaturated, long-chain members of each family.

The long chain saturated fatty acids and unsaturated fatty acids of the n-3, n-6, n-7 and n-9 series can be synthesized from myristic acid (C14:0) and palmitic acid (C16:0). Long-chain fatty acids of the n-6 and n-3 series can also be synthesized from precursors obtained from dietary precursors to elongation (ELOVL) and desaturation steps as indicated in these pathways. **Abbreviations**. ACC: acetyl-CoA carboxylase; ELOVL: elongase of very
long chain fatty acid; FASN: fatty acid synthase; FADS: fatty acid desaturase; SCD: stearoyl-CoA desaturase.

Figure 4. Lipids derived from arachidonic acid (AA), eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA).

AA is metabolized by cyclooxygenases (COX) 1/2 to prostaglandins(PGs) and thromboxanes(TXs) and by 5-lipoxygenase (5-LOX) to leukotrienes (LTs) which are involved in the initiation of the inflammatory response (red colour).

Hydroxyeicosatetraenoic acids (HETEs) and lipoxins are also synthesized from

arachidonic acid and here 5-, 12- and 15-LOX and cytochrome (Cyt) P450 are involved.

Eicosapentaenoic acid is metabolized to 3-series PGs by COX and 3-series TXs by 5-LOX (weak

pro-inflammatory properties, pink colour) and to E-series resolvins (anti-inflammatory and pro-

resolving actions, green colour) by CYP450 and 5-LOX.

Resolvins of the D-series, protectins and maresins are derived from docosahexaenoic acid.

Lipoxins, resolvins, protectins and maresins have anti-inflammatory and pro-resolving actions.

Author	Population	Biological	Technique	Findings in NASH relative to NAFL
	(n)	Sample		
Puri 2007 ⁵	Non-cirrhotic	Liver	TLC	↓ PC
	NASH(n=9)			↑lysoPC
	NAFL(n=9)			↓DHA/EPA/AA
	Control(n=9)			↑ω-6/ω-3 PUFA ratio
				↑FC

Table 1. Main lipidomic studies in NAFLD patients

Puri 2009 ⁶	Non-cirrhotic	Plasma	TLC	↓DHA//DPA ratio
	NASH(n=50)			↓ MUFAs
	NAFL(n=25)			↓ plasmalogens
	Control(n=50)			↑5-, 8-, 11-, 15-HETE
Gorden	Cirrhosis (n=20)	Plasma	LC-MS	↓ lysoPE
2015 ⁷	NASH(n=20)			↑ PE
	NAFL(n=17)			↑ Cer, DH-Cer and DH-deoxyCer
	Control(n=31)			↓ Sph
Zhou 2016 ⁸	NASH(n=69)	Plasma	UPLC-MS	↑SFA(14:0/16:0/18:0)
	NAFL(n=117)			↑MUFA(44:1/54:1)
	Non-NASH(n=249)			↓ sphingomyelin
	Non-NAFL(n=132)			↓lysoPC
Chiappini	NASH(n=15)	Liver	LC-MS	↑SFA(14:0/16:0/18:0)
2017 ¹⁰	NAFL(n=39)			↑MUFA(16:1/18:1)
	Control(n=7)			↑ω-6/ω-3 PUFA ratio
				↑Cer(C16/C18)
				↓phospholipids (PC/PE/PI/PS)
				↑ sphingomyelin

Abbreviations: AA: arachidonic acid; TLC: thin-layer chromatography; FC: free cholesterol; PC:

phosphatidylcholine; PE: phosphatidylethanolamine; PI: phosphatidylinositol; PS:

phosphatidylserine; Cer: ceramide; DH: dihydro; DHA: docosahexanoic acid; DPA:

Docosapentaenoic acid; EPA: Eicosapentanoic acid; LysoPC: lysophosphatidylcholine; HETE:

hydroxyeicosatetraenoic acid; LC: liquid chromatography; MS: mass spectrometry;

Sph: sphingosine; UPLC: ultra-performance liquid chromatography.

Molecular	Agent	Biological effect	Furthest de	evelopmental stage in
target			NASH	
			Preclinical	Clinical
				(RCT ID number)
LXR	25HC3S	LXR/SREBP-1c inhibition	+	Ib:
		NF-KB inhibition		ACTRN12615000267
				550
FXR	Obeticholic	FXR activation	+	III:
	acid			Clinicaltrials.gov ID:
				NCT02548351
	Px-104	FXR activation	+	lla:
				Clinicaltrials.gov ID:
				NCT0199910
	LMB763	FXR activation	+	IIa:
				Clinicaltrials.gov ID:
				NCT02913105
		EVD activation		Hat
	Gs-9674	TAK activation	T	11a.
				Clinicaltrials.gov ID:
				NCT02854605
PPAR-α/δ	Elafibranor	PPAR-α/δ activation	+	IIa:
				Clinicaltrials.gov ID:
				NCT01694849
	MRV	DDAD Spatiantian	Droolinical	
ГГАК-0	WIDA-	rrak-o activation	ricciniicai	-
	8025(selade			

Table 2. Molecular targets and pharmacological agents targeting lipotoxicity in NASH

	par)			
ΡΡΑΚ-α/γ	Saroglitazar	PPAR- α/γ activation	+	IIa:
				CTRI/2010/091/00010
				8
PPAR-α/δ/γ	IVA337	PPAR- $\alpha/\delta/\gamma$ activation	preclinical	-
	(Lanifibran			
	or)			
ACC1/2	GS-0976	ACC1/2 inhibitor	+	IIa
				Clinicaltrials.gov ID:
				NCT02856555
DGAT-1	Pradigastat	DGAT-1 inhibitor	+	IIa
				Clinicaltrials.gov ID:
				NCT01811472
PLA2	ASB14780,	PLA2 inhibition	Preclinical	-
	AVX001,			
	AVX002			
Autotaxin	PF8380	Autotaxin inhibitor	Preclinical	-
Ceramide	fenretinide	DES-1 and CeS5 inhibitor	Preclinical	-
	ST1072	CeS(4,6) inhibitor	Preclinical	-
	myriocin,	SPT inhibitor	Preclinical	-
	FTY720			
	Gly-MCA	Intestinal FXR inhibition	Preclinical	-
	FIASMAs:	Inhibitors of ceramide	Preclinical	-
	TCAs,	hydrolysis from sphingomyelin		
	SSRIs,	by SMase		

	CCBs			
S1P	fingolimod	S1P type 1 receptor (SIPR1)	Preclinical	-
		antagonist		
	VPC23019	S1PR(1,3)) antagonist	Preclinical	-
ω-3 PUFA	ω-3 fish oil	ω-3 PUFA supplementation	+	IIa*:
				ClinicalTrials.gov:
				NCT00681408,
				NCT01154985,
				NCT01992809
SCD-1	aramchol	SCD-1 inhibitor	+	IIa:
				NCT01094158,
				NCT02279524
5-LOX and	tipelukast	5-LOX inhibitor, LTD4 receptor	+	IIa:
LTD4		antagonist and PDE 3/4 inhibitor		NCT02681055
LTB4	CP105696	LTB4 receptor antagonist	Preclinical	-
Resolvin D1	BDA-RvD1	RvD1 analogues	Preclinical	-
	Lipo-RvD1			
Resolvin E1	RX10045	RvE1 analogue	Preclinical	-

Abbreviations: ACC: Acetyl-CoA carboxylase; TCA: tricyclic antidepressants; SSRI: selective serotonin-reuptake inhibitors; CCBs: calcium channel blockers; 25HC3S: 25-Hydroxycholesterol-3-sulfate; PLA2: phospholipase A2; CeS: ceramide synthase; DES: Dihydroceramide desaturase; S1P: Sphingosine-1-Phosphate; SPT: serine palmitoyltransferase; SMase: sphingomyelinase; SCD-1: Stearoyl Coenzyme A Desaturase-1; LOX: lipoxygenase; LT: leukotriene; PDE: phosphodiesterase,

* only RCT with post-treatment liver histology are reported.

Table 3. Involvement of Specialized Proresolving Mediators(SPMs) in the different steps of

 fibrosis resolution in NASH

Step 1: Cessation of chronic injury to the liver (allowing hepatocyte recovery and modulating the microenvironment)					
SPM	Molecular mechanism	Cellular pathway	Ref		
MaR1	Hepatocyte:	\downarrow JNK activation $\rightarrow \downarrow$ apoptosis	112		
RvD1	\downarrow PA- and hypoxia-induced ER	\downarrow CHOP activation $\rightarrow \downarrow$ apoptosis	146		
	stress	↑ antiapoptotic miRNA signature			
	↓ SREBP-1c activation	↑ mitochondrial β-oxidation \rightarrow ↓ steatosis	143		
	\uparrow PPAR-α, and PGC-1α	↓ <i>de novo</i> lipogenesis→↓steatosis			
	activation \rightarrow \uparrow mitochondrial				
	function				
	↑AMPK phosphorylation				
	↑ Akt phosphorylation				
	↑ autophagy				
RvD1-D6	Adipocyte:	↓ secretion of pro-inflammatory adipokines	112		
RvE1-E2-	↑AMPK phosphorylation	(IL-1, TNF-α, IL-6)	144		
ЕЗ,	↑ Akt phosphorylation	↑ secretion of adiponectin	145		
MaR1,	↑ autophagy	\uparrow insulin sensitivity and FFA oxidation			
PD1	\uparrow PPAR- γ expression	\downarrow lipolysis $\rightarrow \downarrow$ flow of toxic SFA to the liver			
	↑ IRS-1/IRS-2 expression				
	↑ GLUT-2/-4 expression				
Step 2. Intra	ahepatic balance switch from pro-in	flammatory to restorative(phenotypic adjustments of in	nmune cells,		
especially induction	of restorative macrophages),				
SPM	Molecular mechanism	Cellular pathway	Ref		
MaR1	Kupffer cell:	↑efferocytosis of apoptotic cells and cellular	112		
	\downarrow PA- and hypoxia-induced ER	debris			
	stress	\downarrow LTB4 and TNF- α secretion			
LXA4	Macrophage:	↑ MAPK/HSP27 activation \rightarrow ↑ IL-10	138		
15-epi-	Switch from a M1 pro-	secretion	139		
LXA4	inflammatory to a M2 pro-resolving	↑PI3K/Akt signaling→↑ efferocytosis			
RvD1	phenotype	\downarrow MCP-1, TNF- α , IL-1 secretion			
RvE1	↑ autophagy				
PD1					

RvD1-D2	Adaptive immune T cells	Switch CD4 ⁺ T cell differentiation from a	140
MaR1	Shift from a pro-inflammatory to a	pro-inflammatory $T_h 1/T_h 17$ phenotype to a	
	pro-resolving phenotype	pro-resolving T _{reg} phenotype	
		\downarrow TNF- α and IL-17 secretion	
		↑ IL-22 secretion	
Step 3. Deac	tivation of myofibroblasts (by senescence, a	apoptosis and inactivation)	
SPM	Molecular mechanism	Cellular pathway	Ref
LXA4,	↓TGF-β1/Smad2/3-axis activation	\downarrow epithelial-to-mesenchimal transition	149
RvD1,		(EMT) and HSC activation	150
RvE1,			151
MaR1			152
	Step 4. Extracellula	ar matrix degradation	
SPM	Molecular mechanism	Cellular pathway	Ref
RvD1,	Kupffer cells, macrophages,	↓TIMP-1 secretion and ↑ MMP-9/12 activity	148
RvE1,	HSCs	→↑ ECM degradation	151
PD1	restored balance between matrix stabilizing and matrix degrading factors/enzymes		152

Abbreviations: ACC: acetyl-CoA carboxylase; FAS: fatty acid synthase; PA: palmitate; CHOP: CCAAT/enhancer-binding protein homologous protein; SREBP-1, \uparrow PPAR- α , and PGC-1 α ; CPT-1a: carnitine palmitoyltransferase, ACOX-1: acyl-coenzyme A oxidase JNK: c-Jun N-terminal kinase;

REFERENCES

1.Chalasani N, Younossi Z, Lavine JE, et al. The diagnosis and management of nonalcoholic fatty liver disease: Practiceguidance from the American Association for the Study of Liver Diseases.Hepatology. 2018;67:328-357

2.Musso G, Cassader M, Gambino R.Non-alcoholicsteatohepatitis: emerging molecular targets and therapeutic strategies. Nat Rev Drug Discov. 2016;15:249-74

3.Goldberg D, Ditah IC, Saeian K, et al. Changes in the Prevalence of Hepatitis C Virus Infection, Non-alcoholicSteatohepatitis, and Alcoholic LiverDiseaseAmong Patients with Cirrhosis or Liver Failure on the Waitlist for Liver Transplantation. Gastroenterology.2017;152:1090-1099

4. Yamaguchi K, Yang L, McCall S, et al. Inhibiting triglyceride synthesis improves hepatic steatosis but exacerbates liver damage and fibrosis in obese mice with nonalcoholic steatohepatitis. Hepatology 2007;45:1366-1374.

5.Puri P, Baillie RA, Wiest MM, et al. A Lipidomic Analysis of Nonalcoholic Fatty Liver Disease. Hepatology 2007;46:1081-1090

6.Puri P, Wiest MM, Cheung O, et al. The plasma lipidomic signature of nonalcoholic steatohepatitis. Hepatology 2009;50:1827–1838.

7.Gorden DL, Myers DS, Ivanova PT, et al. Biomarkers of NAFLD progression: a lipidomics approach to an epidemic.J Lipid Res. 2015;56:722-36.

8.Zhou Y, Oresic M, Leivonen M, et al. Noninvasive Detection of Nonalcoholic Steatohepatitis Using Clinical Markers and Circulating Levels of Lipids and Metabolites. Clin Gastro Hep 2016;14:1463–1472

9.Chiappini F, Desterke C, Bertrand-Michel J, et al. Hepatic and serum lipid signatures specific tononalcoholic steatohepatitis in murine models. Sci Rep. 2016;6:31587

10.Chiappini F, Coilly A, Kadar H, et al. Metabolism dysregulation induces a specific lipid signature of nonalcoholic steatohepatitis in patients.Sci Rep. 2017;7:46658.

11Serhan CN. Treating inflammation and infection in the 21st century: new hints from decoding resolution mediators and mechanisms.FASEB J. 2017;31:1273-1288.

12.Musso G, Gambino R, Cassader M. Cholesterol metabolism and the pathogenesis of nonalcoholic steatohepatitis.Prog Lipid Res. 2013;52:175-91

13.Donnelly KL, Smith CI, Schwarzenberg SJ, et al. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. J Clin Invest. 2005;115: 1343–1351.

14.Stremmel W, StafferS, Wannhoff A, et al. Plasma membrane phospholipase A2 controls hepatocellular fatty acid uptake and is responsive to pharmacological modulation: implications for nonalcoholic steatohepatitis. FASEB J. 2014; 28: 3159–3170.

15.Win S, Than TA, Zhang J, et al. New insights into the role and mechanism of c-Jun-N-terminal kinase signaling in the pathobiology of liver diseases.Hepatology. 2017 Nov 30. doi:

10.1002/hep.29689.[Epub ahead of print]

16.Vernia S, Cavanagh-Kyros J, Garcia-Haro L, et al. et al. The PPARalpha-FGF21 hormone axis contributes to metabolic regulation by the hepatic JNK signaling pathway. Cell Metab. 2014; 20: 512–525.

17.Cazanave SC, Wang X, Zhou H, Rahmani M, Grant S, Durrant DE, et al. Degradation of Keap1 activates BH3-only proteins Bim and PUMA during hepatocyte lipoapoptosis. Cell Death Differ 2014;21:1303-1312.

18.Luo W, Xu Q, Wang Q, et al. Effect of modulation of PPAR-γ activity on Kupffer cells M1/M2 polarization in the development of non-alcoholic fatty liver disease.Sci Rep. 2017;7:44612.
19.Musso G, Gambino R, Cassader M. Musso G Recent insights into hepatic lipid metabolism in non-alcoholic fatty liver disease (NAFLD).Prog Lipid Res. 2009;48:1-26.

20.Higuchi N, Kato M, Shundo Y, et al. Liver X receptor in cooperation with SREBP-1c is a major lipid synthesis regulator in nonalcoholic fatty liver disease. Hepatol Res. 2008;38:1122-9

21.Ren S, Ning Y. Sulfation of 25-hydroxycholesterol regulates lipid metabolism, inflammatory responses, and cell proliferation. Am J Physiol Endocrinol Metab 2014;306:E123–30

22.Kim MJ and Lin WQ. DUR-928, an endogenous regulatory molecule, exhibits antiinflammatory and antifibrotic activity in a mouse model of NASH. Poster session presented at: AASLD's Emerging Trends in NAFLD; 2017 Mar 17-18

23.Neuschwander-Tetri B, et al. Farnesoid X nuclear receptor ligand obeticholic acid for noncirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial.Lancet. 2015;385:956-65

24.Haczeyni F, Wang H, Barn V, et al. The selective peroxisome proliferator–activated receptordelta agonist seladelpar reverses nonalcoholic steatohepatitis pathology by abrogating lipotoxicityin diabetic obese mice. Hepatol Comm 2017;1: 663-674

25.Staels B, Rubenstrunk A, Noel B, et al. Hepatoprotective effects of the dual peroxisome proliferator-activated receptor alpha/delta agonist, GFT505, in rodent models of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis.Hepatology. 2013;58:1941-52.

26.Ratziu V, Harrison SA, Francque S, et al. Elafibranor, an Agonist of the Peroxisome Proliferator-Activated Receptor-α and -δ, Induces Resolution of Nonalcoholic Steatohepatitis Without Fibrosis Worsening. Gastroenterology.2016;150:1147-1159.

27.Bays HE, Schwartz S, Littlejohn T, et al. MBX-8025, a novel peroxisome proliferator receptordelta agonist: lipid and other metabolic effects in dyslipidemic overweight patients treated with and without atorvastatin.J Clin Endocrinol Metab. 2011;96:2889-97 28.Tsukamoto H, She H, Hazra S, et al. Anti-adipogenic regulation underlies hepatic stellate cell transdifferentiation.JGastroenterolHepatol. 2006;21(S3):S102-5.

29.Joshi SR. Saroglitazar for the treatment of dyslipidemia in diabetic patients.Expert Opin Pharmacother. 2015;16:597-606

30.Wettstein G, Luccarini JM, Poekes L, et al. The new-generation pan-peroxisome proliferatoractivated receptor agonist IVA337 protects the liver from metabolic disorders and fibrosis. HepatolComm 2017; 1: 524-537

31.Banshi D. To Assess the Effect of 4mg Saroglitazar on Patients of Diabetes Dyslipidemia with Nonalcoholic Fatty Liver Disease for 24 Weeks. American Diabetes Association 75th Scientific Sessions, June 5 - 9, 2015, Boston, Massachusetts Abstract 712-P

32.Harriman G, Greenwood J, Bhat S, et al. Acetyl-CoA carboxylase inhibition by ND-630 reduces hepatic steatosis, improves insulin sensitivity, and modulates dyslipidemia in rats.Proc Natl AcadSci U S A. 2016;113:E1796-805.

33.Lawitz EJ, Poordad F, Coste A, et al. Acetyl-CoA carboxylase (ACC) inhibitor GS-0976 leads tosuppression of hepatic de novo lipogenesis and significant improvements in MRI-PDFF, MRE,and markers of fibrosis after 12 weeks of therapy in patients with NASH. J Hep 2017; 66(S1): S34

34.Irshad Z, Dimitri F, Christian M, et al. Diacylglycerol acyltransferase 2 linksglucoseutilization to fatty acidoxidation in the brown adipocytes. J Lipid Res. 2017;58:15-30.

35.Stone SJ, Myers HM, Watkins SM, et al. "Lipopenia and skin barrier abnormalities in DGAT2deficient mice". J Biol Chem 2004;279:11767–76.

36.Yamamoto Y, Utsunomiya H, Miyake T, et al. Upregulated absorption of dietary saturated fatty acids with changes of intestinal fatty acid transporters in non-alcoholic steatohepatitis. Hepatology 2015;62(S1):1252A-1253A

37.Sanyal AJ, Cusi K, Patel S, et al. Effect of Pradigastat, a Diacylglycerol Acyltransferase 1 Inhibitor, on Liver Fat Content in Nonalcoholic Fatty Liver Disease. Hepatology 2015;62(S1): 1253A

38.Li C, Li L, Lian J, et al. Roles of Acyl-CoA:Diacylglycerol Acyltransferases 1 and 2 in Triacylglycerol Synthesis and Secretion in Primary Hepatocytes. Arterioscler Thromb Vasc Biol. 2015;35:1080-91

39.Toriumi K, Horikoshi Y, YoshiyukiOsamuraR, et al. Carbon tetrachloride-induced hepatic injury through formation of oxidized diacylglycerol and activation of the PKC/NF-κB pathway. Lab Invest. 2013;93:218-29.

40.Takekoshi S, Kitatani K, Yamamoto Y. Roles of Oxidized Diacylglycerol for Carbon Tetrachloride-induced Liver Injury and Fibrosis in Mouse. Acta Histochem Cytochem. 2014;47:185-94.

41.Kakisaka K, Cazanave SC, Fingas CD, et al. Mechanisms of lysophosphatidylcholine-induced hepatocyte lipoapoptosis. Am. J. Physiol. Gastrointest. Liver Physiol. 2012;302:G77–G84.

42.Hollie NI, Cash JG, Matlib MA, et al. Micromolar changes in lysophosphatidylcholine concentration cause minor effects on mitochondrial permeability but major alterations in function.BiochimBiophysActa. 2014;1841:888-95.

43.Hirsova P, Ibrahim SH, Krishnan A, et al. Lipid-Induced Signaling Causes Release of Inflammatory Extracellular Vesicles From Hepatocytes. Gastroenterology. 2016;150:956-67

44.Ibrahim SH, Hirsova P, Tomita K, et al. Mixed lineage kinase 3 mediates release of C-X-C motif ligand 10-bearing chemotactic extracellular vesicles from lipotoxichepatocytes.Hepatology. 2016 ;63:731-44 45.Kaffe E, Katsifa A, Xylourgidis N, et al.Hepatocyteautotaxin expression promotes liver fibrosis and cancer.Hepatology. 2017;65:1369-1383

46.Li Z, Agellon LB, Allen TM, et al.The ratio of phosphatidylcholine to phosphatidylethanolamine influences membrane integrity and steatohepatitis.CellMetab. 2006;3:321-31.

47.Walker AK,Jacobs RL, Watts JL, et al. A conserved SREBP-1/phosphatidylcholine feedback circuit regulates lipogenesis in metazoans. Cell 2011; 147:840–852.

48.Payne F, Lim K, Girousse A, et al. Mutations disrupting the Kennedy phosphatidylcholine pathway in humans with congenital lipodystrophy and fatty liver disease. Proc Natl AcadSci U S A. 2014;111:8901-6.

49.Kanai S, Ishihara K, Kawashita E, et al. ASB14780, an Orally Active Inhibitor of Group IVA Phospholipase A2, Is a Pharmacotherapeutic Candidate for Nonalcoholic Fatty Liver Disease. J Pharmacol Exp Ther. 2016;356:604-14.

50.Rodriguez-Cuenca S, Pellegrinelli V, Campbell M, et al. Sphingolipids and glycerophospholipids - The "ying and yang" of lipotoxicity in metabolic diseases.Prog Lipid Res. 2017;66:14-29

51.Beckmann N, Sharma D, Gulbins E, **et al**. Inhibition of acid sphingomyelinase by tricyclic antidepressants and analogons. Front Physiol. 2014;5:331.

52Jiang C, Xie C, Lv Y, et al.: Intestine-selective farnesoid X receptor inhibition improves obesity-related metabolic dysfunction. Nat Commun 2015; 6: 10166.

53.Holland W, Bikman BT, Wang LP, et al.Lipid-induced insulin resistance mediated by the proinflammatory receptor TLR4 requires saturated fatty acid-induced ceramide biosynthesis in mice.JClinInvest. 2011;121:1858-70.

54.Raichur S, Wang ST, Chan PW, et al. CerS2 haploinsufficiencyinhibits b-oxidation and confers susceptibility to diet-inducedsteatohepatitis and insulin resistance. Cell Metab. 2014; 20: 687–695

55.Turpin SM, Nicholls HT, Willmes DM, et al. Obesity-induced CerS6-dependent C16:0 ceramide productionpromotes weight gain and glucose intolerance. Cell Metab.2014; 20: 678–686
56.Liu Z, Xia Y, Li B, et al. Induction of ER stress-mediated apoptosis by ceramide via disruption of ERCa(2+) homeostasis in human adenoid cystic carcinoma cells. Cell Biosci. 2014;4:71.
57.Cinar R, Godlewski G, Liu J, et al. Hepatic cannabinoid-1 receptors mediate diet-induced insulin resistance by increasing *de novo* synthesis of long-chain ceramides. Hepatology 2014;59:143-53
58.Fucho R, Martínez L, Baulies A, et al. ASMase regulates autophagy and lysosomal membrane permeabilization and its inhibition prevents early stage non-alcoholic steatohepatitis. J Hep 2014; 60: 1126-34.

59.Harvald EB, Olsen AS, Færgeman NJ. Autophagy in the light of sphingolipid metabolism. Apoptosis. 2015; 20:658-70..

60.Lu S, Natarajan SK, Mott JL, et al. Ceramide Induces Human Hepcidin Gene Transcription through JAK/STAT3 Pathway. PLoS One. 2016;11:e0147474.

61.Moles A, Tarrats N, Morales A, et al. Acidic sphingomyelinase controls hepatic stellate cell activation and in vivo liver fibrogenesis. Am J Pathol 2010;177:1214–1224.

62Adada M, Luberto C, Canals D. Inhibitors of the sphingomyelin cycle: Sphingomyelin synthases and sphingomyelinases.Chem Phys Lipids. 2016;197:45-59.

63.Ussher J. R., Koves TR., CadeteVJJ, et al. Inhibition of de novo ceramide synthesis reverses diet-induced insulin resistance and enhances whole-body oxygen consumption. Diabetes 2010;59, 2453–2464

64.Kasumov T, Li L, Li M, et al. Ceramide as a mediator of non-alcoholic Fatty liver disease and associated atherosclerosis. PLoS One. 2015;10:e0126910.

65.Zhao L., Spassieva S, Gable K., et al. Elevation of 20-carbon long chain bases due to a mutation in serine palmitoyltransferase small subunit bresultsinneurodegeneration.Proc.Natl.Acad. Sci.USA 2015;112:12962–12967 66.Wegner MS, Schiffmann S, Parnham MJ, et al. The enigma of ceramide synthase regulation in mammalian cells.Prog Lipid Res. 2016;63:93-119

67.Schiffmann S, Hartmann D, Fuchs S, et al. Inhibitors of specific ceramide synthases.Biochimie. 2012;94:558-65.

68.Novgorodov SA, Riley CL, Keffler JA, et al. SIRT3deacetylates ceramide synthases: implications for mitochondrialdysfunction and brain injury. J. Biol. Chem. 2016; 291:1957–1973 69.Garić D, De Sanctis JB, Wojewodka G, et al. Fenretinide differentially modulates the levels of long- and very long-chain ceramides by downregulating Cers5 enzyme: evidence from bench to bedside.JMol Med (Berl). 2017;95:1053-1064 Jul 10.

70.Mcilroy GD, Tammireddy SR, Maskrey BH, Fenretinide mediated retinoic acid receptor signalling and inhibition of ceramide biosynthesis regulates adipogenesis, lipid accumulation, mitochondrial function and nutrient stress signalling in adipocytes and adipose tissue.Biochem Pharmacol. 2016;100:86-97.

71.Liu Y, Sen S, Wannaiampikul S, et al.Metabolomic profiling in liver of adiponectin knockout mice uncovers lysophospholipid metabolism as an important target of adiponectin action.Biochem J. 2015 469:71-82.

72.Xia J. Targeted Induction of Ceramide Degradation Reveals Roles for Ceramides in Nonalcoholic Fatty Liver Disease and Glucose Metabolism. American Diabetes Association 75th Scientific Sessions, June 5 - 9, 2015, Boston, Massachusetts Abstract 265-LB American Diabetes Association 75th Scientific Sessions, June 5 - 9, 2015, Boston, Massachusetts

73.Rohrbach T, Maceyka M, Spiegel S. Sphingosine kinase and sphingosine-1-phosphate in liver pathobiology. Crit Rev Biochem Mol Biol. 2017 Jun 15:1-11.

74Alvarez SE, Harikumar KB, Hait NC, et al. Sphingosine-1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2.Nature. 2010;465:1084-8.

75.Maceyka M, Sankala H, Hait NC, et al. SphK1 and SphK2, sphingosine kinase isoenzymes with opposing functions in sphingolipid metabolism. J BiolChem 2005;280: 37118–37129.

76.Geng T, Sutter A, Harland MD, et al. SphK1 mediates hepatic inflammation in a mouse model of NASH induced by high saturated fat feeding and initiates proinflammatory signaling in hepatocytes. J Lipid Res. 2015;56:2359-71

77.Li C, Zheng S, You H, et al. Sphingosine 1-phosphate (S1P)/S1P receptors are involved in human liver fibrosis by action on hepatic myofibroblastsmotility.J Hepatol. 2011;54:1205-13.
78.Wang J, Badeanlou L, Bielawski J, et al.Sphingosine kinase 1 regulates adipose proinflammatory responses and insulin resistance.Am J Physiol Endocrinol Metab. 201;306:E756-68.

79.Lee SY, Hong IK, Kim BR,, et al. Activation of sphingosine kinase 2 by endoplasmic reticulum stress ameliorates hepatic steatosis and insulin resistance in mice. Hepatology. 2015;62:135-46.
80.Chen J, Wang W, Qi Y, et al. Deletion of sphingosine kinase 1 ameliorates hepatic steatosis in diet-induced obese mice: Role of PPARγ.BiochimBiophysActa. 2016;1861:138-47.
81.Mauer AS, Hirsova P, Maiers JL, et al. Inhibition of sphingosine 1-phosphate signaling ameliorates murine nonalcoholic steatohepatitis.Am J Physiol Gastrointest Liver Physiol.

2017;312:G300-G313.

82.Fadel FA, Fayyaz S, Japtok L, et al Involvement of Sphingosine 1-Phosphate in Palmitate-Induced Non-Alcoholic Fatty Liver Disease. Cell Physiol Biochem 2016;40:1637-1645
83.Xiu L, Chang N, Yang L, et al. Intracellular sphingosine 1-phosphate contributes to collagen expression of hepatic myofibroblasts in human liver fibrosis independent of its receptors. Am J Pathol. 2015;185:387-98

84.Ding BS, Liu CH, Sun Y, et al. HDL activation of endothelial sphingosine-1-phosphate receptor-1 (S1P1) promotes regeneration and suppresses fibrosis in the liver.JCI Insight.2016;1:e87058.
85.Lee SY, Lee HY, Song JH, et al. Adipocyte-Specific Deficiency of de novo Sphingolipid
Biosynthesis Leads to Lipodystrophy and Insulin Resistance.Diabetes. 2017;66:2596-2609.

~_

86.Alshaker H1,,,Wang Q, Srivats S, et al. New FTY720-docetaxel nanoparticle therapy overcomes FTY720-induced lymphopenia and inhibits metastatic breast tumour growth.Breast Cancer Res Treat. 2017;165:531-543.

87.Akazawa Y, Cazanave S, Mott JL, et al. .Palmitoleate attenuates palmitate-induced Bim and PUMA up-regulation and hepatocyte lipoapoptosis. J. Hepatol. 2010;52:586–593.

88.Listenberger LL, Han X, Lewis SE, et al. Triglycerideaccumulationprotectsagainstfatty acidinduced lipotoxicity. Proc Natl Acad Sci U S A. 2003;100:3077-82.

89.Gormaz, J.G.; Rodrigo, R.; Videla, L.A.; et al. Biosynthesis and bioavailability of long-chain polyunsaturated fatty acids in non-alcoholic fatty liver disease. Prog. Lipid Res. 2010; 49:407–19.
90.Nobili V, Alisi A, Musso G, et al.Omega-3 fatty acids: Mechanisms of benefit and therapeutic effects in pediatric and adult NAFLD. Crit Rev Clin Lab Sci. 2016;53:106-20

91.Das UN. A defect in the activity of Delta6 and Delta5 desaturases may be a factor predisposing to the development of insulin resistance syndrome. ProstaglandinsLeukot Essent Fatty Acids 2005;72:343–50.

92.Sui YH, Luo WJ, Xu QY, et al. Dietary saturated fatty acid and polyunsaturated fatty acid oppositely affect hepatic NOD-like receptor protein 3 inflammasome through regulating nuclear factor-kappa B activation. World J Gastroenterol. 2016;22:2533-44.

93.López-Vicario C1, González-Périz A, Rius B, et al. Molecular interplay between $\Delta 5/\Delta 6$ desaturases and long-chain fatty acids in the pathogenesis of non-alcoholic steatohepatitis. Gut.2014;63:344-55.

94.Depner CM, Philbrick KA, Jump DB. Docosahexaenoic acid attenuates hepatic inflammation, oxidative stress, and fibrosis without decreasing hepatosteatosis in a Ldlr(-/-) mouse model of western diet-induced nonalcoholic steatohepatitis. JNutr.2013;143:315-23.

95.Lytle KA, Depner CM, Wong CP, et al. Docosahexaenoic acid attenuates Western diet-induced hepatic fibrosis in Ldlr-/- mice by targeting the TGFβ-Smad3 pathway. J Lipid Res. 2015;56:1936-

46.

96.Walle P, Takkunen M, Männistö V, et al. Fatty acid metabolism is altered in non-alcoholic steatohepatitis independent of obesity.Metabolism. 2016;65:655-66.

97.Gromovsky AD, Schugar RC1, Brown AL, et al. Δ-5 Fatty Acid Desaturase FADS1 Impacts Metabolic Disease by Balancing Proinflammatory and Proresolving Lipid Mediators.Arterioscler Thromb Vasc Biol. 2017 Oct 26. pii: ATVBAHA.117.309660.

doi: 10.1161/ATVBAHA.117.309660.

98.Yashiro H, Takagahara S, Tamura YO, et al. A Novel Selective Inhibitor of Delta-5 Desaturase Lowers Insulin Resistance and Reduces Body Weight in Diet-Induced Obese C57BL/6J Mice. PLoS One. 2016;11:e0166198.

99.Slagsvold JE, Thorstensen K, Kvitland M, et al. Regulation of desaturase expression in HL60 cells. Scand J Clin Lab Invest. 2007;67:632–642.

100.Cho HP, Nakamura M, Clarke SD. Cloning, expression, and fatty acid regulation of the human delta-5 desaturase. J Biol Chem. 1999;274:37335–37339

101.Wang L, Athinarayanan S, Jiang G, et al.Fatty acid desaturase 1 gene polymorphisms control human hepatic lipid composition.Hepatology. 2015;61:119-28.

102.Okada LSDRR, Oliveira CP, Stefano JT, et al. Omega-3 PUFA modulate lipogenesis, ER stress, and mitochondrial dysfunction markers in NASH - Proteomic and lipidomic insight. ClinNutr. 2017 Sep 7. pii: S0261-5614(17)30314-X. doi: 10.1016/j.clnu.2017.08.031. [Epub ahead of print]

103.He XX, Wu XL, Chen RP, et al. Effectiveness of Omega-3 Polyunsaturated Fatty Acids in Non-Alcoholic Fatty Liver Disease: A Meta-Analysis of Randomized Controlled Trials. PLoS One. 2016;11:e0162368.

104.Freire M, Dalli J, Serhan CN, et al. Neutrophil Resolvin E1 Receptor Expression and Function in Type 2 Diabetes. J Immunol. 2017;198:718-728.

105. Uto Y. Recent progress in the discovery and development of stearoylCoA desaturase inhibitors. Chem Phys Lipids. 2016;197:3-12.

106.Kurikawa N, Takagi T, Wakimoto S, et al. A novel inhibitor of stearoyl-CoAdesaturase-1 attenuates hepatic lipid accumulation, liver injury and inflammation in model of nonalcoholic steatohepatitis. Biol Pharm Bull. 2013;36: 259-67.

107.Safadi R, Konikoff FM, Mahamid M, et al. The fatty acid-bile acid conjugate Aramchol reduces liver fat content in patients with nonalcoholic fatty liver disease.Clin Gastroenterol Hepatol. 2014;12:2085-91

108.Chen C, Shah YM, Morimura K, et al. Metabolomics reveals that hepatic stearoyl-CoA desaturase 1 downregulation exacerbates inflammation and acute colitis. Cell Metab. 2008;7:135-47.

109.Musso G, Cassader M, De Michieli F, et al.. Nonalcoholic steatohepatitis versus steatosis: adipose tissue insulin resistance and dysfunctional response to fat ingestion predict liver injury and altered glucose and lipoprotein metabolism.Hepatology. 2012;56: 933-42

110.Araújo AC, Wheelock CE, Haeggström JZ. The Eicosanoids, Redox-Regulated Lipid Mediators in Immunometabolic Disorders. Antioxid Redox Signal. 2017 Dec 11. doi: 10.1089/ars.2017.7332. [Epub ahead of print]

111.Horrillo R, Gonzalez-Periz A, Martinez-Clemente M, et al. 5-lipoxygenase activating protein signals adipose tissue inflammation and lipid dysfunctionin experimental obesity. J Immunol 2010; 184:3978–3987.

112.Rius B, Duran-Güell M, Flores-Costa R, et al. The specialized pro-resolving lipid mediator maresin 1 protects hepatocytes from lipotoxic and hypoxia-induced endoplasmic reticulum stress. FASEB J. 2017;31:5384-5398.

113.Sun GY, Chuang DY, Zong Y, et al. Role of cytosolic phospholipase A2 in oxidative and inflammatory signaling pathways in different cell types in the central nervous system. Mol Neurobiol. 2014;50:6-14.

114.Omland SH, Habicht A, Damsbo P, et al. A randomized, double-blind, placebo-controlled, dose-escalation first-in-man study (phase 0) to assess the safety and efficacy of topical cytosolic phospholipase A2 inhibitor, AVX001, in patients with mild to moderate plaque psoriasis. J Eur Acad Dermatol Venereol. 2017;31:1161-1167.

115.Chan PC, Hsiao FC, Chang HM, et al.Importance of adipocyte cyclooxygenase-2 and prostaglandin E2-prostaglandin E receptor 3 signaling in the development of obesity-induced adipose tissueinflammation and insulin resistance.FASEB J. 2016;30:2282-97

116.Chung MY, Mah E, Masterjohn C, et al.Green Tea Lowers Hepatic COX-2 and Prostaglandin E2 in Rats with Dietary Fat-Induced Nonalcoholic Steatohepatitis.J Med Food. 2015;18:648-55.

117.Ishihara K, Kanai S, Tanaka K, et al. Group IVA phospholipase A(2) deficiency prevents

CCl4-induced hepatic cell death through the enhancement of autophagy.Biochem Biophys Res Commun. 2016;471:15-20.

118.Peña L, Meana C, Astudillo AM, et al. Critical role for cytosolic group IVA phospholipase A2 in early adipocyte differentiation and obesity. Biochim Biophys Acta. 2016;1861(9 Pt A):1083-95. 119.Hu X, Cifarelli V, Sun S, et al. Major role of adipocyte prostaglandin E2 in lipolysis-induced macrophage recruitment.J Lipid Res. 2016;57:663-73.

120.Henkel J, Frede K, Schanze N, et al.Stimulation of fat accumulation in hepatocytes by PGE_2 dependent repression of hepatic lipolysis, β -oxidation and VLDL-synthesis. Lab Invest. 2012;92:1597-606.

121.Wu J, Chen C, Hu X, et alSuppressing cyclooxygenase-2 prevents nonalcoholic and inhibits apoptosis of hepatocytes that are involved in the Akt/p53 signal pathway.Biochem Biophys Res Commun. 2016;469:1034-40.

122.Henkel J, Gärtner D, Dorn C, et al. Oncostatin M produced in Kupffer cells in response to PGE2: possible contributor to hepatic insulin resistance and steatosis.Lab Invest. 2011;91:1107-17.

123.Tian F, Zhang YJ, Li Y, et al. Celecoxib ameliorates non-alcoholic steatohepatitis in type 2 diabetic rats via suppression of the non-canonical Wnt signaling pathway expression.PLoS One. 2014;9:e83819.

124.Hu X, Cifarelli V, Sun S, et al. Major role of adipocyte prostaglandin E2 in lipolysis-induced macrophage recruitment.J Lipid Res. 2016;57:663-73.

125.Loo TM, Kamachi F, Watanabe Y, et al. Gut Microbiota Promotes Obesity-Associated Liver Cancer through PGE2-Mediated Suppression of Antitumor Immunity. Cancer Discov. 2017;7:522-538.

126.Luan B, Yoon YS, Le Lay J, et al. CREB pathway links PGE2 signaling with macrophage polarization. Proc Natl AcadSci U S A. 2015;112:15642-7.

127.Manferdini C, Paolella F, Gabusi E, et al. Adipose stromal cells mediated switching of the proinflammatory profile of M1-like macrophages is facilitated by PGE2: in vitro evaluation.Osteoarthritis Cartilage. 2017;25:1161-1171.

128.Fanelli A, Ghisi D, Aprile PL, et al Cardiovascular and cerebrovascular risk with nonsteroidal anti-inflammatory drugs and cyclooxygenase 2 inhibitors: latest evidence and clinical implications.TherAdv Drug Saf. 2017;8:173-182

129.Titos E, Clària J, Planagumà A, et al. Inhibition of 5-lipoxygenase induces cell growth arrest and apoptosis in rat Kupffer cells: implications for liver fibrosis. FASEB J.2003 ;17:1745-7.

130.Ferre' N, Martinez-Clemente M, Lopez-Parra M, et al. Increased susceptibility

to exacerbated liver injury in hypercholesterolemic ApoE-deficient mice:

potential involvement of oxysterols. Am J Physiol Gastrointest Liver Physiol 2009;296:G553–G562 131.Martinez-Clemente M, Ferre´N, Gonzalez-Periz A, et al. 5-lipoxygenase deficiency reduces hepatic inflammation and tumor necrosis factor alpha-induced hepatocyte damage in hyperlipidemia-prone ApoE-null mice. Hepatology 2010;51:817–827.

57

132.Titos E, Ferre´ N, Lozano JJ, et al. Protection from hepatic lipid accumulation and inflammation by genetic ablation of 5-lipoxygenase.Prostaglandins Other Lipid Mediat 2010;92:54–61.

133.Matsuda K, Iwak Y; MN-001 (tipelukast), a novel, orally bioavailable drug, reduces fibrosis and inflammation and down-regulates TIMP-1, collagen Type 1 and LOXL2 mRNA overexpression in an advanced NASH (nonalcoholic steatohepatitis) model. Hepatology 2014; 60(S1): 1283A 134.American Association for the Study of Liver Diseases (AASLD) and Industry Colloquium -March 20, 2015, Durham, California, US. http://globenewswire.com/newsrelease/2015/02/25/709915/10122037/en/MediciNova-s-MN-001-tipelukast-NASH-with-Advanced-Fibrosis

135.Li P, Oh DY, Bandyopadhyay G, et al. LTB4 promotes insulin resistance in obese mice by acting on macrophages, hepatocytes and myocytes.Nat Med. 2015;21:239-47.

136.Serhan CN, Clish CB, Brannon J., et al. .Novel functional sets of lipid-derived mediators with anti-inflammatory actions generated from omega-3fatty acids via cyclooxygenase 2-nonsteroidal anti-inflammatory drugs and transcellular processing.J.Exp.Med. 2000;192:1197–1204.

137.Bannenberg G, Serhan CN, Specialized pro-resolvinglipid mediators in the inflammatory response: an update. Biochim.Biophys. Acta 2010;1801:1260–1273.

138.Kang JW, Lee SM. Resolvin D1 protects the liver from ischemia/reperfusion injury by enhancing M2 macrophage polarization and efferocytosis.Biochim Biophys Acta. 2016;1861(9 Pt A):1025-35

139Prieto P, Rosales-Mendoza CE, Terrón V, et al. Activation of autophagy in macrophages by pro-resolving lipid mediators. Autophagy. 2015;11:1729-44.

140.<u>Chiurchiù V</u>, <u>Leuti A</u>, <u>Dalli J</u>, et al.Proresolving lipid mediators resolvin D1, resolvin D2, and maresin 1 are critical in modulating T cell responses. <u>SciTransl Med.</u> 2016;8:353ra111.

141.Clària J,Nguyen BT, MadenciAL, et al. Diversity of lipid mediators in human adipose tissue depots.Am. J. Physiol.Cell Physiol. 2013;304:C1141–C1149.

142.Börgeson E, Johnson AM, Lee YS, et al Lipoxin A4 Attenuates Obesity-Induced Adipose Inflammation and Associated Liver and Kidney Disease. Cell Metab. 2015;22:125-37.

143. Laiglesia LM, Lorente-Cebrián S, Martínez-Fernández L, et al Maresin 1 mitigates liver

steatosis in ob/ob and diet-induced obese mice. Int J Obes (Lond).2017 Sep 12. doi:

10.1038/ijo.2017.226. [Epub ahead of print]

144.Martínez-Fernández L, González-Muniesa P, Laiglesia LM, et al. Maresin 1 improves insulin sensitivity and attenuates adipose tissue inflammation in ob/ob and diet-induced obese mice. FASEB J.2017;31:2135-2145.

145.Rius B, Titos E, Morán-Salvador E, et al. Resolvin D1 primes the resolution process initiated by calorie restriction in obesity-induced steatohepatitis.FASEB J. 2014;28:836-48.

146.Jung TW1, Hwang HJ1, Hong HC, et al. Resolvin D1reducesERstress-inducedapoptosis and triglyceride accumulation through JNK pathway in HepG2 cells. Mol Cell Endocrinol. 2014;39:30-40.

147.Li H, Hao Y, Zhang H, et al. Posttreatment with Protectin DX ameliorates bleomycin-induced pulmonary fibrosis and lung dysfunction in mice. Sci Rep. 2017;7:46754.

148.Yatomi M, Hisada T, Ishizuka T, et al.17(R)-resolvinD1amelioratesbleomycin-induced pulmonary fibrosis in mice.Physiol Rep. 2015;3. pii: e12628.

149.Qiu W, Guo K, Yi L, et al. ResolvinE1reduceshepaticfibrosis in mice with Schistosoma japonicuminfection.ExpTher Med. 2014;7:1481-1485.

150.Lee HJ, Park MK, Lee EJ, et al. ResolvinD1 inhibits TGF-β1-induced epithelial mesenchymal transition of A549 lung cancer cells via lipoxin A4 receptor/formyl peptide receptor 2 and GPR32.Int J Biochem Cell Biol. 2013;45:2801-7.

151.Wang Y, Li R, Chen L, et al.Maresin 1InhibitsEpithelial-to-Mesenchymal Transitionin Vitro and Attenuates Bleomycin Induced Lung Fibrosis in Vivo.Shock. 2015;44:496-502.

59

152.Tang S, Gao C, Long Y, et al. Maresin 1 Mitigates High Glucose-Induced Mouse Glomerular Mesangial Cell Injury by Inhibiting Inflammation and Fibrosis.Mediators Inflamm. 2017;2017:2438247.

153.Orr SK, Colas RA, Dalli J, et al. Proresolving actions of a new resolvin D1 analog mimetic qualifies as an immunoresolvent.Am J Physiol Lung Cell Mol Physiol. 2015;308:L904-11

154.Kain V, Ingle KA, Colas RA, et al. Resolvin D1 activates the inflammation resolving response at splenic and ventricular site following myocardial infarction leading to improved ventricular function. J Mol Cell Cardiol. 2015; 84: 24-35.

155.Musso G, Cassader M, Paschetta E, et al. Thiazolidinediones and Advanced Liver Fibrosis in Nonalcoholic Steatohepatitis: A Meta-analysis.JAMAInternMed. 2017;177:633-640

156.Stiuso P, Scognamiglio I, Murolo M, et al..Serum oxidative stress markers and lipidomic profile to detect NASH patients responsive to an antioxidant treatment: a pilot study. Oxid Med Cell Longev. 2014; 2014:169216.

157.Scorletti E, West AL, Bhatia L, et al. Treating liver fat and serum triglyceride levels inNAFLD, effects of PNPLA3 and TM6SF2 genotypes: results from the WELCOME trial. J Hepatol.2015 ;63:1476-83

158.Kalia HS, GaglioPJ. The Prevalence and Pathobiology of Nonalcoholic Fatty Liver Disease in Patients of Different Races or Ethnicities. Clin Liver Dis. 2016;20:215-24.

159.BasuRay S, Smagris E, Cohen JC, et al. The PNPLA3 variant associated with fatty liver disease (I148M) accumulates on lipid droplets by evading ubiquitylation. Hepatology. 2017;66:1111-1124. 160.Bruschi FV, Claudel T, Tardelli M, et al. The PNPLA3 I148M variant modulates the fibrogenic phenotype of human hepatic stellate cells. Hepatology. 2017;65:1875-1890

Online supplementary table

Table 1. Nuclear transcription factors Liver X receptor(LXR),FXR, SREBF-2/miRNA-33a, PXR,

PPAR- α , -PPAR- δ in the pathogenesis and treatment of NAFLD.

Liver X receptor(LXR)-a			
Cells	Molecular targets	Biological effect	
Hepatocyte,	↑ SREBP-1c and ChREBP activity	↑ hepatic <i>de novo</i> lipogenesis,	
enterocyte,		steatosis and large VLDL secretion	
macrophage	↑ CYP7A1 expression	↑ cholesterol conversion to bile acids	
	↓ NPC1L1expression	↓ intestinal cholesterol absorption	
	\uparrow macrophage, intestinal and hepatic	\uparrow cholesterol reverse transport and	
	ABCG5/G8 expression	excretion into bile and gut lumen	
	↑ macrophage and hepatic ABCA1 expression	↑ cholesterol efflux to acceptor apoA-	
		I to form HDL-C	
	↑ hepatic CEH>↑ FC availability for ABC	↑ cellular free cholesterol efflux	
	transporters		
	↑ hepatocyte CD36 expression	\uparrow uptake of plasma oxLDLs and FFAs	
	↑ LDLR degradation	↓ uptake of plasma LDL	
	\uparrow Angptl3 secretion \rightarrow LPL inhibition	↓ VLDL catabolism	
	↓ apoA-V secretion	VLDL catabolism	
Macrophage	↓ secretion of proinflammatory cytokines IL-	↓ hepatic inflammation	
, KC	1/TNF-α		
HSC	↓ HSC activation	↓ hepatic fibrosis	
	Farnesoid X Receptor	1	

Cell type	Molecular mechanism	Biological action
hepatocyte	↓ SREBP-1c-mediated lipogenesis	Reduced hepatic steatosis
	↑ PPAR-α-mediated FFA β-oxidation	
	↓ gluconeogenesis	
	\uparrow IRS-1 phosphorylation and coupling with the	Enhanced insulin sensitivity
	PI-3K	
	↑AdipoR2 expression	
	↑ CYP7A1 andABCG5/G810 expression	Enhanced bile acid synthesis and
		cholesterol excretion into bile
	↓ hepatic lipase activity and ApoC-III/apoA-1	Reduced plasma HDL-C
	synthesis	
	\downarrow VLDL secretion and HDL-Csynthesis	Reduced plasma TG
	↑ApoC-II synthesis and VLDLR-mediated	
	uptake of VLDL	
	↓NF-κB activation	Reduced inflammation
Marophage,	↓ NF-κB activation \rightarrow ↓ MCP-1 and TGF-ß	Reduced inflammation and
КС	secretion	fibrogenesis
HSC	↓ TGF-ß-R expression	Reduced fibrogenesis
Adipocyte	↑ PPAR-γ expression	Improved adipose tissue
	↑ adiponectin and AdipoR2 expression	dysfunction
	\downarrow TNF- α secretion	
1		

Enterocyte	Enhanced gut barrier function and	Reduced bacterial endotoxemia
	secretion of antibacterial factors angiogenin,	Increased bile acid synthesis, EE
	iNOS, IL-18	and fat oxidation
	Enhanced FGF-19 secretion	
	PPAR-α	
Cell type	Molecular mechanism	Biological action
Hepatocyte,	↑ expression of : acyl-CoA synthetase, CPT1A,	Increased mitochondrial and
Miocyte,	VLCAD/LCAD/MCAD, acyl-CoA	peroxisomal ß-oxidation
Adipocyte	dehydrogenase, trifunctional protein HADHB,	Increased ω-oxidation
	ACOX1, L-bifunctional protein EHHADH	Increased ketogenesis
	↑ CYP4A and HMGCA activity	
	↑ FATP, CD36, L-FABP activity	Enhanced FFA uptake
	↑ LPL activity and reduced apoC-III	Enhanced lipolysis of TG
	↑ apo-AI/apo-AII synthesis	Increased HDL-C levels
Hepatocyte	\uparrow p65 binding to NF-κB response element of C3	
	promoter \rightarrow reduced complement C3 secretion	Reduced inflammatory response and
	↓ NF-κB activation	endothelial dysfunction
	↑ FGF21 expression	
	\downarrow expression of IL-6 , IL-1,TNF-a, ICAM-1,	↑ metabolic effects of PPAR- α
	VCAM-1	
	Increased catalase activity	Enhanced H ₂ O ₂ detoxification
	ΡΡΑ R- β(δ)	
Cell type	Molecular mechanism	Biological action

Hepatocyte	\uparrow mitochondrial β-oxidation	Improved hepatic steatosis and
		insulin resistance
	↑ ABCA1 expression	Increased HDL-C levels
Macrophage	↓ M1/M2 phenotype ratio	Reduced inflammatory and
КС	\downarrow NF- κ B activation and TGF- β 1 secretion	fibrogenesis
Adipocyte,	\uparrow PGC-1 α -mediated mitochondrial biogenesis	Enhanced fat oxidation and EE
miocyte	and ß-oxidation	
	↑ mitochondrial UCP-1/3 expression	
	↑ LPL expression	Reduced plasma TG
Enterocyte	↓ NPC1L1 expression and cholesterol	Reduced cholesterol accumulation
	reabsorption from bile and intestine	
	ΡΡΑR-γ	L
Cell type	Molecular mechanism	Biological action
Cell type Adipocyte	Molecular mechanism ↑ subcutaneous preadipocyte differentiation and	Biological action Enhanced insulin sensitivity and
Cell type Adipocyte	Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced
Cell type Adipocyte	Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes ↑ insulin signalling and adiponectin secretion	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation
Cell type Adipocyte	Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes ↑ insulin signalling and adiponectin secretion ↑ GLUT1/GLUT4 translocation	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic
Cell type Adipocyte	Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes ↑ insulin signalling and adiponectin secretion ↑ GLUT1/GLUT4 translocation ↑ release of pro-inflammatory FFAs and	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic inflammation
Cell type Adipocyte	 Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes ↑ insulin signalling and adiponectin secretion ↑ GLUT1/GLUT4 translocation ↑ release of pro-inflammatory FFAs and cytokines TNF-α, resistin and IL-6 	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic inflammation
Cell type Adipocyte Skeletal	 Molecular mechanism ↑ subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes ↑ insulin signalling and adiponectin secretion ↑ GLUT1/GLUT4 translocation ↑ release of pro-inflammatory FFAs and cytokines TNF-α, resistin and IL-6 ↑ GLUT1/GLUT4 expression and translocation 	Biological actionEnhanced insulin sensitivity andsubcutaneous fat storage Enhancedglucose disposal and FFA oxidationReduced systemicinflammationEnhanced insulin sensitivity, glucose
Cell type Adipocyte Skeletal miocyte,	Molecular mechanism \uparrow subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes \uparrow insulin signalling and adiponectin secretion \uparrow GLUT1/GLUT4 translocation \uparrow release of pro-inflammatory FFAs and cytokines TNF- α , resistin and IL-6 \uparrow GLUT1/GLUT4 expression and translocation to the cell surface	Biological actionEnhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic inflammationEnhanced insulin sensitivity, glucose disposal and FFA oxidation
Cell type Adipocyte Skeletal miocyte, hepatocyte	Molecular mechanism \uparrow subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes \uparrow insulin signalling and adiponectin secretion \uparrow GLUT1/GLUT4 translocation \uparrow release of pro-inflammatory FFAs and cytokines TNF- α , resistin and IL-6 \uparrow GLUT1/GLUT4 expression and translocation to the cell surface \downarrow gluconeogenesis (adponectin-mediated)	Biological action Enhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic inflammation Enhanced insulin sensitivity, glucose disposal and FFA oxidation
Cell type Adipocyte Skeletal miocyte, hepatocyte Macrophage	Molecular mechanism \uparrow subcutaneous preadipocyte differentiation and apoptosis of insulin-resistant visceral adipocytes \uparrow insulin signalling and adiponectin secretion \uparrow GLUT1/GLUT4 translocation \uparrow release of pro-inflammatory FFAs and cytokines TNF- α , resistin and IL-6 \uparrow GLUT1/GLUT4 expression and translocation to the cell surface \downarrow gluconeogenesis (adponectin-mediated)M2 phenotype switching	Biological actionEnhanced insulin sensitivity and subcutaneous fat storage Enhanced glucose disposal and FFA oxidation Reduced systemic inflammationEnhanced insulin sensitivity, glucose disposal and FFA oxidationAnti-inflammatory and pro-resolving

HSC	Inhibition of TGFβ-1/Smad3-signaling pathway	Anti-fibrotic effects
	Induction of cell apoptosis	

Abbreviations: ABC. ATP-binding cassette; CYP7A1: Sterol 7a hydroxylase; 27OHC: 27-

hydroxycholesterol; 7OHC: 7-hydroxycholesterol; 22(R)OHC: 22(R)-hydroxycholesterol, 24(S): 24(S)-hydroxycholesterol,

27OHC: 27-hydroxycholesterol 24(S), 25OOC: 24(S), 25-epoxycholesterol 25OHC3S:25-

hydroxycholesterol-3-sulfate, 22(S)OHC: 22(S)-hydroxycholesterol;

SAA: serum amyloid A protein; LE: late endosome; LY: lysosome;

PM: plasma membrane; IL: interleukin; HMGCoAR: 3-hydroxy-3-methyl-glutaryl-CoA reductase;

NPC1, NPC2: Niemann-Pick C1, C2; NPC1L1: Niemann-Pick C1-like 1; oxLDL: oxidized low density lipoproteins;PPAR: peroxisome proliferator-activated receptor; SHP: small heterodimeric partner; StARD4:Steroidogenic acute regulatory protein D4; LE/LY: late endosomes/lysosomes; PEPCK: phosphoenol-pyruvate carboxykinase; IRS-1: insulin-receptor substrate-1; TGF: transforming growth factor; CEH: cholesteryl ester hydrolase; LDLR: low density lipoprotein receptor; FFA: free fatty acids; HSC:hepatic stellate cells; VLDL: very low density lipoproteins

Online supplementary Table 2. Role of group IVA phospholipase A2 and cyclooxygenase (COX)-2-generated mediators in the pathogenesis of NASH and hepatic fibrosis

group IVA phospholipase A2			
15-deoxy-PGI2 and PGE2-mediated			
Cell type and molecular pathways	Cellular effect	Biological effect	

		1	
Hepatocyte, adipocyte:			
→↓ cAMP synthesis →↓ HSL activity	Lipid storage		
↓ VLDL-TG secretion		Steatosis	
\uparrow SREBP-1c activation \rightarrow \uparrow <i>de novo</i> lipogenesis		Obesity	
\uparrow DGAT-2 activation \rightarrow \uparrow TG synthesis			
↓ constitutive autophagy	cell death	Necrosis	
Kupffer cells, HSCs:			
\uparrow MCP-1 secretion \rightarrow monocyte recruitment to the liver	monocyte recruitment	Inflammation	
↑ TGF- β secretion → HSC activation	HSC activation	Fibrosis	
NOX activation \rightarrow \uparrow ROS production			
Lysophosphatidylcholine-mediated			
Hepatocyte, adipocyte:			
\downarrow mitochondrial oxidative phosphorylation	Cell apoptosis	necrosis	
↑ ER stress			
↑ JNK activation			
HSC:			
↑ activation by autotaxin-generated LPA	HSC activation	fibrogenesis	
Cyclooxygenase(COX)-2			
Cell type and molecular pathways	Cellular effect	Biological effect	

		1
Hepatocyte:	↓ mitochondrial β-	Steatosis
\downarrow PGC1 $\alpha \rightarrow \downarrow$ CPT1 activity	oxidation	
↓ adipose triglyceride lipase		
↓ MTTP synthesis	↓ TG hydrolysis	
	↓ VLDL secretion	
↓ Akt /↑ p53 signaling	↑ anontosis	Necro-
↑ JNK expression		110010-
	↑ proinflammatory	inflammation
	cytokine secretion	
Kupffer cells:	↓ Akt activation	Insulin ressitance
↑ oncostatin M \rightarrow ↑ SOCS3 secretion by hepatocytes	μ mitochondrial FFA β-	Steatosis
$\rightarrow \downarrow$ CPT1 in hepatocytes		Steatosis
	oxidation	
HSCs:	↓ antitumor immunity	HCC progression
↑ PGE2 secretion \rightarrow ↓ regulatory T cells (Treg) activation		
Adipocyte:	macrophage recruitment	Enhanced
↑ pro-inflammatory cytokine and chemokine MCP-1		induction and
secretion		
	resolution of inflammation	resolution of
↑ macrophage polarization switch from a M1 to a M2		inflammation
phenotype		

Abbreviations: ABCA1: ATP-binding cassette transporters A1; ACC: acetyl-CoA carboxylase;

AMPK: adenosine-monophosphate kinase; **Ccl3 : chemokine(C-C motif ligand 3 ;** CD36: cluster of differentiation-36; CHOP: C/EBP homologous protein; CPP: calciprotein particles; CPT-1: carnitine palmitoyltransferase-I; ; ER: endoplasmic reticulum; FAS: fatty acid synthase; FFA: free fatty acids; FXR: farnesoid X-receptor; GLUT: glucose transporter; HMG-CoAR: 3-hydroxy-3-methylglutaryl-coenzyme A reductase; IL: interleukin; 11β-HSD1: 11β-hydroxysteroid dehydrogenase type 1; IRE1 α : inositol requiring element 1 α ; IRS-1: insulin receptor substrate-1; KLF: Kruppel-like factor; LDL: low-density lipoprotein; LDL-R: low-density lipoprotein receptor; LPA: lysophosphatidic acid; MCP-1: monocyte chemotactic protein-1; NO: nitric oxide; NOX:

NADPH oxidase; OCA: obeticholic acid; PCSK9 : proprotein convertasesubtilisin kexin 9 ; PGC-1 α : peroxisome proliferator-activated receptor- γ coactivator-1 α ; ROS: reactive oxygen species; SCD-1: stearoyl-CoA desaturase-1; SOD2: superoxide dimutase-2; SR-A1: scavenger receptor-A1; SOCS3: suppressor of cytokine signaling 3; SR-B1: scavenger receptor-B1; SREBP: sterolresponsive element binding protein; STAT3: signal transducer and activator of transcription; TGF- β : transforming growth factor- β ; TLR: toll-like receptor; TNF: tumor necrosis factor; VLDL: very low density lipoprotein; VSCMs: vascular smooth muscle cells;

Acius (FOFAS)			
Hepatocyte			
Type of	Transcription factor/cellular	Cellular molecular mechanism	Biological effect
PUFA	pathway		
EPA,	↓ LXR/SREBP-1c activation	↓ <i>de novo</i> lipogenesis	↓ hepatic triglyceride
DHA			
EPA,	↓ ChREBP activation	↓ de novo lipogenesis	↓ hepatic triglyceride
DHA		↓ glycolysis	
EPA,	↓ SREBP- activation	↓ cholesterol synthesis	↓ hepatic cholesterol
DHA			

 \uparrow fatty acid β-oxidation

 \downarrow caspase-1 activation

 \downarrow IL-1 secretion

↓ pro-inflammatory IL-1 secretion

↓ cholesterol and FA synthesis

 \downarrow apoptosis

Online Supplementary Table 3. Main pathways and biological effect of Polyunsaturated Fatty Acids (PUFAs)

EPA

DHA

DHA

DHA.

EPA

 \uparrow PPAR- α activation

 \downarrow NF- κ B activation ⁴⁶

inflammasome

NLRP3

↑ FXR activation

activation⁴⁶

.

↓ hepatic triglyceride

↓ necro-inflammation

hepatic triglyceride

inflammation

and cholesterol

T

.....

hepaticnecro-

DHA,	\downarrow HNF-4 α activation	\downarrow VLDL assembly and secretion	↓ plasma triglyceride
EPA			
DHA	↓Tolllike receptor-2/4/9	↓proinflammatory cytokine	↓necroinflammation
	48	secretion	
DHA	\downarrow NOX expression ⁴⁸	↓ superoxide/hydrogen peroxide	\downarrow oxidative stress
		production	
Macrophage/Kupffer cells			
DHA	\uparrow PPAR-γ activation	Polarization to a M2, proresolving	Anti-inflammation
		phenotype	Fibrosis resolution
Hepatic Stellate Cell (HSC)			
DHA	\downarrow TGF β -Smad3-Col1A1 axis	↓ collagen deposition	↓ fibrosis
	activation		

Abbreviations: SREBP-1c: sterol regulatory element-binding protein 1c; ChREBP: carbohydrate response element binding protein; Col1A1: collagen 1A1; TGF β : transforming growth factor- β ; mothers against decapentaplegic homolog (Smad)3; NF- κ B: nuclear factor-kappa B; NOD-like receptor protein 3 (NLRP3) inflammasome, LXR: liver X receptor












Acetil-CoA



Palmitoleic acid



Malonil-CoA

$$CH_3 - (CH_2)_n - C \sim O$$

OH

Miristic Acid (n=12); 14:0 Palmitic Acid (n=14); 16:0 Stearic Acid (n=16); 18:0

$$CH_3 - (CH_2)_7 - CH = CH - (CH_2)_7 - C \sim OH$$

Oleic acid



$$\begin{array}{c} H \\ H - C - O \\ H - C \\ H - C - O \\ H - C - O \\ H - C \\ H - C - O \\ H - C \\ H \\ H - C \\ H - C \\ H \\ H - C \\ H - C \\ H \\ H - C \\ H - C \\ H \\$$

Diacylglycerol

Triacylglycerol

ω -3 fatty acids

$$CH_3 - (CH_2 - CH = CH)_3 - (CH_2)_7 - C_{OH}$$

 $CH_{3} - (CH_{2} - CH = CH)_{5} - (CH_{2})_{3} - C$

 $CH_3 - (CH_2 - CH = CH)_6 - (CH_2)_2 - C$

 α -Linolenic acid

Eicosapentaenoic acid

Docosahexaenoic acid

 ω -6 fatty acids

$$CH_3 - (CH_2)_4 - (CH = CH - CH_2)_4 - (CH_2)_2 - C \bigvee_{OH}^{O}$$

Arachidonic acid





Phosphocholine



Sphingosine



Sphingosine-1phosphate



Ceramide



Leukotriene A



Leukotriene A5



Prostaglandin E2



Resolvin D1



Resolvin E1



